

# Energy and Momentum in General Relativity

M. Sharif \*

Department of Mathematics, University of the Punjab,  
Quaid-e-Azam Campus Lahore-54590, PAKISTAN.

## Abstract

The energy and momentum for different cosmological models using various prescriptions are evaluated. In particular, we have focused our attention on the energy and momentum for gravitational waves and discuss the results. It is concluded that there are methods which can provide physically acceptable results.

**Keywords:** Energy-momentum problem, gravitational waves

---

\*E-mail: hasharif@yahoo.com

# 1 Introduction

The notion of energy has been one of the most thorny and important problems in Einstein's theory of General Relativity (GR). There have been many attempts [1,2,3] to get a well defined expression for local or quasi-local energy and momentum. However, there is still no generally accepted definition known. As a result, different people have different points of view. Cooperstock [4] argued that in GR, energy and momentum are localized in regions of the non-vanishing energy and momentum tensor and consequently gravitational waves are not carriers of energy and momentum in vacuum. The gravitational waves, by definition, have zero stress-energy tensor. Thus the existence of these waves was questioned. However, the theory of GR indicates the existence of gravitational waves as solutions of Einstein's field equations [5]. Infact this problem arises because energy is not well defined in GR.

The problem for gravitational waves was resolved by Ehlers and Kundt [6], Pirani [7] and Weber and Wheeler [8] by considering a sphere of test particles in the path of the waves. They showed that these particles acquired a constant momentum from the waves. Qadir and Sharif [9] presented an operational procedure, embodying the same principle, to show that gravitational waves impart a momentum. Rosen [10] investigated whether or not cylindrical gravitational waves have energy and momentum. He used the energy-momentum pseudo tensors of Einstein and Landau Lifshitz and carried out calculations in cylindrical polar coordinates. However, he arrived at the conclusion that the energy and momentum density components vanish. These results supported the conjecture of Scheidegger [11] that physical system cannot radiate gravitational energy. Later, he pointed out [12] that the energy and momentum densities turn out to be non-vanishing and reasonable if the calculations are performed in Cartesian coordinates. Rosen and Virbhadra [13] explicitly evaluated these quantities in the Einstein's prescription by using Cartesian coordinates and found them finite and well defined. Virbhadra [14] then used Tolman, Landau-Lifshitz and Papapetrou's prescriptions to evaluate the energy and momentum densities and found that the same results turn out in all these prescriptions.

Energy and momentum density are usually defined by a second rank tensor  $T_a^b$ . The conservation of energy and momentum are described by the requirement that the tensor's divergence is zero. However, in GR, the partial derivative in the usual conservation equation  $T_{a,b}^b = 0$  is replaced by a covariant derivative.  $T_a^b$  then represents the energy and momentum of matter and

all non-gravitational fields and no longer satisfies  $T_{a,b}^b = 0$ . A contribution from the gravitational field must be added to obtain an energy-momentum expression with zero divergence. Einstein obtained such an expression and many others such as Landau and Lifshitz, Papapetrou and Weinberg gave similar prescriptions [15]. The expressions they gave are called energy-momentum complexes because they can be expressed as a combination of  $T_a^b$  and a pseudotensor, which is interpreted to represent the energy and momentum of the gravitational field. These complexes have been heavily criticized because they are non-tensorial, i.e. they are coordinate dependent. For the Einstein, Landau-Lifshitz, Papapetrou, Weinberg (ELLPW) energy-momentum complexes, one gets physically meaningful results only in *Cartesian coordinates* [16-18]. Because of this drawback, many others, including Moller [18], Komar [19] and Penrose [1], have proposed coordinate independent definitions. Each of these, however, has its own drawbacks [20,21].

In this paper we gather various results to show that different prescriptions can provide the same result for different cosmological models. Also, we shall see from the analysis that when rotation is included, the problem becomes considerably complicated and the results obtained may not be the same. This has been explained by applying to a class of cylindrical gravitational waves. The paper has been planned as follows. In the next section, we shall describe different prescriptions to evaluate energy and momentum densities. In section three, these methods will be applied to different cosmological models. The section four contains the evaluation of energy and momentum for gravitational waves. Finally, we shall conclude the results.

## 2 Various Prescriptions to Evaluate Energy and Momentum

In this section we describe different methods to calculate energy and momentum in GR.

### (i) Energy and Momentum in Einstein's Prescription

The energy-momentum complex of Einstein [18,20] is given by

$$\Theta_a^b = \frac{1}{16\pi} H_{a,c}^{bc}, \quad (1)$$

where

$$H_a^{bc} = \frac{g_{ad}}{\sqrt{-g}}[-g(g^{bd}g^{ce} - g^{cd}g^{be})]_{,e}, \quad (2)$$

where Latin indices run from 0 to 3 and Greek indices from 1 to 3.  $\Theta_0^0$  is the energy density,  $\Theta_0^\alpha$  are the momentum density components, and  $\Theta_\alpha^0$  are the components of energy current density. The Einstein energy-momentum satisfies the local conservation laws

$$\frac{\partial \Theta_a^b}{\partial x^b} = 0. \quad (3)$$

### (ii) Energy and Momentum in Landau-Lifshitz Prescription

The energy-momentum complex of Landau-Lifshitz [17] is given by

$$L^{ab} = \frac{1}{16\pi} S^{abcd}_{,cd}, \quad (4)$$

where

$$S^{abcd} = -g[g^{ab}g^{cd} - g^{ac}g^{bd}]. \quad (5)$$

$L^{ab}$  is symmetric in its indices.  $L^{00}$  is the energy density,  $L^{0\alpha}$  are the momentum (energy current) density components.  $S^{abcd}$  has symmetries of the Riemann curvature tensor. The energy-momentum complex of Landau and Lifshitz satisfies the local conservation laws

$$\frac{\partial L^{ab}}{\partial x^b} = 0, \quad (6)$$

where

$$L^{ab} = -g(T^{ab} + t^{ab}). \quad (7)$$

$g$  is the determinant of the metric tensor  $g_{ab}$ ,  $T^{ab}$  is the energy-momentum tensor of the matter and all non-gravitational fields, and  $t^{ab}$  is known as Landau-Lifshitz energy-momentum pseudo tensor. Thus the locally conserved quantity  $L^{ab}$  contains contributions from the matter, non-gravitational fields and gravitational fields.

### (iii) Energy and Momentum in Papapetrou's Prescription

The energy-momentum complex of Papapetrou [22] is given by

$$\Omega^{ab} = \frac{1}{16\pi} N^{abcd}_{,cd}, \quad (8)$$

where

$$N^{abcd} = \sqrt{-g}[g^{ab}\eta^{cd} - g^{ac}g^{bd} + g^{cd}\eta^{ab} - g^{bd}g^{ac}], \quad (9)$$

and  $\eta^{ab}$  is the Minkowski spacetime.  $\Omega^{00}$  and  $\Omega^{\alpha 0}$  are the energy and momentum density components respectively. The Papapetrou energy-momentum complex satisfies the local conservation laws

$$\frac{\partial \Omega^{ab}}{\partial x^b} = 0. \quad (10)$$

#### (iv) Energy and Momentum in Weinberg Prescription

The energy-momentum complex of Weinberg [23] is given by

$$W^{ab} = \frac{1}{16\pi} \Delta^{abc}, \quad (11)$$

where

$$\Delta^{abc} = \frac{\partial h_e^e}{\partial x_a} \eta^{bc} - \frac{\partial h_e^e}{\partial x_b} \eta^{ac} - \frac{\partial h^{ea}}{\partial x^e} \eta^{bc} + \frac{\partial h^{eb}}{\partial x^e} \eta^{ac} + \frac{\partial h^{ac}}{\partial x_b} - \frac{\partial h^{bc}}{\partial x_a} \quad (12)$$

and

$$h_{ab} = g_{ab} - \eta_{ab}. \quad (13)$$

The indices on  $h_{ab}$  or  $\frac{\partial}{\partial x_a}$  are raised or lowered with the help of  $\eta$ 's. The Weinberg energy-momentum complex  $W^{ab}$  contains contributions from the matter, non-gravitational and gravitational fields, and satisfies the local conservation laws

$$\frac{\partial W^{ab}}{\partial x^b} = 0. \quad (14)$$

$W^{00}$  and  $W^{\alpha 0}$  are the energy and momentum density components respectively.

#### (v) Energy and Momentum in Möller Prescription

The energy-momentum complex of Moller [18] is given by

$$M_a^b = \frac{1}{8\pi} \chi_{a,c}^{bc}, \quad (15)$$

satisfying the local conservation laws:

$$\frac{\partial M_a^b}{\partial x^b} = 0, \quad (16)$$

where the antisymmetric superpotential  $\chi_a^{bc}$  is

$$\chi_a^{bc} = \sqrt{-g}[g_{ad,e} - g_{ae,d}]g^{be}g^{cd}. \quad (17)$$

$M_0^0$  is the energy density and  $M_\alpha^0$  are the momentum density components.

**(vi) Energy and Momentum in Qadir-Sharif's Prescription**

The energy-momentum complex of Qadir-Sharif [9] is given by

$$p_a = \int F_a, \quad (18)$$

where

$$F_0 = m[\{\ln(A/\sqrt{g_{00}})\}_{,0} - g_{\alpha\beta,0}g_0^{\alpha\beta}/4A], \quad F_i = m(\ln \sqrt{g_{00}})_{,i}, \quad (19)$$

and  $A = (\ln \sqrt{-g})_{,0}$ ,  $g = \det(g_{\alpha\beta})$ . This force formula depends on the choice of frame, which is not uniquely fixed. The quantity, whose proper time derivative is  $F_a$ , is called the momentum four-vector for the test particle. The spatial components of  $p_a$  give the momentum imparted to test particles as defined in the preferred frame (in which  $g_{0\alpha} = 0$ ).

### 3 Application to Various Cosmological Models

**(i) Bianchi Type I Universes**

The Bianchi type I spacetimes are expressed by the line element

$$ds^2 = dt^2 - e^{2l}dx^2 - e^{2m}dy^2 - e^{2n}dz^2, \quad (20)$$

where  $l, m, n$  are functions of  $t$  alone. Using ELLPW prescriptions, it turns out that energy-momentum distribution is zero. This supports the viewpoint of Tyron [24].

**(ii) Axially Symmetric Scalar Field**

It is well known that the Kaluza-Klein and the superstring theories predict the scalar fields as a fundamental interaction in Physics. Scalar fields are fundamental components of the Brans-Dicke theory and of the inflationary models. Also, they are a good candidate for the dark matter in spiral galaxies. Because they interact very weakly with matter we have never seen one but many of the theories containing scalar fields are in good concordance with

measurements in weak gravitational fields. Also, we expect that they can play an important role in strong gravitational fields like at the origin of the universe or in pulsars or black holes. The metric that we consider [25,26] is an axially symmetric solution to the field equations derived from the action for gravity minimally coupled to a scalar field. The solution is

$$ds^2 = \left(1 - \frac{2M}{r}\right)dt^2 - \frac{e^{2k_a}}{1 - \frac{2M}{r}}dr^2 - r^2(e^{2k_a}d\theta^2 + \sin^2\theta d\phi^2), \quad (21)$$

with

$$e^{2k_a} = \left(1 + \frac{M^2 \sin^2\theta}{r^2\left(1 - \frac{2M}{r}\right)}\right)^{\frac{-1}{a^2}}, \quad \phi = \frac{1}{2a} \ln\left(1 - \frac{2M}{r}\right), \quad (22)$$

where  $a$  is a constant of integration and  $\phi$  is the scalar field. This solution is one of the new classes of solutions to the Einstein-Maxwell theory non-minimally coupled to a dilatonic [26]. The metric given by Eq.(21) is almost spherically symmetric and represents a gravitational body (gravitational monopole) with scalar field. The scalar field deforms the spherical symmetry. We observe that when  $a \rightarrow \infty$  we recover the Schwarzschild solution.

When we apply Möller's prescription, it yields

$$E = M. \quad (23)$$

Thus the energy distribution is given by the mass  $M$ . In the case of the Schwarzschild metric we obtain the same result.

### (iii) Charged Regular Black Hole

The Reissner-Nordstrom (RN) metric is the only static and asymptotically flat solution of the Einstein-Maxwell equations and it represents an electrically charged black hole. The metric is given by

$$ds^2 = A(r)dt^2 - B(r)dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (24)$$

where

$$A(r) = B^{-1}(r) = 1 + \frac{2M}{r} + \frac{q^2}{r^2} \quad (25)$$

and  $q$  and  $M$  are the electric charge and the mass of the black hole respectively.

A solution to the coupled system of the Einstein field equations of the nonlinear electrodynamics was recently given by E. Ayon-Beato and A. Garcia (ABG) [27]. This solution represents a regular black hole with mass  $M$

and electric charge  $q$  and avoids thus the singularity problem. Also, the metric asymptotically behaves as the RN solution. The usual singularity of the RN solution, at  $r = 0$ , has been smoothed out and now it simply corresponds to the origin of the spherical coordinates. The line element is given by (24) with

$$A(r) = B^{-1}(r) = 1 - \frac{2M}{r} \left(1 - \tanh\left(\frac{q^2}{2Mr}\right)\right). \quad (26)$$

If the electric charge vanishes we reach the Schwarzschild solution. At large distances (26) resembles to the RN solution and can be written

$$A(r) = B^{-1}(r) = 1 + \frac{2M}{r} + \frac{q^2}{r^2} + \frac{q^6}{12M^2r^4} + O\left(\frac{1}{r^6}\right). \quad (27)$$

Using Einstein's prescription, we get the energy distribution of the ABG black hole given by

$$E(r) = M - \frac{q^2}{2r} + \frac{q^6}{24M^2r^3} - \frac{q^{10}}{240M^4r^5} + O\left(\frac{1}{r^6}\right). \quad (28)$$

This can also be written as

$$E(r) = E_{RN}(r) + \frac{q^6}{24M^2r^3} - \frac{q^{10}}{240M^4r^5} + O\left(\frac{1}{r^6}\right). \quad (29)$$

It follows that if  $q = 0$  we have the energy of a Schwarzschild black hole. The Möller's prescription gives

$$E(r) = M - \frac{q^2}{r} + \frac{q^6}{6M^2r^3} - \frac{q^{10}}{40M^4r^5} + O\left(\frac{1}{r^6}\right). \quad (30)$$

From Eq.(28) it results that in the Einstein prescription the first two terms in the expression of the energy correspond to the Penrose quasi-local mass definition evaluated by Tod [1,28]. The Möller's prescription provides in the expression of the energy (30) a term  $M - \frac{q^2}{r}$  which agrees with the Komar [19] prescription.

#### (iv) Kerr-Newmann Metric

The stationary axially symmetric and asymptotically flat Kerr-Newmann (KN) solution is the most general black hole solution to the Einstein-Maxwell equations. This describes the exterior gravitational and electromagnetic field of a charged rotating object. The KN metric in Boyer-Lindquist coordinates  $(t, \rho, \theta, \phi)$  is expressed by the line element

$$ds^2 = \frac{\Delta}{r_0^2} [dt - a \sin^2 \theta d\phi]^2 - \frac{\sin^2 \theta}{r_0^2} [(\rho^2 + a^2) d\phi - a dt]^2 - \frac{r_0^2}{\Delta} d\rho^2 - r_0^2 d\theta^2, \quad (31)$$

where  $\Delta = \rho^2 - 2M\rho + q^2 + a^2$  and  $r_0^2 = \rho^2 + a^2 \cos^2 \theta$ .  $M, q$  and  $a$  are respectively mass, electric charge and rotation parameters. Aguirregabiria et al. [29] studied the energy-momentum complexes of ELLPW for the KN metric. They showed that these definitions give the same results for the energy and energy current densities. This is given as

$$E_{ELLPW} = M - \frac{q^2}{4\rho} \left[ 1 + \frac{(a^2 + \rho^2)}{a\rho} \arctan\left(\frac{a}{\rho}\right) \right]. \quad (32)$$

Using Möller's prescription, we have

$$E_{M\ddot{o}l} = M - \frac{q^2}{2\rho} \left[ 1 + \frac{(a^2 + \rho^2)}{a\rho} \arctan\left(\frac{a}{\rho}\right) \right]. \quad (33)$$

The result of Möller's agrees with the energy distribution obtained by Cohen and de Felice [30] in Komar's prescription. The second term of the energy distribution differs by a factor of two from that computed by Aguirregabiria et al. using ELLPW complex. The total energy ( $\rho \rightarrow \infty$  in all these energy expressions) give the same result  $M$ .

#### (v) Melvin's Magnetic Universe

Melvin [31] obtained an axially symmetric electrovac solution ( $J^a = 0$ ) describing the Schwarzschild black hole in Melvin's magnetic universe. The spacetime is

$$ds^2 = \Lambda^2 \left[ \left(1 - \frac{2M}{r}\right) dt^2 - \left(1 - \frac{2M}{r}\right)^{-1} dr^2 - r^2 d\theta^2 \right] - \Lambda^{-2} r^2 \sin^2 \theta d\phi^2 \quad (34)$$

and the Cartan components of the magnetic field are

$$H_r = \Lambda^{-2} B_0 \cos \theta, \quad H_\theta = -\Lambda^{-2} B_0 (1 - 2M/r)^{1/2} \sin \theta, \quad (35)$$

where

$$\Lambda = 1 + \frac{1}{4} B_0^2 r^2 \sin^2 \theta. \quad (36)$$

$M$  and  $B_0$  are constants. The ELLP energy distribution gives

$$E = Mc^2 + \frac{1}{6} B_0^2 r^3 + \frac{1}{20} \frac{G}{c^4} B_0^4 r^5 + \frac{1}{140} \frac{G^2}{c^8} B_0^6 r^7 + \frac{1}{2520} \frac{G^3}{c^{12}} B_0^8 r^9. \quad (37)$$

The above result can be expressed in geometrized units (gravitational constant  $G = 1$  and the speed of light in vacuum  $c = 1$ ) as follows

$$E = M + \frac{1}{6} B_0^2 r^3 + \frac{1}{20} B_0^4 r^5 + \frac{1}{140} B_0^6 r^7 + \frac{1}{2520} B_0^8 r^9. \quad (38)$$

The first term  $Mc^2$  is the rest-mass energy of the Schwarzschild black hole, the second term  $\frac{1}{6}B_0^2r^3$  is the well-known classical value of the energy of the magnetic field under consideration, and rest of the terms are general relativistic corrections. For very large  $B_0r$ , the general relativistic contribution dominates over the classical value for the magnetic field energy.

## 4 Application to Gravitational Waves

### (i) Plane Gravitational Waves

The metric for the plane-fronted gravitational waves is [5,32]

$$ds^2 = dt^2 - dx^2 - L^2(t, x)[\exp\{2\beta(t, x)\}dy^2 + \exp\{-2\beta(t, x)\}dz^2], \quad (39)$$

where  $L$  and  $\beta$  are arbitrary functions subject to the vacuum Einstein equations

$$L_{,\alpha\alpha} + L\beta_{,\alpha}^2 = 0, \quad \alpha = 0, 1. \quad (40)$$

Since  $L$  and  $\beta$  are functions of  $u = t - x$ , Eqs.(40) reduce to the single equation

$$L_{uu} + L\beta_u^2 = 0. \quad (41)$$

Using Qadir-Sharif procedure, we obtain

$$F_0 = -m(\ddot{L} + \dot{\beta}^2)L/\dot{L} = 0, \quad F_i = 0, \quad (42)$$

where a dot denotes differentiation with respect to  $t$ . Consequently the momentum four-vector becomes  $p_a = \text{constant}$ . Thus there is a constant energy and momentum. The constant, here, determines the strength of the wave. This exactly coincides with the Ehlers-Kundt method in which they demonstrate that the test particles acquire a constant momentum and hence a constant energy, from the plane gravitational waves.

### (ii) Cylindrical Gravitational Waves

To describe cylindrical gravitational waves one uses cylindrical polar coordinates  $(\rho, \theta, \phi)$  and the time  $t$ , and one takes the line element in the form [8]

$$ds^2 = e^{2\gamma-2\psi}(dt^2 - d\rho^2) - \rho^2e^{-2\psi}d\phi^2 - e^{2\psi}dz^2, \quad (43)$$

where  $\gamma = \gamma(\rho, t)$ ,  $\psi = \psi(\rho, t)$ . To satisfy the Einstein field equations for empty space one takes

$$\psi_{tt} - \psi_{\rho\rho} - \frac{1}{\rho}\psi_{\rho} = 0, \quad (44)$$

$$\gamma_t = 2\rho\psi_\rho\psi_t, \quad \gamma_\rho = \rho(\psi_\rho^2 + \psi_t^2), \quad (45)$$

where subscript denote partial derivatives. Using ELLP prescription, we obtain the energy-momentum densities given by

$$\Theta_0^0 = \frac{1}{8\pi}e^{2\gamma}(\psi_\rho^2 + \psi_t^2), \quad (46)$$

$$\Theta_1^0 = \frac{1}{4\pi\rho}x\psi_\rho\psi_t, \quad (47)$$

$$\Theta_2^0 = \frac{1}{4\pi\rho}y\psi_\rho\psi_t, \quad (48)$$

$$\Theta_0^1 = -e^{2\gamma}\Theta_1^0, \quad (49)$$

$$\Theta_0^2 = -e^{2\gamma}\Theta_2^0, \quad (50)$$

$$\Theta_0^3 = \Theta_3^0 = 0. \quad (51)$$

The energy density of the cylindrical gravitational waves is finite and positive definite, and the momentum density components reflect the symmetry of the spacetime.

### (iii) A Class of Rotating Cylindrical Gravitational Waves

A class of solutions of the gravitational field equations describing vacuum spacetimes outside rotating cylindrical sources is given by the line element of the form [33]

$$ds^2 = e^{2\gamma-2\psi}(dt^2 - d\rho^2) - \mu^2e^{-2\psi}(\omega dt + d\phi)^2 - e^{2\psi}dz^2, \quad (52)$$

in the cylindrical coordinates  $(\rho, \phi, z)$ . Here the metric functions  $\gamma, \mu, \psi$  and  $\omega$  depend on the coordinates  $t$  and  $\rho$  only. When  $\omega = 0$ , the metric represents spacetimes without rotation, in which the polarization of gravitational waves has only one degree of freedom and the direction of polarization is fixed [5]. It is to be noticed that if we take  $\omega = 0$  and  $\mu = \rho$ , the above metric reduces to a special case of cylindrical gravitational waves [8]. Einstein's vacuum field equations for the metric form (52) are given by

$$(\mu\psi_v)_u + (\mu\psi_u)_v = 0, \quad (53)$$

$$\mu_{uv} - \frac{l^2}{8}\mu^{-3}e^{2\gamma} = 0, \quad (54)$$

$$\omega_v - \omega_u = l\mu^{-3}e^{2\gamma}, \quad (55)$$

$$\gamma_u = \frac{1}{2\mu_u}(\mu_{uu} + 2\mu\psi_u^2), \quad (56)$$

$$\gamma_v = \frac{1}{2\mu_v}(\mu_{vv} + 2\mu\psi_v^2), \quad (57)$$

where  $\psi_u = \frac{\partial\psi}{\partial u}$ , etc. The subscripts  $u = t - \rho$  and  $v = t + \rho$  are retarded and advanced times respectively. Here  $l$  is a constant length characteristic of the rotation of the system which is positive and is specifically attributed with rotating gravitational waves. For  $l = 0$  we have  $\omega = \omega(t)$  from Eq.(55) and  $\mu_{tt} = \mu_{\rho\rho}$  from Eq.(54). A simple transformation to a rotating frame reduces the waves to non-rotating generalized Beck spacetimes which have been studied by many authors [32,34,35].

The energy and momentum densities in Einstein's prescription are

$$\Theta_0^0 = \frac{1}{16\pi\mu^2\rho^3}[\mu^2(-\mu + \mu_{\rho\rho} - 2\mu_{\rho\rho}\rho^2 - \mu\omega^2\rho^2 - 2\mu\omega\omega_{\rho\rho} - \mu_{\rho\rho}\omega^2\rho^3) \quad (58)$$

$$+ \rho^2(\mu + 2\mu\gamma_{\rho\rho} - \mu_{\rho\rho})e^{2\gamma} - \mu^4\rho^2(\mu\omega_{\rho}^2 + \mu\omega\omega_{\rho\rho} - 2\mu\omega\omega_{\rho}\gamma_{\rho} + 3\mu_{\rho}\omega\omega_{\rho})e^{-2\gamma}],$$

$$\Theta_1^0 = \frac{1}{16\pi\rho^5}[2\mu\rho^2\dot{\gamma}x - 6\dot{\mu}x^3 - 2\dot{\mu}\rho^2x - 2\dot{\mu}_{\rho}\rho x^3 - \mu\omega^2\rho y \quad (59)$$

$$- \mu^2\rho y(\mu\omega_{\rho} - \mu\omega_{\rho\rho}\rho + 2\mu\omega_{\rho}\gamma_{\rho\rho} - 3\mu_{\rho}\omega_{\rho\rho})e^{-2\gamma}],$$

$$\Theta_2^0 = \frac{1}{16\pi\rho^4}[2\mu\rho\dot{\gamma}y - 2\dot{\mu}_{\rho}\rho^2y + \mu\omega\rho x - \mu\omega_{\rho}\rho^2x - \mu_{\rho}\omega\rho^2x \quad (60)$$

$$+ \mu^2x(\mu\omega_{\rho} - \mu\omega_{\rho\rho}\rho + 2\mu\omega_{\rho}\gamma_{\rho\rho} - 3\mu_{\rho}\omega_{\rho\rho})e^{-2\gamma}],$$

$$\Theta_0^1 = \frac{1}{16\pi\mu^2\rho^3}[\mu^2(-\dot{\mu}x + 2\dot{\mu}_{\rho}\rho x + 2\mu\omega\dot{\omega}\rho^2x - 2\mu\omega\ddot{\gamma}\rho^2y + 2\mu\omega_{\rho}\gamma_{\rho\rho}\rho^2y \quad (61)$$

$$- \mu\omega_{\rho\rho}\rho^2y + 2\mu\omega\gamma_{\rho\rho}\rho^2y + \dot{\mu}\omega^2\rho^2x + 2\mu_{\rho}\omega\gamma_{\rho\rho}\rho^2y - 3\mu_{\rho}\omega_{\rho\rho}\rho^2y - 2\mu_{\rho\rho}\omega\rho^2y) - \rho^2x(2\mu\dot{\gamma} - \dot{\mu})e^{2\gamma} + \mu^4\rho x(\mu\dot{\omega}_{\rho} + \mu\omega\dot{\omega}_{\rho} - 3\mu\omega\omega_{\rho}\dot{\gamma} + 3\dot{\mu}\omega\omega_{\rho})e^{-2\gamma}],$$

$$\Theta_0^2 = \frac{1}{16\pi\mu^2\rho^3}[\mu^2(2\dot{\mu}_{\rho}\rho y + \mu\omega_{\rho\rho}\rho^2y - 2\mu\omega_{\rho}\gamma_{\rho\rho}\rho^2y - 2\mu\omega\gamma_{\rho\rho}\rho^2y + 2\mu\omega\dot{\omega}\rho^2y \quad (62)$$

$$+ 2\mu\dot{\omega}\dot{\gamma}\rho^2x + 2\mu\omega\ddot{\gamma}\rho^2x + \dot{\mu}\omega^2\rho^2y + 2\dot{\mu}\omega\dot{\gamma}\rho^2x + 3\mu_{\rho}\omega_{\rho\rho}\rho^2y - 2\mu_{\rho}\omega\gamma_{\rho\rho}\rho^2y + 2\mu_{\rho\rho}\omega\rho^2y) - \rho^2y(2\mu\dot{\gamma} - \dot{\mu})e^{2\gamma} + \mu^4\rho y(\mu\dot{\omega}_{\rho} + \mu\omega\dot{\omega}_{\rho} - 2\mu\omega\omega_{\rho}\dot{\gamma} + 3\dot{\mu}\omega\omega_{\rho})e^{-2\gamma}],$$

$$\Theta_0^3 = \Theta_3^0 = 0. \quad (63)$$

Now for  $\omega = 0$  and  $\mu = \rho$ , Eqs.(58)-(62) become the energy and momentum densities of cylindrical gravitational waves given by Rosen and Virbhadra [13].

The energy and momentum density components in Papapetrou's prescription is given by

$$\Omega^{00} = \frac{1}{16\pi\mu^2\rho^3}[\mu^2(-\mu + \mu_{\rho\rho} - 2\mu_{\rho\rho}\rho^2 - \mu\omega^2\rho^2 - 2\mu\omega\omega_{\rho}\rho^3 - \mu_{\rho}\omega^2\rho^3) \quad (64)$$

$$+(\mu\rho^2 + 2\mu\gamma_{\rho}\rho^3 - \mu_{\rho}\rho^3)e^{2\gamma}],$$

$$\Omega^{01} = \frac{1}{16\pi\mu^2\rho^3}[\mu^2(-\dot{\mu}x + 2\dot{\mu}_{\rho}\rho x + \mu\omega y - \mu\omega_{\rho}\rho y - \mu\omega_{\rho\rho}\rho^2 y \quad (65)$$

$$+ 2\mu\omega\dot{\omega}\rho^2 x + \dot{\mu}\omega^2\rho^2 x - \mu_{\rho}\omega\rho y - 2\mu_{\rho}\omega_{\rho}\rho^2 y$$

$$- \mu_{\rho\rho}\omega\rho^2 y) - \rho^2 x(2\mu\dot{\gamma} - \dot{\mu})e^{2\gamma}],$$

$$\Omega^{02} = \frac{1}{16\pi\mu^2\rho^3}[\mu^2(\dot{\mu}y + 2\dot{\mu}_{\rho}\rho y + 2\mu\omega\dot{\omega}\rho^2 y + \dot{\mu}\omega^2\rho^2 y - \mu\omega x \quad (66)$$

$$+ \mu\omega_{\rho}\rho x + \mu\omega_{\rho\rho}\rho^2 x + \mu_{\rho}\omega\rho x + 2\mu_{\rho}\omega_{\rho}\rho^2 x$$

$$+ \mu_{\rho\rho}\omega\rho^2 x) - \rho^2 y(2\mu\dot{\gamma} - \dot{\mu})e^{2\gamma}],$$

$$\Omega^{03} = \Omega^{30} = 0. \quad (67)$$

We see that for  $\omega = 0$  and  $\mu = \rho$ , Eqs.(64)-(66) yield the same result as given by Virbhadra [13].

#### (iv) Spherical Gravitational Waves

The gravitational waves with spherical wavefronts are given by the line element of the form [36]

$$ds^2 = e^{-M}(dt^2 - d\rho^2) - e^{-U}(e^V dz^2 + e^{-V} d\phi^2), \quad (68)$$

where the metric functions  $U, V$  and  $M$  depend on the coordinates  $t$  and  $\rho$  only. Einstein's vacuum field equations imply that  $e^{-U}$  satisfies the wave equation

$$(e^{-U})_{tt} - (e^{-U})_{\rho\rho} = 0, \quad (69)$$

and that  $V$  satisfies the linear equation

$$V_{tt} - U_t V_t - V_{\rho\rho} + U_{\rho} V_{\rho} = 0. \quad (70)$$

The remaining equations for  $M$  are

$$U_{tt} - U_{\rho\rho} = \frac{1}{2}(U_t^2 + U_\rho^2 + V_t^2 + V_\rho^2) - U_t M_t - U_\rho M_\rho = 0, \quad (71)$$

$$2U_{t\rho} = U_t U_\rho - U_t M_\rho - U_\rho M_t + V_t V_\rho. \quad (72)$$

It is well known that, if Eqs.(69) and (70) are satisfied, the Eqs.(71) and (72) are automatically integrable.

Using Qadir-Sharif prescription, we obtain

$$F_0 = m\left[\dot{U} + \frac{\ddot{M} + 2\ddot{U}}{\dot{M} + 2\dot{U}} - \frac{3\dot{U}^2 + \dot{V}^2}{\dot{M} + 2\dot{U}}\right], \quad F_1 = -m\frac{M'}{2}, \quad F_2 = 0 = F_3. \quad (73)$$

The corresponding four-vector momentum will become

$$p_0 = m\left[U + \ln(\dot{M} + 2\dot{U}) - \int \frac{3\dot{U}^2 + \dot{V}^2}{\dot{M} + 2\dot{U}} dt\right] + f_1(\rho), \quad (74)$$

$$p_1 = -\frac{m}{2} \int M' dt + f_2(\rho), \quad p_2 = \text{constant} = p_3. \quad (75)$$

where dot denotes differentiation with respect to time and prime with respect to  $\rho$ ,  $f_1$  and  $f_2$  are arbitrary functions of  $\rho$ . Eqs.(74) and (75) provide the general expression of the momentum four-vector for the gravitational waves with spherical wavefronts. As we are interested in evaluating the momentum imparted by gravitational waves we need to calculate the term  $p_1$ . For this purpose, we require the value of  $M$ .

The background region ( $t < \rho$ , Minkowski) is described by the solution  $U = -\ln t - \ln \rho$ ,  $V = \ln t - \ln \rho$  and  $M = 0$ . Substituting these values in Eqs.(74) and (75), we have

$$p_0 = m \ln(-2/\rho) + f_1(\rho), \quad p_\alpha = \text{constant}. \quad (76)$$

The quantity  $p_0$  can be made zero by choosing  $f_1(\rho) = -m \ln(-2/\rho)$  and the momentum term  $p_\alpha$  will be zero for a particular choice of an arbitrary constant as zero. Thus the four-vector momentum vanishes in the background region (Minkowski) as was expected.

The solution on the wavefront ( $t = \rho$ ) can be written in the form  $U = -2 \ln t$ ,  $V = 0$ ,  $M = 0$ . Using these values in Eqs.(74) and (75), it follows that

$$p_0 = m \ln(-4) + f_1(\rho), \quad p_\alpha = \text{constant}. \quad (77)$$

We see that the momentum turns out to be constant which can be made zero if we choose constant as zero.

The solution in the wave region ( $t > \rho$ ) can be found by solving Eqs.(69) and (70) and is given in the form [35,36]

$$U = -\ln t - \ln \rho, \quad V = \ln t - \ln \rho + \tilde{V}(t, \rho). \quad (78)$$

The case of a single component gives

$$\tilde{V}(t, \rho) = a_k (t\rho)^k H_k \left( \frac{t^2 + \rho^2}{2t\rho} \right) \quad (79)$$

for some  $k \geq \frac{1}{2}$  and constant  $a_k$ . In this case,  $M$  is given by

$$M = \frac{1}{2k} a_k (t^2 - \rho^2) (t\rho)^{k-1} H_{k-1} - \frac{1}{2k} (t\rho)^{2k} a_k^2 \left[ k^2 H_k^2 - \frac{(t^2 - \rho^2)^2}{4t^2 \rho^2} H_{k-1}^2 \right]. \quad (80)$$

Notice that the dimension of  $a_k$  is  $L^{-2k}$ . For the purpose of simplicity, we take a special case when  $k = 1$  for which  $M$  takes the form

$$M = \frac{1}{2} a_1 (t^2 - \rho^2) H_0 - \frac{1}{2} (t\rho)^2 a_1^2 \left[ H_1^2 - \frac{(t^2 - \rho^2)^2}{4t^2 \rho^2} H_0^2 \right], \quad (81)$$

where  $H_0 = \ln(t/\rho)$ ,  $H_1 = \frac{1}{2}[(t/\rho + \rho/t) \ln(t/\rho) - (t/\rho - \rho/t)]$ . After taking derivative of  $M$  with respect to  $\rho$ , we substitute it in Eq.(75) and after a tedious integration, we obtain

$$p_1 = m \frac{a_1}{4} \left[ \frac{2}{3} (t^3/\rho) - 6t\rho + 4t\rho \ln(t/\rho) + a_1 \left\{ \frac{1}{5} (t^5/\rho) + \frac{2}{27} t^3 \rho + 5t\rho^3 \right. \right. \quad (82) \\ \left. \left. - 4 \left( \frac{5}{9} t^3 \rho + t\rho^3 \right) \ln(t/\rho) + \frac{4}{3} t^3 \rho (\ln(t/\rho))^2 \right\} \right] + f_2(\rho).$$

This gives the momentum imparted to test particles by gravitational waves with spherical wavefronts. The quantity  $p_1$  can be made zero for the  $t \rightarrow 0$  limit by choosing  $f_2 = 0$ . However, it immediately indicates the presence of singularity when  $\rho = 0$  and this singularity at  $\rho = 0$  acts as a source of the gravitational waves inside the wave region. This coincides with the result evaluated by using Möller's prescription [18]. This is a physically reasonable expression for the momentum imparted by gravitational waves. The interpretation of  $p_0$  in the  $e\psi N$  formalism is given elsewhere [37]. It can also be shown that, near the wavefront as  $t \rightarrow \rho$

$$\tilde{V} \sim a_k (t + \rho)^{-1} (t - \rho)^{1+2k}, \quad M \sim \left( \frac{1 + 2k}{2k} \right) a_k (t - \rho)^{2k}. \quad (83)$$

Using Eqs.(75) and (83), it follows that

$$p_1 = \frac{1}{4k}m(1 + 2k)a_k(t - \rho)^{2k} + f_2(\rho). \quad (84)$$

This gives the momentum near the spherical wavefront. We see that for a particular choice of  $f_2$ , it reduces to the momentum expression given by Eq.(77) on the wavefront. We remark that our results exactly coincide with those evaluated by using Möller's prescription for the background region and on the wavefront. For the wave region, these two can be equated for a particular choice of an arbitrary function  $f_2$ . We have seen that in all the three cases we obtain a physically reasonable expression for the momentum.

## 5 Discussion

It is usually believed that different energy-momentum complexes could give different results for a given geometry . Keeping this point in mind, we have applied various prescriptions to different cosmological models. We have also extended this analysis to gravitational waves. For many spacetimes, it is found that various methods could give the same result which is physically well-defined.

The Bianchi type I metric gives zero energy-distribution in ELLPW prescriptions. For the axially symmetric scalar field, using Möller's method, the energy distribution becomes the mass  $M$  which coincides with the Schwarzschild metric. We also obtain the physically interesting results for the charged regular black hole, KN metric and Melvin's magnetic universe as given in the analysis.

Further, we have evaluated energy and momentum distribution for gravitational waves by different prescriptions. The Ehler-Kundt method gives the physically reasonable result that plane gravitational waves impart a constant energy and momentum to test particles in their path. However, it does not provide a simple formula that can be applied to other cases. We have obtained the same result using Qadir-Sharif formalism. For cylindrical gravitational waves, using ELLP energy-momentum distribution, we have obtained the similar and physically acceptable result. We have also calculated the energy-momentum distribution for a class of cylindrical gravitational waves using EP prescriptions. It can be seen that the energy and momentum densities for a class of rotating gravitational waves are finite and well-defined in

both the prescriptions. It follows from Eqs.(58-63) and (64-67) that though the energy-momentum complexes of Einstein and Papapetrou are not exactly the same but are similar upto certain terms. However, it is interesting to note that both the results reduce to the same energy and momentum densities of a special case of cylindrical gravitational waves as given in [13,14].

Finally, we have applied the Qadir-Sharif prescription to spherical gravitational waves. It is interesting to note that this provide physical acceptable result. This result supports the result evaluated by using Möller's prescription. We can conclude that different prescriptions can provide the same meaningful result.

## Acknowledgment

The author would like to thank MOST for funding this work.

## References

- [1] Penrose, R.: Proc. Roy. Soc. London **A381**(1982)53.
- [2] Misner, C.W. and Sharp, D.H. Phys. Rev. **136**(1964)B571.
- [3] Bondi, H.: Proc. Roy. Soc. London **A427**(1990)249.
- [4] Cooperstock, F.I.: Found. Phys. **22**(1992)1011; in *Topics in Quantum Gravity and Beyond: Papers in Honor of L. Witten* eds. Mansouri, F. and Scario, J.J. (World Scientific, Singapore, 1993) 201; in *Relativistic Astrophysics and Cosmology*, eds. Buitrago et al (World Scientific, Singapore, 1997)61; Annals Phys. **282**(2000)115.
- [5] Misner, C.W., Thorne, K.S. and Wheeler, J.A.: *Gravitation* (W.H. Freeman, San Francisco, 1973).
- [6] Ehlers, J. and Kundt, W.: *Gravitation: An Introduction to Current Research*, ed. L. Witten (Wiley, New York, 1962)49.
- [7] Pirani, F.A.E.: *Gravitation: An Introduction to Current Research*, ed. L. Witten (Wiley, New York, 1962)199.
- [8] Weber, J. and Wheeler, J.A.: Rev. Mod. Phy. **29**(1957)509;  
Weber, J.: *General Relativity and Gravitational Waves*, (Interscience, New York, 1961).
- [9] Qadir, A. and Sharif, M.: Physics Letters **A 167**(1992)331.
- [10] Rosen, N: Helv. Phys. Acta. Suppl. **4**(1956)171.
- [11] Scheidegger, A.E.: Rev. Mod. Phys. **25**(1953)451.
- [12] Rosen, N.: Phys. Rev. **291**(1958)110.

- [13] Rosen, N. and Virbhadra, K.S.: Gen. Rel. and Grav. **26**(1993)429.
- [14] Virbhadra, K.S.: Pramana-J. Phys. **45**(1995)215.
- [15] Brown, J.D. and York, J.w.: Phys. Rev. D **47**(1993)1407.
- [16] Tolman, R.C.: *Relativity, Thermodynamics and Cosmology* (Oxford University Press, Oxford).
- [17] Landau, L.D. and Lifshitz, E.M.: *The Classical Theory of Fields* (Pergamon Press, Oxford 1987).
- [18] Möller, C.: Ann. Phys. (NY) **4**(1958)347.
- [19] Komar, A.: Phys. Rev. **113**(1959)934.
- [20] Möller, C.: Ann. Phys. (NY) **12**(1961)118;
- [21] Bergqvist, G.: Class. Quantum Grav. **9**(1992)1753; Bernstein, D.H. and Tod, K.P.: Phys. Rev. **D 49**(1994)2808.
- [22] Papapetrou, A.: Proc. R. Irish. Acad. **A52**(1948)11.
- [23] Weinberg, S.: *Gravitation and Cosmology: Principles and Applications of General Theory of Relativity* (John Wiley and Sons, Inc., New York, 1972).
- [24] Tyron, E.P: Nature **246**(1973)396.
- [25] Matos, T., Nunez, D. and Quevedo, H.: Phys. Rev. **D 51**(1995)310.
- [26] Matos, T., Nunez, D. and Rios, M.: Class. Quantum Grav.**17**(2000)3917.
- [27] E. Ayon-Beato and A. Garcia: Phys. Lett. **B464**(1999)25.
- [28] Tod, K. P.: Proc. R. Soc. London **A388**(1983)457.
- [29] Aguirregabiria, J.M., Chamorro, A. and Virbhadra, K.S.: Gen Rel. Grav. **28**(1996)1393.
- [30] Cohen, J.M. and de Felice, F.: J. Math. Phys. **25**(1984)992.
- [31] Ernst, F.J.: J. Math. Phys. **15**(1975)1409; **17**(1976)54.

- [32] Kramer, D., Stephani, H. , MacCallum, M. and Herlt, E.: *Exact Solutions of Einstein's Field Equations* (Cambridge University Press, Cambridge, 1980).
- [33] Mashhoon, Bahram , McClune, James C. and Quevedo, Hernando: Class. Quantum Grav.**17**(2000)533.
- [34] Verdaguer, E.: Phys. Rep. **229**(1993)1;  
Griffths, J.B.: *Colliding Plane Waves in General Relativity* (Oxford University Press, Oxford, 1991).
- [35] Alekseev, G.A. and Griffths, J.B.: Class. Quantum Grav. **13**(1996)2191.
- [36] Alekseev, G.A. and Griffths, G.B.: Class. Quantum Grav.**12**(1995)L13.
- [37] Qadir, Asghar, Sharif, M. and Shoaib, M.: Nuovo Cimento **B 13**(2000)419.