Semiclassical States in Quantum Cosmology: Bianchi I Coherent States

Brett Bolen,^{1, 2, *} Luca Bombelli,^{1, 3, †} and Alejandro Corichi^{1, 4, ‡}

¹Department of Physics and Astronomy, University of Mississippi University, MS 38677-1848, U.S.A. ²Department of Physics, Rhodes College Memphis, TN 38112-1690, U.S.A. ³Perimeter Institute for Theoretical Physics

35 King Street North, Waterloo, Ontario, Canada N2J 2W9

⁴Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México

Apartado Postal 70-543, México D.F. 04510, México

(Dated: 9 June 2004)

Abstract

We study coherent states for Bianchi type I cosmological models, as examples of semiclassical states for time-reparametrization invariant systems. This simple model allows us to study explicitly the relationship between exact semiclassical states in the kinematical Hilbert space and corresponding ones in the physical Hilbert space, which we construct here using the group averaging technique. We find that it is possible to construct good semiclassical physical states by such a procedure in this model; we also discuss the sense in which the original kinematical states may be a good approximation to the physical ones, and the situations in which this is the case. In addition, these models can be deparametrized in a natural way, and we study the effect of time evolution on an "intrinsic" coherent state in the reduced phase space, in order to estimate the time for this state to spread significantly.

PACS numbers: 04.60.Ds, 98.80.Qc, 03.65.Sq.

^{*}Electronic address: brett@phy.olemiss.edu

[†]Electronic address: luca@phy.olemiss.edu

[‡]Electronic address: corichi@nuclecu.unam.mx

I. INTRODUCTION

Bianchi cosmological models have long been used as toy models in both classical general relativity and quantum gravity. The small but non-trivial number of degrees of freedom they possess makes their dynamics interesting from the point of view of shedding light on features of the full theory, even though it reduces to that of a relativistic particle in a low-dimensional space. These models are homogeneous cosmologies, in which the symmetries are imposed on the canonical action or Hamiltonian before the dynamics is obtained. This procedure is, in general, not equivalent to imposing the symmetries on the full dynamical equations, but in a classical theory this can only potentially give some extra solutions with respect to the symmetric ones for the full theory, and is not a problem [1]. Thus, the classical Bianchi models can be very fruitfully used in exploring issues such as the evolution of anisotropies in simplified versions of the early universe, or the chaotic nature of the evolution of the metric close to the initial singularity.

In the quantum theory, the difference between any physical prediction of a minisuperpace model and the "full theory" may be more substantial [2], but it is generally believed that some qualitative features of the models are still useful indications of what to expect from the full theory when different approaches to its quantization are used. In particular, minisuperspace models are time-reparametrization invariant, thus exhibiting the much celebrated "problem of time" [3]; they represent then a fruitful playground in which to explore quantization methods, in the hope of getting some insight on how to approach this problem in full quantum gravity. It is not surprising then that minisuperpaces have been analyzed from the Euclidean path integral approach to quantum gravity [4], consistent histories [5] and refined algebraic quantization [6] perspectives.

It is in this spirit that we study here coherent states of minisuperspace models. The aim is to understand the role of coherent states in defining semiclassical states for timereparametrization invariant systems. In recent years, an increasing amount of work has been devoted to understanding the phenomenology of semiclassical states in quantum gravity, by which here we mean that sector of the physical states in which the gravitational degrees of freedom are sharply peaked around values satisfying the classical constraint equations, because it is expected that these states will possess quasi-classical properties in certain regimes [7]. The study of the semiclassical limit of quantum cosmological models is of course not new; however, most of the treatments that we are aware of deal with the WKB method, where the wave function is assumed to be of a special form and some component of it is shown to satisfy the Hamilton-Jacobi equation (for a review, see Ref [8]).

In this paper, for simplicity, we consider vacuum Bianchi type I models, whose classical description can be reduced to that of a relativistic particle moving in a 3-dimensional, linear configuration space, subject to the Hamiltonian constraint of general relativity, which in this case reduces to the free particle Hamiltonian. These models have been extensively studied, but the type of questions we are concerned with here, however, have been only recently considered within the context of constrained systems in Ref [7], where a general framework for the construction of semiclassical states for a class of constrained theories is discussed. Here, we follow much of that approach in spirit, but the constraint and physical observables that arise in these Bianchi models are not of the same type and we have to adapt the general constructions to the present situation, and the fact that we are dealing with cosmological models allows us to ask additional physically interesting questions.

Despite the simplicity of the classical model, the quantum counterpart is non-trivial, be-

cause of the presence of the constraint. First, we consider coherent states in the kinematical Hilbert space, peaked around a point on the constraint surface in phase space but without taking the quantum constraint into account, and characterized by a set of "squeezing" parameters. These states are only meant to be approximations to physical coherent states, i.e., solutions to the quantum constraint, which we find by applying the group averaging procedure of the Refined Algebraic Quantization program (RAQ) [6] to the kinematical coherent states; one expects these to be the ones that capture the "essential" features of semiclassical Bianchi I models. Finally, we consider the reduced phase space, described by true Dirac observables, and define coherent states. The main purpose of this article is to compare these three classes of states, in terms of describing semiclassical situations, assess how good the kinematical coherent states are as approximations to the physical ones, and use the "intrinsic" ones to get an insight into evolution issues.

The structure of the paper is the following. In Sec. II we recall some basic notions about Bianchi models. In Sec. III we recall the notion of coherent states, and discuss what we mean by semiclassical states for a system. Sec. IV considers the kinematical coherent states for Bianchi I models, and in Sec. V the Physical coherent states. In Sec. VI we study the time evolution of the "intrinsic coherent states," and in particular some estimates of the time it would take for a coherent state to lose its peakedness properties. We end with a discussion in Sec. VII. Throughout the paper we set c = G = 1, but keep \hbar dimensionful.

II. PRELIMINARIES ON BIANCHI MODELS AND THEIR OBSERVABLES

In this section, we will provide some background information on Bianchi models, the variables used to describe them, and their physical observables; readers who are familiar with these models should be able to skim through it quickly.

Bianchi models are spatially homogeneous cosmologies: they describe spacetimes which can be foliated by spacelike hypersurfaces Σ_t (all diffeomorphic to the "space manifold" Σ), on each of which a group G of isometries acts transitively, and they are classified by the Lie algebra of the group G.

If one uses three linearly independent left-invariant forms σ^i on Σ as a basis for its cotangent space, the spacetime metric can be written in the form $g = -N^2(t) dt \otimes dt + q_{ij}(t) \sigma^i \otimes \sigma^j$, where in some Bianchi models the matrix g_{ij} can be diagonalized and written, in the Misner parametrization [9], as

$$q_{ij} = e^{2\beta^0} \operatorname{diag}(e^{2(\beta^+ + \sqrt{3}\beta^-)}, e^{2(\beta^+ - \sqrt{3}\beta^-)}, e^{-4\beta^+}); \qquad (2.1)$$

we thus see that $\sqrt{\det q} = e^{3\beta^0}$, and $\beta^0(t)$ characterizes the volume of Σ_t , while the $\beta^{\pm}(t)$ characterize its anisotropy; all three functions take values in $(-\infty, +\infty)$. We will take as configuration variables the β^i , with i = 0, +, -; the configuration space is thus $\mathcal{C} = \mathbb{R}^3$. The dynamics of these models then reduces (before imposing the constraints) to that of a relativistic particle in 3 dimensions, and is governed by a Hamiltonian constraint of the form

$$H = \frac{1}{2} \eta^{ij} p_i p_j + U(\beta^i) , \qquad (2.2)$$

in the vacuum case, where p_i is the canonical momentum conjugate to β^i , and η_{ij} a flat metric on C, with line element $\eta_{ij} d\beta^i d\beta^j = -(d\beta^0)^2 + (d\beta^+)^2 + (d\beta^-)^2$, and the form of the potential U depends on the model under consideration. Here we should point out that even when the flat metric resembles that of 3D Minkowski spacetime, it is actually conformally related to the De Witt supermetric on C. As with all general relativistic models, the Hamiltonian is also a constraint, in that allowed classical phase space points have to satisfy the scalar constraint $H(p_i, \beta^i) = 0$; this is the only constraint for these models, since the metric (2.1) satisfies the vector constraint identically.

In this paper, we restrict ourselves to begin with to the simplest models, the vacuum ones of type I (with homogeneity group U(1)³, or the additive \mathbb{R}^3). In this case, $U(\beta^i)$ vanishes identically; this allows us to not only solve the quantum constraint easily, but also to conceptually isolate the issue of comparing the different coherent states on a linear phase space with a relatively simple constraint, from those related to the structure of phase space or the form of the potential. In the canonical quantum theory, following the Dirac prescription, states have to be annihilated by the operator version \hat{H} of the constraint, i.e., be invariant under the group $e^{-i\lambda\hat{H}}$ it generates, and it is therefore very advantageous, to use the *p*-representation, because the Hamiltonian is then just a multiplication operator. We will also rescale the variables by $p_i \mapsto p_i/\hbar$, which makes the new p_i dimensionless (like the β^i); we can think of this alternatively as a rescaling of the action or Hamiltonian, or as a way to give the value of p_i in terms of the quantum unit \hbar , which is useful when characterizing the extent to which a fluctuation is small or large in the classical sense. For simplicity of notation, in the rest of the paper we will use only subscripts, and denote $\beta_i \equiv \beta^i$ (that is, the index is not lowered with the metric η_{ij}).

In order to evaluate candidate semiclassical states by comparing fluctuations of observables, we need to identify the proper set of physical observables to consider. It is natural to look for a sufficient number of Dirac observables, in the sense that their values constitute a full characterization of a (classical, gauge-invariant) state of the system. Since we will be using the *p*-representation, to simplify the terminology it is convenient to call the space $\tilde{\mathcal{C}} = \mathbb{R}^3$ of p_i 's our new configuration space. Classically, configuration observables are then functions $f(p_i)$ on $\tilde{\mathcal{C}}$, and "momentum" observables, specifically linear functions of the canonically conjugate β_i , can be thought of as vector fields u on $\tilde{\mathcal{C}}$, since there is a canonical assignment of a linear momentum observable P_u everywhere in the phase space Γ to each such vector field; we can then associate to these observables quantum operators \hat{f} and \hat{u} acting on $\psi(p)$, defined respectively by

$$(\hat{f}\psi)(p) := f(p)\psi(p) , \qquad (\hat{u}\psi)(p) := i\left(\mathcal{L}_u + \frac{1}{2}\operatorname{div}_{\mu}u\right)\psi(p) , \qquad (2.3)$$

where $\operatorname{div}_{\mu} u$ is the divergence of u with respect to some volume element $d\mu(p)$ on $\tilde{\mathcal{C}}$.

We know that the reduced phase space has four dimensions, so we need four independent observables $\{O_{\alpha}\}_{\alpha=1,\ldots,4}$, in addition to the constraint itself. The first observation is that the variables p_{\pm} are Dirac observables, since the Hamiltonian constraint in the new variables,

$$H(\beta_i, p_i) = \frac{1}{2} \left(-p_0^2 + p_+^2 + p_-^2 \right), \qquad (2.4)$$

depends on the p_i only. The not so obvious task is that of finding a suitable pair of linear momentum observables, or vector fields on $\tilde{\mathcal{C}}$, to represent the conjugate variables, since the β_i , for which the CCR with the p_i are of the form $[\hat{\beta}_i, \hat{p}_j] = i \, \delta^i{}_j$, are not physical observables; in fact, one can easily see that no function of the β_i alone is a constant of the motion. As Dirac "momentum" observables for this model we will use instead the combinations $v_{\pm} := p_0 \beta_{\pm} + p_{\pm} \beta_0$ which, together with $v_0 := p_+ \beta_- - p_- \beta_+$ are associated with the vector fields (denoted here by the same symbols)

$$v_{+} = p_{0} \frac{\partial}{\partial p_{+}} + p_{+} \frac{\partial}{\partial p_{0}} , \qquad v_{-} = p_{0} \frac{\partial}{\partial p_{-}} + p_{-} \frac{\partial}{\partial p_{0}} , \qquad v_{0} = p_{+} \frac{\partial}{\partial p_{-}} - p_{-} \frac{\partial}{\partial p_{+}} , \qquad (2.5)$$

i.e., the two boosts and the spatial rotation that generate the (2+1) Lorentz group [10] on the 'minisuperspace' \tilde{C} .

We may note that in general there is a possible ambiguity in the class of observables, consisting in adding to each one a term proportional to the constraint; such an addition does not change the value of the observable on the constraint surface. One can implement a general strategy to remove the ambiguity [7], but in the present case no such change would preserve the property that all observables are linear in the configuration variables.

III. SEMICLASSICAL AND COHERENT STATES

In this section, we will specify our criteria for a quantum state to be semiclassical, following the general discussion of Ref [7], recall the basic definition of the most common example of such a state, a coherent state, and specify the setup and goals for the rest of the paper, specializing the discussion to Bianchi I cosmology. (For other references on semiclassical states for systems with constraints, see also Refs [13, 14].)

Given a point (\bar{q}_i, \bar{p}_i) in a classical phase space, a complete set of physical observables $\{O_\alpha\}_{\alpha=1,\dots,4}$ for the theory (completeness will be the only requirement, the observables do not need to generate an algebra; they may be overcomplete), and a pair of parameter values $(\epsilon_\alpha, \delta_\alpha)$ for each O_α , we will consider a state ψ to be semiclassical if for each α

$$|\langle \hat{O}_{\alpha} \rangle_{\psi} - O_{\alpha}(\bar{q}_i, \bar{p}_i)| < \epsilon_{\alpha} , \qquad (\Delta \hat{O}_{\alpha})_{\psi} < \delta_{\alpha} .$$

$$(3.1)$$

In other words, we are assigning tolerances; the $(\epsilon_{\alpha}, \delta_{\alpha})$ give a measure of how close the expectation values must be to the classical values, and how narrowly peaked a state must be, for the latter to be considered semiclassical. One often takes the O_{α} to be the canonical variables (\bar{q}_i, \bar{p}_i) themselves, but this is not necessary; in some cases, as we will see, one is forced to make other choices, and in general the choice can affect in a crucial way which states are considered to be semiclassical.

If the system has a set of constraints $H_I(q_i, p_i) = 0$ (assumed here to be first class), a couple of additional points need to be addressed. First of all, the observables need to be physical in the Dirac sense of commuting (weakly) with the constraints, $\{O_{\alpha}, H_I\} \approx 0$, but they need only be complete in the reduced phase space—if one knows how to characterize it. Secondly, if the state does not belong to the physical Hilbert space, we specify the value of an additional pair of tolerance parameters (ϵ_I, δ_I) for each constraint, and require that $\forall I$

$$|\langle \hat{H}_I \rangle_{\psi}| < \epsilon_{\alpha} , \qquad (\Delta \hat{H}_I)_{\psi} < \delta_I . \tag{3.2}$$

An important issue, which in fact underlies the motivation for the current work, arises when one has a constrained system for which one prefers, or is forced, to work with kinematical states. One may want to know in that case whether the kinematical semiclassical states are, in an appropriate sense, good approximations to the physical ones, so that one may extract from the former useful information on the latter, for example regarding quantum fluctuations. The general formulation of this issue deserves further study; here we will just mention that two possible definitions of when $\psi_{\rm kin}$ is a good approximation to $\psi_{\rm phy}$ are: (i) The two states are semiclassical with respect to the same (\bar{q}_i, \bar{p}_i) , with the same tolerances; and the more restrictive definition (ii) For all O_{α}

$$\frac{|\langle \hat{O}_{\alpha} \rangle_{\rm phy} - \langle \hat{O}_{\alpha} \rangle_{\rm kin}|}{|\langle \hat{O}_{\alpha} \rangle_{\rm phy}|} , \frac{|(\Delta \hat{O}_{\alpha})^2_{\rm phy} - (\Delta \hat{O}_{\alpha})^2_{\rm kin}|}{(\Delta \hat{O}_{\alpha})^2_{\rm phy}} \ll 1 .$$
(3.3)

Qualitatively, the first definition simply aims at identifying situations in which the spreads of measured values for the O_{α} in $\psi_{\rm kin}$ and $\psi_{\rm phy}$ are both within the same tolerances, while the second one may be suited for situations in which one would like all quantities which can be calculated from the spread of values of O_{α} with the two states to nearly coincide.

It would be natural to ask at this point whether the values of the tolerances $(\epsilon_{\alpha}, \delta_{\alpha})$ can be arbitrarily chosen. Although the definition above is consistent for arbitrary (positive) values, if the δ_{α} are too small no state will satisfy the conditions. For example, in the case of a pair of conjugate variables (p, q), choosing δ_p and δ_q such that $\delta_p \delta_q < \hbar/2$ would be too restrictive; in general, quantum uncertainty relations will translate into some (macroscopically) mild conditions that set lower bounds for combinations of δ_{α} 's. The ϵ_{α} in principle can be arbitrary, but in practice they may be restricted by the way the states are constructed, as mentioned below, and by their interpretation. A physically motivated choice would be to set the value of each ϵ_{α} to be close to, or smaller than, that of the corresponding δ_{α} , with the latter being smaller than the experimental resolution for O_{α} .

We should also comment here that it may seem like the introduction of the parameters ϵ_{α} is unnecessary, since one can usually choose the states in such a way that the expectation values of the \hat{O}_{α} agree with their classical values. This is correct, in principle; however, some of the physical states we will use are constructed (using group averaging) out of states with the right expectation values for our choice of physical observables, but do not have this property themselves. Besides, it may sometimes simplify calculations to choose states which don't have exactly the classical expectation values for the chosen \hat{O}_{α} .

Coherent states are among the most commonly used states with a semiclassical interpretation; they are relatively easy to work with, are often associated with realizable physical situations, and satisfy minimum uncertainty relations. However, those relations only refer to the product of uncertainties for pairs of conjugate variables, and not individual uncertainties, so given some set of $\{(O_{\alpha}, \epsilon_{\alpha}, \delta_{\alpha})\}_{\alpha=1,\dots,4}$, not all coherent states may satisfy our definition of semiclassical states. To look at this question, let us summarize the freedom available in defining coherent states for a theory.

Consider a theory with a linear phase space $\Gamma = \mathbf{R}^{2N} = \{(q_i, p_i)\}_{i=1,\dots,N}$, where the phase space coordinates are dimensionless. Then a coherent state centered at $(\bar{q}_i, \bar{p}_i) \in \Gamma$ in the *p*-representation is (the square root of) a Gaussian wave function [15],

$$\psi_{\bar{p},\bar{q},\sigma}(p) = \prod_{i=1}^{N} \mathcal{N}_{i} \exp\left\{-(p_{i}-\bar{p}_{i})^{2}/4\sigma_{i}^{2}-\mathrm{i}\,\bar{q}_{i}\,(p_{i}-\bar{p}_{i})\right\},\tag{3.4}$$

which depends on the N parameters σ_i , the widths of the Gaussian in the various p_i directions; $\mathcal{N}_i = (\sqrt{2\pi}\sigma_i)^{-1/2}$ is a normalization factor. The values of the σ_i are the freedom we have (once we fix the classical phase space point (\bar{q}_i, \bar{p}_i)) to ensure that the state is semiclassical. Since the Fourier transform of a Gaussian of width σ_i is another Gaussian, of width $1/2\sigma_i$, we expect that for each choice of observables and tolerances $\{(O_\alpha, \epsilon_\alpha, \delta_\alpha)\}$ there will be a finite range of values of the σ_i for which the state $\psi_{\bar{p},\bar{q},\sigma}$ is semiclassical, provided the tolerances are large enough to be consistent with the uncertainty relations.

In the familiar case of a simple harmonic oscillator, where the Hamiltonian contains a pair of parameters (m_i, ω_i) for each degree of freedom, one normally takes $\sigma_i = \sqrt{m_i \omega_i \hbar/2}$, and ends up with the coherent states commonly used in that case, while other choices for the σ_i lead to squeezed states. Here, we don't have such dimensionful parameters available, and it might seem that this fact will make the process of defining coherent states somewhat difficult. In reality, this is not the case; we don't have the same natural choice for the σ_i , but we have instead the freedom of choosing values that will make our states satisfy our semiclassicality conditions.

The situation for Bianchi I models is then as follows. Our observables O_{α} are the p_{\pm} and v_{\pm} , and we have a single constraint H; since the two quantities in each of the two observable pairs have a similar physical interpretation, we will assign the same tolerances to them. Thus, given a classical state $(\bar{p}_i, \bar{\beta}_i) \in \bar{\Gamma}$ and a set of tolerance parameters $(\epsilon_p, \epsilon_v, \delta_p, \delta_v, \epsilon_H, \delta_H)$, a quantum state ψ will be called semiclassical if it satisfies the conditions

$$|\langle \hat{p}_{\pm} \rangle_{\psi} - \bar{p}_{\pm}| < \epsilon_p , \qquad |\langle \hat{v}_{\pm} \rangle_{\psi} - \bar{v}_{\pm}| < \epsilon_v , \qquad (\Delta \hat{p}_{\pm})_{\psi} < \delta_p , \qquad (\Delta \hat{v}_{\pm})_{\psi} < \delta_v , \qquad (3.5)$$

and, if ψ is not a solution of the quantum constraint,

$$|\langle \hat{H} \rangle_{\psi}| < \epsilon_H , \qquad (\Delta \hat{H})_{\psi} < \delta_H .$$

$$(3.6)$$

Notice that we could have chosen the observables p_0 and v_0 as part of our set, and assigned tolerances to them as well. In this model, our results would not have been very different, although in general one cannot even assume that observables which are functions of other operators that have small quantum fluctuations have small fluctuations themselves.

In the kinematical Hilbert space, based on a linear configuration space, we can use coherent states of the form (3.4). Then, given a set of tolerances we will have more inequalities than free parameters (the σ_i or a set of equivalent ones); although it is reasonable to expect that solutions do exist for sufficiently large ($\epsilon_{\alpha}, \delta_{\alpha}$), one of our goals is to check that this is true for "small" tolerances as well. In the physical Hilbert space these states cannot be used directly, but we will follow the group averaging procedure to set up appropriately modified coherent states peaked around points of the reduced phase space, check if they are semiclassical as well, and whether the original kinematical states can be considered good approximations. In the concluding section, we will mention models for which the configuration space is not linear, in which case one needs to replace the coherent states defined above by a different choice. The specific form of the states used will depend on the characteristics of the model, and we will discuss some options later.

IV. KINEMATICAL COHERENT STATES

The first step in the quantization procedure is to find a representation of the CCR on the relevant Hilbert space. In our case, given the form of the Hamiltonian, the most convenient choice for the kinematical Hilbert space is to set $\mathcal{H}_{kin} = L^2(\tilde{\mathcal{C}}, dp_j)$. In this representation, the \hat{p}_j operators act by multiplication and the $\hat{\beta}_i$ act as $\hat{\beta}_i = i \partial/\partial p_i$; this is the standard Schrödinger representation of the CCR. As for the additional observables v_i , since the measure μ on \mathcal{H}_{kin} is trivial, we simply get div_{μ} $v_{\pm} = \text{div}_{\mu} v_0 = 0$, and

$$\hat{v}_{\pm}\psi = i(p_0 \partial_{\pm} + p_{\pm} \partial_0)\psi, \qquad \hat{v}_0\psi = i(p_+ \partial_- - p_- \partial_+)\psi,$$
(4.1)

where $\partial_i := \partial/\partial p_i$.

Given a point $\overline{P} = (\overline{\beta}_i, \overline{p}_i)$ on the constraint surface $\overline{\Gamma}$ in the classical phase space Γ , we construct a coherent state centered at \overline{P} in the above representation with the usual prescription (3.4), as

$$\psi_{\rm kin}(p_i) = \prod_{i=0,\pm} \mathcal{N}_i \exp\left\{-(p_i - \bar{p}_i)^2 / 2\tau_i - i(p_i - \bar{p}_i)\bar{\beta}_i\right\},\tag{4.2}$$

where the free parameters τ_i measure the width of the coherent state in momentum space (in the natural Planck units), analogously to the σ_i in (3.4), and the normalization factors are $\mathcal{N}_i = (\pi \tau_i)^{-1/4}$. In this state, we recover the usual properties of coherent states, namely that the expectation values of the canonical variables are

$$\langle \hat{\beta}_i \rangle_{\rm kin} = \bar{\beta}_i , \qquad \langle \hat{p}_i \rangle_{\rm kin} = \bar{p}_i , \qquad (4.3)$$

and their fluctuations have the following expressions

$$(\Delta \hat{\beta}_i)_{\rm kin}^2 = 1/2\tau_i , \qquad (\Delta \hat{p}_i)_{\rm kin}^2 = \tau_i/2 , \qquad (4.4)$$

so that $(\Delta \hat{\beta}_i)(\Delta \hat{p}_i) = \frac{1}{2}$ for each *i* (no summation over *i*). In practice, given the similar interpretation of β_+ and β_- , we will often set $\tau_+ = \tau_-$, for simplicity. For the moment, we shall leave the three τ_i unspecified, but we will assume that the state is narrowly peaked in the sense that the τ_i are small, $\tau_i/\bar{p}_0^2 \ll 1$ for all *i*.

Although we are not imposing the quantum constraint on this state, since we will compare it with a corresponding physical coherent state, the quantities we need to calculate are the expectation values and fluctuations of the physical operators we have identified, in the state $\psi_{\rm kin}$. We already have those of \hat{p}_{\pm} ; for the remaining ones (among which we include v_0 , although strictly speaking it is not necessary), the expectation values are

$$\langle \hat{v}_{\pm} \rangle_{\rm kin} = \int d^3 p \; \psi^*_{\rm kin}(p_i) \, i \, (p_0 \, \partial_{\pm} + p_{\pm} \, \partial_0) \, \psi_{\rm kin}(p_i) = \bar{p}_0 \bar{\beta}_{\pm} + \bar{p}_{\pm} \bar{\beta}_0 \;, \tag{4.5}$$

$$\langle \hat{v}_0 \rangle_{\rm kin} = \int d^3 p \; \psi^*_{\rm kin}(p_i) \, i \, (p_+ \partial_- - p_- \partial_+) \, \psi_{\rm kin}(p_i) = \bar{p}_+ \bar{\beta}_- - \bar{p}_- \bar{\beta}_+ \; , \qquad (4.6)$$

and for the fluctuations $(\Delta \hat{O}_{\alpha})^2_{\rm kin} = \langle \hat{O}^2_{\alpha} \rangle_{\rm kin} - \langle \hat{O}_{\alpha} \rangle^2_{\rm kin}$ we obtain

$$(\Delta \hat{v}_{\pm})_{\rm kin}^2 = \frac{1}{2} + \frac{1}{4} \left(\frac{\tau_0}{\tau_{\pm}} + \frac{\tau_{\pm}}{\tau_0} \right) + \frac{1}{2} \left(\frac{\bar{p}_{\pm}^2}{\tau_0} + \frac{\bar{p}_0^2}{\tau_{\pm}} \right) + \frac{1}{2} \left(\bar{\beta}_0^2 \tau_{\pm} + \bar{\beta}_{\pm}^2 \tau_0 \right) , \qquad (4.7)$$

$$(\Delta \hat{v}_0)_{\rm kin}^2 = -\frac{1}{2} + \frac{1}{4} \left(\frac{\tau_+}{\tau_-} + \frac{\tau_-}{\tau_+} \right) + \frac{1}{2} \left(\frac{\bar{p}_+^2}{\tau_-} + \frac{\bar{p}_-^2}{\tau_+} \right) + \frac{1}{2} \left(\bar{\beta}_-^2 \tau_+ + \bar{\beta}_+^2 \tau_- \right) \,. \tag{4.8}$$

In addition, for the Hamiltonian constraint operator we get

$$\langle \hat{H} \rangle_{\rm kin} = \frac{1}{4} \left(-\tau_0 + \tau_+ + \tau_- \right)$$
(4.9)

$$(\Delta \hat{H})_{\rm kin}^2 = \frac{1}{2} \left(\bar{p}_0^2 \tau_0 + \bar{p}_+^2 \tau_+ + \bar{p}_-^2 \tau_- \right) + \frac{1}{8} \left(\tau_0^2 + \tau_+^2 + \tau_-^2 \right) \,, \tag{4.10}$$

where we have used the classical constraint $-\bar{p}_0^2 + \bar{p}_+^2 + \bar{p}_-^2 = 0$. Thus, as expected on general grounds [7], the expectation values of all relevant operators agree with the classical values, except for the (quadratic) Hamiltonian. In addition, for the kinematical coherent state to be a good semiclassical state according to our choice of tolerances, the above fluctuations need to satisfy the conditions (3.5) and (3.6).



FIG. 1: The region in the parameter space of (τ_0, τ) where the inequalities for a semiclassical state are satisfied is shown together with several of the curves that bound the region.

Specifically, the only nontrivial condition arising from the expectation values of observables is the one from the constraint in (3.6),

$$|\tau_{+} + \tau_{-} - \tau_{0}| < 4 \epsilon_{H} , \qquad (4.11)$$

and the conditions on the uncertainties in the \hat{p}_{\pm} , the \hat{v}_{\pm} and \hat{H} give, respectively

$$\tau_{\pm} < 2\,\delta_p^2 \tag{4.12}$$

$$\frac{1}{2} + \frac{1}{4} \left(\frac{\tau_0}{\tau_\pm} + \frac{\tau_\pm}{\tau_0} \right) + \frac{1}{2} \left(\frac{\bar{p}_\pm^2}{\tau_0} + \frac{\bar{p}_0^2}{\tau_\pm} \right) + \frac{1}{2} \left(\bar{\beta}_0^2 \tau_\pm + \bar{\beta}_\pm^2 \tau_0 \right) < \delta_v^2 \tag{4.13}$$

$$\frac{1}{2}\left(\bar{p}_{0}^{2}\tau_{0} + \bar{p}_{+}^{2}\tau_{+} + \bar{p}_{-}^{2}\tau_{-}\right) + \frac{1}{8}\left(\tau_{0}^{2} + \tau_{+}^{2} + \tau_{-}^{2}\right) < \delta_{H}^{2}$$

$$(4.14)$$

These inequalities could certainly always be satisfied if all tolerances were considered as free parameters, for example by choosing large enough values of the δ_{α} , but what we are interested in is finding, for fixed $\{(\epsilon_{\alpha}, \delta_{\alpha})\}$, a range of values for the τ_i which satisfy the inequalities. We will show explicitly how to do this in the case $\tau_+ = \tau_- =: \tau$, which simplifies both the calculations and the visualization of the results.

For example, Eq (4.14) becomes now a simple quadratic relation between τ and τ_0 ,

$$\tau_0^2 + 4\,\bar{p}_0^2\tau_0 + (4\,\bar{p}_0^2\tau + 2\,\tau^2 - 8\,\delta_H^2) < 0 , \qquad (4.15)$$

which can also be written in a form showing explicitly that (τ, τ_0) must be inside an ellipse centered at $(-\bar{p}_0^2, -2\bar{p}_0^2)$. The full set of inequalities is

$$0 < \tau < 2\,\delta_p^2 \tag{4.16}$$

$$2\tau - 4\epsilon_H < \tau_0 < 2\tau + 4\epsilon_H \tag{4.17}$$

$$(\tau_0 + 2\,\bar{p}_0^2)^2 + 2\,(\tau + \bar{p}_0^2)^2 < 8\,\delta_H^2 + 6\,\bar{p}_0^4 \tag{4.18}$$

$$\left(\bar{\beta}_{0}^{2}\tau + \bar{\beta}_{\pm}^{2}\tau_{0}\right)\tau_{0}\tau + \left(1 - 2\,\delta_{v}^{2}\right)\tau_{0}\tau + \frac{1}{2}\left(\tau_{0}^{2} + \tau^{2}\right) + \left(\bar{p}_{\pm}^{2}\tau + \bar{p}_{0}^{2}\tau_{0}\right) < 0 , \qquad (4.19)$$

where the last inequality is equivalent to (4.13) for non-vanishing τ and τ_0 .

Fig. 1 illustrates roughly the range of allowed (τ, τ_0) values, for the choice of parameter values $\bar{p}_0 = 1$, $\bar{p}_{\pm} = 1/\sqrt{2}$, $\bar{\beta}_0 = 1$, $\bar{\beta}_{\pm} = 1/\sqrt{2}$, $\delta_p = 0.2$, $\delta_v = 10$, $\epsilon_H = 0.1$, $\delta_H = 0.6$. Strictly speaking, many values in the "allowed region" shown do not satisfy the conditions under which our small- τ/\bar{p}_0^2 approximation is valid, but the plot gives a useful qualitative description of the effect of each inequality in the τ_0 - τ plane.

The following argument tells us that the first three inequalities can in practice be replaced by one of them. Notice that, given that we are assuming \bar{p}_0 to be large, and the relationship between H and the p_i 's, it is reasonable to set for simplicity

$$\epsilon_H \approx \delta_H \approx \bar{p}_0 \, \delta_p \,, \qquad \text{so} \qquad \delta_p^2 \ll \epsilon_H, \qquad \delta_H \ll \bar{p}_0^2 \,.$$

$$(4.20)$$

Consider first the inequality (4.17). It defines a strip of allowed (τ, τ_0) values around the $\tau_0 = 2 \tau$ line, and this strip intersects the τ and τ_0 axes, respectively, at

$$\tau = 2 \epsilon_H , \qquad \tau_0 = 4 \epsilon_H . \tag{4.21}$$

With the choice of parameters we just made, $2 \epsilon_H > 2 \delta_p^2$, so the first of the two inequalities in (4.17) becomes redundant, provided that (4.16) is satisfied. However, if we now look at (4.18), it is easy to see that the ellipse intersects the axes at, respectively,

$$\tau = \sqrt{\bar{p}_0^4 + 4\,\delta_H^2} - \bar{p}_0^2 \approx 2\,\delta_p^2 \,, \qquad \tau_0 = \sqrt{4\,\bar{p}_0^4 + 8\,\delta_H^2} - 2\,\bar{p}_0^2 < 4\,\epsilon_H \,, \tag{4.22}$$

where we have used the fact that \bar{p}_0 is large in the inequalities. Thus, kinematical coherent states are semiclassical if (τ, τ_0) lie inside the portion of the ellipse (4.18) that defines a neighborhood of the origin in the first quadrant,

$$0 < \tau_0 < 2\,\bar{p}_0^2 \left(\sqrt{1 + \frac{2\,\delta_H^2}{\bar{p}_0^4}} - 1 \right), \qquad 0 < \tau < \bar{p}_0^2 \left(\sqrt{1 - \frac{\tau_0^2 + 4\,\bar{p}_0^2\tau_0 - 8\,\delta_H^2}{2\bar{p}_0^2}} - 1 \right), \quad (4.23)$$

and satisfy the single pair of additional inequalities (4.19).

We now need to find out whether τ and τ_0 can be chosen to meet these requirements. Clearly, not all points in the region (4.23) satisfy (4.19). This can be seen setting $\tau_0 = 0$, in which case we get $\frac{1}{2}\tau^2 + \bar{p}_{\pm}^2\tau < 0$, which has no positive solutions for τ ; in fact, if we consider (4.19) as a second-order inequality for τ for a given (positive) value of τ_0 ,

$$\left(\bar{\beta}_{0}^{2}\tau_{0} + \frac{1}{2}\right)\tau^{2} + \left[\bar{\beta}_{\pm}^{2}\tau_{0}^{2} + \left(1 - 2\,\delta_{v}^{2}\right)\tau_{0} + \bar{p}_{\pm}^{2}\right]\tau + \left(\frac{1}{2}\,\tau_{0}^{2} + \bar{p}_{0}^{2}\tau_{0}\right) < 0 , \qquad (4.24)$$

we see that there are positive solutions for τ only if the coefficient of the linear term is a sufficiently large negative number. We can also see directly from (4.13) that we must have $\delta_v^2 \gg 1$. For any positive τ_0 , as δ_v becomes very large, the range of values of τ which satisfy (4.24) becomes the full positive half-line $(0, \infty)$, so all we need to do is pick a large enough δ_v for all conditions to be satisfied in a region of the τ - τ_0 plane. The fact that δ_v must be large for suitable states to exist is an example of the uncertainty relations putting bounds on products of tolerances, and a consequence of the fact that we are requiring δ_p to be small.

We finish this section by giving a set of parameter values (different from the ones in Fig. 1) which satisfy the relationships mentioned above, and allow us to choose smaller values for τ and τ_0 , for use in later parts of the paper. If for simplicity we set $\bar{p}_+ = \bar{p}_-$ and $\bar{\beta}_+ = \bar{\beta}_-$ (the two inequalities (4.24) or their equivalents become a single one), it is natural to make the choices

$$\bar{p}_{\pm} = 1$$
, $\bar{p}_0 = \sqrt{2}$, $\bar{\beta}_{\pm} = 1$, $\bar{\beta}_0 = 1$ (4.25)

for the classical phase space point (notice that all inequalities are invariant under an appropriate rescaling of the variables involved; this can be used for example to assign any desired value to \bar{p}_0 , which we think of as an arbitrary 'global scale'), and

$$\delta_p = 0.15 , \quad \delta_v = 12 , \quad \epsilon_H = \delta_H = 0.25$$
(4.26)

for the relevant tolerances. Then, a pair of width parameters that solves the inequalities is

$$\tau = \tau_0 = 0.025 . \tag{4.27}$$

V. GROUP-AVERAGED COHERENT STATES

In the Dirac approach to the quantization of first-class constrained systems, a *physical state* Ψ_{phy} is one which is annihilated by the Hamiltonian constraint, $\hat{H} \cdot \Psi_{\text{phy}} = 0$. It is a general feature of constrained systems that physical states will *not* belong to the kinematical Hilbert space \mathcal{H}_{kin} , and the Refined Algebraic Quantization program has been developed to deal with such instances [16]. The general strategy is the following. One first identifies a dense subspace Φ of the Hilbert space \mathcal{H}_{kin} , where the Hamiltonian is well defined (that is, Φ is contained in the domain of \hat{H}). The physical states will then belong to the dual Φ^* , which satisfies the inclusion relation $\Phi \subset \mathcal{H}_{\text{kin}} \subset \Phi^*$. The choice of the dense subset Φ is a delicate matter, and normally has some physical implications [16], but for our purposes we are only interested in looking at coherent states which have some "nice" properties that make them well behaved, and those states belong to most choices of the space Φ . The procedure we will follow for implementing the Dirac requirement that physical quantum states be annihilated by the constraint operators is the so-called group averaging, a particularly popular approach within the general RAQ program.

The intuitive idea is to start with a state $\psi \in \Phi$ and average it over the group generated by the constraint. If all goes well, one will end up with an element of Φ^* satisfying the constraint. Our strategy will be very naive. Starting with the kinematical coherent state (4.2), group averaging gives us the physical coherent state $\psi_{\text{phy}} = N^{-1} \int d\lambda e^{-i\lambda \hat{H}} \psi_{\text{kin}}$, where N is a normalization factor. Since in p_i -space the constraint is a multiplication operator, the integration is straightforward and gives a delta function $\delta(p_0^2 - p_+^2 - p_-^2)$, so we can identify $\psi_{\text{phy}}(p_{\pm})$ with $\psi_{\text{kin}}((p_+^2 + p_-^2)^{1/2}, p_{\pm})$, up to a new normalization factor to be determined,¹

$$\psi_{\rm phy}(p_{\pm}) = \mathcal{N} e^{-[(p_{\pm}^2 + p_{\pm}^2)^{1/2} - \bar{p}_0]^2 / 2\tau_0 - (p_{\pm} - \bar{p}_{\pm})^2 / 2\tau_{\pm} - (p_{\pm} - \bar{p}_{\pm})^2 / 2\tau_{\pm}} \times e^{-i[(p_{\pm}^2 + p_{\pm}^2)^{1/2} - \bar{p}_0]\bar{\beta}_0 - i(p_{\pm} - \bar{p}_{\pm})\bar{\beta}_{\pm} - i(p_{\pm} - \bar{p}_{\pm})\bar{\beta}_{\pm}}} .$$
(5.1)

¹ The group average procedure, in a strict sense, will yield also a contribution from the past null cone, but since we are considering Gaussian wave functions that are almost zero there, the contribution from that null cone is negligible.

Of course, strictly speaking what one is doing is defining a function on the future null cone $(p_0^2 - p_+^2 - p_-^2 = 0, p_0 > 0)$, with coordinates p_{\pm} . The inner product induced by the group averaging procedure is then of the form

$$(\psi|\phi\rangle = \int \frac{\mathrm{d}p_{+}\,\mathrm{d}p_{-}}{2\,(p_{+}^{2}+p_{-}^{2})^{1/2}}\,\psi_{\mathrm{phy}}^{*}(p_{\pm})\,\phi_{\mathrm{phy}}(p_{\pm})\;.$$
(5.2)

Again, we want to calculate the expectation values and fluctuations of the physical observables \hat{p}_{\pm} and \hat{v}_i , and compare them to the values computed with the kinematical coherent states. The strategy here will be to represent the Dirac observables in the physical Hilbert space making use of the measure (5.2) induced by the group average and the general form of the operators given by (2.3). The momentum observables act as multiplication operators; to find the action of the configuration-like ones we use again (2.3), where now $\mu(p_{\pm}) = \frac{1}{2} (p_{+}^2 + p_{-}^2)^{-1/2}$ is the measure appearing in (5.2). This gives, on physical states,

$$\hat{v}_{\pm}\psi(p_{\pm}) = i\sqrt{p_{+}^{2} + p_{-}^{2}} \partial_{\pm}\psi(p_{\pm}) , \qquad \hat{v}_{0}\psi(p_{\pm}) = i(p_{+}\partial_{-} - p_{-}\partial_{+})\psi(p_{\pm}) .$$
 (5.3)

The first step, before calculating expectation values and fluctuations, is to calculate \mathcal{N} . However, because of the presence of the expression $(p_+^2 + p_-^2)^{1/2}$ in the exponent of ψ_{phy} , as well as in the measure, the relevant integrals cannot be calculated exactly, and we need to use an approximation.

If the state ψ_{phy} is narrowly peaked in the p_{\pm} directions, we can make use of the assumption $\tau \ll \bar{p}_0^2$ in the approximation, and keep only the first few terms in the expansion of every result in powers of τ/\bar{p}_0^2 . To see how to implement this idea, let us parametrize the deviation of p_{\pm} from the classical values \bar{p}_{\pm} by a new pair of variables,

$$x := \frac{p_+}{\bar{p}_0} \cos \bar{\theta} + \frac{p_-}{\bar{p}_0} \sin \bar{\theta} - 1 , \qquad y := -\frac{p_+}{\bar{p}_0} \sin \bar{\theta} + \frac{p_-}{\bar{p}_0} \cos \bar{\theta} , \qquad (5.4)$$

where θ is the angle between the vector (\bar{p}_+, \bar{p}_-) and the positive p_+ -axis, and the origin in the (x, y) coordinates corresponds to the point where the Gaussian is peaked. We can, equivalently solve

$$p_{+} = (x+1)\bar{p}_{+} - y\bar{p}_{-}, \qquad p_{-} = (x+1)\bar{p}_{-} + y\bar{p}_{+}.$$
(5.5)

Intuitively, x and y are the deviations from \bar{p}_{\pm} along the radial direction in the p_{\pm} plane and along the $p_{+}^{2} + p_{-}^{2} = \bar{p}_{0}^{2}$ circle, respectively, both expressed in units of \bar{p}_{0} . Using these definitions we get, e.g., that $p_{+}^{2} + p_{-}^{2} = \bar{p}_{0}^{2} [(1+x)^{2} + y^{2}]$, and in terms of the variables x and y the physical state can be written as

$$\psi_{\text{phy}}(x,y) = \mathcal{N} e^{-\{[(1+x)^2+y^2]^{1/2}-1\}^2 \bar{p}_0^2/2\tau_0 - (x^2+y^2)\bar{p}_0^2/2\tau}} \times e^{-i\{[(1+x)^2+y^2]^{1/2}-1\}\bar{p}_0\bar{\beta}_0 - i(\bar{p}_+x-\bar{p}_-y)\bar{\beta}_+ - i(\bar{p}_-x+\bar{p}_+y)\bar{\beta}_-};$$
(5.6)

in the remainder of this paper, we will assume for simplicity that $\tau_+ = \tau_- =: \tau$. Notice also that we will always assume that the parameters satisfy the classical constraint $\bar{p}_0^2 = \bar{p}_+^2 + \bar{p}_-^2$.

Our strategy for obtaining expressions for \mathcal{N} , as well as all expectation values and fluctuations, will be the following. All calculations can be reduced to integrals of the form

$$\int \frac{\mathrm{d}x\,\mathrm{d}y}{[(1+x)^2+y^2]^{1/2}}\,\psi(x,y)^*\,\hat{O}\,\psi(x,y)\,\,,\tag{5.7}$$

where \hat{O} may be a combination of multiplication and derivative operators. Since $\psi^* \hat{O} \psi$ always contains the Gaussian factor $\exp\{-(x^2 + y^2)\bar{p}_0^2/\tau\}$ which is peaked at small values of x and y, we will expand all remaining factors in the integrand, including the measure, in powers of x and y, and calculate the integral term by term in the expansion. Each non-vanishing term is a Gaussian integral of the form

$$\int_{-\infty}^{+\infty} \mathrm{d}x \, x^{2n} \,\mathrm{e}^{-x^2 \bar{p}_0^2/\tau} \int_{-\infty}^{+\infty} \mathrm{d}y \, y^{2m} \,\mathrm{e}^{-y^2 \bar{p}_0^2/\tau} = \frac{\pi \,(2n-1)!! \,(2m-1)!!}{2^{n+m}} \left(\frac{\tau}{\bar{p}_0^2}\right)^{n+m+1}, \qquad (5.8)$$

with a coefficient that depends on the parameters in ψ and, most importantly, contains one inverse power of τ for each derivative in \hat{O} ; if we want to keep terms up to a certain order in τ/\bar{p}_0^2 when adding the integrals of the type (5.8), we need to take this into account. With the exception of $(\Delta \hat{H})_{\rm kin}^2$ and $(\Delta \hat{v}_i)_{\rm kin}^2$, the expectation values and fluctuations of our operators in the kinematical states contain a single power of τ ; therefore, in order to compare those results with the corresponding ones for physical states, for the latter in practice we need to keep terms in our expansions up to the two leading orders in τ/\bar{p}_0^2 .

The normalization factor \mathcal{N} is the simplest expression to calculate. Since

$$\mathcal{N}^{-2} = \frac{\bar{p}_0}{2} \int \frac{\mathrm{d}x \,\mathrm{d}y}{[(1+x)^2 + y^2]^{1/2}} \,|\psi_{\rm phy}(x,y)|^2 = \frac{\pi\tau}{2\bar{p}_0} \left[1 - \left(\frac{\bar{p}_0^2}{\tau_0} - \frac{1}{2}\right) \frac{\tau}{2\bar{p}_0^2} + \mathcal{O}(\tau^2) \right], \quad (5.9)$$

the result is

$$\mathcal{N}^2 = \frac{2\bar{p}_0}{\pi\tau} \left[1 + \left(\frac{\bar{p}_0^2}{\tau_0} - \frac{1}{2}\right) \frac{\tau}{2\bar{p}_0^2} \right] + \mathcal{O}(\tau) .$$
 (5.10)

With \mathcal{N} , we can now calculate other expectation values. It is useful to first calculate the auxiliary quantities

$$\langle x \rangle = -\frac{\tau}{2\bar{p}_0^2} + \left(\frac{\bar{p}_0^2}{\tau_0} - 1\right) \frac{\tau^2}{4\bar{p}_0^4} + \mathcal{O}(\tau^3) , \qquad \langle y \rangle = 0 ,$$

$$\langle x^2 \rangle = \frac{\tau}{2\bar{p}_0^2} - \left(\frac{\bar{p}_0^2}{\tau_0} - 1\right) \frac{\tau^2}{2\bar{p}_0^4} + \mathcal{O}(\tau^3) , \qquad \langle y^2 \rangle = \frac{\tau}{2\bar{p}_0^2} - \frac{\tau^2}{4\bar{p}_0^4} + \mathcal{O}(\tau^3) , \qquad (5.11)$$

with which we find for the \hat{p}_{\pm} ,

$$\langle \hat{p}_{\pm} \rangle_{\rm phy} = \int \frac{\mathrm{d}p_{+}\mathrm{d}p_{-}}{2(p_{+}^{2}+p_{-}^{2})^{1/2}} p_{\pm} |\psi_{\rm phy}|^{2}$$

= $\bar{p}_{\pm} \left(\langle \hat{x} \rangle + 1 \right) = \bar{p}_{\pm} \left[1 - \frac{\tau}{2\bar{p}_{0}^{2}} + \left(\frac{\bar{p}_{0}^{2}}{\tau_{0}} - 1 \right) \frac{\tau^{2}}{4\bar{p}_{0}^{4}} + \mathcal{O}(\tau^{3}) \right],$ (5.12)

and for the fluctuations of the same operators,

$$\langle \hat{p}_{\pm}^{2} \rangle_{\rm phy} = \bar{p}_{\pm}^{2} \left(1 + 2\langle \hat{x} \rangle + \langle \hat{x}^{2} \rangle \right) + \bar{p}_{\mp}^{2} \langle \hat{y}^{2} \rangle$$

$$= \bar{p}_{\pm}^{2} + \frac{\bar{p}_{\mp}^{2} - \bar{p}_{\pm}^{2}}{2\bar{p}_{0}^{2}} \tau - \frac{\bar{p}_{\mp}^{2} \tau^{2}}{4\bar{p}_{0}^{4}} + \mathcal{O}(\tau^{3})$$

$$\left(\Delta \hat{p}_{\pm} \right)_{\rm phy}^{2} = \langle \hat{p}_{\pm}^{2} \rangle_{\rm phy} - \langle \hat{p}_{\pm} \rangle_{\rm phy}^{2} = \frac{\tau}{2} + \left[\frac{\bar{p}_{\pm}^{2}}{\bar{p}_{0}^{2}} - \left(\frac{1}{2} + \frac{\bar{p}_{\pm}^{2}}{\tau_{0}} \right) \right] \frac{\tau^{2}}{2\bar{p}_{0}^{2}} + \mathcal{O}(\tau^{3}) .$$

$$(5.13)$$

As for the Hamiltonian constraint, the exact physical state ψ_{phy} of course gives $\langle \hat{H} \rangle_{\text{phy}} = 0$ and $(\Delta \hat{H})^2_{\text{phy}} = 0$, by definition.

The \hat{v}_i operators contain derivatives with respect to p_{\pm} ; to calculate their expectation values it is useful to recast those in terms of (x, y), and we find, from (5.3) and (5.5),

$$\hat{v}_{\pm}\psi(x,y) = \mathrm{i}\sqrt{(1+x)^2 + y^2} \left(\frac{\bar{p}_{\pm}}{\bar{p}_0}\partial_x \mp \frac{\bar{p}_{\mp}}{\bar{p}_0}\partial_y\right)\psi$$
$$\hat{v}_0\psi(x,y) = \mathrm{i}\left[-y\,\partial_x + (1+x)\,\partial_y\right]\psi.$$
(5.14)

Then for the expectation values we find

$$\langle \hat{v}_{\pm} \rangle_{\rm phy} = \frac{i}{2} \int dx \, dy \, \psi_{\rm phy}^* \left(\frac{\bar{p}_{\pm}}{\bar{p}_0} \, \partial_x \mp \frac{\bar{p}_{\mp}}{\bar{p}_0} \partial_y \right) \psi = (\bar{p}_0 \bar{\beta}_{\pm} + \bar{p}_{\pm} \bar{\beta}_0) - (\bar{p}_0 \bar{\beta}_{\pm} + 2 \, \bar{p}_{\pm} \bar{\beta}_0) \frac{\tau}{4\bar{p}_0^2} + \mathcal{O}(\tau^2) , \langle \hat{v}_0 \rangle_{\rm phy} = \frac{i}{2} \int \frac{dx \, dy}{[(1+x)^2 + y^2]^{1/2}} \, \psi_{\rm phy}^* \Big(-y \, \partial_x + (1+x) \, \partial_y \Big) \psi_{\rm phy} = (\bar{p}_{\pm} \bar{\beta}_{-} - \bar{p}_{-} \bar{\beta}_{\pm}) \left(1 - \frac{\tau}{2\bar{p}_0^2} + \mathcal{O}(\tau^2) \right);$$
(5.15)

similarly, the fluctuations of these operators are, respectively,

$$(\Delta \hat{v}_{\pm})_{\rm phy}^2 = \frac{\bar{p}_0^2}{2\tau} + \frac{1}{4} \left(1 - \frac{\bar{p}_{\pm}^2}{\bar{p}_0^2} + \frac{2\bar{p}_{\pm}^2}{\tau_0} \right) + \mathcal{O}(\tau) ,$$

$$(\Delta \hat{v}_0)_{\rm phy}^2 = \frac{\bar{p}_0^2}{2\tau} - \frac{1}{4} + \mathcal{O}(\tau) .$$
(5.16)

We now have all the results we need to write down the conditions that the physical states be semiclassical, but we can also bypass this step by a direct comparison between the expectation values for the O_{α} in the two sets of states,

$$\langle \hat{p}_{\pm} \rangle_{\rm phy} - \bar{p}_{\pm} = -\frac{\tau}{2\bar{p}_0^2} + \mathcal{O}(\tau^2)$$
 (5.17)

$$\langle \hat{v}_{\pm} \rangle_{\rm phy} - \bar{v}_{\pm} = -(\bar{p}_0 \bar{\beta}_{\pm}^2 + 2 \, \bar{p}_{\pm} \bar{\beta}_0^2) \, \frac{\tau}{4\bar{p}_0^2} + \mathcal{O}(\tau^2) \,,$$
 (5.18)

and between the corresponding fluctuations,

$$(\Delta \hat{p}_{\pm})_{\rm phy}^2 - (\Delta \hat{p}_{\pm})_{\rm kin}^2 = -\left(\frac{\bar{p}_{\pm}^2}{\tau_0} + \frac{1}{2} - \frac{\bar{p}_{\pm}^2}{\bar{p}_0^2}\right)\frac{\tau^2}{\bar{p}_0^2} + \mathcal{O}(\tau^3)$$
(5.19)

$$(\Delta \hat{v}_{\pm})^{2}_{\rm phy} - (\Delta \hat{v}_{\pm})^{2}_{\rm kin} = -\frac{1}{4} \left(1 + \frac{\tau_{0}}{\tau} + \frac{\tau}{\tau_{0}} \right) - \frac{\bar{p}^{2}_{\pm}}{4 \bar{p}^{2}_{0}} - \bar{\beta}^{2}_{\pm} \tau_{0} + \mathcal{O}(\tau) .$$
(5.20)

The first pair of relationships tell us that the physical coherent states have expectation values for the O_{α} that are close (to order τ) to the classical values; that fact, together with the observation that the right-hand sides of the last two equations are negative (provided that $\tau_0 < 2 \bar{p}_0^2$), shows that the physical states are also semiclassical, with respect to the same tolerances used for the kinematical state and values for ϵ_p and ϵ_v that can be chosen to be small of order τ . In particular, if we choose as reference classical state the one specified by (4.25), and our criteria for a semiclassical state include the choices (4.26) plus, for example,

$$\epsilon_p = 0.01 , \quad \epsilon_v = 0.02 , \qquad (5.21)$$

then the state $\psi_{\rm phy}$ constructed using the width parameters given in (4.27) is semiclassical.

Thus, with this choice of widths, the kinematical state is compatible with the physical one in the less restrictive of the two possible definitions mentioned in Sec III. In the more restrictive sense, however, it is in general not a good approximation, because the right-hand side of (5.19) is not necessarily small compared to $(\Delta \hat{p}_{\pm})^2_{\rm phy}$, nor that of (5.20) compared to $(\Delta \hat{v}_{\pm})^2_{\rm phy}$. For the former to happen, the widths would have to satisfy $\tau \ll \tau_0$ (or \bar{p}_{\pm} would have to be very small, but this cannot be true for both signs), while for the latter to happen, they would have to satisfy (either $\tau \approx \tau_0$ or) $\tau \ll \tau_0 \ll \bar{p}_0^2$.

VI. TIME EVOLUTION

In the previous sections we have considered two kinds of coherent states, kinematical and "group averaged" ones. The first kind are coherent states defined on the full phase space and peaked around a point of the constraint surface. They do not satisfy the quantum constraint. The second kind of states are constructed out of the first kind by averaging over the orbit generated by the constraint. Even though the second kind of states are in a sense *dynamical* since one can think of the constraint (for totally constrained systems, such as this one) as generating time evolution, in a strict sense they are 'frozen'. Once the quantum state belongs to the physical Hilbert space, the notion of 'gauge' time evolution is lost. There is no notion of time evolution, no dynamics. Within this perspective, the observables that we have considered, namely Dirac observables, are constant along the orbits of the constraint and are therefore constants of the motion according to the same dynamical interpretation. This is the well known "frozen dynamics formalism" for time-reparametrization invariant constraint systems.

Is there a way of recovering dynamics from such a formalism? Again the answer is in the affirmative and several solutions have been known for a while [10, 17]. We shall consider one of those possibilities. For this we shall proceed in two steps. In the first part of the remainder of this section we recast the quantum constraint equation in such a way that it resembles a Schrödinger equation. In the second part we consider observables (different from the ones already considered) that capture the intuitive notion of time evolution and that will allow us to ask dynamically meaningful questions such as: given a coherent state with a small spread in the anisotropies at, say, the Plank time, we would like to know how much time later (as measured by a proper measure of the total volume) the state will spread. Based on the previous results obtained so far, where we found a range of parameters that produce acceptable coherent states, we shall try to give answers to these type of questions.

A. Schrödinger Equation

The idea here is to consider states at the kinematical level that are in the p representation for the \pm sector but are functions of the spatial volume and will therefore be diagonal on β_0 . That is, we will consider functions of the form $\Psi(\beta_0, p_{\pm})$. We want to consider the quantum constraint $\hat{H} = -\frac{1}{2} \left(\hat{p}_0^2 + \hat{p}_+^2 + \hat{p}_-^2 \right) = 0$, rewritten as

$$\hat{H} = -\frac{1}{2} \left(\hat{p}_0 + \sqrt{\hat{p}_+^2 + \hat{p}_-^2} \right) \left(\hat{p}_0 - \sqrt{\hat{p}_+^2 + \hat{p}_-^2} \right) \,, \tag{6.1}$$

and look for solutions to the quantum constraint that are of the form

$$(\hat{p}_0 - \sqrt{\hat{p}_+^2 + \hat{p}_-^2}) \cdot \Psi(\beta_0, p_\pm) = 0.$$
(6.2)

The choice of relative sign in (6.2) is equivalent to the choice we made earlier of the future (as opposed to past) light cone in *p*-space, in the context of group-averaged coherent states. In a sense we are following the original "already parametrized" program of ADM.

Note that this strategy is equivalent to considering the reduced phase space, described by true Dirac observables, and defining wave functions of them that have a "time dependence" as they are also functions of β_0 . Among these states, we can look for coherent states peaked around 'physical' phase space points. We start therefore with the reduced phase space Γ_r for which we will use the coordinates $\Gamma_r = \{(p_I, \beta_I), I = \pm\}$, and the Hilbert space $\mathcal{H}_r = L^2(\mathbb{R}^2, dp_+ dp_-)$. The sector of the solutions to the Hamiltonian constraint now takes the form of a Schrödinger equation,

$$i \frac{\partial}{\partial \beta_0} \Psi(\beta_0, p_{\pm}) = \sqrt{p_+^2 + p_-^2} \Psi(\beta_0, p_{\pm}) .$$
 (6.3)

That is, any (normalizable) function $\psi(p_{\pm})$ gives rise to a solution to the quantum constraint via the "time evolution equation" (6.3), whose solution can easily be written as

$$\Psi(\beta_0, p_{\pm}) = e^{-i\sqrt{p_{\pm}^2 + p_{-}^2} \beta_0} \psi(p_{\pm}) .$$
(6.4)

The strategy now is to start with a coherent state "intrinsic" to the p_{\pm} plane (as a coordinatization of the reduced configuration space) and then construct the corresponding physical state. This choice is motivated by the fact that the quantities β_{\pm} have a clear space-time interpretation as the anisotropies of the cosmological model. Let us then consider, as initial state,

$$\psi_0(p_{\pm}) = \prod_{i=\pm} \mathcal{N}_i \, \exp[-(p_i - \bar{p}_i)^2 / 2\tau_i - i \, (p_i - \bar{p}_i)\bar{\beta}_i] \,, \tag{6.5}$$

which is a coherent state peaked around the point $(\bar{p}_{\pm}, \bar{\beta}_{\pm})$ on the reduced phase space, and with spread dictated by the parameters τ_i . We get then

$$\Psi_{0}(\beta_{0}, p_{\pm}) = e^{-i\sqrt{p_{\pm}^{2} + p_{-}^{2}}\beta_{0}}\psi_{0}(p_{\pm})$$

=
$$\prod_{i=\pm} \mathcal{N}_{i} \exp\left[-(p_{i} - \bar{p}_{i})^{2}/2\tau_{i} - i(p_{i} - \bar{p}_{i})\bar{\beta}_{i}\right] e^{-i\sqrt{p_{\pm}^{2} + p_{-}^{2}}\beta_{0}}.$$
 (6.6)

Let us note that the general solution should be a function of $(\beta_0 - t_0)$, for t_0 the "initial value" of the parameter β_0 . Now, when $\beta_0 = 0$, the *classical* volume element is given by $\sqrt{\det q} = 1$ in Planck units. Thus, the "Planck epoch" is given by the values of $\beta_0 = 0$ (recall that the "singularity" is at $\beta_0 \to -\infty$). We shall then, in the remainder of this section, take $t_0 = 0$, and consider the "initial wave function" at $t_0 = 0$ as being defined at the Planck epoch.

B. Time Evolution of Observables

The next step is to consider observables. On the one hand we know from the general theory of constrained systems that physical observables are those operators that preserve the space of physical states. Properly represented Dirac observables have that property and that is why we have considered them in previous sections. On the other hand, we are interested in disentangling dynamical information from the particular (Schrödinger-like) representation of this part. This means that we need observables that capture the notion of time evolution. Such observables exist and are sometimes referred to as evolving constants of the motion [17]. The physical states considered in this part are solutions to the equation (6.3) so we have to define observables that preserve the space of such solutions. If we were to consider the operator $\hat{\beta}_0$ (that naturally acts as a multiplication operator), we would be thrown out of the physical Hilbert space. It is not a physical observable. What we need to do is to consider the following situation: let us assume that the phase space function β_0 is an (internal) time parameter, and that we can consider a new kind of observables 'defined at time' $\beta_0 = t$. That is, we fix a $\beta_0 = t$ slice in the configuration space (p_{\pm}, β_0) and consider the restriction of the wave function to that slice, $\Psi(\beta_0 = t, p_{\pm})$. We can then act with an operator (that might be ' β_0 -dependent') and consider the time evolution of the resulting wave function, thus obtaining again a physical state. This is nothing but the ordinary prescription for defining Heisenberg observables in ordinary quantum mechanics.

Consider an observable that depends explicitly on β_0 and does not commute with the constraint (for instance the volume form $V(\beta_0) = e^{3\beta_0}$). The strategy is to define a oneparameter family of observables $V(t) = V(\beta_0 = t)$ as Heisenberg observables and then consider the corresponding operators. Note that $\hat{V}(t_1) \neq \hat{V}(t_2)$ if $t_1 \neq t_2$, as physical operators on the physical Hilbert space. The observables that we will be interested in are of two kinds. On the one hand, we shall consider the 'configuration observables' \hat{p}_{\pm} that are constants of the motion since they commute with \hat{H} , and on the other hand those operators that measure the anisotropies, namely the operators $\hat{\beta}_{\pm}$. These are not explicitly 'time dependent', but we expect both expectation values and dispersions to be changing quantities.²

Let us then compute the expectation values and quantum fluctuations for the operators $\hat{\beta}_{\pm} = i \partial/\partial p_{\pm}$. The expectation value will take the form

$$\langle \hat{\beta}_{+} \rangle_{t} = \mathcal{N}^{2} \left[\int \mathrm{d}^{2} p_{\pm} \, \mathrm{e}^{-(p_{+}-\bar{p}_{+})^{2}/\tau} \, \frac{p_{+}}{\sqrt{p_{+}^{2}+p_{-}^{2}}} \right] t + \bar{\beta}_{+} \, .$$
 (6.7)

Note that there will be a shift of the 'peak' of the wave function, since the expectation value will have a linear t dependence. This is to be expected since classically we know that the system evolves in a way that the β_{\pm} depend linearly on time. In order to compute the coefficient we need to make some approximations as in previous sections. Making use of the same assumptions, we arrive for the integral to the following expression,

$$\langle \hat{\beta}_{+} \rangle_{t} = \bar{\beta}_{+} + t \,\mathcal{N}^{2} \bar{p}_{0}^{2} \int \mathrm{d}x \,\mathrm{d}y \,\mathrm{e}^{-\bar{p}_{0}^{2}(x^{2}+y^{2})/\tau} \,\frac{(1+x)\bar{p}_{+} - y\bar{p}_{-}}{\bar{p}_{0} \left[(1+x)^{2} + y^{2}\right]^{1/2}} \,. \tag{6.8}$$

² Let us qualify the statement: We expect that the one-parameter family of numbers $\langle \hat{\beta}_{\pm} \rangle_t$, corresponding to the one-parameter family of operators $\hat{\beta}_{\pm,t}$, will depend on the value of t.

This can be approximated, as a power series expansion in τ , by the expression

$$\langle \hat{\beta}_{\pm} \rangle_t = \bar{\beta}_{\pm} + \frac{\bar{p}_{\pm}}{\bar{p}_0} \left[1 - \frac{1}{4} \frac{\tau}{\bar{p}_0^2} - \frac{3}{8} \frac{\tau^2}{\bar{p}_0^4} + \mathcal{O}(\tau^3) \right] t .$$
 (6.9)

Note that at zeroth order in τ the speed of the peak coincides precisely with the classical value. Let us now compute the fluctuations of the operators and consider the expectation value $\langle \hat{\beta}_{+}^{2} \rangle$, to second order in τ as,

$$\langle \hat{\beta}_{+}^{2} \rangle_{t} = \frac{1}{2\tau} + \bar{\beta}_{+}^{2} + 2\bar{\beta}_{+} \frac{\bar{p}_{+}}{\bar{p}_{0}} \left[1 - \frac{1}{4} \frac{\tau}{\bar{p}_{0}^{2}} - \frac{3}{8} \frac{\tau^{2}}{\bar{p}_{0}^{4}} \right] t + \frac{\bar{p}_{+}^{2}}{\bar{p}_{0}^{2}} \left[1 - \frac{1}{2} \frac{\tau}{\bar{p}_{0}^{2}} + \frac{33}{4} \frac{\tau^{2}}{\bar{p}_{0}^{4}} \right] t^{2} .$$
 (6.10)

From here we can then compute the fluctuations of the anisotropy observables. We have then,

$$(\Delta\beta_{\pm})_t^2 = \langle \hat{\beta}_{\pm}^2 \rangle_t - \langle \hat{\beta}_{\pm} \rangle_t^2 = \frac{1}{2\tau} + \frac{143}{16} \frac{\bar{p}_{\pm}^2}{\bar{p}_0^6} \tau^2 t^2 , \qquad (6.11)$$

where we keep terms that are at most quadratic in τ . We see that, as in the case of a non-relativistic free particle, the spread of the Gaussian grows quadratically with 'time' t. The next step is to estimate the value t_{max} of the maximum allowed value for t before the wave packet spreads considerably. For that we use the values of the parameters \bar{p}_{\pm} , \bar{p}_0 and τ chosen in (4.25) and (4.27). The reason for this is that we expect that the conditions imposed on the expectation values and fluctuations of the relevant operators yield realistic values for the parameters defining the coherent states; even though in this part we start with intrinsic coherent states at 'Planck time', the resulting wave function is closely related to the physical coherent states of Sec. V, and it seems reasonable to employ the parameters used there. If we define the time for the wave packet to spread significantly as the value for t_{max} such that $(\Delta \beta_{\pm})_{t_{\text{max}}}^2$ is about twice as large as $(\Delta \beta_{\pm})_{t_0}^2 \approx 20$, with the values of the parameters given in Sec IV it is easy to see that $t_{\text{max}} \approx 170$. That is, the time for which the packet spreads to about twice its size is within 200 times the fundamental time unit, the Planck time. With a different choice of parameters that yield smaller values of τ , one might have slightly longer spread times, but within the same order of magnitude. Another possibility would be to take the values used in Fig. 1 and choose $\tau = .03$, which is near the high end of the range of allowed values. In this case the time $t_{\rm max} \approx 65$, again of the same order of magnitude as with the previous choice.

One might ask about the possibility of having semiclassical states that approximate the β 's more accurately, which means smaller initial fluctuations (of the order of $1/\tau$). This would require large values of τ , for which the expansion in powers of τ would not be justified. In that case, one would need to expand expectation values and fluctuations of observables in inverse powers of τ , and use those in the set of inequalities that express the semi-classicality conditions. With this choice, one might be able to construct states that take longer to spread. We shall not explore this possibility further. Let us end this section with a remark. Even though we have found a 'time' (as measured in terms of the volume element) for which the initial wave-packet spreads, we would not like to suggest that this has a direct significance for the spread of the wave-function in realistic cosmological models, since we have neglected any matter and the effect that this may have in possible decoherence effects on the wave function.

VII. CONCLUSION AND OUTLOOK

In this paper we have only considered the simplest type of anisotropic cosmological models, the vacuum ones of Bianchi type I. For our purposes, the simplicity of these models lies in the fact that their phase space is a vector space, their only constraint is quadratic, and it depends on the p variables only; in this sense, these models are similar to the general class of models treated in Ref [7], in which kinematical and physical coherent states are also discussed. However, on the one hand our Hamiltonian and the set of physical observables we chose are not of the type used there, and on the other hand the fact that the Hamiltonian itself is the constraint raises the usual issue of time evolution in generally covariant systems, and led us to the construction of a third type of state that we called "intrinsic", with which physical predictions on the evolution of physical observables can be attempted. The relevance of our findings in this simple model for the ultimate goal of constructing realistic semi-classical states in full quantum gravity still needs to be explored. However, we hope that these first steps might shed some light on the program.

To explore further the relationships between kinematical and physical coherent states as candidate semiclassical states for gravity models, a natural extension of this work would be to consider other Bianchi type models. As soon as one starts doing that, one is faced with the fact that the potential $U(\beta^i)$ does not vanish, which implies that the quantum Hamiltonian is no longer a multiplication operator. Fortunately, in some cases this difficulty can be circumvented. As is well known, the dynamics of a system on a configuration space with a given metric subject to a potential U, can be mapped into that of a system with a new metric (the Jacobi metric) and vanishing potential. In our case, the transformation of a Hamiltonian like (2.2) into an equivalent one describing a free particle follows the standard procedure, but what one usually ends up with is a curved Jacobi metric replacing η_{ij} . What is not so obvious is the fact that in the case of many homogeneous cosmologies, known as diagonal, intrinsically multiply transitive (DIMT) models, the new metric is still flat [10]. These models include the following cases: Bianchi types I and II; sub-families of Bianchi types III, VIII and IX defined by $\beta^- = 0$ (the Taub model); Kantowski-Sachs spacetimes; and Bianchi type V models with $\beta^+ = 0$. For Bianchi type I and II, the minisuperspaces are 3-dimensional, parametrized by β^0 , β^+ and β^- in the Misner scheme; for the remaining DIMT models they are 2-dimensional, parametrized by an appropriate subset of the β^i [11]. Note that even though the type III and V models are class B, which means that in general the Hamiltonian description is more subtle [12], these particular cases do not suffer from the problems described in Refs [1, 12].

Thus, DIMT models are natural candidates for a first extension of our work. When one carries out the transformation to a new set of phase space variables $(\tilde{\beta}^i, \tilde{p}_i)$ in which the potential term in (2.2) vanishes and the metric again has the form η_{ij} , the difference between the various DIMT models shows up in the fact that the allowed ranges of the \tilde{p}_i are not the whole real line [10]. Therefore, those variables cannot be regarded as momenta in the regular sense of elements of the cotangent space, and in the quantum theory the corresponding operators in the $\tilde{\beta}_i$ -representation cannot be the usual derivative operators. (Note: this feature, which also occurs in full quantum gravity, has been addressed in that context by Klauder [18] and Loll [19].) We have then one more reason to set up the quantum theory in the \tilde{p} -representation [10]. This time, however, the "configuration space" for these models is a proper subset of 2+1 Minkowski space, with restrictions on the values of the \tilde{p}_i depending on the model under consideration, and the constraint surface is the intersection of the (future) light cone of the origin with this subset.

With this setup, one can define suitable generalizations of coherent states in the configuration spaces of the various DIMT models, which now are non-trivial spaces (for example, a half-line rather than the whole real line), and analyze their properties as candidate semiclassical states. One possible strategy is to construct such states as suggested by Klauder's work [20], and search for a proper subset of states that are consistent with our physical requirements, but we shall leave that investigation for a future publication. As a further step, one can envision extending the work to models which cannot be formulated as free particles on portions of Minkowski space, which will require new techniques to handle the quantum constraint (it would be interesting, for example, to study the connection with Kiefer's use of the "principle of constructive interference" to build wave packets in minisuperspace [21] and other models for the emergence of classical Friedmann-Robertson-Walker spacetimes [22]), and then possibly some non-homogeneous, midisuperspace models. In terms of our original goal of understanding the relationship between the kinematical and physical semiclassical states for quantum gravity, the hope is that at some point one may see a pattern that will allow us to make statements of a more general nature.

Acknowledgements

We are grateful to Abhay Ashtekar for many discussions, which provided the motivation for this work, and for suggestions. We also thank Michael Ryan for comments and David Sanders for help with the figure. Partial support for this work was provided by NSF grant PHY-0010061, CONACyT grant J32754-E and DGAPA-UNAM grant 112401.

- Fels M E and Torre C G 2002 "The principle of symmetric criticality in general relativity," Class. Quantum Grav. 19 641, arXiv:gr-qc/0108033
- [2] Kuchař K V and Ryan M P 1989 "Is minisuperspace quantization valid? Taub in Mixmaster," Phys. Rev. D 40 3982; Kuchař K V and Ryan M P 1986 "Can minisuperspace quantization be justified?, in *Gravitational Collapse and Relativity*, edited by H. Sato and Nakamura (World Scientific, Singapore)
- Butterfield J and Isham C J 1999 "On the emergence of time in quantum gravity," arXiv:gr-qc/9901024; Kuchař K V 1992 "Time and interpretations of quantum gravity," in G Kunstatter et al General relativity and Relativistic Astrophysics (World Scientific 1992) 211-314
- [4] Hawking S W 1979 "The path integral approach to quantum gravity," in S W Hawking and W Israel General Relativity: An Einstein Centenary Survey (Cambridge University Press) 746-789; Hartle J B and Hawking S W 1983 "Wave function of the universe," Phys. Rev. D 28 2960; Marolf D 1996 "Path integrals and instantons in quantum gravity," Phys. Rev. D 53 6979, arXiv:gr-qc/9602019
- [5] For a recent account, see Halliwell J J 2002 "The interpretation of quantum cosmology and the problem of time" arXiv:gr-qc/0208018
- [6] Marolf D 1995 "Quantum observables and recollapsing dynamics," Class. Quantum Grav. 12 1199, arXiv:gr-qc/9404053; "Almost ideal clocks in quantum cosmology: A brief derivation of time," Class. Quantum Grav. 12 2469, arXiv:gr-qc/9412016

- [7] Ashtekar A, Bombelli L and Corichi A 2004 "Coherent and semiclassical states for constrained systems", preprint
- [8] Kiefer C 1994 "The Semiclassical approximation to quantum gravity" Canonical Gravity: From Classical to Quantum J Ehlers and H Friedrich eds (Springer, Berlin), arXiv:gr-qc/9312015
- [9] Misner C W 1972 "Minisuperspace" in J R Klauder, ed Magic Without Magic: J A Wheeler (Freeman, San Francisco) 441–473
- [10] Ashtekar A, Tate R and Uggla C 1993 "Minisuperspaces: Observables and quantization" Int. J. Mod. Phys. D2 15-50, arXiv:gr-qc/9302017
- [11] Ryan M P 1972 Hamiltonian Cosmology (Springer-Verlag)
- [12] Ryan M P and Waller S M 1997 "On the Hamiltonian formulation of class B Bianchi cosmological models" arXiv:gr-qc/9709012
- [13] Ashworth M C 1998 "Coherent state approach to time-reparametrization invariant systems" Phys. Rev. A 57 2357, arXiv:quant-ph/9611026
- [14] Date G and Singh P 2001 "Semi-classical states in the context of constrained systems", arXiv:quant-ph/0109127
- [15] Schiff L I 1968 Quantum Mechanics 3rd edition (McGraw-Hill)
- [16] Marolf D 1995 "Refined algebraic quantization: Systems with a single constraint" arXiv:gr-qc/9508015; "Group averaging and refined algebraic quantization: Where are we now?" arXiv:gr-qc/0011112; Giulini D and Marolf D 1999 "On the generality of refined algebraic quantization" Class. Quantum Grav. 16 2479, arXiv:gr-qc/9812024
- [17] Rovelli C 1991 "What is observable in classical and quantum gravity?" Class. Quantum Grav. 8 297; Rovelli C 1991 "Time in quantum gravity: a hypothesis" Phys. Rev. D 43 442
- [18] Klauder J 2003 "Affine quantum gravity" Int. J. Mod. Phys. D12 1769–1774, arXiv:gr-qc/0305067
- [19] Loll R 1997 "Imposing det E > 0 in discrete quantum gravity," Phys. Lett. **B399** 227, arXiv:gr-qc/9703033
- [20] Klauder K 1986 "Global, uniform semiclassical approximations for quantum systems on the half-line" Phys. Rev. A 34 4486–4489; Watson G and Klauder J R 2000 "Generalized affine coherent states: A natural framework for the quantization of metric-like variables" J. Math. Phys. 41 8072–8082, arXiv:quant-ph/0001026
- [21] Kiefer C 1988 "Wave packets in minisuperspace" Phys. Rev. D 38 1761–1772
- [22] Kim S P, Ji J Y, Shin H S and Soh K S 1997 "Coherence and emergence of classical spacetime" Phys. Rev. D 56 3756–3758, arXiv:gr-qc/9703064