arXiv:gr-qc/0404015v2 23 Jul 2004

Small Black Holes on Branes: Is the horizon regular or singular?

D. Karasik,* C. Sahabandu, P. Suranyi, and L. C. R. Wijewardhana

Department of Physics, University of Cincinnati, Cincinnati, Ohio

We investigate the following question: Consider a small mass, with ϵ (the ratio of the Schwarzschild radius and the bulk curvature length) much smaller than 1, that is confined to the TeV brane in the Randall-Sundrum I scenario. Does it form a black hole with a regular horizon, or a naked singularity? The metric is expanded in ϵ and the asymptotic form of the metric is given by the weak field approximation (linear in the mass). In first order of ϵ we show that the iteration of the weak field solution, which includes only integer powers of the mass, leads to a solution that has a singular horizon. We find a solution with a regular horizon but its asymptotic expansion in the mass also contains half integer powers.

I. INTRODUCTION

Black holes in theories with extra dimensions have been studied widely. Myers and Perry [1] found Schwarzschild type solutions (MPS) in *D*-dimensional asymptotically flat space. Black hole solutions were also found in asymptotically AdS space [2, 3]. No non-trivial closed form black hole solutions, other than the black string solution [4], which extends in a uniform manner from the brane into the extra dimension, have been found in three-brane theories of the Randall-Sundrum type [5, 6]. Given that there is considerable interest surrounding the production of black holes at accelerators [7], and in collisions of cosmic rays [8], it is important to develop approximate methods to find black hole solutions in Randall-Sundrum brane world theories.

Some initial attempts at finding black hole solutions centered on deriving the induced metric on the brane by solving the Hamiltonian constraint conditions [9]. Some of the induced solutions do not arise from matter distributions confined to the brane [10]. Linearized solutions about RS backgrounds [11] as well as numerical solutions [12, 13] have also been derived.

In a recent paper [14] we have studied the metric around a small mass that is confined to the TeV brane in Randall-Sundrum type I scenario (RSI)[5]. The configuration is characterized by two length scales; the Schwarzschild radius μ , which is related to the mass $\mu = \sqrt{8G_5M/(3\pi)}$, and the curvature length of the bulk ℓ , which is related to the cosmological constant and the brane tension. We study the case $\mu \ll \ell$ where the metric can be expanded in the dimensionless parameter, $\epsilon = \mu/\ell \ll 1$. There are two ways to make $\epsilon \ll 1$: (1) $\mu \to 0$ while ℓ is finite. This limit is just the original Randall-Sundrum scenario in the absence of matter. Keeping ℓ finite and expanding in μ (or actually in M) is called the weak field approximation. First order in M is linearized gravity. Linearized gravity is valid at distances much larger than the Schwarzschild radius, $r \gg \mu$. (2) $\ell \to \infty$ while μ is finite. This limit is a five dimensional black hole in an asymptotically flat background. The metric for this configuration is the Myers and Perry solution (MPS). We studied the expansion in ϵ with μ kept finite. We call the first order expansion in ϵ - the ϵ solution. The ϵ solution is valid at distances much smaller than the bulk curvature length $r \ll \ell$.

On one hand, the ϵ solution is needed to study the horizon and the thermodynamics of the black hole, for which the linearized solution is unsuitable. On the other hand, the ϵ solution cannot be fixed uniquely without satisfying certain boundary conditions. We need the linearized solution to identify the mass and to satisfy the junction conditions on the Planck brane because the distance between the branes in RSI is of the same order as the bulk curvature length $d \sim \ell$.

Since we need both the ϵ solution and the linearized solution we look at the region where both solutions are valid, $\mu \ll r \ll \ell$. In this region we expand the linearized solution to first order in ℓ^{-1} and the ϵ solution to first order in M. We require that these two expansions will coincide. Using this method, we are able to incorporate the asymptotic characteristics into the short-ranged ϵ solution. In other words, the linearized solution is the asymptotic boundary condition for the ϵ solution.

In this paper we study the regularity of the horizon. In section II we summarize the results from [14]. In section III we analyze the horizon and formulate the conditions such that the ϵ solution will be regular on the horizon. We also calculate the thermodynamics parameters of the black hole assuming it is regular. In section IV we represent the ϵ solution with Legendre functions. We assume that in the asymptotic region the metric is given by the linearized

^{*}Electronic address: karasik@physics.uc.edu

solution (linear in the mass). Another reasonable assumption is that when iterating the linearized solution, the post linearized solution includes only integer powers of the mass. This assumption is motivated by the post Newtonian behavior of 4 dimensional black holes, where the asymptotic solution is an expansion in integer powers of G_4M/r . We show that under this assumption the horizon is singular. In section V we show that one can construct a regular solution provided that the post linearized solution includes half integer powers of the mass. We find it interesting that the first law of black hole thermodynamics, dM = T dS, eliminates the terms of order $M^{3/2}$ from the asymptotic expansion.

II. THE ϵ SOLUTION

We use the following ansatz for the metric

$$ds^{2} = a(\rho,\psi) \left[-B(\rho,\psi)dt^{2} + \frac{A(\rho,\psi)}{B(\rho,\psi)}d\rho^{2} + 2V(\rho,\psi)d\rho\,d\psi + \rho^{2}U(\rho,\psi)d\psi^{2} + \rho^{2}\sin^{2}\psi\,d\Omega_{2}^{2} \right] , \tag{1}$$

where $d\Omega_2^2$ is the metric on the unit 2-sphere. The conformal factor is taken from RSI as $a(\rho, \psi) = (1 - \epsilon \rho | \cos \psi |)^{-2}$. The TeV brane is located at $\psi = \pi/2$ as can be read off of the conformal factor. We work only in the interval $0 \le \psi \le \pi/2$, and assume Z_2 symmetry about the brane. The point mass is located at $\rho = 0$. The ϵ solution is just MPS solution with ϵ corrections

$$B(\rho, \psi) = 1 - \frac{1}{\rho^2} - \epsilon B_1(\rho, \psi) ,$$
 (2a)

$$A(\rho, \psi) = 1 + \epsilon A_1(\rho, \psi) , \qquad (2b)$$

$$U(\rho,\psi) = 1 + \epsilon U_1(\rho,\psi) , \qquad (2c)$$

$$V(\rho, \psi) = \epsilon V_1(\rho, \psi) . \tag{2d}$$

We have rescaled the coordinates with μ such that $\rho = 1$ corresponds to the MPS horizon at the Schwarzschild radius. The gauge is partially fixed by the coefficient of $d\Omega_2^2$. There is still a gauge freedom associated with the coordinate transformation $\rho \to \rho(1 - \epsilon F)$, $\psi \to \psi + \epsilon F \tan \psi$.

The ϵ solution can be written in terms of the gauge function $F(\rho, \psi)$ and the wave function $H(\rho, \psi)$. The solution is [20]

$$B_1 = -\frac{2}{\rho^2} F(\rho, \psi) ,$$
 (3a)

$$A_{1} = 8\rho(\rho^{2} - 1)\cos\psi - 12H(\rho,\psi) - 4\tan\psi H_{,\psi}(\rho,\psi) + 2F(\rho,\psi) + 2\rho F_{,\rho}(\rho,\psi) , \qquad (3b)$$

$$U_{1} = \frac{2\tan^{2}\psi}{\rho} \left[6\rho H(\rho,\psi) - (2\rho^{2} - 1)H_{,\rho}(\rho,\psi) - \rho F(\rho,\psi) - \rho \cot\psi F_{,\psi}(\rho,\psi) \right] , \qquad (3c)$$

$$V_{1} = -2\rho^{2}(2\rho^{2} - 1)\sin\psi + 2\rho^{2}\tan\psi H_{,\rho}(\rho,\psi) + \frac{\rho^{2}F_{,\psi}(\rho,\psi)}{\rho^{2} - 1} - \rho^{2}\tan\psi F_{,\rho}(\rho,\psi) + \frac{\rho^{3}F_{,\psi}(\rho,\psi)}{\rho^{2} - 1}.$$
(3d)

The wave function $H(\rho, \psi)$ satisfies the differential equation

$$(\rho^2 - 1)\left(H_{,\rho\rho} - \frac{1}{\rho}H_{,\rho}\right) + H_{,\psi\psi} + 2\frac{\cos^2\psi + 1}{\sin\psi\cos\psi}H_{,\psi} = 0.$$
(4)

Junction conditions on the brane. Israel junction conditions [15] on the brane are simply $B_{1,\psi}(\rho, \pi/2) = 0$, $A_{1,\psi}(\rho, \pi/2) = 0$, $V_1(\rho, \pi/2) = 0$, and $U_1(\rho, \pi/2) < \infty$. These conditions imply the following

$$F_{,\psi}(\rho,\pi/2) = 0$$
, (5a)

$$(3H + \tan \psi H_{,\psi})_{,\psi}\Big|_{\psi=\pi/2} = -2\rho(\rho^2 - 1) , \qquad (5b)$$

$$H_{,\psi}(\rho,\pi/2) = 0$$
, (5c)

$$F(\rho, \pi/2) = 6H(\rho, \pi/2) - \frac{2\rho^2 - 1}{\rho}H_{,\rho}(\rho, \pi/2) .$$
(5d)

$$H(\rho,\psi) = \delta_0 + \frac{\rho^3}{16\sin^3\psi} (4\psi - \sin 4\psi) + \frac{\rho}{64\sin^3\psi} (4\sin 2\psi - 8\psi\cos 2\psi - 12\psi + 3\sin 4\psi) + \frac{1}{1024\rho\sin^3\psi} (1024\delta + 8\psi + 32\psi\cos 2\psi + 12\psi\cos 4\psi - 16\sin 2\psi - 5\sin 4\psi) + \mathcal{O}(\rho^{-(1+2\alpha)}), \quad (6)$$

where δ and δ_0 are arbitrary constants. The asymptotic form of the gauge function is

$$F(\rho,\psi) = \frac{\rho}{2} \left(\cos\psi + \frac{\psi}{\sin\psi} \right) + 6\delta_0 + \frac{F_2(\psi)}{\rho} + \mathcal{O}(\rho^{-(1+2\alpha)}) , \qquad (7)$$

where $F_2(\pi/2) = 8\delta - 5\pi/64$ and $F_2''(\pi/2) = 47\pi/64 - 24\delta$.

In the functions (6) and (7) we have neglected corrections of order $\rho^{-(1+2\alpha)}$ where $\alpha > 0$. The reason is that such terms will contribute to order $M^{1+\alpha}$, which is beyond the linearized solution.

A reasonable assumption is that if one will go beyond the linearized solution, only integer powers of M will appear in the metric. This assumption will force the functions $H(\rho, \psi)$ and $F(\rho, \psi)$ to be odd functions of ρ , therefore one must set $\delta_0 = 0$ and the neglected terms are of order ρ^{-3} .

III. ANALYZING THE HORIZON

The configuration is called a black hole if there exists a Killing horizon, i.e. a null surface with a Killing field normal to the surface. The ansatz (1) is static, and the static Killing vector becomes null on the surface B = 0. The static Killing vector is $\xi^{\mu} = \delta_0^{\mu}$, it is normalized such that asymptotically $\xi^2 = -1$.

A vector normal to the surface B = 0 must be proportional to $n_{\mu} = B_{,\mu}$. If the surface B = 0 is null, then the normal vector must be null on B = 0.

$$n^{\mu}n_{\mu}|_{B=0} = \frac{(B_{,\psi})^2}{\rho^2 a U} = 0.$$
(8)

Therefore, the derivative $B_{,\psi}$ must vanish on the surface B = 0. This implies that the surface $B(\rho, \psi) = 0$ is actually defined as $\rho = \rho_H = \text{constant}$. The ansatz for $B(\rho, \psi)$ is

$$B(\rho,\psi) = \left(1 - \frac{\rho_H^2}{\rho^2}\right) b(\rho,\psi) , \qquad (9)$$

where $b(\rho, \psi)$ does not vanish at $\rho = \rho_H$, but it is regular such that $\lim_{\rho \to \rho_H} (\rho - \rho_H) b(\rho, \psi) = 0$.

The surface gravity, κ , for the ansatz (1) is defined as

$$\kappa^{2} = -\frac{1}{2}g^{\mu\nu}g^{\lambda\sigma}\xi_{\mu;\lambda}\xi_{\nu;\sigma}\Big|_{B=0} = \frac{(B_{,\rho})^{2}}{4A} \ . \tag{10}$$

According to a theorem by Rácz and Wald [16], if the surface gravity is not constant on the horizon, the horizon is singular. This can be verified in the ansatz (1) by calculating the Kretchmann scalar around the horizon. Provided that the function $B(\rho, \psi)$ is given by Eq.(9), the Kretchmann scalar can be expanded in powers of B

$$R^{\mu\nu\lambda\sigma}R_{\mu\nu\lambda\sigma} = \frac{1}{B(\rho,\psi)} \frac{32(\kappa,\psi)^2}{\rho^2 a^2 U} + \text{regular terms} .$$
(11)

Clearly, at any point where $\kappa_{,\psi} \neq 0$, the surface B = 0 is singular. If the surface gravity is not constant but vanishing at some points on the horizon, these points will be regular. An ansatz for the function $A(\rho, \psi)$, which is compatible with Eqs.(9, 10), is

$$A(\rho,\psi) = \frac{b(\rho_H,\psi)^2}{\rho_H^2 \kappa^2} + \left(1 - \frac{\rho_H^2}{\rho^2}\right) \alpha(\rho,\psi) , \qquad (12)$$

where $\lim_{\rho \to \rho_H} (\rho - \rho_H) \alpha(\rho, \psi) = 0.$

 ϵ expansion on the horizon. Take the ansatz (1) together with Eqs.(9, 12) and expand in ϵ . Zero order in ϵ should be MPS solution, therefore, the following ϵ expansions should be used

$$b(\rho,\psi) = 1 + \epsilon b_1(\rho,\psi) , \qquad (13a)$$

$$\alpha(\rho,\psi) = \epsilon a_1(\rho,\psi) , \qquad (13b)$$

$$\rho_H = 1 + \epsilon \zeta , \qquad (13c)$$

$$\kappa = 1 + \epsilon \chi . \tag{13d}$$

Comparing Eqs.(13) with Eqs.(2), one can solve for b_1 and a_1

$$b_1(\rho,\psi) = \frac{2\zeta - \rho^2 B_1(\rho,\psi)}{\rho^2 - 1} , \qquad (14)$$

$$a_1(\rho,\psi) = \frac{\rho^2 \left[A_1(\rho,\psi) - 2b_1(1,\psi) + 2\zeta + 2\chi\right]}{\rho^2 - 1} .$$
(15)

The expansion in ϵ on the horizon is possible only if the functions b_1 and a_1 are finite at $\rho = 1$. Therefore, the numerators in Eqs.(14, 15) must vanish at $\rho = 1$, and one can calculate ζ , $b_1(1, \psi)$, and χ

$$\zeta = \lim_{\rho \to 1} \frac{1}{2} B_1(\rho, \psi) , \qquad (16)$$

$$b_1(1,\psi) = -\lim_{\rho \to 1} \frac{\left(\rho^2 B_1(\rho,\psi)\right)_{,\rho}}{2\rho} , \qquad (17)$$

$$\chi = \lim_{\rho \to 1} \frac{1}{2} A_1(\rho, \psi) - b_1(1, \psi) - \zeta .$$
(18)

Both ζ and χ are constants, therefore Eqs.(16, 18) put restrictions on the solution (3). Equation (16) implies that

$$\zeta = -F(1,\psi) = \text{constant} \quad \Rightarrow \quad F_{,\psi}(1,\psi) = 0 \ . \tag{19}$$

Substitute the solution (3) in Eq.(18) to calculate the correction to the surface gravity

$$\chi = 2\left(3H(1,\psi) + \tan\psi H_{,\psi}(1,\psi)\right) = \text{constant} .$$
⁽²⁰⁾

So, the combination $3H(1,\psi) + \tan \psi H_{\psi}(1,\psi)$ must be constant.

Another constraint comes from Eq.(3d). The function V_1 must be finite at $\rho = 1$ (just like b_1 and a_1). This implies that

$$\lim_{\rho \to 1} (\rho^2 - 1) V_1(\rho, \psi) = \tan^2 \psi H_{,\psi}(1, \psi) + F_{,\psi}(1, \psi) = 0 .$$
⁽²¹⁾

Since $F_{,\psi}(1,\psi) = 0$ we deduce that $H_{,\psi}(1,\psi) = 0$. The functions H and F are constants on the horizon and using Eq.(5d) can be evaluated as

$$H(1,\psi) = H(1,\pi/2) , F(1,\psi) = F(1,\pi/2) = 6H(1,\pi/2) - H_{,\rho}(1,\pi/2) .$$
(22)

To summarize, if the ϵ expansion is valid on the horizon and the horizon is regular then

$$F_{,\psi}(1,\psi) = 0$$
, (23a)

$$H_{,\psi}(1,\psi) = 0$$
, (23b)

$$\rho_H = 1 + \epsilon \zeta = 1 + \epsilon \left[-6H(1, \pi/2) + H_{,\rho}(1, \pi/2) \right], \qquad (23c)$$

$$\kappa = 1 + \epsilon \chi = 1 + 6\epsilon H(1, \pi/2) . \tag{23d}$$

Thermodynamics. If the horizon is regular then one can talk about thermodynamics of the black hole. The zeroth law of black hole thermodynamics states that the surface gravity is constant on the horizon [16]. The temperature, T, associated with the black hole is

$$\frac{1}{T} \equiv \frac{2\pi\mu}{\kappa} = 2\pi\mu \left(1 - 6\epsilon H(1, \pi/2)\right) \ . \tag{24}$$

One should remember that the physical dimension of the surface gravity is length⁻¹ therefore it is rescaled with μ^{-1} . The entropy of the black hole is related to the area of the horizon

$$S \equiv \frac{A_H}{4G_5} = \frac{2\mu^3}{4G_5} \int_0^{\pi/2} d\psi \int_0^{\pi} d\theta \int_0^{2\pi} d\phi \sqrt{g_{\psi\psi}g_{\theta\theta}g_{\phi\phi}} \Big|_{\rho=1+\epsilon\zeta} = \frac{2\pi\mu^3\rho_H^3}{G_5} \int_0^{\pi/2} d\psi \sin^2\psi \frac{1+\epsilon/2U_1(1,\psi)}{(1-\epsilon\cos\psi)^3} \,. \tag{25}$$

Using the solution (3) and the constraints (22, 23), one can verify that on the horizon $U_1(1,\psi) = 2 \tan^2 \psi \left[H_{,\rho}(1,\pi/2) - H_{,\rho}(1,\psi)\right]$. The entropy is

$$S = \frac{2\pi^2 \mu^3}{4G_5} \left[1 + \epsilon \left\{ 3\zeta + \frac{4}{\pi} + \frac{4}{\pi} \int_0^{\pi/2} d\psi \sin^2 \psi \tan^2 \psi \left[H_{,\rho}(1,\pi/2) - H_{,\rho}(1,\psi) \right] \right\} \right] .$$
(26)

The integral in Eq.(26) can be simplified using integration by parts as

$$\int_{0}^{\pi/2} d\psi \sin^{2} \psi \tan^{2} \psi \left[H_{,\rho}(1,\pi/2) - H_{,\rho}(1,\psi)\right] = \int_{0}^{\pi/2} d\psi \frac{\sin\psi}{\cos^{2}\psi} \sin^{3} \psi \left[H_{,\rho}(1,\pi/2) - H_{,\rho}(1,\psi)\right]$$
$$= \left[\frac{\sin^{3}\psi}{\cos\psi} \left\{H_{,\rho}(1,\pi/2) - H_{,\rho}(1,\psi)\right\}\right]_{0}^{\pi/2} - \int_{0}^{\pi/2} \frac{d\psi}{\cos\psi} \left\{\sin^{3}\psi \left[H_{,\rho}(1,\pi/2) - H_{,\rho}(1,\psi)\right]\right\}_{,\psi}$$
$$= H_{,\rho\psi}(1,\pi/2) - \frac{3\pi}{4}H_{,\rho}(1,\pi/2) + \int_{0}^{\pi/2} \frac{d\psi}{\cos\psi} \left[\sin^{3}\psi H_{,\rho}(1,\psi)\right]_{,\psi} , \qquad (27)$$

where in the last step we have used L'Hopital's rule to evaluate the boundary term at $\psi = \pi/2$. Using Equations (27) and (23c), the entropy (26) can be written as

$$S = \frac{2\pi^2 \mu^3}{4G_5} \left[1 + \epsilon \left\{ -18H(1,\pi/2) + \frac{4}{\pi} \left[1 + H_{,\rho\psi}(1,\pi/2) + \int_0^{\pi/2} \frac{d\psi}{\cos\psi} \left[\sin^3\psi H_{,\rho}(1,\psi) \right]_{,\psi} \right] \right\} \right] .$$
(28)

The first law of black hole thermodynamics states that $T^{-1} = \partial S/\partial M$. In all known black hole solutions the first law is satisfied as a result of Einstein equations. In the ϵ solution it is not trivially satisfied, but should be imposed as a boundary condition. The mass appears in the entropy only through $\mu = \sqrt{8G_5M/(3\pi)}$ and $\epsilon = \mu/\ell$, so

$$\frac{1}{T} = \frac{\partial S}{\partial M} = 2\pi\mu \left[1 + \epsilon \left\{ -24H(1,\pi/2) + \frac{16}{3\pi} \left[1 + H_{,\rho\psi}(1,\pi/2) + \int_0^{\pi/2} \frac{d\psi}{\cos\psi} \left[\sin^3\psi H_{,\rho}(1,\psi) \right]_{,\psi} \right] \right\} \right] .$$
(29)

Equations (29) and (24) must be consistent, therefore, there is another constraint on the function H

$$H(1,\pi/2) = \frac{8}{27\pi} \left[1 + H_{,\rho\psi}(1,\pi/2) + \int_0^{\pi/2} \frac{d\psi}{\cos\psi} \left[\sin^3\psi H_{,\rho}(1,\psi) \right]_{,\psi} \right] \,. \tag{30}$$

IV. THE SINGULAR HORIZON

A detailed study of the horizon in the ϵ solution requires a specific representation of the function $H(\rho, \psi)$, which solves the differential equation (4) and the boundary conditions (5, 6, 23).

A possible representation for the solution is a combination of Associated Legendre functions in ρ and Hypergeometric functions in ψ

$$H(\rho,\psi) = \int d\lambda R(\rho;\lambda)\Psi(\psi;\lambda)$$
(31a)

$$R(\rho;\lambda) = \rho \sqrt{\rho^2 - 1} \left[a(\lambda) Q_{\frac{\lambda-1}{2}}^1 (2\rho^2 - 1) + b(\lambda) P_{\frac{\lambda-1}{2}}^1 (2\rho^2 - 1) \right]$$
(31b)

$$\Psi(\psi;\lambda) = c(\lambda) \,_2F_1(\frac{1-\lambda}{2}, \frac{1+\lambda}{2}, \frac{5}{2}, \sin^2\psi) + \frac{d(\lambda)}{\sin^3\psi} \,_2F_1(\frac{-2-\lambda}{2}, \frac{-2+\lambda}{2}, -\frac{1}{2}, \sin^2\psi) \tag{31c}$$

A lengthy discussion about the expansion (31) appears in [14]. There we assumed that the asymptotic solution beyond the linearized solution includes only integer powers of M. Therefore, the function H is odd in ρ and is given by

$$H(\rho,\psi) = \rho\sqrt{\rho^{2}-1} \left[-\frac{2}{3\pi}Q_{-1/2}^{1}(2\rho^{2}-1)\frac{3[\sin(2\psi)-2\psi\cos(2\psi)]}{8\sin^{3}\psi} + Q_{1/2}^{1}(2\rho^{2}-1)\left(-\frac{1}{3\pi}\frac{3[4\psi-\sin(4\psi)]}{32\sin^{3}\psi} + \frac{d_{2}}{\sin^{3}\psi} - \frac{\pi}{3}\frac{3[8\psi+4\psi\cos(4\psi)-3\sin(4\psi)]}{32\pi^{2}\sin^{3}\psi}\right) + \frac{\pi}{3}\left(P_{1/2}^{1}(2\rho^{2}-1) + \frac{2}{\pi^{2}}\frac{\partial Q_{n-1/2}^{1}(2\rho^{2}-1)}{\partial n}\Big|_{n=1}\right)\frac{3[4\psi-\sin(4\psi)]}{32\sin^{3}\psi} + \sum_{n=2}^{\infty}a(\lambda=2n)Q_{n-1/2}^{1}(2\rho^{2}-1)\frac{3[n\cos(2n\psi)\sin(2\psi)-\cos(2\psi)\sin(2n\psi)]}{8n(1-n^{2})\sin^{3}\psi}\right].$$
(32)

The set $a(\lambda = 2n)_{n>2}$ is fixed by the junction condition (5b)

$$a(\lambda = 2n)_{n \ge 2} = \frac{16}{3\pi(\lambda^2 - 4)} .$$
(33)

The parameter d_2 is undetermined.

The problem appears when one tries to apply the conditions (20, 23) to the solution (32).

• The associated Legendre functions of the second kind are not analytic at $\rho = 1$

$$\rho \sqrt{\rho^2 - 1} Q_{\nu}^1 (2\rho^2 - 1) \sim -\frac{1}{2} - \nu (\nu + 1)(\rho^2 - 1) \ln |\rho^2 - 1| + \mathcal{O}(\rho^2 - 1) .$$
(34)

As a result, one can evaluate the function $H(1, \psi)$ but not the derivative $H_{,\rho}(1, \psi)$.

• The surface gravity, (20), is not constant. The combination $3H + \tan \psi H_{,\psi}$ is not constant on the horizon

$$\kappa = 1 + 2\epsilon \left[3H(1,\psi) + \tan\psi H_{,\psi}(1,\psi) \right] = 1 + 2\epsilon \left[\frac{2\psi\sin\psi}{\pi} + \frac{3\cos\psi}{2\pi} - \sum_{n=2}^{\infty} a(\lambda = 2n) \frac{3\sin(2n\psi)}{4n\sin\psi} \right] = 1 + 2\epsilon \sin\psi .$$
(35)

As was mentioned earlier, according to the theorem by Rácz and Wald [16], if the surface gravity is not constant on the horizon, the horizon is singular. One may notice that on the brane $\kappa_{,\psi}|_{\psi=\pi/2} = 0$. So, according to Eq.(11), although the horizon is singular, the singularity is naked from the bulk but it is covered on the brane.

V. GIVE UP THE POST LINEARIZED ASSUMPTION

We give up the assumption that the asymptotic solution can be expanded in integer powers of M, but still keep the linearized solution as the asymptotic boundary condition. According to Eq.(6) we can include terms of order $\rho^{-(1+2\alpha)}$. It is convenient to work with the Legendre expansion (31). The asymptotic behavior of the associated Legendre functions of the first (second) kind is $\rho \sqrt{\rho^2 - 1} P(Q)^1_{(\lambda-1)/2} (2\rho^2 - 1) \sim \rho^{1+(-)\lambda}$. Therefore, we can include associated Legendre functions of the second kind with $\lambda > 2$ in the expansion. Actually, we have already included even λ terms in the expansion (32). We first try to include also odd λ terms.

$$H(\rho,\psi) = \delta_{0} + \rho \sqrt{\rho^{2} - 1} \left[-\frac{2}{3\pi} Q_{-1/2}^{1} (2\rho^{2} - 1) \frac{3[\sin(2\psi) - 2\psi\cos(2\psi)]}{8\sin^{3}\psi} + Q_{1/2}^{1} (2\rho^{2} - 1) \left(-\frac{1}{3\pi} \frac{3[4\psi - \sin(4\psi)]}{32\sin^{3}\psi} + \frac{d_{2}}{\sin^{3}\psi} - \frac{\pi}{3} \frac{3[8\psi + 4\psi\cos(4\psi) - 3\sin(4\psi)]}{32\pi^{2}\sin^{3}\psi} \right) \\ + \frac{\pi}{3} \left(P_{1/2}^{1} (2\rho^{2} - 1) + \frac{2}{\pi^{2}} \frac{\partial Q_{n-1/2}^{1} (2\rho^{2} - 1)}{\partial n} \bigg|_{n=1} \right) \frac{3[4\psi - \sin(4\psi)]}{32\sin^{3}\psi} \\ + \sum_{\lambda=3}^{\infty} a(\lambda) Q_{(\lambda-1)/2}^{1} (2\rho^{2} - 1) \frac{3[\lambda\cos(\lambda\psi)\sin(2\psi) - 2\cos(2\psi)\sin(\lambda\psi)]}{2\lambda(4 - \lambda^{2})\sin^{3}\psi} \right].$$
(36)

The free parameters are d_2 , δ_0 , and $a(\lambda = 2k + 1)_{k \ge 1}$. Our goal is to make the surface gravity constant at $\rho = 1$. Adding the odd λ terms does not change the junction condition (5b) and therefore the coefficients $a(\lambda = 2n)$ are the same as in Eq.(33). We will try to fix the coefficients δ_0 , $a(\lambda = 2k + 1)$ such that the combination $3H + \tan \psi H_{,\psi}$, which appears in the surface gravity (20), is constant. From Eq.(36) we find that

$$3H(1,\psi) + \tan\psi H_{,\psi}(1,\psi) = 3\delta_0 + \sin\psi - \sum_{k=1}^{\infty} a(\lambda = 2k+1) \frac{3\sin(2k+1)\psi}{2(2k+1)\sin\psi} \equiv 3H(1,\pi/2) .$$
(37)

To solve for a(2k+1), one can multiply (37) by $\sin \psi$ and expand in the set $\{\sin(2k+1)\psi\}_{k=0,1...}$, which is complete and orthogonal over the interval $[0, \pi/2]$ provided that the boundary conditions are f(0) = 0 and $f'(\pi/2) = 0$. The coefficients are

$$\delta_0 = H(1, \pi/2) - \frac{8}{9\pi} , \qquad (38)$$

$$a(\lambda = 2k+1)_{k \ge 1} = \frac{-16}{3\pi(\lambda^2 - 4)} .$$
(39)

Since we have multiplied (37) with $\sin \psi$ one should verify that (37) holds for $\psi = 0$, and indeed it is.

0.7

At this stage we have to verify the consistency of; (i) the representation of the function H, Eq.(36), (ii) the zeroth law of thermodynamics, Eqs.(38, 39), and (iii) the first law of thermodynamics, Eq.(30).

We use the following expansions for the Legendre functions around $\rho = 1$

$$\rho \sqrt{\rho^2 - 1} P_{\nu}^1 (2\rho^2 - 1) = \nu(\nu + 1)(\rho^2 - 1) + \frac{\nu^2(\nu + 1)^2}{2} (\rho^2 - 1)^2 + \mathcal{O}(\rho^2 - 1)^3, \qquad (40)$$

$$\rho \sqrt{\rho^2 - 1} Q_{\nu}^1 (2\rho^2 - 1) = -\frac{1}{2} + \frac{\nu(\nu + 1)}{2} (\rho^2 - 1) \left[1 - \ln(\rho^2 - 1) - 2\gamma_E - 2\bar{\psi}(\nu + 1) \right]$$

$$+ \frac{\nu^2(\nu + 1)^2}{4} (\rho^2 - 1)^2 \left[\frac{5\nu^2 + 5\nu + 2}{2\nu(\nu + 1)} - \ln(\rho^2 - 1) - 2\gamma_E - 2\bar{\psi}(\nu + 1) \right] + \mathcal{O}(\rho^2 - 1)^3, \qquad (41)$$

where γ_E is Euler's constant and $\bar{\psi}(z)$ is the logarithmic derivative of the Gamma function, $\bar{\psi}(z) = d/dz \ln \Gamma(z)$. First, we evaluate Eq.(36) at $\rho = 1$, $\psi = \pi/2$

$$H(1,\pi/2) = \delta_0 + \frac{1}{8} + \frac{1}{32} - \frac{d_2}{2} + \frac{3}{32} - \frac{1}{2} \sum_{\lambda=3}^{\infty} \frac{(-1)^{\lambda} 16}{3\pi (\lambda^2 - 4)} \frac{3\sin(\lambda\pi/2)}{\lambda(4 - \lambda^2)}$$
$$= \delta_0 + \frac{1}{4} - \frac{d_2}{2} - \frac{8}{\pi} \sum_{k=1}^{\infty} \frac{(-1)^k}{(2k+1)((2k+1)^2 - 4)^2} = \delta_0 + \frac{8}{9\pi} - \frac{d_2}{2}.$$
 (42)

Comparing Eq.(42) with the result of the first law, Eq.(38), we conclude that $d_2 = 0$.

Second, we would like to evaluate the first law, Eq.(30). For this we need to evaluate $H_{,\rho}(1,\psi)$. We use the expansions (40, 41) in Eq.(36) and find that

$$H_{,\rho}(1,\psi) = \frac{h_0(\psi)}{\sin^3\psi} \left[\ln(\rho^2 - 1) + 2\gamma_E + 2\bar{\psi}(1/2) \right] + h_1(\psi) .$$

$$(43)$$

$$-7\psi + 4\psi\cos 2\psi + 3\psi\cos 4\psi - 2\sin 2\psi + \sin 4\psi$$

$$h_0(\psi) = \frac{32\pi}{32\pi} + \frac{2}{\pi} \sum_{\lambda=3}^{\infty} \frac{(-1)^{\lambda} (\lambda^2 - 1)}{\lambda (\lambda^2 - 4)^2} \left[\lambda \cos \lambda \psi \sin 2\psi - 2 \cos 2\psi \sin \lambda \psi\right] .$$

$$(44)$$

$$h_{1}(\psi) = \frac{5\psi - 2\sin 4\psi + 3\psi\cos 4\psi}{8\pi\sin^{3}\psi} + \frac{4}{\pi}\sum_{\lambda=3}^{\infty}\frac{(-1)^{\lambda}(\lambda^{2} - 1)}{\lambda(\lambda^{2} - 4)^{2}} \left[\bar{\psi}\left(\frac{\lambda + 1}{2}\right) - \bar{\psi}\left(\frac{1}{2}\right)\right]\frac{\lambda\cos\lambda\psi\sin 2\psi - 2\cos 2\psi\sin\lambda\psi}{\sin^{3}\psi} .$$
(45)

Let us deal with the function $h_0(\psi)$, which multiplies the diverging term $\ln(\rho^2 - 1)$. One can verify that $h_0(\psi = \pi/2) = 0$ and $h_0(\psi = 0) = 0$. Therefore, one can expand the function $h_0(\psi)$ in the set $\{\sin 2n\psi\}_{n=1,2...}$, which is complete and orthogonal over the interval $[0, \pi/2]$. It is easy to verify that for $n \ge 1$

$$\int_0^{\pi/2} h_0(\psi) \sin 2n\psi \, d\psi = 0 \ . \tag{46}$$

So, the function $h_0(\psi) = 0$ and the first derivative of the function H is finite on the horizon. The first law, Eq.(30), together with Eq.(38) can be used to find the value of δ_0

$$\delta_0 = \frac{8}{27\pi} \left[-2 + h_1'(\pi/2) + \int_0^{\pi/2} \frac{d\psi}{\cos\psi} \left[h_1(\psi) \sin^3\psi \right]_{,\psi} \right] \,. \tag{47}$$

From Eq.(45) one can deduce the following

$$\left[h_{1}(\psi)\sin^{3}\psi\right]_{,\psi} = \frac{5-5\cos4\psi - 12\psi\sin4\psi}{8\pi} - \frac{4}{\pi}\sum_{\lambda=3}^{\infty}\frac{(-1)^{\lambda}(\lambda^{2}-1)}{\lambda(\lambda^{2}-4)}\left[\bar{\psi}\left(\frac{\lambda+1}{2}\right) - \bar{\psi}\left(\frac{1}{2}\right)\right]\sin2\psi\sin\lambda\psi \ . (48)$$

$$h_1'(\pi/2) = \left[h_1(\psi) \sin^3 \psi \right]_{,\psi} \Big|_{\psi=\pi/2} = 0 .$$
(49)

$$\int_{0}^{\pi/2} \frac{d\psi}{\cos\psi} \left[h_1(\psi) \sin^3\psi \right]_{,\psi} = \frac{5}{\pi} + \frac{2}{\pi} \sum_{n=2}^{\infty} \frac{(-1)^n}{(n^2 - 1)} \left[\bar{\psi} \left(n + 1/2 \right) - \bar{\psi} \left(1/2 \right) \right] = 2 .$$
(50)

Now, substitute in Eq.(47) to find $\delta_0 = 0$.

To summarize, if we give up the post linearized assumption and allow for asymptotic behavior of $M^{n/2}$ then we can satisfy the requirement that the surface gravity is constant and the solution appears to be regular on the horizon. The function H and the first derivative $H_{,\rho}$ are finite on the horizon. The derivatives $H_{,\psi}$ and $H_{,\psi\psi}$ are vanishing there. Therefore, based on Eq.(4) we can conclude that $H_{,\rho\rho}$ is finite on the horizon as well. This is a sufficient condition for the metric to be C^1 . In general this is not sufficient to ensure that the the metric is not singular. To double check, we look at the Kretchmann scalar up to first order in ϵ

$$K = R^{abcd} R_{abcd} = \frac{72}{\rho^8} - \epsilon \left[\frac{576}{\rho^8} \left(F(\rho, \psi) + \rho(\rho^2 - 1)\cos\psi \right) + \frac{96}{\rho} \left(\frac{3H(\rho, \psi) + \tan\psi H_{,\psi}(\rho, \psi)}{\rho^6} \right)_{,\rho} \right] .$$
(51)

As one can see, the Kretchmann scalar depends only on the first derivative $H_{,\rho}(\rho,\psi)$, and therefore it is finite. An ambiguity is left for the second order in ϵ . The Kretchmann scalar at the second order in ϵ depends on higher derivatives of H, which might be divergent. However, second order in ϵ in the Kretchmann scalar depends on the first and second orders in the metric. So nothing can be done at this stage before finding the second order corrections to the metric. These should cancel the divergence part coming from the first order in the metric. We should emphasize that the situation with the surface gravity is different. It is true that if the surface gravity is not constant in first order in ϵ , it will affect the Kretchmann scalar only at second order in ϵ . However, this contribution cannot be compensated by second order corrections to the metric. Therefore, the surface gravity must be constant at all orders.

What about the asymptotic behavior of the metric? The inclusion of half integer powers of the mass changes the post Newtonian potential. For large ρ , the terms with $a(\lambda = 2k+1)_{k\geq 1}$ contribute to the function H (and throughout Eq.(5d) to the function F) terms of order ρ^{-2k} . As a result, the gravitational potential, g_{tt} , acquires terms of the form

$$\epsilon \frac{a(\lambda = 2k+1)}{\rho^{2k+2}} \sim \frac{a(\lambda = 2k+1)(G_5M)^{k+3/2}}{\bar{\rho}^{2k+2\ell}} , \qquad (52)$$

where $\bar{\rho} = \mu \rho$ is a dimensionful coordinate. The term with δ_0 would have contributed a term of the form $\delta_0 (G_5 M)^{3/2} \bar{\rho}^{-2} \ell^{-1}$. But, since $\delta_0 = 0$ the half integer terms start only from $M^{5/2}$.

We learn that, in general, the asymptotic solution should be expanded in powers of $(G_5M)^{1/2}$. The lowest order must be $(G_5M)^1$ and not $(G_5M)^{1/2}$ otherwise there will not be a well defined conserved mass. One might expect that by iterating the lowest order there will be only integer powers of M. However, the boundary conditions on the horizon require also half integer powers of the mass. In particular it is the zeroth law of black hole thermodynamics (constant surface gravity) that forced us to include the half integer terms. The coefficients of the terms $(G_5M)^{k/2}$, k = 5, 7..., which are $a(\lambda = 2k + 1)_{k\geq 1}$, are completely determined by the zeroth law (39). The coefficient of $(G_5M)^{3/2}$, which is δ_0 , is not fixed by the requirement that the surface gravity is constant. The actual value of the surface gravity depends on δ_0 . The first law of black hole thermodynamics, dM = T dS, forces δ_0 to vanish, and therefore eliminates the term of order $(G_5M)^{3/2}$ from the asymptotic expansion.

The dominant part of the post Newtonian potential remains $(G_5 M)^2/(r^2 \ell^2)$, which is responsible for the precession of perihelion calculations. Since we don't know of any experimental evidence for higher order terms in the potential, it looks like the half integer terms cannot be detected experimentally.

VI. SUMMARY AND DISCUSSION

We study a small black hole located on the TeV brane in Randall-Sundrum I (two branes) scenario. We expand the metric in ϵ , which is the ratio between the Schwarzschild radius and the bulk curvature length. We find the solution up to first order in ϵ . The solution satisfies Einstein's equations in the bulk and Israel junction conditions on the TeV brane. The asymptotic form of the metric is fixed by the weak field approximation (linearized gravity).

In Randall-Sundrum II (single brane scenario)[6] it was conjectured that there were no large static black holes localized on the brane[17]. However, small black holes might exist and one can apply our method of ϵ expansion to find them. Although the linearized solution for a single brane is very different from that for two branes, in first order of ϵ they are similar. The only difference is $\epsilon \to -\epsilon$. So, the results for the ϵ solution and the discussion of regularity are equivalent in the two scenarios.

A crucial issue is the post linearized form of the asymptotic solution. For Four dimensional black holes the post Newtonian behavior is given in terms of an expansion in integer powers of G_4M/r (the only dimensionless combination). The equivalent in RS models would be an expansion in integer powers of $G_5M/(r\ell)$. However, if we assume that the post linearized metric includes only integer powers of the mass then the surface gravity on the horizon is not constant. This means that the horizon is singular, and the configuration of a small mass on the TeV brane describes a naked singularity. As we mentioned earlier, the singularity is naked from the bulk but it is covered on the brane.

In section V we have shown that there is a solution with a regular horizon, i.e. the surface gravity is constant and the Kretchmann scalar is finite up to first order in ϵ . However, this requires a different asymptotic form of the metric, as one must include half integer powers of the mass. In general, the expansion parameter for the asymptotic solution is $(G_5M)^{1/2}$, but still the lowest order is $(G_5M)^1$ and not $(G_5M)^{1/2}$ otherwise there will not be a well defined conserved mass. Surprisingly, the first law of black hole thermodynamics, dM = T dS, eliminates the term of order $(G_5M)^{3/2}$, as well, from the asymptotic expansion. The leading term in the post Newtonian potential remains $(G_5M)^2/(r^2\ell^2)$, which is the term responsible for the precession of perihelion calculations in the Schwarzschild metric. The next order term is $(G_5M)^{5/2}$. This term does not exist in the four dimensional Schwarzschild metric, but it is very hard to detect such a term in any asymptotic measurement.

As we mentioned earlier, these results are also valid for small black holes in RS single brane scenario. In principle, the post Newtonian behavior can be verified numerically. However, the asymptotic behavior of the numerical solution of [13] is not well known [18]. Future numerical studies with a regular horizon might detect the fractional power behavior in the post Newtonian potential.

The configuration discussed in this paper, of a small (few TeV) mass on the brane in RS scenario, can be the final state of high energy particle collisions. In some sense such a collision is similar to the collapse of matter in ordinary four dimensional gravity, it might end up as a black hole or generate a naked singularity[19]. If the mass of the object is somewhat bigger than the Planck mass (TeV) the behavior of these two possibilities is very different. The black hole will radiate thermally. It will mainly radiate ordinary particles on the brane. The naked singularity has no temperature, it evaporates in an explosive fashion in the order of Planck time. It will mainly radiate gravitons into the bulk (where the singularity is naked) and not on the brane (where it is covered). further investigations of this issue will be carried out elsewhere.

Acknowledgments

This work is supported in part by the U.S. Department of Energy Grant No. DE-FG02-84ER40153. We thank Richard Gass and Cenalo Vaz for fruitful discussions.

- [1] R.C. Myers and M.J. Perry, Annals Phys. 172:304 (1986).
- [2] S.W. Hawking and D.N. Page, Commun. Math. Phys. 87:577 (1983).
- [3] D. Birmingham, Class. Quant. Grav. 16:1197-1205 (1999).
- [4] A. Chamblin, S.W. Hawking, H.S. Reall, Phys. Rev. D61:065007 (2000).
- [5] L. Randall and R. Sundrum, Phys. Rev. Lett. 83:3370-3 (1999).
- [6] L. Randall and R. Sundrum, Phys. Rev. Lett. 83:4690-3 (1999).
- [7] S. Giddings and E. Katz J. Math. Phys. 42:3082-3102 (2001);
 - T. Banks and W. Fischler, *Preprint* hep-th/9906038;
 - S. Dimopoulos and G. Landsberg, Phys. Rev. Lett. 87:161602 (2001);

S.B. Giddings and S. Thomas, *Phys. Rev.* D65:056010 (2002);

- D.M. Eardley and S.B. Giddings, *Phys. Rev.* D66:044011 (2002) gr-qc/0201034.
- [8] L.A. Anchordoqui, J.L. Feng, H. Goldberg and A.D. Shapere, *Phys. Rev.* D65:124027 (2002) hep-ph/0112247.
- [9] N. Dadhich et. al. Phys. Lett. B487:1 (2000);
 - C. Germani and R. Maartens, Phys. Rev. D64:124010 (2001);
 - R. Casadio, et.al. Phys. Rev. D65:084040 (2002);
 - T. Shiromizu, K. I. Maeda and M. Sasaki, Phys. Rev. D62:024012 (2000);
- R. Casadio and L. Mazzacurati, Mod. Phys. Lett. A18: 651-660 (2003) gr-qc/0205129.
- [10] P. Kanti and K. Tamvakis, Phys. Rev. D65:084010 (2002) hep-th/0110298;
- P. Kanti, I. Olasagasti and K. Tamvakis *Phys. Rev.* D68:124001 (2003) hep-th/0307201.
 [11] S.B. Giddings *et.al. JHEP*0003:023 (2000);
- J. Garriga and T. Tanaka, Phys. Rev. Lett. 84:2778 (2000).
- [12] A. Chamblin et.al. Phys. Rev. D63:064015 (2001);
- T. Wiseman, *Phys. Rev.* D**65**:124007 (2002).
- [13] H. Kudoh, T. Tanaka, and T. Nakamura Phys. Rev. D68:024035 (2003)
- [14] D. Karasik, C. Sahabandu, P. Suranyi, and L.C.R. Wijewardhana, Phys. Rev. D69:064022 (2004). gr-qc/0309076.
- [15] W. Israel, Nuovo Cimento B44:1 (1966); B48:463 (1966).
- [16] I. Rácz and R. M. Wald, Class. Quant. Grav. 9:2643-2656 (1992).
- [17] T. Tanaka, Prog. Theor. Phys. Suppl 148:307 (2003);
 R. Emparan, A. Fabbri and N. Kaloper, JHEP 08:043 (2002)
- [18] Private communication with the authors of [13].
- [19] T. Harada et. al., Phys. Rev. D64:041501 (2001);
 T.P. Singh, C. Vaz, Phys. Lett. B481:74-78 (2000);
 T.P. Singh, C. Vaz, Phys. Rev. D61:124005 (2000);
 C. Vaz, L. Witten, Phys. Lett.B442:90-96 (1998);
 S. Barve, T.P. Singh, C. Vaz, L. Witten, Nucl. Phys. B532:361-375 (1998)
- [20] This solution is obtained from the solution in [14] using redefinition of $A \to A/B$ and redefining the gauge function $F \to F 6H_0 + (2\rho \rho^{-1})H'_0$.