Two-dimensional description of *D*-dimensional static black holes with pointlike sources

Mariano Cadoni*

Dipartimento di Fisica, Università di Cagliari, and INFN, Sezione di Cagliari, Cittadella Universitaria 09042 Monserrato, Italy

Salvatore Mignemi[†]

Dipartimento di Matematica, Università di Cagliari, viale Merello 92, 09123 Cagliari, Italy, and INFN, Sezione di Cagliari

We derive two-dimensional (2D) solutions of a generic dilaton gravity model coupled with matter, which describe D-dimensional static black holes with pointlike sources. The equality between the mass M of the D-dimensional gravitational solution and the mass m of the source can be preserved also at the level of the 2D gravity model.

Two-dimensional (2D) models of gravity have been used in various contexts as an effective description of *D*-dimensional gravity. In the case of pure gravity the 2D dilaton gravity model (see e.g. Ref. [1] and references therein) can be integrated exactly and it can be used to give an effective description of the radial modes of *D*-dimensional gravity at the classical, semiclassical and full quantum level. It is almost impossible to give a comprehensive list of all the problems of black hole physics that have been tackled using 2D dilaton gravity models. The Hawking radiation and the information puzzle [2] for black holes, the microscopic interpretation of the black hole entropy [3] and the construction of acoustic analogues of gravitational black holes [4] are only a few of many examples.

Unfortunately, many nice features of the 2D model are spoiled by the presence of matter. The existence and uniqueness of the solutions are consequence of a 2D Birkhoff's theorem [5, 6], which has been shown to hold for pure 2D dilaton gravity but in general does not hold when matter is present. This is a strong limitation of 2D models. It is therefore important to look for general solutions of matter-coupled dilaton gravity models.

In this letter we will consider a generic 2D dilaton gravity model coupled with a pointlike source. The model gives an effective description of *D*-dimensional static black holes with pointlike sources. Moreover, gravitational models in which the matter contribution is represented by δ -like terms emerge in brane world scenarios [7, 8]. We will derive the general solution of the model and show that choosing appropriately the integration constants, the equality between gravitational and inertial mass can be preserved at the 2D level.

Let us consider generic D-dimensional Einstein gravity coupled to matter fields and a pointlike source of mass m. The action is [9]

$$A_D = \frac{1}{16\pi G_D} \int d^D x \sqrt{-g} \ R - \int d^D x \sqrt{-g} \ \mathcal{L}_{MF} - m \int dt \sqrt{-g_{ij}} \frac{dx^i}{dt} \frac{dx^j}{dt},\tag{1}$$

where G_D is the *D*-dimensional Newton constant and \mathcal{L}_{MF} is the Lagrangian for the matter fields (gauge fields, scalar fields etc).

It is well known that in absence of the pointlike source *D*-dimensional spherically symmetric gravity allows for an effective two-dimensional description in terms of a 2D dilaton gravity model. The 2D model is obtained by retaining only the radial modes in the action (1). After a suitable Weyl rescaling of the metric one gets the 2D action $A_2 = \int d^2x \sqrt{-g} \left(\phi R + \lambda^2 V(\phi)\right)$. The scalar field ϕ (the dilaton) parametrizes the volume of the transverse (D-2)-dimensional sphere, λ is a parameter with dimensions of length⁻¹ and the dilaton potential $V(\phi)$ depends both on the dimension of the spacetime *D* and on the form of the matter field action \mathcal{L}_{MF} (We shall only consider the case in which the matter fields can be treated, from the 2D point of view, as constant background solutions).

When the pointlike source is present the situation becomes more involved and it is not a priori evident that a 2D description of the gravity model (1) exists. Two main problems arise. First, the *D*-dimensional stress-energy tensor for the pointlike source is proportional to $\delta^D(x - x(\tau))$. Solving the *D*-dimensional field equations in the case of spherical symmetry is essentially equivalent to find the Green's functions of the Laplace operator in D-1 dimensional Green's functions is required. Assuming that this correspondence between (D-1)-dimensional and one-dimensional Green's functions is required. Assuming that this correspondence has been found, one is faced with a second problem. Even though the mass M of the D-dimensional solution equals the mass m of the the source, one may end up with a 2D solution for which $M \neq m$.

Progress in solving the first problem has been achieved in Ref. [10] (see also Refs. [11, 12, 13, 14]). In that paper the one-dimensional Green's functions have been given as functions of |r|, r being the spacelike coordinate of the 2D spacetime, but unfortunately, the solutions of the 2D gravity model are characterized by $m \neq M$. Retaining only the radial modes in the action (1) and after a Weyl rescaling of the 2D metric needed to get rid of the kinetic terms for the dilaton, one obtains a 2D model characterized by a dilaton potential $V(\phi)$ and a coupling function $W(\phi)$,

$$A = A_G + A_M = \frac{1}{2} \int d^2x \sqrt{-g} \left(\phi R + \lambda^2 V(\phi)\right) - m \int dt \, W(\phi) \sqrt{-g_{\mu\nu} \frac{dx^{\mu}}{dt} \frac{dx^{\nu}}{dt}} \,. \tag{2}$$

The form of the coupling function $W(\phi)$ depends both on the dimensionality D of the spacetime and on the Weyl rescaling performed on the 2D metric.

The field equations for the 2D dilaton gravity model (2) are (we do not consider the motion of the source)

$$R = -\lambda^2 \frac{dV}{d\phi} + \frac{2m}{\sqrt{-g}} \frac{dW}{d\phi} \int d\tau \, \delta^2(x - x(\tau)),$$

$$-(\nabla_\mu \nabla_\nu \phi - g_{\mu\nu} \nabla^2 \phi) - \frac{1}{2} g_{\mu\nu} \lambda^2 V = T^{(M)}_{\mu\nu},$$
(3)

where τ is the proper time and $T^{(M)}_{\mu\nu}$ is the stress energy tensor for the pointlike matter source,

$$T^{(M)}_{\mu\nu} = \frac{m}{\sqrt{-g}} W(\phi) \int d\tau \, \delta^2(x - x(\tau)) u_\mu u_\nu, \tag{4}$$

with u^{μ} the 2-velocity of the particle.

If we consider the field equations in vacuum $T^{(M)}_{\mu\nu} = 0$, we can define a scalar function [15]

$$M = \frac{1}{2\lambda} \left(\lambda^2 \int V(\phi) d\phi - (\nabla \phi)^2 \right), \tag{5}$$

which is constant by virtue of the field equations and gives the mass of the gravitational solution.

Adopting static coordinates and the Schwarzschild gauge,

$$ds^{2} = -U(r)dt^{2} + U^{-1}(r)dr^{2},$$
(6)

and considering a source particle at rest at the origin, the field equations (3) become,

$$\frac{d^2 U}{dr^2} = \lambda^2 \frac{dV}{d\phi} - 2m \frac{dW}{d\phi} \,\delta(r),$$

$$\frac{dU}{dr} \frac{d\phi}{dr} = \lambda^2 V,$$

$$2U \frac{d^2 \phi}{dr^2} + \frac{dU}{dr} \frac{d\phi}{dr} = \lambda^2 V - 2m W \,\delta(r),$$
(7)

The general solution of Eqs. (7) is a generalization of that found in Ref. [10]. In two spacetime dimensions the coordinate r has not the meaning of a radial coordinate, so that one can take $-\infty < r < \infty$. The delta function singularity at r = 0 can be generated writing the solution as function of |r|,

$$U = \frac{J(\phi)}{\sigma^2} - \gamma, \quad \phi = \sigma \lambda |r| + \beta, \tag{8}$$

where $J(\phi) = \int V(\phi) d\phi$ and σ, β, γ are integration constants. Inserting Eqs. (8) into Eqs. (7), one easily finds that (8) is a solution of the system (7) if the integration constants satisfy

$$J(\beta) - \sigma^2 \gamma = -\frac{m\sigma}{2\lambda} W(\beta), \quad V(\beta) = -\frac{m\sigma}{\lambda} \frac{dW}{d\phi}(\beta).$$
(9)

In general the mass M of the solution, calculated using Eq. (5) away from the source in the r > 0 region, is not equal to the mass m of the source. However, in 2D an additional freedom is present compared to higher dimensions. In fact, the solution (8) is parametrized by three integration constants constrained by only two equations (9). The remaining freedom can be fixed requiring m = M. The origin of this additional freedom can be traced back to the peculiarity of 2D gravity: the radial coordinate has no natural normalization in two dimensions. More precisely, the field equations (7) in absence sources are invariant under the rescaling $r \to \Omega r$, $U \to \Omega^2 U$, with constant Ω . This is analogous to the possibility of freely rescaling the time coordinate in generic dimension. The invariance implies the presence in the solution (8) of an integration constant, σ , with no higher-dimensional analogue, that can be fixed arbitrarily. When sources are added, one must of course rescale also m to preserve the invariance.

Requiring m = M and using equation (8) into Eq. (5) one finds

$$\gamma = \frac{2m}{\lambda\sigma^2}.$$
(10)

As an example, we consider the case $V = a\phi^h$, $W = b\phi^k$ (a, b, h, k are arbitrary real constants). Using equations (8), (9) and (10) one finds the solution

$$U = \frac{a}{\sigma^2(h+1)} (\sigma\lambda|r| + \beta)^{h+1} - \gamma, \qquad \phi = \sigma\lambda|r| + \beta, \tag{11}$$

with

$$\beta = B^{\frac{1}{(h+1)}} \left(\frac{m}{\lambda a}\right)^{\frac{1}{h+1}}$$

$$\sigma = -\frac{1}{bk} B^{\frac{h-k+1}{h+1}} \left(\frac{m}{\lambda a}\right)^{-\frac{k}{h+1}}$$

$$\gamma = 2ab^2 k^2 B^{\frac{2(k-h-1)}{h+1}} \left(\frac{m}{\lambda a}\right)^{\frac{2k+h+1}{h+1}}.$$
(12)

where B = (4k(h+1)/(2k-h-1)). The solution exists for every B > 0.

A particularly interesting special case of the power-law potentials discussed above is the dimensional reduction of the four-dimensional (4D) Schwarzschild solution. Consider the 4D Einstein action coupled to a point particle of mass m given by Eq. (1) with $\mathcal{L}_{MF} = 0$, and substitute the ansatz

$$ds_{(4)}^2 = ds_{(2)}^2 + \frac{2}{\lambda^2} \phi \, d\Omega_2^2. \tag{13}$$

After a Weyl rescaling of the 2D metric,

$$g \to \frac{1}{\sqrt{2\phi}} g,$$
 (14)

one finds a 2D model of the form (2) with

$$V(\phi) = (2\phi)^{-1/2}, \quad W(\phi) = (2\phi)^{-1/4}.$$
 (15)

Using Eqs. (11) and (12), with $h = -\frac{1}{2}, k = -\frac{1}{4}, a = \frac{1}{\sqrt{2}}, b = (2)^{-1/4}$, one easily finds for r > 0

$$U = \sigma^{-2}\sqrt{2\phi} - \gamma, \quad \phi = \sigma\lambda r + \beta, \quad \sigma = 2\sqrt{\frac{m}{\lambda}}, \quad \beta = \frac{m^2}{2\lambda^2}, \quad \gamma = \frac{1}{2}.$$
 (16)

It is also straightforward to check that after going back to the original Weyl frame, $g_s = g/\sqrt{2\phi}$, changing coordinates

$$t' = \frac{t}{\sigma}, \quad R^2 = \frac{2}{\lambda^2}\phi = \frac{2}{\lambda^2}\left(\sigma\lambda r + \beta\right),\tag{17}$$

and setting $G = \lambda^{-2}$, the 2D solution (16) reduces to the 4D Schwarzschild solution, $ds^2 = -(1 - 2Gm/R)dt'^2 + (1 - 2Gm/R)^{-1}dR^2 + R^2 d\Omega_2^2$.

In the same way one may discuss the dimensional reduction of more general solutions like Reissner-Nordström or Schwarzschild-anti de Sitter.

 $^{^{\}ast}$ Electronic address: mariano.cadoni@ca.infn.it

 $^{^{\}dagger}$ Electronic address: smignemi@vaxca1.unica.it

- [1] D. Grumiller, W. Kummer and D. V. Vassilevich, Phys. Rept. 369 (2002) 327 [arXiv:hep-th/0204253].
- [2] C. G. Callan, S. B. Giddings, J. A. Harvey and A. Strominger, Phys. Rev. D 45 (1992) 1005 [arXiv:hep-th/9111056].
- [3] M. Cadoni and S. Mignemi, Nucl. Phys. B 557 (1999) 165 [arXiv:hep-th/9902040].
- [4] M. Cadoni, Class. Quant. Grav. 22 (2005) 409 [arXiv:gr-qc/0410138].
- [5] D. Louis-Martinez and G. Kunstatter, Phys. Rev. D 49 (1994) 5227.
- [6] M. Cavaglia, V. de Alfaro and A. T. Filippov, Phys. Lett. B 424 (1998) 265 [arXiv:hep-th/9802158].
- [7] L. Randall and R. Sundrum, Phys. Rev. Lett. 83 (1999) 4690 [arXiv:hep-th/9906064].
- [8] P. Brax, C. van de Bruck and A. C. Davis, Rept. Prog. Phys. 67 (2004) 2183 [arXiv:hep-th/0404011].
- [9] We adopt the notations of S. Weinberg, Gravitation and Cosmology, New York 1972.
- [10] M. Melis and S. Mignemi, arXiv:gr-qc/0411136.
- [11] R. B. Mann, A. Shiekh and L. Tarasov, Nucl. Phys. B **341** (1990) 134.
- [12] R. B. Mann, M. S. Morris and S. F. Ross, Class. Quant. Grav. 10 (1993) 1477 [arXiv:hep-th/9202068].
- [13] M. M. Leite and V. O. Rivelles, Phys. Lett. B 392 (1997) 305 [arXiv:hep-th/9608049].
- [14] C. G. Boehmer and P. Bronowski, arXiv:gr-qc/0502030.
- [15] R. B. Mann, Phys. Rev. D 47 (1993) 4438 [arXiv:hep-th/9206044].