

# Essential Constants in General Relativity

G.O. Papadopoulos\*

*National & Kapodistrian University of Athens, Physics Department  
Nuclear & Particle Physics Section  
Panepistimioupolis, Ilisia GR 157-71, Athens, Hellas*

**Dedication** *The author wishes to dedicate this work to his nieces.*

## Abstract

The issue of the essential constants, within the context of General Relativity, is studied in the present short work. A rather commonly accepted definition as well as a conclusive way to pinpoint them from a set of constants in a given metric tensor field, which may or may not be a solution to the Einstein's Field Equations, are given. Although the problem of distinction between essential and spurious constants is classic, no particular progress can be found in the literature so far. It is this gap that the present work comes to fill.

**MSC-Class** (2000): 83C05;83C15;83C20

**Keywords:** Essential Constants, Spurious Constants

## 1 Introduction

As it is well known, the Einstein's Field Equations (EFE's) are a set of highly nonlinear, mixed, partial differential equations of the second order. Typically and according to the standard theory of Differential Equations, in order to solve them, one needs initial and/or boundary conditions. Of course in practice, there are a lot of problems ensuing from the nonlinear and hyperbolic character of the EFE's and the existence of constraints (due to the Bianchi Identities); e.g., the causal structure of an arbitrary Einstein space-time can have some severe pathologies, which usually can be avoided by postulating the existence of a spacelike Cauchy hypersurface endowed

---

\*E-Mail: gpapado@cc.uoa.gr

with the property, that any causal curve intersects it at precisely one point (*globally hyperbolic space-times*) [1].

It is a common practice that by the integration procedure of the EFE's, either no initial and/or boundary conditions are given or, taking into account the complexity of the EFE's, some Ansätze concerning the metric tensor field are employed. In any case, one ends up with a metric tensor field containing a set of constants. Some of them correspond to the lack of initial and/or boundary conditions, while some others are related to the employed Ansätze.

Let an Einstein space-time which is described by a pair  $(\mathcal{M}, g)$ , where  $\mathcal{M}$  is a 4-dimensional manifold and  $g$  is an Einstein metric tensor field on it; a non degenerate, covariant tensor field of order 2, with the property that at each point of  $\mathcal{M}$  one can choose a frame of 4 vectors  $\{e_0, e_1, e_2, e_3\}$ , such that:  $g(e_\alpha, e_\beta) = \eta_{\alpha\beta}$ , where  $\eta$  is a diagonal matrix with entries  $\{-1, 1, 1, 1\}$  (i.e., the Minkowskian metric)<sup>1</sup>. In principle, this metric tensor field will depend not only on the space-time points  $x^\mu$  but also on a set of constants  $\{\lambda^{(1)}, \dots, \lambda^{(n)}\}$ . In a local coordinate system, it is<sup>2</sup>:

$$g_{\alpha\beta} = g_{\alpha\beta}(x^\mu; \lambda^{(i)}) \quad (1.1)$$

There are two cases: this metric tensor field is either a solution to the EFE's or not; then it simply defines a geometry without any particular physical meaning. Among these constants, some are integration constants, if the metric tensor field (1.1) is a solution to the EFE's, and some are (obviously) spurious –since they can be imposed by hand (e.g., a changing in the scale and the origin of the  $x^0 \equiv t$  (time) coordinate:  $t \rightarrow c_1 t + c_2$ ). Logically, one would like to dispose of the redundant constants.

In the first case (the metric tensor field is a solution to the EFE's), and if the integration procedure of the EFE's as well as the employed Ansätze –if any– are known, then in each and every step of this procedure, the residual coordinate freedom of the given metric tensor field can be determined and used in order to eliminate some constants –if possible. The problem here is twofold: one must use brute force i.e., try to find (usually, by inspection or by hypothesis) those general coordinate transformations (GCT's) which eliminate the constant(s) under discussion but if one is not able to find them, then one can not claim that the constant(s) is(are) essential –since, in general, there is no way to assure that there are not such GCT's.

In the second case (the metric tensor field is not a solution to the EFE's), one must guess which of the constants may be spurious and try to use an eliminating procedure in terms of general coordinate transformations. The previously mentioned twofold problem occurs here as well.

---

<sup>1</sup>Greek indices take values on the interval  $\{0, \dots, 3\}$ .

<sup>2</sup>Latin indices take values on the interval  $\{0, \dots, n\}$ .

## 2 Essential & Spurious Constants

It seems that, the only case in which one can cope with the aforementioned difficulties is when Homotheties exist (see e.g., [2]): indeed, if a metric tensor field admits an homothetic vector field, then an overall constant in front of the metric tensor field can be eliminated and this constant turns out to be spurious; a well known result [2].

In order to find a systematic and compact way to attack the problem of the essential constants, one must return to the metric tensor field (1.1) and pose a naturally arisen question:

*Which of these constants describe, at least locally, the inner properties of the geometry of the pair  $(\mathcal{M}, g)$ ?*

It is well known that the geometry of an arbitrary space-time—in fact of any Riemannian space— can be described, at least locally, by its *Curvature Invariant Relations*; functionally independent relations among scalars, which are constructed from the Riemann tensor and its covariant derivatives up to a given order (although this scheme fails in the case of the pp-waves; see e.g., [2], Chapter 9, for a relative discussion on these issues). Barring degeneracies, such as pp-waves where all above mentioned scalars vanish, curvature invariant relations have one very important property: they do not depend on space-time points  $x^\mu$ ; their forms are invariant statements (i.e., they have the same form in all coordinate systems). Consequently if they depend on some parameters/constants, that means these constants, which clearly are some from those in the metric tensor field, are not affected by GCT's. For, if a constant of the metric tensor field could be eliminated by a GCT, then the metric tensor field in the new coordinate system as well as all the curvature invariant relations based on it would lack this particular constant; an invariant statement, since the curvature invariant relations are form invariant. Therefore, only the non absorbable constants will appear in the curvature invariant relations and in this sense, they will describe, indirectly, the inner properties of the geometry of the pair  $(\mathcal{M}, g)$ .

These thoughts lead to the following:

**Definition 2.1.** A constant  $\lambda^{(i)}$ , for a given  $i$ , contained in the metric tensor field of an Einstein space-time, is said to be *essential* if and only if it can not be eliminated by a GCT. Every constant which is not essential is said to be *spurious*.

This definition stands to reason, independently of the above mentioned geometrical considerations, and is thus rather commonly accepted<sup>3</sup>.

---

<sup>3</sup>Evidently, *essential* and *spurious* are complementary notions.

Based on this definition, one can deduce a conclusive criterion which will help one to find —if any— the spurious constants among the totality of the constants contained in a metric tensor field.

Let the metric tensor field (1.1) and a variation  $\delta\lambda^{(j)}$  of the  $j$ -th constant  $\lambda^{(j)}$ , while the rest of them are kept fixed. It is:

$$\begin{aligned}\delta g_{\alpha\beta}(x^\mu; \lambda^{(i)}) \Big|_{\lambda^{(i \neq j)} = \text{const.}} &\equiv g_{\alpha\beta}(x^\mu; \lambda^{(j)} + \delta\lambda^{(j)}) - g_{\alpha\beta}(x^\mu; \lambda^{(j)}) \\ &= \frac{\partial g_{\alpha\beta}(x^\mu; \lambda^{(i)})}{\partial \lambda^{(j)}} \Big|_{\lambda^{(i \neq j)} = \text{const.}} \delta\lambda^{(j)}\end{aligned}\quad (2.1)$$

[no summation condition]

up to the first order and when  $\delta\lambda^{(j)} \rightarrow 0$ . Obviously such a variation does change the form of the metric tensor field.

According to the definition (2.1), the varied constant  $\lambda^{(j)}$  is spurious if and only if it is induced by a GCT —since a spurious constant can be not only eliminated but also introduced by a GCT.

So, if one demands for this alteration to be induced by a GCT, then there must exist an infinitesimal GCT, which depends on the varied constant  $\lambda^{(j)}$  (seen as a parameter) and is described in terms of a vector field  $\mathbf{\Gamma}$  (where  $\Gamma^\alpha$  are the components of the field  $\mathbf{\Gamma}$  ( $\mathbf{\Gamma} = \Gamma^\alpha \frac{\partial}{\partial x^\alpha}$ ) in a coordinate system):

$$x^\mu \rightarrow \tilde{x}^\mu : \tilde{x}^\mu = x^\mu - \delta\lambda^{(j)} \Gamma^\mu \quad (2.2)$$

such that, the variation of the metric tensor field to be:

$$\delta g_{\alpha\beta}(x^\mu) \equiv \tilde{g}_{\alpha\beta}(\tilde{x}^\mu) - g_{\alpha\beta}(\tilde{x}^\mu) = \delta\lambda^{(j)} (\mathcal{L}_{\mathbf{\Gamma}} g)_{\alpha\beta} \quad (2.3)$$

again, up to the first order and when  $\delta\lambda^{(j)} \rightarrow 0$ .

These lead to the following:

**Criterion 2.1.** A constant  $\lambda^{(i)}$ , for a given  $i$ , contained in the metric tensor field of an Einstein space-time, is *spurious* if and only if there exists a vector field  $\mathbf{\Gamma}$ , such that the Lie derivative  $\mathcal{L}_{\mathbf{\Gamma}} g$  of the metric tensor field with respect to it equals the variation  $\frac{\delta g}{\delta \lambda^{(i)}}$  of the metric tensor field with respect to the constant under discussion:

$$\mathcal{L}_{\mathbf{\Gamma}} g = \frac{\delta g}{\delta \lambda^{(i)}} \quad (2.4)$$

This is a tensorial equation and in a local coordinate system, condition (2.4) assumes the form:

$$\Gamma_{\alpha;\beta} + \Gamma_{\beta;\alpha} = \frac{\partial g_{\alpha\beta}(x^\mu; \lambda^{(i)})}{\partial \lambda^{(j)}} \Big|_{\lambda^{(i \neq j)} = \text{const.}} \quad (2.5)$$

The semi column symbol (;) denotes covariant differentiation with respect to the metric tensor field (1.1). Existence of a solution (even of a partial one) implies that the constant under discussion is spurious, since the integral curves of the corresponding vector field  $\Gamma$  can serve as that particular GCT, which eliminates/introduces the constant (see interchange between the active and the passive character of a transformation and the example of the integral curves of a Killing field). On the other hand inconsistency of the system (2.5) is tantamount to the statement that the constant under consideration is essential. Each time, only one constant  $\lambda^{(i)}$  can be checked with this algorithm, but the entire method, based on partial differential equations of the first order, is conclusive and has no exceptions at all. Substraction of the spurious constants from the totality of the constants contained in (1.1) results in the totality of the essential constants contained in this metric tensor field. It is clear that, in the above analysis, the dimension of the manifold  $\mathcal{M}$ , the signature of the metric tensor field as well as its “nature” (i.e., whether it is a solution to the EFE’s or not) do not play any role. So, the criterion (2.1) holds for any metric tensor field on any manifold, but it is of importance for the case of the Einstein space-times.

An immediate application:

Consider a metric tensor field, containing one constant, of the form:

$$g_{\alpha\beta}(x^\mu; \lambda^{(1)}) = \lambda^{(1)} \gamma_{\alpha\beta}(x^\mu) \quad (2.6)$$

where the metric tensor field  $\gamma$  contains no constants. Then, since:

$$\frac{\partial g_{\alpha\beta}(x^\mu; \lambda^{(1)})}{\partial \lambda^{(1)}} = \gamma_{\alpha\beta}(x^\mu) \quad (2.7)$$

application of the condition (2.5) of the criterion (2.1) results in:

$$\Gamma_{\alpha;\beta} + \Gamma_{\beta;\alpha} = \frac{1}{\lambda^{(1)}} g_{\alpha\beta}(x^\mu; \lambda^{(1)}) \quad (2.8)$$

and the constant  $\lambda^{(1)}$  is spurious (i.e., absorbable) if and only if the metric tensor field  $g$  has an homothety; a classical result is thus regained [2].

Another classical example concludes this section: Consider the 2-sphere:

$$g_{\alpha\beta}(\theta, \phi; \lambda^{(1)}) = (\lambda^{(1)})^2 \begin{pmatrix} 1 & 0 \\ 0 & \sin^2(\theta) \end{pmatrix} \quad (2.9)$$

in a local coordinate system  $\{\theta, \phi\}$ . This time, application of the condition (2.5) results in an inconsistent system (i.e., the 2-sphere has no homothety), so the parameter  $\lambda^{(1)}$ , which here stands for the radius, is essential –as it should.

Although the previous examples concern cases where the constant under discussion is an overall factor, criterion (2.1) is applicable to any metric tensor field of the general form (1.1) –as one can check.

### 3 Discussion

In the present work a twofold problem, that of defining and pinpointing the essential constants contained in a given metric tensor field of an Einstein space-time (or of a generic metric tensor field on any manifold), is studied.

With the help of some geometrical notions (i.e., description of the inner properties of a geometry in terms of its curvature invariant relations) the wanted definition is hinted. The proposed definition is simple and logical, and it seems to have its own “*raison d’être*”, independently of the initial geometrical considerations.

Using this very definition as a basis, a conclusive criterion which serves to identify the spurious constants, and consequently the complementary notion of the essential constants, is established.

The problem studied in the present work is the “dual” of another problem, which can take the form of the following question:

*For a given Einstein space-time, how many essential constants is expected to be contained in the metric tensor field, which is a solution to the EFE’s?*

No general answer to this question exists so far; only partial step towards the solution to this problem have been made. None the less, the case of the spatially homogeneous Einstein space-times (these are used to define the so called “Spatially Homogeneous Cosmological Models” see e.g., [2]) has been treated and a concrete, general answer to this question is given to the literature [3] (and the references therein). Indeed, in that case there exists an algorithm, which, taking into account the simply transitively action of an Isometry Group on the Cauchy hypersurface, gives the number of the essential constants expected to be contained in a general solution to the EFE’s [3]. On the other hand, the merits of the above analysis are the generality of the application, since there are no exceptions at all (holds for any metric tensor field on any manifold), and the extreme simplicity of the proposed method (which is based on partial differential equations of the first order).

#### Acknowledgements

This work was supported by the National & Kapodistrian University of Athens, Special Account for the Research Grant-No. 70/4/5000. The author is indebted to Associate Professor T. Chistodoulakis for his encouragement, and for his helpful comments, which contributed to a clearer exposition of the essence of this work.

## References

- [1] Y. Bruhat, “*The Cauchy Problem*”, in *Gravitation: An Introduction to Current Research*, ed. by L. Witten, John Wiley & Sons, Inc., N.Y. (1962);  
R.M. Wald, *General Relativity*, Chicago Univ. Press, Chicago (1984);  
S. Weinberg *Gravitation & Cosmology*, John Wiley & Sons, Inc., N.Y. (1972)
- [2] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Hertl, *Exact Solutions of Einstein’s Field Equations*, 2nd Edition, Cambridge University Press, (2003)
- [3] T. Christodoulakis, G. Kofinas, E. Korfiatis, G.O. Papadopoulos and A. Paschos, *Time-dependent automorphism inducing diffeomorphisms in vacuum Bianchi cosmologies and the complete closed form solutions for types II and V*, J. Math. Phys. **42** (2001), no. 8, 3580–3608;  
T. Christodoulakis, S. Hervik and G.O. Papadopoulos, *Essential constants for spatially homogeneous Ricci-flat manifolds of dimension  $4+1$* , J. Phys. A **37** (2004), no. 13, 4039–4058