Can the Acceleration of Our Universe Be Explained by the Effects of Inhomogeneities?

Akihiro Ishibashi[†] and Robert M. Wald^{†‡}

Enrico Fermi Institute[†] and Department of Physics[‡] The University of Chicago, Chicago, IL 60637, USA

November 26, 2024

Abstract

No. It is simply not plausible that cosmic acceleration could arise within the context of general relativity from a back-reaction effect of inhomogeneities in our universe, without the presence of a cosmological constant or "dark energy." We point out that our universe appears to be described very accurately on all scales by a Newtonianly perturbed FLRW metric. (This assertion is entirely consistent with the fact that we commonly encounter $\delta \rho / \rho > 10^{30}$.) If the universe is accurately described by a Newtonianly perturbed FLRW metric, then the back-reaction of inhomogeneities on the dynamics of the universe is negligible. If not, then it is the burden of an alternative model to account for the observed properties of our universe. We emphasize with concrete examples that it is not adequate to attempt to justify a model by merely showing that some spatially averaged quantities behave the same way as in FLRW models with acceleration. A quantity representing the "scale factor" may "accelerate" without there being any physically observable consequences of this acceleration. It also is *not* adequate to calculate the second-order stress energy tensor and show that it has a form similar to that of a cosmological constant of the appropriate magnitude. The second-order stress energy tensor is gauge dependent, and if it were large, contributions of higher perturbative order could not be neglected. We attempt to clear up the apparent confusion between the second-order stress energy tensor arising in perturbation theory and the "effective stress energy tensor" arising in the "shortwave approximation."

1 Introduction

The apparent acceleration of our universe is one of the most striking cosmological observations of recent times. In the context of Friedmann-Lemaitre-Robertson-Walker (FLRW) models in general relativity, the acceleration of our universe would require either the presence of a cosmological constant or a new form of matter ("dark energy") with large negative pressure. However, since there does not appear to be any natural explanation for the presence of a cosmological constant of the necessary size, nor does there appear to be any natural candidate for dark energy, it is tempting to look for alternative explanations. In recent years, there have been at least two approaches that have attempted to account for the observed acceleration of our universe within the framework of general relativity as being a consequence of deviations of our universe from exact FLRW symmetry, without invoking the presence of a cosmological constant or dark energy.

One approach notes that the mass density of our universe is, in fact, extremely inhomogeneous on scales much smaller than the Hubble radius. In order to get an effective homogeneous, isotropic universe, one needs to average and/or smooth out the inhomogeneities on some appropriate choice of spatial slicing. In such an "averaged" (or effective) FLRW universe, one can then define "effective cosmological parameters" [1, 2, 3]. One then finds that the equations of motion for these effective cosmological parameters differ, in general, from the equations satisfied by these parameters in FLRW models. If they differ in a way that corresponds to adding a cosmological constant (or dark energy) of the right magnitude, then one may hope to have explained the acceleration of our universe in the context of general relativity, without invoking the presence of a cosmological constant or dark energy [4, 5, 6, 7, 8, 9, 10].

A second approach has attempted to account for the acceleration of our universe as a back-reaction effect of long-wavelength cosmological perturbations. Here one constructs an "effective energy-momentum tensor" for these perturbations from second-order perturbation theory, and adds this as a source term in Einstein's equation [11, 12, 13, 14]. If this effective energy-momentum tensor has a form similar to that of a cosmological constant (or, at least, provides a negative pressure term) and is of the appropriate magnitude, then one may hope that this could explain the acceleration of our universe.

Given that our universe appears to be very accurately described by a FLRW model, with very small deviations from homogeneity and isotropy, it would seem extremely implausible that cosmologically important effects could result from the second (or higher) order corrections produced by these small departures from a FLRW model. The main purposes of this paper are to explain this point with somewhat more precision and to point out some significant flaws in the arguments that have been made in the context of the above two approaches. In particular, we emphasize that one cannot justify a model by merely showing that spatially averaged quantities behave the same way in FLRW models with acceleration; rather one must show that all of the predictions of the model are compatible with observations. We illustrate this point by showing that spatial averaging can yield "acceleration" for the case of a universe that consists of two disconnected components, each of which is decelerating! We also show that spatial averaging can yield acceleration for suitably chosen slices of Minkowski spacetime. With regard to the back-reaction effects of long-wavelength perturbations, we note that the effective energy-momentum tensor is highly gauge dependent, as has previously been pointed out by Unruh [15]. Even in the long-wavelength limit, we show that one can get essentially any answer one wishes for the effective energymomentum tensor, even though one cannot get any new physical phenomena beyond those already present in FLRW models. We also emphasize the difference between the use of an effective energy-momentum tensor for gravitational perturbations in the context of second-order perturbation theory and the use of a similar effective energy-momentum tensor in the context of the "shortwave approximation" [16, 17]. The former is highly gauge dependent (and, thus, not easily interpreted) and must be "small" if higher order perturbative corrections are to be neglected. The latter is essentially gauge independent and need not be "small." However, the "shortwave approximation" clearly is not valid for analyzing long-wavelength cosmological perturbations.

In the next section, we point out that our universe appears to be very accurately described on all scales by a Newtonianly perturbed FLRW metric, despite the presence of large density contrasts. If this is correct, then higher order corrections to this metric resulting from inhomogeneities would be negligible. Thus, any model that attempts to explain the acceleration of our universe as a consequence of higher order effects of inhomogeneities will have to overcome the seemingly impossible burden of explaining why the universe appears to be so well described by a model that has only very small departures from a FLRW metric.¹ In section 3, we illustrate that spatial averaging can produce an entirely spurious "acceleration" that is not associated with any physical observations. In section 4, we emphasize the distinction between second-order perturbation theory and the shortwave approximation, and we analyze the gauge dependence of the effective energy-momentum tensor arising in second-order perturbation theory.

2 The Newtonianly Perturbed FLRW Metric

By a Newtonianly perturbed FLRW metric, we mean a metric of the form

$$d\tilde{s}^{2} = -(1+2\Psi)dt^{2} + a^{2}(t)(1-2\Psi)\gamma_{ij}dx^{i}dx^{j}.$$
 (1)

¹ For example, rotation of the cosmic matter may produce acceleration effects [5] but these acceleration effects must be negligible in view of the observed isotropy of our universe [18, 19, 20].

where γ_{ij} denotes the metric of a space of constant curvature (3-sphere, flat, or hyperboloid), and Ψ satisfies

$$|\Psi| \ll 1, \quad \left|\frac{\partial\Psi}{\partial t}\right|^2 \ll \frac{1}{a^2} D^i \Psi D_i \Psi, \quad (D^i \Psi D_i \Psi)^2 \ll (D^i D^j \Psi) D_i D_j \Psi, \qquad (2)$$

where D_i denotes the derivative operator associated with γ_{ij} . Suppose that the stress-energy content of this spacetime consists of fluid components that are very nearly homogeneously and isotropically distributed but may have arbitrary equation of state (such as radiation, dark energy, and/or a cosmological constant) together with components that may be very inhomogeneously distributed on scales small compared with the Hubble radius but are nearly pressureless and move with velocity much smaller than light relative to the Hubble flow (such as ordinary matter and dark matter). The stress-energy of the smoothly distributed components take the form

$$T_{ab}^{(s)} \approx \rho^{(s)}(t)dt^2 + P^{(s)}(t)a^2(t)\gamma_{ij}dx^i dx^j , \qquad (3)$$

where $P^{(s)} = P^{(s)}(\rho^{(s)})$ is arbitrary, whereas the inhomogeneously distributed components have a stress-energy of the "dust" form

$$T_{ab}^{(m)} \approx \rho^{(m)}(t, x^i) dt^2 \,. \tag{4}$$

If one plugs the metric form eq. (1) into Einstein's equation and uses eqs. (2), the spatial average yields the usual FLRW equations for the scale factor a with stress-energy source consisting of the sum of eq. (3) and the spatial average of eq. (4),

$$3\left(\frac{\dot{a}}{a}\right)^2 = \kappa^2 \left(\rho^{(s)} + \bar{\rho}^{(m)}\right) - 3\frac{K}{a^2}, \qquad (5)$$

$$3\frac{\ddot{a}}{a} = -\frac{\kappa^2}{2} \left(\rho^{(s)} + \bar{\rho}^{(m)} + 3P^{(s)} \right) , \qquad (6)$$

where $\bar{\rho}$ denotes the spatial average of ρ (taken on a t = const. time slice with respect to the underlying FLRW metric with $K = \pm 1, 0$). The dominant remaining terms in Einstein's equation then yield

$$\frac{1}{a^2}\Delta_{(3)}\Psi = \frac{\kappa^2}{2}\delta\rho\,,\tag{7}$$

where $\Delta_{(3)} \equiv \gamma^{ij} D_i D_j$, and where $\delta \rho = \rho^{(m)} - \bar{\rho}^{(m)}$ denotes the deviation of the density from the spatial average. In the following we assume the metric γ_{ij} to be a flat spatial metric, i.e., K = 0.

It should be emphasized that the above discussion does *not* constitute a *derivation* that a metric form, eq. (1), is a good approximation to a solution to Einstein's equation for stress-energy eq. (3) and (4) when eqs (5)–(7) hold. Rather, all that has been shown is that if one *assumes* a metric of the form eq. (1) with eqs. (2)holding and with the stress-energy given by eqs. (3) and (4), then eqs. (5), (6)and (7) must hold. The above discussion is thus analogous to the usual textbook "derivations" of the ordinary Newtonian limit of general relativity, where one also postulates a spacetime metric of a suitable "Newtonian form" and assumes that the matter distribution is approximately of the form (4) (see, e.g., section 4.4a of [21]). One can derive the ordinary Newtonian limit more systematically by considering one-parameter families of solutions to Einstein's equation with suitable limiting properties (see, e.g., [22] and [23] and references cited therein). It would be more difficult to provide an analogous analysis here, but we see no reason to doubt that eq. (1) is a good approximation to a solution to Einstein's equation when eqs (5)-(7) hold, provided, of course, that conditions (2) are satisfied.

We now assert that the metric, eq. (1), appears to very accurately describe our universe on *all scales*, except in the immediate vicinity of black holes and neutron stars. The basis for this assertion is simply that the FLRW metric appears to provide a very accurate description of all phenomena observed on large scales, whereas Newtonian gravity appears to provide an accurate description of all phenomena observed on small scales. The metric (1) together with eqs (2), predicts that large scale phenomena will be accurately described by a FLRW model, whereas the metric (1) together with eq. (7) predicts that Newtonian gravity will hold in regions small compared with the Hubble radius in which the stress-energy (4) dominates over (3) (see [24]).

Note that the validity of eq. (1) for accurately describing phenomena on *small* scales holds despite the fact that the density contrast of matter is commonly quite large [25, 26]

$$\frac{\delta\rho}{\rho} \gg 1$$
. (8)

Indeed, for the solar system, galaxies, clusters of galaxies, we can estimate respectively, $\delta \rho / \rho \approx 10^{30}$, $\approx 10^5$, $\approx 10^2 \gg 1$. Nevertheless, in all of these cases, we have $\Psi \approx 10^{-6} \sim 10^{-5} \ll 1$, and the other conditions appearing in eq. (2) also hold. Even for neutron stars, $\Psi \approx 10^{-1}$, so the metric eq. (1) is probably not too bad an approximation even in the vicinity of neutron stars.

The key point of this section is that if our assertion is correct that a metric of the form of eq. (1) accurately describes our universe, and if it also is true that conditions (2) hold, then the nonlinear correction terms occurring in eqs. (5) and (6) are negligibly small. It therefore is manifest that nonlinear corrections² to

² We take this opportunity to comment upon one misconception related to the validity of the Newtonianly perturbed FLRW metric. It is commonly stated that when $\delta\rho/\rho \gg 1$, one enters a "nonlinear regime." This might suggest that the validity of eqs. (1) and/or (7) would be questionable whenever $\delta\rho/\rho \gg 1$. However, this is not the case; the proper criteria for the validity of the metric eq. (1) are conditions (2), not $\delta\rho/\rho \ll 1$. It is true that nonlinear effects become important for the motion of matter when $\delta\rho/\rho \gg 1$. This follows simply from the fact

the dynamics of the universe will be negligible, i.e., there will be no important "back-reaction" effects of the inhomogeneities on the observed expansion of the universe on large scales. In particular, accelerated expansion cannot occur if the smoothly distributed matter satisfies the strong energy condition. However, our assertion that the metric, eq. (1), very accurately describes our universe is merely an assertion, and we cannot preclude the possibility that other models (e.g., with large amplitude, long-wavelength gravitational waves or with matter density inhomogeneities of a different type) might also fit observations. Our main point of this paper, however, is that if one wishes to propose an alternative model, then it is necessary to show that all of the predictions of this model are compatible with observations such as the observed redshift-luminosity relation for type Ia supernovae and the various observed properties of the cosmological microwave background (CMB) radiation. As we shall illustrate in the next two sections, it does not suffice to show merely that the spatially averaged scale factor behaves in a desired way or that an effective stress-energy tensor is of a desired form.

3 Cosmic Acceleration via Averaging

The type of spatial averaging in the context of Newtonianly perturbed FLRW models that was done to derive eqs. (5) and (6) above is not problematical. The metric very nearly has FLRW symmetry, so there is a natural choice of spatial slices on which one can take spatial averages. Since $\Psi \ll 1$, it makes negligible difference if one uses the spacetime metric (1) or the corresponding "background" FLRW metric (i.e., eq. (1) with Ψ set equal to zero) to define the averaging.

If one has a metric that does not nearly have FLRW symmetry, one can, of course, still define spatial averaging procedures. However, these will, in general, be highly dependent on the choice of spatial slicing, and the results obtained from spatial averaging need not be interpretable in a straightforward manner. We now illustrate these comments with concrete examples.

For simplicity and definiteness, we consider an inhomogeneous universe with

that self-gravitation is a nonlinear effect, and self-gravitational effects on the motion of matter cannot be ignored when $\delta \rho / \rho > 1$. But this does not mean that one must include nonlinear corrections to the metric form, eq. (1), or to eq. (7) in order to get a good approximation to the spacetime metric. Indeed, if one is trying to describe the solar system in the context of the ordinary Newtonian limit of general relativity, one must include "nonlinear effects" to obtain the correct motion of the planets; they would move on geodesics of the flat metric rather than the Newtonianly perturbed metric if not for these "nonlinear effects." However, the corrections to the spacetime metric of the solar system arising from nonlinear terms in Einstein's equation are entirely negligible.

irrotational dust. In the comoving synchronous gauge,³ the metric takes the form

$$ds^{2} = -dt^{2} + q_{ij}(t, x^{m})dx^{i}dx^{j}.$$
(9)

Let Σ denote a hypersurface of constant t, let \mathcal{D} denote a compact region of Σ and let ϕ be a scalar field on Σ . The average, $\langle \phi \rangle_{\mathcal{D}}$, of ϕ over the domain $\mathcal{D} \subset \Sigma$ may be defined by

$$\langle \phi \rangle_{\mathcal{D}} \equiv \frac{1}{V_{\mathcal{D}}} \int_{\mathcal{D}} \phi d\Sigma \,, \tag{10}$$

where $V_{\mathcal{D}}$ denotes the volume of \mathcal{D} and $d\Sigma$ is the proper volume element of Σ .⁴ Define the averaged scale factor, $a_{\mathcal{D}}$, by

$$a_{\mathcal{D}} \equiv (V_{\mathcal{D}})^{1/3} \,. \tag{11}$$

We "time evolve" \mathcal{D} by making it be comoving with the dust, i.e., the (comoving) coordinates of the boundary of \mathcal{D} remain constant with time. Following Buchert [1], one then obtains from Einstein's equation the following equations of motion for $a_{\mathcal{D}}$,

$$3\frac{\ddot{a}_{\mathcal{D}}}{a_{\mathcal{D}}} = -\frac{\kappa^2}{2} \langle \rho \rangle_{\mathcal{D}} + Q_{\mathcal{D}} , \qquad (12)$$

$$3\left(\frac{\dot{a}_{\mathcal{D}}}{a_{\mathcal{D}}}\right)^2 = \kappa^2 \langle \rho \rangle_{\mathcal{D}} - \frac{1}{2} \langle \mathcal{R} \rangle_{\mathcal{D}} - \frac{1}{2} Q_{\mathcal{D}} , \qquad (13)$$

together with

$$\left(a_{\mathcal{D}}^{6}Q_{\mathcal{D}}\right)^{\cdot} + a_{\mathcal{D}}^{4}\left(a_{\mathcal{D}}^{2}\langle\mathcal{R}\rangle_{\mathcal{D}}\right)^{\cdot} = 0.$$
(14)

Here \mathcal{R} is the scalar curvature of Σ and

$$Q_{\mathcal{D}} \equiv \frac{2}{3} \left(\langle \theta^2 \rangle_{\mathcal{D}} - \langle \theta \rangle_{\mathcal{D}}^2 \right) - \langle \sigma_{ij} \sigma^{ij} \rangle_{\mathcal{D}} , \qquad (15)$$

In deriving these equations, it is important to bear in mind that the averaging of the time derivative $\langle \dot{\phi} \rangle_{\mathcal{D}} \equiv \langle \partial \phi / \partial t \rangle_{\mathcal{D}}$ of a locally defined quantity ϕ differs in general from the time derivative of the averaged quantity $\langle \phi \rangle_{\mathcal{D}} \equiv \partial \langle \phi \rangle_{\mathcal{D}} / \partial t$, since the volume element and $V_{\mathcal{D}}$ may depend on the time t. Indeed, we have

$$\langle \phi \rangle^{\cdot}{}_{\mathcal{D}} = \langle \dot{\phi} \rangle_{\mathcal{D}} + \langle \theta \phi \rangle_{\mathcal{D}} - \langle \theta \rangle_{\mathcal{D}} \langle \phi \rangle_{\mathcal{D}} , \qquad (16)$$

³ If the universe is filled with irrotational dust, then the comoving synchronous gauge defines a natural choice of slicing, namely the slices orthogonal to the world lines of the dust. However, for an inhomogeneous universe, this gauge choice typically will break down on timescales much shorter than cosmological timescales, due to formation of caustics. For example, synchronous coordinates defined in a neighborhood of the Earth would typically break down on a timescale of order the free fall time to the center of the Earth, i.e., ~ 1 hour.

⁴ For a different type of averaging procedure than that given by eq. (10) and its application to cosmology, see e.g., [27, 28].

where θ denotes the expansion of the world lines of the dust.

It is immediately seen from (12) that "averaged acceleration" $\ddot{a}_{\mathcal{D}} > 0$ is achieved if

$$Q_{\mathcal{D}} > \frac{\kappa^2}{2} \langle \rho \rangle_{\mathcal{D}} \,. \tag{17}$$

Buchert [29] has discussed cosmological implications of the condition (17).

A number of authors (see, e.g., Refs. [6, 9, 10]) have sought to account for the observed acceleration of our universe by means of inhomogeneous models that satisfy eq. (17). However, our main point of this section is that even if our universe (or a suitable spatial domain of our universe) satisfies eq. (17) and thus has $\ddot{a}_{\mathcal{D}} > 0$, this does *not* imply that the model will possess any physically observable attributes of an accelerating FLRW model. Indeed Nambu and Tanimoto [9] have shown that in a cosmological model in which \mathcal{D} can be written as a union of regions each of which is locally homogeneous and isotropic, we have

$$a_{\mathcal{D}}^{2}\ddot{a}_{\mathcal{D}} = a_{1}^{2}\ddot{a}_{1} + a_{2}^{2}\ddot{a}_{2} + \dots + \frac{2}{a_{\mathcal{D}}^{3}}\sum_{i\neq j}a_{i}^{3}a_{j}^{3}\left(\frac{\dot{a}_{i}}{a_{i}} - \frac{\dot{a}_{j}}{a_{j}}\right)^{2}, \qquad (18)$$

where a_i denotes the locally defined scale factor in the *i*-th patch, and, in this case, $a_{\mathcal{D}} \equiv (a_1^3 + a_2^3 + \cdots)^{1/3}$. Consider, now, a model where at time *t* the universe consists of two disconnected(!) dust filled FLRW models, one of which is expanding and the other of which is contracting. Both components of the universe are, of course, decelerating, i.e., $\ddot{a}_1 < 0$, $\ddot{a}_2 < 0$. Nevertheless, it is not difficult to see from eq. (18) that $\ddot{a}_{\mathcal{D}} > 0$ can easily be satisfied. For example, if we take $a \equiv a_1 = a_2$ and $\dot{a}_1 = -\dot{a}_2$ we obtain

$$a_{\mathcal{D}}^2 \ddot{a}_{\mathcal{D}} = 2a^3 \left\{ \frac{\ddot{a}}{a} + 4\left(\frac{\dot{a}}{a}\right)^2 \right\} = \frac{7}{3}\kappa^2 a^3 \rho > 0.$$
⁽¹⁹⁾

We thereby obtain a very simple model where the universe is accelerating according to the definition eq. (17), but all observers see only deceleration. This graphically illustrates that satisfaction of eq. (17) in a model is far from sufficient to account for the physically observed effects of acceleration in our universe. We see no reason to believe that the spatially averaged acceleration found, e.g., in the models of [9, 10] directly corresponds to any physical effects of acceleration such as would be observed in type Ia supernovae data. The only way to tell if a model displays physically observable effects of acceleration is to calculate these effects.

As already mentioned above, the averaging procedure defined by eq. (10) also has ambiguities both with regard to the choice of time slicing and the choice of domain \mathcal{D} . One may also artificially produce an averaged cosmic acceleration as a result of a suitably chosen time-slicing. To show this explicitly, we give an example of accelerated expansion in Minkowski spacetime. We note first that for a general inhomogeneous universe (i.e., with no assumption concerning the form of the stress-energy), the equation of motion for $a_{\mathcal{D}}$ can be expressed as

$$3\frac{\ddot{a}_{\mathcal{D}}}{a_{\mathcal{D}}} = -\langle \mathcal{R} \rangle_{\mathcal{D}} - 6\left(\frac{\dot{a}_{\mathcal{D}}}{a_{\mathcal{D}}}\right)^2 + \langle (G_{ab} + \frac{1}{2}g_{ab}G^c{}_c)t^at^b \rangle_{\mathcal{D}}, \qquad (20)$$

where G_{ab} denotes the Einstein tensor. Therefore, for any vacuum spacetime, if there is a domain \mathcal{D} such that $-\langle \mathcal{R} \rangle_{\mathcal{D}} > 6 \left(\dot{a}_{\mathcal{D}} / a_{\mathcal{D}} \right)^2$, then \mathcal{D} describes an accelerated expansion insofar as $a_{\mathcal{D}}$ is concerned.

To construct an accelerating \mathcal{D} in Minkowski spacetime, we start with two hyperboloidal slices, one of which corresponds to an expanding time-slice in the Milne chart covering the future of the origin

$$ds^{2} = -da^{2} + a^{2}(d\xi^{2} + \sinh^{2}\xi d\Omega^{2}), \qquad (21)$$

and the other of which is a similar hyperboloidal slice that is contracting. The idea of the construction is to join these two hyperboloids at some radius, smooth out the join region, and choose \mathcal{D} so that $\dot{a}_{\mathcal{D}} = 0$ (see Figure 1). Since the hyperboloids have negative scalar curvature, we thereby have $\mathcal{R} < 0$ except near the radius where the hyperboloids are joined. However, we can show that this construction can be done so that the contribution from the join region can be made arbitrarily small. Consequently, we obtain $3\ddot{a}_{\mathcal{D}}/a_{\mathcal{D}} = -\langle \mathcal{R} \rangle_{\mathcal{D}} > 0$. Details of this construction are given in the appendix. Since Minkowski spacetime does not display any physical effects associated with accelerated expansion, this example shows quite graphically that "acceleration" as defined by the above averaging procedure can easily arise as a gauge artifact produced by a suitable choice of time slicing.

4 Cosmological Back-reaction in the Long-wavelength Limit

A number of authors have considered the back-reaction effects of perturbations of FLRW models, particularly with regard to modes whose wavelength is comparable to or larger than the Hubble radius [11, 12, 13, 14]. The basic strategy has been to compute the second order terms in Einstein's equation arising from these perturbations, thereby obtaining an "effective stress-energy tensor" for the perturbations. If the form of this effective stress-energy tensor corresponds to that of a positive cosmological constant of the correct magnitude, then one might hope to have provided a mechanism for obtaining the observed acceleration of our universe as a back-reaction effect of long-wavelength perturbations, without the need to introduce a cosmological constant or dark energy.⁵

 $^{^{5}}$ The motivation in [11, 12] was actually to use back-reaction effects to attempt to cancel the presence of a large cosmological constant rather than to use back-reaction to directly produce acceleration.



Figure 1: An accelerating domain \mathcal{D} (thick line) in Minkowski spacetime (t, r) with the angular coordinates suppressed. The domain \mathcal{D} is constructed by cutting portions out of the two hyperboloids (thin dashed lines), $t = \sqrt{a^2 + r^2}$ and $t = 2\sqrt{a^2 + r_0^2} - \sqrt{a^2 + r^2}$, and joining them at $r = r_0$. The matching can be done in a smooth manner, as explained in Appendix. The boundary radius $r = r_{\mathcal{D}}$ is chosen so that $\dot{a}_{\mathcal{D}} = 0$.

We comment, first, that, even without extensive analysis, there is an intrinsic implausibility to this type of explanation. If one considers perturbations of wavelength less than the Hubble radius, it is hard to imagine that the perturbations could be so small that we do not notice any significant deviations from homogeneity and isotropy, yet so large that their *second order* effects produce very significant changes to the dynamics of our universe. On the other hand, if one goes to the long-wavelength limit, then the perturbation should correspond closely to a perturbation to a spatially homogeneous cosmological model. But, given the severe constraints on anisotropy arising from CMB observations, the perturbation should, in fact, correspond to a perturbation towards another FLRW model. Thus, one should not be able to obtain any new phenomena (such as acceleration without a cosmological constant or dark energy) that are not already present in FLRW models.

At least part of the confusion with regard to the calculation of the backreaction effects of cosmological perturbations appears to stem from the fact that the notion of an "effective stress-energy tensor" for perturbations arises in two quite different contexts, namely (i) ordinary perturbation theory and (ii) the "shortwave approximation." We now explain this distinction. For simplicity, we restrict consideration in the following discussion to the vacuum case; cosmological perturbations for the Einstein-scalar-field system will be considered later in this section.

Ordinary perturbation theory (see, e.g., section 7.5 of [21]) arises by considering a one-parameter family of metrics $g_{ab}(\alpha)$ that is jointly analytic in its dependence on α and the spacetime point. We refer to $g_{ab}^{(0)} \equiv g_{ab}(0)$ as the "background metric." Roughly speaking, as $\alpha \to 0$, $g_{ab}(\alpha)$ differs from $g_{ab}^{(0)}$ by a perturbation that becomes of arbitrarily small amplitude but maintains a fixed profile. One expands $g_{ab}(\alpha)$ in a power series in α about $\alpha = 0$

$$g_{ab}(\alpha) = \sum_{n} \frac{1}{n!} \alpha^n g_{ab}^{(n)} \,. \tag{22}$$

The perturbation equations for $g_{ab}^{(n)}$ for the vacuum Einstein equation,

$$G_{ab} = 0, (23)$$

are then obtained by differentiating the Einstein tensor, $G_{ab}(\alpha)$, of $g_{ab}(\alpha)$ n times with respect to α at $\alpha = 0$. The zeroth order equation is just Einstein's equation for $g_{ab}^{(0)}$

$$G_{ab}[g^{(0)}] = 0. (24)$$

The first order equation is

$$G_{ab}^{(1)}[g^{(1)}] = 0, \qquad (25)$$

where $G_{ab}^{(1)}$ denotes the linearized Einstein tensor off of the background metric $g_{ab}^{(0)}$. The second order equation is

$$G_{ab}^{(1)}[g^{(2)}] = -G_{ab}^{(2)}[g^{(1)}], \qquad (26)$$

where $G_{ab}^{(2)}[g^{(1)}]$ denotes the second-order Einstein tensor constructed from $g_{ab}^{(1)}$.

As can be seen from eq. (26), minus the second-order Einstein tensor (divided with the gravitational constant), $-\kappa^{-2}G_{ab}^{(2)}[g^{(1)}]$, plays the role of an "effective stress-energy tensor" associated with the perturbation $g_{ab}^{(1)}$ in the sense that it acts as a source term for the second-order metric perturbation $g_{ab}^{(2)}$. However, this does not mean that one can treat $-\kappa^{-2}G_{ab}^{(2)}[g^{(1)}]$ as though it were a new form of matter stress-energy that can be inserted into the right side of the exact Einstein equation (23) as opposed to the right side of eq. (26). For one thing, the second-order Einstein tensor is highly gauge dependent (as we shall illustrate explicitly below), so it is not straightforward to interpret its meaning. The key point, however, is that eq. (26) arises only in the context of perturbation theory. If $G_{ab}^{(2)}[g^{(1)}]$ is very small, then its effects on the spacetime metric can be reliably calculated from eq. (26). But if $G_{ab}^{(2)}[g^{(1)}]$ is large enough to produce cosmologically interesting effects (such as acceleration), then the third and higher order contributions to $g_{ab}(\alpha)$ will also be large, and one cannot reliably compute back-reaction effects from second-order perturbation theory.

This situation occurring in perturbation theory contrasts sharply with the situation that arises when one uses the "shortwave approximation" [17, 30, 31]. Here, one wishes to develop a formalism in which the self-gravitating effects of gravitational radiation—and the consequent effects on the spacetime metric on scales much larger than the wavelength of the radiation—can be reliably obtained, even when these effects are "large." Again, one considers a one-parameter family of metrics $g_{ab}(\beta)$ that has a continuous limit to the metric $g_{ab}^{(0)} \equiv g_{ab}(0)$. Thus, as in ordinary perturbation theory, as $\beta \to 0$, $g_{ab}(\beta)$ differs from $g_{ab}^{(0)}$ by a perturbation of arbitrarily small amplitude. However, one now requires $g_{ab}(\beta)$ to be such that, roughly speaking, as $\beta \to 0$, the ratio of the amplitude to the wavelength of the perturbation goes to a finite, non-zero limit; see [31] for a precise statement of what is required in this limit. Thus, in this scheme, the dominant terms in Einstein's equation as $\beta \to 0$ are actually the linear terms in the second derivatives of the first order perturbation, which diverge as $1/\beta$. One thereby obtains

$$G_{ab}^{(1)}[g_{ab}^{(1)}] = 0. (27)$$

The quadratic terms in the first order perturbation are of zeroth order in β , so they make a contribution to the Einstein tensor that is comparable to that of $g_{ab}^{(0)}$. The linear terms in the second-order perturbation also contribute to this order, but these contributions can be eliminated by averaging over a spacetime region that is large compared with the wavelength of the perturbation. One thereby obtains

$$G_{ab}[g^{(0)}] = \langle -G_{ab}^{(2)}[g^{(1)}] \rangle , \qquad (28)$$

where the brackets on the right side of eq. (28) denote a suitably defined spacetime average. It can be shown that $\langle -G_{ab}^{(2)}[g^{(1)}]\rangle$ is gauge invariant in a suitably defined sense. We refer to [31] for further details of the derivation and meaning of these equations.

Although eq. (28) is quite similar in form to eq. (26), the meaning and range of validity of these equations are quite different. In contrast to eq. (26), it should be possible to use eq. (28) to calculate the back-reaction effects of gravitational radiation even when these effects are large. The catch, however, is that eq. (28) can be used only when the wavelength of the perturbation is much smaller than the curvature lengthscale of the background spacetime. Thus, if the universe were filled with gravitational radiation of wavelength much smaller than the Hubble radius, then it should be possible to use eq. (28) to reliably calculate the backreaction effects of this radiation, even if this radiation is the dominant form of "matter" in the universe. However, eq. (28) manifestly *cannot* be used to calculate the back-reaction effects of long-wavelength perturbations.

We conclude this section by deriving an explicit formula for the gauge dependence of the second-order "effective stress-energy tensor" arising in ordinary perturbation theory for long-wavelength scalar-type perturbations of a FLRW universe containing a scalar field. By doing so, we will see that one can get essentially any answer one wishes for this effective stress-energy tensor by making appropriate gauge transformations. This graphically shows that one cannot draw any physical conclusions merely by examining the form of the effective stress-energy tensor arising in second-order perturbation theory. ⁶

Consider a background flat FLRW universe

$$ds^{2} = -dt^{2} + a^{2}(t)\gamma_{ij}dx^{i}dx^{j}, \qquad (29)$$

filled with a scalar field whose energy momentum tensor is given by

$$T_{ab} = \nabla_a \phi \nabla_b \phi - \frac{1}{2} g_{ab} \{ \nabla^c \phi \nabla_c \phi + 2U(\phi) \}.$$
(30)

The unperturbed background equations of motion for a and ϕ , which are functions of only t, are given by

$$H^{2} \equiv \left(\frac{\dot{a}}{a}\right)^{2} = \frac{\kappa^{2}}{3} \left(\frac{1}{2}\dot{\phi}^{2} + U\right), \qquad (31)$$

$$\ddot{\phi} + 3H\dot{\phi} + \frac{\partial U}{\partial \phi} = 0, \qquad (32)$$

where in these equations and hereafter the dot denotes the derivative with respect to t.

We focus on the scalar-type perturbations. The general form of a scalar-type metric perturbation is

$$d\tilde{s}^{2} = -(1+2A\mathbb{S})dt^{2} - 2aB\mathbb{S}_{i}dtdx^{i} + a^{2}\{(1+2H_{L}\mathbb{S})\gamma_{ij} + 2H_{T}\mathbb{S}_{ij}\}dx^{i}dx^{j}, \quad (33)$$

and the scalar field perturbation is given by

$$\tilde{\phi} = \phi + \delta \phi \mathbb{S} \,. \tag{34}$$

Here S denotes a plane wave on flat 3-space with wavevector \mathbf{k} , and \mathbb{S}_i and \mathbb{S}_{ij} are the divergence-free vector and transverse-traceless tensor defined by

$$\mathbb{S}_{i} = -\frac{1}{k} D_{i} \mathbb{S}, \quad \mathbb{S}_{ij} = \frac{1}{k^{2}} \left(D_{i} D_{j} - \frac{1}{3} \gamma_{ij} \Delta_{(3)} \right) \mathbb{S}, \quad (35)$$

with D_i being the derivative operator associated with the 3-space metric γ_{ij} as in (2), and $k^2 = \mathbf{k} \cdot \mathbf{k}$. Here and in the following, perturbation variables are understood as corresponding Fourier expansion coefficients—hence functions merely of *t*—and we omit the index **k** unless otherwise stated.

⁶ Discussion of the characterization of the back-reaction effects of perturbations in terms of physical variables can be found in [32, 33].

Under infinitesimal gauge transformations of the scalar-type;

$$t \to t + T\mathbb{S}, \quad x^i \to x^i + L\mathbb{S}^i,$$
 (36)

the perturbation variables, A, B, H_L , H_T , $\delta\phi$, change as

$$A \rightarrow A - \dot{T},$$
 (37)

$$B \rightarrow B + a\dot{L} + \frac{k}{a}T,$$
 (38)

$$H_L \rightarrow H_L - \frac{k}{3}L - HT$$
, (39)

$$H_T \rightarrow H_T + kL$$
, (40)

$$\delta \phi \rightarrow \delta \phi - \phi T$$
. (41)

In particular, it follows from eqs. (38) and (40) that the following combinations

$$X_T \equiv \frac{a}{k} \left(\frac{a}{k} \dot{H}_T - B \right) , \quad X_L \equiv -\frac{1}{k} H_T , \qquad (42)$$

change as

$$X_T \to X_T - T$$
, $X_L \to X_L - L$. (43)

Hence, by inspection of eqs. (37), (39) and (41), one can immediately obtain gauge-invariant perturbation variables [34, 35];

$$\Psi \equiv A - \dot{X_T}, \quad \Phi \equiv H_L - \frac{k}{3}X_L - HX_T, \quad \Delta\phi \equiv \delta\phi - \dot{\phi}X_T.$$
(44)

Any scalar-type gauge-invariant perturbation quantity can be expressed as a linear combinations of the gauge-invariant variables Ψ , Φ , and $\Delta \phi$, and their time derivatives.

It follows from the linearized Einstein equations that the gauge-invariant variables defined above satisfy the following equations [35]

$$\frac{k^2}{a^2}(\Psi + \Phi) = 0, \qquad k\left[2(\dot{\Phi} - H\Psi) + \kappa^2 \dot{\phi} \Delta \phi\right] = 0.$$
(45)

which correspond, respectively, to the trace-free part of the space-space component and the time-space component of the linearized Einstein equations. For $k^2 \neq 0$, we obtain from eq. (45) the following relations between Φ , Ψ , and $\Delta \phi$;

$$\Phi = -\Psi, \quad \frac{\Delta\phi}{\dot{\phi}} = -\frac{1}{\dot{H}} \left(\dot{\Psi} + H\Psi \right) \,. \tag{46}$$

It then also follows from Einstein equations that Ψ is governed by

$$\ddot{\Psi} + \left(H - \frac{\ddot{H}}{\dot{H}}\right)\dot{\Psi} + \left(2\frac{\dot{H}}{H} - \frac{\ddot{H}}{\dot{H}}\right)H\Psi + \frac{k^2}{a^2}\Psi = 0.$$
(47)

We therefore have found that all of the scalar-type perturbations variables are given in terms of the variables Ψ , X_T , and X_L by

$$A = \Psi + \dot{X}_T, \qquad (48)$$

$$B = -a\dot{X}_L - \frac{k}{a}X_T, \qquad (49)$$

$$H_L = -\Psi + HX_T + \frac{k}{3}X_L, \qquad (50)$$

$$H_T = -kX_L, (51)$$

$$\frac{\delta\phi}{\dot{\phi}} = -\frac{1}{\dot{H}}\left(\dot{\Psi} + H\Psi\right) + X_T.$$
(52)

The variable Ψ is gauge invariant and satisfies eq. (47). On the other hand, the gauge transformation law (43) implies that the functions X_T and X_L are completely arbitrary, i.e., they may be chosen to take any values that one wishes. Thus, the specification of X_T and X_L in terms of Ψ corresponds to fixing the gauge freedom. For example, the choice

$$X_T = X_L = 0 \tag{53}$$

corresponds to the Poisson gauge (or the longitudinal gauge), in which the metric, eq. (33), takes precisely the form of eq. (1). Another example is the choice

$$X_T = -\int_{t_*}^t \Psi(t')dt' + C_1(k), \qquad (54)$$

$$X_L = k \int_{t_*}^t \left\{ \int_{t'_*}^{t'} \Psi(t'') dt'' \right\} \frac{dt'}{a^2(t')} - kC_1(k) \int_{t_*}^t \frac{dt'}{a^2(t')} + C_2(k) , \quad (55)$$

where t_* denotes some reference time and C_1 and C_2 are arbitrary constants. This choice corresponds to the synchronous gauge, A = B = 0, in which C_1 and C_2 parameterize the residual gauge freedom in this gauge.

The second-order effective stress-energy tensor for the Einstein-scalar-field system is defined by

$${}^{(\text{eff})}T_{ab} \equiv -\frac{1}{\kappa^2} G^{(2)}_{ab}[g^{(1)}] + T^{(2)}_{ab}[\delta\phi, g^{(1)}], \qquad (56)$$

where $G_{ab}^{(2)}$ denotes the second order Einstein tensor and $T_{ab}^{(2)}$ is the similarly defined second order contribution to T_{ab} , eq. (30), arising from the first order perturbation $(\delta\phi, g^{(1)})$. We now calculate the second-order effective stress-energy tensor in order to explicitly demonstrate its gauge dependence. It is very convenient to express ^(eff) T_{ab} in terms of the variables Ψ , X_T , and X_L , since any dependence of ^(eff) T_{ab} on X_T , and X_L will explicitly show its gauge dependence. Clearly, since ^(eff) T_{ab} is quadratic in the first order perturbation, it must consist of a part, ${}^{(\text{eff}:\Psi)}T_{ab}$, that is quadratic in Ψ , a part, ${}^{(\text{eff}:X)}T_{ab}$, that is quadratic in (X_T, X_L) , and a part, ${}^{(\text{eff}:\Psi,X)}T_{ab}$, containing the "cross-terms" between Ψ and X_T, X_L . The quantity ${}^{(\text{eff}:\Psi)}T_{ab}$ is gauge invariant, ⁷ but both ${}^{(\text{eff}:X)}T_{ab}$ and ${}^{(\text{eff}:\Psi,X)}T_{ab}$ are gauge dependent. Thus, ${}^{(\text{eff})}T_{ab}$ will be gauge invariant if and only if these latter pieces vanish.

It is easy to verify that both ${}^{(\text{eff}:X)}T_{ab}$ and ${}^{(\text{eff}:\Psi,X)}T_{ab}$ are nonvanishing. To see explicitly that ${}^{(\text{eff}:X)}T_{ab}$ is nonvanishing, it suffices to consider the case where we impose the additional restriction

$$X_L = -k \int_{t_*}^t \frac{X_T(t')}{a^2(t')} dt', \qquad (57)$$

which ensures that B = 0, thereby considerably simplifying the calculation. We also focus attention on the long-wavelength limit. A brute force calculation then yields

$$\kappa^{2 \text{ (eff:}X)}T_{00} = \left[-(\dot{H} + 3H^{2})\dot{X}_{T}^{2} + (\ddot{H} + 6H\dot{H} + 12H^{3})X_{T}\dot{X}_{T} + \left(3\dot{H}^{2} + 12\dot{H}H^{2} + 3H\ddot{H} + \ddot{H} - \frac{3}{4}\frac{\ddot{H}^{2}}{\dot{H}} \right)X_{T}^{2} \right] + O(k^{2})(58)$$

$$\kappa^{2 \text{ (eff:}X)}T_{ij} = g_{ij} \left[2H\dot{X}_{T}\ddot{X}_{T} - 2(\dot{H} + 2H^{2})X\ddot{X} - (\dot{H} + H^{2})\dot{X}_{T}^{2} - (3\ddot{H} + 22H\dot{H} + 12H^{3})X_{T}\dot{X}_{T} - (\ddot{H} + H^{2})\dot{X}_{T}^{2} + \frac{1}{2}\ddot{H} \right)X_{T}^{2} \right] + O(k^{2}). \quad (59)$$

Thus, even in the case of a pure gauge perturbation, $\Psi = 0$, we can obtain a nonvanishing effective stress-energy tensor for long-wavelength perturbations. Indeed, since X_T is entirely arbitrary, we see that we can get essentially any answer one wishes for ${}^{(\text{eff})}T_{ab}$. For example, if one wishes to have a pure gauge perturbation in which ${}^{(\text{eff})}T_{ab}$ takes the form of a cosmological constant, one would merely have to solve the second-order ordinary differential equation for X_T that results when one equates the right side of eq. (58) to minus the right side of eq. (59). This manifestly demonstrates that one cannot derive any physical consequences by merely examining the form of the second-order effective stress-energy tensor.

⁷ In fact, $(\text{eff}:\Psi)T_{ab}$ is precisely the "effective energy-momentum tensor for cosmological perturbations" of [11, 12]. However, contrary to the claims of [11, 12], this effective energymomentum tensor is gauge-invariant only in the trivial sense that any gauge-dependent quantity can be viewed as gauge invariant once a gauge has been completely fixed. In the variations taken in [11, 12] to obtain their effective energy-momentum tensor, X_T and X_L were implicitly assumed to be independent of Ψ , corresponding to the choice of the Poisson (longitudinal) gauge $X_T = X_L = 0$. However, different specifications of X_T and X_L in terms of Ψ —i.e., different choices of gauge—would lead to *different* expressions for the effective energy-momentum tensor in terms of Ψ .

Finally, we comment that we derived the above "long-wavelength limit" form of ${}^{(\text{eff})}T_{ab}$ for scalar-type perturbations by considering perturbations with $k^2 \neq 0$ and then taking the limit as $k^2 \to 0$. Alternatively, we could have directly considered scalar-type perturbations with $k^2 = 0$. It is not immediately obvious that this would give equivalent results, since when $k^2 = 0$, the quantities \mathbb{S}_i and \mathbb{S}_{ij} do not exist, so B and H_T are not defined and the x^i coordinate freedom in eq. (36) does not exist. Furthermore, eqs. (45) become trivial, and therefore the relation, eq. (46) need not hold. Thus, it is not entirely straightforward to make a physical correspondence between perturbations with $k^2 = 0$ and the $k^2 \to 0$ limit of perturbations with $k^2 \neq 0$. Nevertheless, such a one-to-one, onto correspondence does exist⁸, and can be explicitly achieved by using the gauge freedom available when $k^2 \neq 0$ to set $B = H_T = 0$ and using the gauge freedom available when $k^2 = 0$ to set $A = -H_L$. Further discussion of the relationship between perturbations in the long-wavelength limit and exactly homogeneous perturbations can be found in Refs. [36, 37, 38].

Since scalar-type perturbations with $k^2 = 0$ manifestly correspond to perturbations to other FLRW spacetimes, it is clear that one cannot find any new physical phenomena that are not already present in FLRW models by studying long-wavelength perturbations and dropping all terms that are $O(k^2)$. For example, consider a FLRW model which contains two matter components, such as dust and radiation or two scalar fields. In such a model, there exist nontrivial gauge-invariant perturbations even in the $k \to 0$ limit, and implications of such perturbations to the cosmological back-reaction problem have been discussed in [8, 32, 33]. However, in the $k \to 0$ limit such perturbations merely correspond to perturbations to other FLRW models; in the above examples, they would correspond to changing the proportion of dust and radiation or changing the initial conditions of the scalar fields. These perturbations cannot give rise to any new phenomena—such as physically measurable acceleration—that are not already present in exact FLRW models.

5 Summary

In this paper, we have argued that the attempts to explain cosmic acceleration by effects of inhomogeneities, without invoking a cosmological constant or dark energy, are, at best, highly implausible. A Newtonianly perturbed FLRW metric appears to describe our universe very accurately on all scales. In this model, the back-reaction effects of inhomogeneities on the cosmological dynamics are negligible even though the density contrast may be very large on small scales.

We focused much of our attention on exposing the flaws in two types of attempts to explain acceleration by effects of inhomogeneities. (i) Starting from

⁸ It also is worth pointing out that, although the gauge freedom is different, the effective stress-energy tensor for $k^2 = 0$ perturbations remains highly gauge dependent.

an inhomogeneous model, one can obtain an effective FLRW universe by spatial averaging. This effective FLRW universe may display acceleration. However, we showed explicitly via concrete examples that acceleration of the effective FLRW universe may occur in situations where no physically observable effects of acceleration actually occur. (ii) The back-reaction effects of a perturbation of a FLRW universe are described at second order by an effective stress-energy tensor constructed from the first order perturbation. In particular cases, this effective stress-energy tensor may take the form of a cosmological constant, thereby suggesting that it could produce acceleration. However, we pointed out that (unlike the effective stress-energy tensor arising in the shortwave approximation), the effective stress-energy tensor arising in second order perturbation theory is highly gauge dependent and must be small in order to justify neglecting higher order corrections. We explicitly evaluated the second-order effective stress-energy tensor for pure gauge scalar-type perturbations of an Einstein-scalar field model, and showed that it can take essentially any form one wishes, including the form of a cosmological constant.

Acknowledgments

This research was supported by NSF grant PHY 00-90138 to the University of Chicago.

Appendix

Here we provide some details of the construction of the accelerating domain \mathcal{D} in Minkowski spacetime that was described below eq. (20). Let t and r be, respectively, the standard time and radial coordinates in Minkowski spacetime. Let a > 0 and $r_0 > \epsilon > 0$. Let f be a smooth, monotone decreasing function of one variable such that f(x) = 1 for all $x \leq 1/2$, f(x) = 0 for all $x \geq 1$. Define

$$\psi(r) = f\left(\frac{r-r_0}{2\epsilon} + 1\right). \tag{60}$$

Then $\psi(r) = 1$ whenever $r \leq r_0 - \epsilon$ and $\psi(r) = 0$ whenever $r \geq r_0$. Furthermore, there exists a constant C > 0, independent of ϵ , such that $\epsilon |\psi'| < C$ for all $0 < \epsilon < r_0$, where $\psi' \equiv \partial \psi / \partial r$. Define

$$F(r) \equiv \{\psi(r) - \psi(-r + 2r_0)\} \left(\sqrt{r^2 + a^2} - \sqrt{r_0^2 + a^2}\right) + \sqrt{r_0^2 + a^2}.$$
 (61)

Then F is smooth and the hypersurface Σ defined by t = F(r) also is smooth. For $r < r_0 - \epsilon$, Σ is the expanding hyperboloid $t = \sqrt{r^2 + a^2}$, whereas for $r > r_0 + \epsilon$, Σ

is the contracting hyperboloid $t = -\sqrt{r^2 + a^2} + 2\sqrt{r_0^2 + a^2}$. The local expansion rate $H \equiv \dot{a}/a$ of Σ smoothly changes from 1/a to -1/a in the junction interval $(r_0 - \epsilon, r_0 + \epsilon)$. Nowhere does Σ display an accelerated expansion locally.

Now let us take our domain \mathcal{D} to be a ball of radius $r_{\mathcal{D}}$ on Σ , where $r_{\mathcal{D}}$ is chosen so that $\dot{a}_{\mathcal{D}}$ vanishes. It is always possible to find such an $r_{\mathcal{D}}$ since, by construction, $V_{\mathcal{D}}$, hence $\dot{a}_{\mathcal{D}}$, is a smooth function of r, and the local expansion rate $H = \dot{a}/a$ is positive when $r < r_0 - \epsilon$, whereas it is negative when $r_0 + \epsilon < r$. Since the third term of eq. (20) vanishes for Minkowski spacetime, if $\langle \mathcal{R} \rangle_{\mathcal{D}}$ is negative in \mathcal{D} , then eq. (20) shows an acceleration $3\ddot{a}_{\mathcal{D}}/a_{\mathcal{D}} = -\langle \mathcal{R} \rangle_{\mathcal{D}} > 0$. However, apart from the junction region $(r_0 - \epsilon, r_0 + \epsilon)$, Σ is intrinsically a hyperbolic space with a negative scalar curvature $\mathcal{R} = -6/a^2$. Furthermore, the following calculation shows that the contribution to $\langle \mathcal{R} \rangle_{\mathcal{D}}$ from the junction region can be made negligibly small. The induced metric on Σ is

$$ds^{2} = (1 - F'^{2}) dr^{2} + r^{2} d\Omega^{2}, \qquad (62)$$

so the scalar curvature of the junction region is given by

$$\mathcal{R} = -\frac{4rF'F'' + 2(F')^2 \left\{1 - (F')^2\right\}}{r^2 \left\{1 - (F')^2\right\}^2}.$$
(63)

Using the formula,

$$F' = -\frac{r_0}{\sqrt{r_0^2 + a^2}} (\psi + \epsilon \psi') + O(\epsilon) , \qquad (64)$$

which is obtained from the properties of ψ in the junction interval, one finds

$$\int_{r_0-\epsilon}^{r_0+\epsilon} dr r^2 \sqrt{1-(F')^2} \mathcal{R} = \int_{r_0-\epsilon}^{r_0+\epsilon} dr \left\{ 2\frac{2-(F')^2}{\sqrt{1-(F')^2}} - 4\frac{\partial}{\partial r} \left(\frac{r}{\sqrt{1-(F')^2}}\right) \right\} \\
= O(\epsilon),$$
(65)

which can be made arbitrarily small by taking $\epsilon \to 0$. Thus, the contribution to $\langle \mathcal{R} \rangle_{\mathcal{D}}$ from the junction region can indeed be made arbitrarily small, so that $-\langle \mathcal{R} \rangle_{\mathcal{D}} \approx 6/a^2 > 0$. Thus, one obtains an accelerating domain $\mathcal{D} \subset \Sigma$ in Minkowski spacetime through the volume averaging process.

References

- [1] Buchert, T., 2000, Gen. Rel. Grav. **32**, 105.
- [2] Buchert, T., 2001, Gen. Rel. Grav. 33, 1381.
- [3] Buchert, T. and Carfora, M., 2003 Phys. Rev. Lett. **90**, 031101.

- [4] Rasanen, S., 2004, JCAP 0402 003.
- [5] Kolb, E.W., Matarrese, S., Notari, A., and Riotto, A., hep-th/0503117.
- [6] Kolb, E.W., Matarrese, S., and Riotto, A., astro-ph/0506534.
- [7] Barausse, E., Matarrese, S., and Riotto, A., 2005 Phys. Rev. D 71, 063537.
- [8] Nambu, Y., 2005 Phys. Rev. D **71**, 084016.
- [9] Nambu, Y. and Tanimoto, M., gr-qc/0507057.
- [10] Moffat, J.W., astro-ph/0505326.
- [11] Mukhanov, V.F. Abramo, L.R.W., and Brandenberger, R.H., 1997 Phys. Rev. Lett. 78, 1624.
- [12] Abramo, L.R.W., Brandenberger, R.H., and Mukhanov, V.F., 1997 Phys. Rev. D 56, 3246.
- [13] Nambu, Y., 2002 Phys. Rev. D 65, 104013.
- [14] Brandenberger, R.H. and Lam, C.S., hep-th/0407048.
- [15] Unruh, W., 1998 astro-ph/9802323.
- [16] Brill, D. and Hartle, J., 1964 Phys. Rev. **135**, B271.
- [17] Isaacson, R., 1968 Phys. Rev. 166, 1272.
- [18] Flanagan, E.E., 2005 Phys. Rev. D 71, 103521.
- [19] Hirata, C.M. and Seljak, U., 2005 astro-ph/0503582.
- [20] Geshnizjani, G., Chung, D.J.H., and Afshordi, N., 2005 Phys. Rev. D 72, 023517.
- [21] Wald, R.M., 1984 General Relativity, University of Chicago Press: Chicago.
- [22] Futamase, T. and Schutz, B.F., 1983 Phys. Rev. D 28, 2363.
- [23] Ehlers, J., 1997 Class. Quant. Grav. 14, A119.
- [24] Holz, D.E. and Wald, R.M., 1998 Phys. Rev. D 58, 063501.
- [25] Barrow, J.D., 1988 Quart. J. Roy. astr. Soc., **30**, 163.
- [26] Futamase, T., 1989 Mon. Not. R. astr. Soc. 237, 187.
- [27] Zalaletdinov, R.M., 1997 Bull. Astron. Soc. India 25, 401.

- [28] Coley, A.A., Pelavas, N. and Zalaletdinov, R.M., gr-qc/0504115.
- [29] Buchert, T., gr-qc/0507028.
- [30] Misner, C. W., Thorne, K. S. and Wheeler, J.A., 1973 *Gravitation* (Freeman, San Francisco).
- [31] Burnett, G.A., 1989 J. Math. Phys. **30**, 90.
- [32] Abramo, L.R. and Woodard, R.P., 2002 Phys. Rev. D 65, 043507.
- [33] Geshnizjani, G. and Brandenberger, R.H., 2005 JCAP 04, 006.
- [34] Bardeen, J.M., 1980 Phys. Rev. D 22, 1882.
- [35] Kodama, H. and Sasaki, M., 1984 Prog. Theor. Phys. Supple. 78, 1.
- [36] Nambu, Y. and Taruya, A, 1998 Class. Quant. Grav. 15, 2761.
- [37] Kodama, H. and Hamazaki, T., 1998 Phys. Rev. D 57, 7177.
- [38] Sasaki, M. and Tanaka, T., 1998 Prog. Theor. Phys. 99, 763.