

# ON SPACETIMES WITH CONSTANT SCALAR INVARIANTS

ALAN COLEY, SIGBJØRN HERVIK, NICOS PELAVAS

ABSTRACT. We study Lorentzian spacetimes for which all scalar invariants constructed from the Riemann tensor and its covariant derivatives are constant (*CSI* spacetimes). We obtain a number of general results in arbitrary dimensions. We study and construct warped product *CSI* spacetimes and higher-dimensional Kundt *CSI* spacetimes. We show how these spacetimes can be constructed from locally homogeneous spaces and *VSI* spacetimes. The results suggest a number of conjectures. In particular, it is plausible that for *CSI* spacetimes that are not locally homogeneous the Weyl type is *II*, *III*, *N* or *O*, with any boost weight zero components being constant. We then consider the four-dimensional *CSI* spacetimes in more detail. We show that there are severe constraints on these spacetimes, and we argue that it is plausible that they are either locally homogeneous or that the spacetime necessarily belongs to the Kundt class of *CSI* spacetimes, all of which are constructed. The four-dimensional results lend support to the conjectures in higher dimensions.

## 1. INTRODUCTION

Let  $(\mathcal{M}, g)$  denote a differentiable manifold, either of Riemannian or Lorentzian signature. We are interested in all spacetimes for which all scalar invariants constructed from the Riemann tensor and its covariant derivatives are constant (denoted as *CSI* spacetimes). In the case of a Riemannian manifold, if it is *CSI* then it is locally homogeneous [1]. This theorem is false for Lorentzian manifolds (there are a number of counterexamples; e.g., the non-homogeneous *VSI* spacetimes [2]).

There are a number of *CSI* spacetimes that arise from homogeneous spacetimes and simple warped products (see section 3). Some examples of *CSI* spacetimes that arise as simple *tensor sum* generalizations of *VSI* spacetimes with a cosmological constant are given in [3] (some higher dimensional versions are given in [4]). Simple examples of *fibred products* were given in [5] and [6] (see also [7]). It is also straightforward to construct *CSI* spacetimes as *warped products*. All of these spacetimes belong to the class  $CSI_R$  described below.

We begin with some definitions, and a brief discussion of the four-dimensional case.

**1.1. Notation.** Let us denote by  $\mathcal{I}_k$  the set of all scalar invariants constructed from the curvature tensor and its covariant derivatives up to order  $k$ .

**Definition 1.1** (*VSI<sub>k</sub>* spacetimes).  $\mathcal{M}$  is called *VSI<sub>k</sub>* if for any invariant  $I \in \mathcal{I}_k$ ,  $I = 0$  over  $\mathcal{M}$ .

**Definition 1.2** (*CSI<sub>k</sub>* spacetimes).  $\mathcal{M}$  is called *CSI<sub>k</sub>* if for any invariant  $I \in \mathcal{I}_k$ ,  $\partial_\mu I = 0$  over  $\mathcal{M}$ .

Moreover, if a spacetime is *VSI<sub>k</sub>* or *CSI<sub>k</sub>* for all  $k$ , we will simply call the spacetime *VSI* or *CSI*, respectively. The set of all locally homogeneous spacetimes will be denoted by  $H$ . Clearly  $VSI \subset CSI$  and  $H \subset CSI$ .

**Definition 1.3** (*CSI<sub>R</sub>* spacetimes). Let us denote by  $CSI_R$  all reducible *CSI* spacetimes that can be built from *VSI* and  $H$  by (i) warped products (ii) fibred products, and (iii) tensor sums (defined more precisely later).

**Definition 1.4** (*CSI<sub>F</sub>* spacetimes). Let us denote by  $CSI_F$  those spacetimes for which there exists a frame with a null vector  $\ell$  such that all components of the Riemann tensor and its covariants derivatives in this frame have the property that (i) all positive boost weight components (with respect to  $\ell$ ) are zero and (ii) all zero boost weight components are constant.

Note that  $CSI_R \subset CSI$  and  $CSI_F \subset CSI$ . (There are similar definitions for  $CSI_{F,k}$  etc. [8]).

**Definition 1.5** (*CSI<sub>K</sub>* spacetimes). Finally, let us denote by  $CSI_K$ , those *CSI* spacetimes that belong to the (higher-dimensional) Kundt class (defined later); the so-called Kundt *CSI* spacetimes.

In particular, we shall study the relationship between  $CSI_R$ ,  $CSI_F$ ,  $CSI_K$  and especially with  $CSI \setminus H$ . We note that by construction  $CSI_R$  is at least of Weyl type *II* (i.e., of type *II*, *III*, *N* or *O* [9]), and by definition  $CSI_F$  and  $CSI_K$  are at least of Weyl type *II* (more precisely, at least of Riemann type *II*). In 4D,  $CSI_R$ ,  $CSI_F$  and  $CSI_K$  are closely related, and it is plausible that  $CSI \setminus H$  is at least of Weyl type *II* (see section 7).

In four dimensions the Weyl classification and the Petrov classification are closely related [9]. However, due to the fact that the Weyl classification contains many more possibilities in dimensions higher than four we will *restrict the term Petrov classification to four dimensions only*.

**1.2. 4D CSI.** We are particularly interested in the four-dimensional *CSI* spacetimes. For a Riemannian manifold every *CSI* is homogeneous ( $CSI \equiv H$ ). This is not true for Lorentzian manifolds. However, for every *CSI* with particular constant invariants there is a homogeneous spacetime (not necessarily unique) with precisely the same constant invariants. This suggests that *CSI* can be “built” from *H* and *VSI* (e.g.,  $CSI_R$ ).

In addition, there is a relationship between the *CSI* conditions and (i) the rank of the Riemann tensor and its holonomy class [12] (ii) the existence of curvature collineations [12] (iii) the condition of non-complete backsolvability<sup>1</sup> (NCB) [11], in addition to (iv) curvature homogeneity<sup>2</sup> and (v) sectional curvature [12]. The relationship between *CSI* and curvature homogeneity in 4D is studied in [13]. In locally homogeneous spacetimes all of the sectional curvatures (Gaussian curvatures) are constant. With the exception of certain special plane wave and constant curvature spacetimes (and all vacuum spacetimes), the sectional curvature uniquely determines the spacetime metric [12].

It is clear that *CSI* consists of  $CSI_R$  and, possibly, some other very special spacetimes. Let us present a *heuristic argument* for this in 4D. Let us suppose that the spacetime is *CSI*. For spacetimes that are completely backsolvable (CB) there is a special invariant frame such that all components of the Riemann tensor are constant. With respect to this invariant frame there exist smooth vector fields  $\zeta_i$  ( $i = 1, 2, 3, 4$ ) that act transitively on the manifold whose directional derivatives leave the Riemann tensor invariant (i.e.,  $\zeta_i$  form a Lie algebra of curvature collineations that span  $\mathcal{M}$ ). In general, every curvature collineation is a homothetic vector, so as a result there is a homothetic group acting transitively on  $\mathcal{M}$ . Therefore, this then implies that in general the *CSI* spacetime is locally homogeneous. The exceptions are those spacetimes that are not CB (NCB [11]) and those spacetimes for which a curvature collineation is not a homothetic vector [12], and these spacetimes are related to the *VSI* spacetimes.

**1.3. Overview.** In this paper we shall study *CSI* spacetimes. In the next section we shall begin by summarizing some important results. In section 3 we construct a subclass of  $CSI_R$  spacetimes that arise as warped products of a homogeneous space and a *VSI* spacetime. In section 4 we consider the higher-dimensional Kundt class. In section 5 examples of  $CSI_K$  spacetimes in higher dimensions are given. We discuss the results in section 6, and present a number of conjectures. In section 7 we summarize the results in 4D in detail.

There are three appendices. In Appendix A, we present all of the 3-dimensional *VSI* metrics, which is necessary for the explicit determination of the metrics in the 4-dimensional  $CSI_R$  spacetimes. In Appendix B, 4-dimensional *CSI* spacetimes

---

<sup>1</sup>Complete backsolvability (CB) refers to when all of the components of the Riemann tensor can be determined from the CZ set [11] of zeroth order curvature invariants, once all the remaining frame freedom has been used to fix components of the Riemann tensor.

<sup>2</sup> $\mathcal{M}$  is curvature homogeneous (of order zero) if there exists a frame with respect to which the Riemann tensor has constant components.

are given explicitly, and the relationship between  $CSI_R$ ,  $CSI_F$ , and  $CSI_K$  in 4D is discussed. Finally, in Appendix C, we write down the metric for higher dimensional  $VSI$  spacetimes in a canonical form.

## 2. SPACETIMES WITH CONSTANT SCALAR INVARIANTS

**2.1. Riemannian case.** Let us first consider the Riemannian case where a great deal about these spacetimes is known. The reason the Riemannian case is easier to deal with is because the orthogonal group  $O(d)$  is compact, and hence, the orbits of its group action are also compact.

**Theorem 2.1.** *If  $\mathcal{M}$  is a Riemannian spacetime, then:*

- (1)  $VSI_0 \Rightarrow VSI$ .
- (2) *There exists a  $k \in \mathbb{N}$  such that  $CSI_k \Rightarrow CSI$ .*

*Proof.* (1): See proof below. (2): See [1] □

**Theorem 2.2** (Riemannian  $VSI$ ). *A Riemannian space is  $VSI$  if and only if it is flat.*

*Proof.* Consider the invariant  $R_{ABCD}R^{ABCD}$ . Using an orthonormal frame, this invariant is a sum of squares; hence,  $R_{ABCD}R^{ABCD} = 0 \Leftrightarrow R_{ABCD} = 0$ . □

**Theorem 2.3** (Riemannian  $CSI$ ). *A Riemannian space is  $CSI$  if and only if it is locally homogeneous. Moreover, the set of curvature invariants,  $\mathcal{I}$ , uniquely determines the metric up to isometries.*

*Proof.* See Prüfer-Tricerri-Vanhecke [1]. □

**Theorem 2.4** (Singer). *If  $\mathcal{M}$  is curvature homogeneous of order  $k \geq \frac{1}{2}n(n-1)$ , then  $\mathcal{M}$  is locally homogeneous.*

*Proof.* See Singer [14]. □

**2.2. Lorentzian case.** The Lorentzian case is more difficult to deal with and only partial results are known. However, in the  $VSI$  case, it was shown in [15] that  $VSI_2$  implies  $VSI$ . A similar result in the  $CSI$  case is not known; however, in this case we believe that an  $n$  exists such that  $CSI_n$  implies  $CSI$ .

In the Riemannian case  $\mathcal{M}$  is locally homogeneous if and only if  $\mathcal{M}$  is curvature homogeneous to all orders (Theorem 2.4). In the Lorentzian case, we have that

**Theorem 2.5** (Podesta and Spiro). *There is an integer  $K_{p,q}$  such that if  $(M, g)$  is a complete, simply connected pseudo Riemannian manifold of type  $(p, q)$  which is  $K_{p,q}$ -curvature homogeneous, then  $\mathcal{M}$  is locally homogeneous.*

*Proof.* See Podesta and Spiro [16]. □

We argued above that in 4D Lorentzian manifolds for every  $CSI$  with particular constant invariants there is a locally homogeneous spacetime with the same constant invariants. It is plausible that this is true in higher dimensions.

**Conjecture 2.6.** *Assume that a Lorentzian spacetime  $\mathcal{M}$  is a  $CSI$  spacetime with curvature invariants  $\mathcal{I}$ . Then there exists a locally homogeneous space  $\widetilde{\mathcal{M}}$  with curvature invariants  $\widetilde{\mathcal{I}} = \mathcal{I}$ .*

**2.3. Boost-weight decomposition.** Consider an arbitrary covariant tensor  $T$  and a null frame  $\{\ell_A, n_A, m_A^{\hat{\mu}}\}$ ; i.e.,

$$\ell_A n^A = 1, \quad m_A^{\hat{\mu}} m^{\hat{\nu}A} = \delta^{\hat{\mu}\hat{\nu}}, \quad \ell_A \ell^A = n_A n^A = \ell_A m^{\hat{\mu}A} = n_A m^{\hat{\mu}A} = 0.$$

Consider now a boost in the  $\ell^A - n^A$ -plane:

$$(1) \quad \{\tilde{\ell}_A, \tilde{n}_A, \tilde{m}_A^{\hat{\mu}}\} = \{e^\lambda \ell_A, e^{-\lambda} n_A, m_A^{\hat{\mu}}\}.$$

We can consider the vector-space decomposition of the tensor  $T$  in terms of the boost weight with respect to the above boost (see [10]):

$$(2) \quad T = \sum_b (T)_b,$$

where  $(T)_b$  denotes the projection of the tensor  $T$  onto the vector space of boost-weight  $b$ . The components of the tensor  $(T)_b$  with respect to the frame will transform according to:

$$(3) \quad (T)_b{}_{AB\dots} = e^{-b\lambda} (T)_b{}_{\tilde{A}\tilde{B}\dots}.$$

Furthermore, we note that

$$(4) \quad (T \otimes S)_b = \sum_{b'+b''=b} (T)_{b'} \otimes (S)_{b''}.$$

Moreover, the connection  $\nabla$  can similarly be decomposed according to whether it raises, lowers or preserves the boost weight:

$$(5) \quad \nabla = (\nabla)_{-1} + (\nabla)_0 + (\nabla)_1,$$

where  $(\nabla)_b$  is defined by

$$(\nabla)_{b\mathbf{X}} \equiv \nabla_{(\mathbf{X})_b},$$

for all vectors  $\mathbf{X}$ . We note that for the metric,  $g = (g)_0$ ; hence, raising or lowering tensor indices preserves the boost weight.

An invariant of a tensor,  $T$ , is necessarily  $SO(1, n)$ -invariant; in particular, for any full contraction of  $T$ :

$$(6) \quad \text{Cont}[T] = \text{Cont}[(T)_0].$$

This property can now be utilized to construct spaces with constant curvature invariants. In the  $VSI$  case, there exists a null-frame such that the Riemann tensor,  $R$ , has the property:

$$(7) \quad (R)_b = 0, \quad b \geq 0.$$

This property alone implies that  $\mathcal{I}_0 = 0$ , regardless of  $(R)_b$  ( $b < 0$ ). If the curvature tensor is of the form:

$$(8) \quad (R)_b = 0, \quad b > 0, \quad \text{and } (R)_b{}_{ABCD,E} = 0, \quad b = 0,$$

in a frame for which  $g_{AB,E} = 0$ , it is necessarily a  $CSI_0$  spacetime. In general, further restrictions must be put on  $\nabla$  in order to make it  $CSI$ . We note that spacetimes can be classified according to their boost-weight components  $(R)_b$ . In particular, a classification using the Weyl tensor  $C$ , which generalises the Petrov classification in 4D, has been employed based on the components  $(C)_b$  [9, 10].

**2.4. Kundt metrics.** In the 4D *VSI* spacetimes, all of the positive boost weight spin coefficients (e.g.,  $\kappa, \rho$  and  $\sigma$ ) are zero, and hence it follows that  $\ell$  is geodesic, non-expanding, shear-free and non-twisting and the spacetime is Kundt  $K$  [17]. This is also true for *CSI* spacetimes, in the sense that if the *CSI* spacetime is not locally homogeneous, then in general it belongs to  $K$ . That is, it is plausible that in 4D,  $CSI \equiv H \cup CSI_K$ .

In higher dimensions it was also shown that  $\ell$  is geodesic, non-expanding, shear-free and non-twisting (i.e.,  $L_{ij} = 0$ ) in *VSI* spacetimes [15]. In locally homogeneous spacetimes, in general there exists a null frame in which the  $L_{ij}$  are constants. We therefore anticipate that in *CSI* spacetimes that are not locally homogeneous,  $L_{ij} = 0$ . For higher dimensional *CSI* spacetimes with  $L_{ij} = 0$ , the Ricci and Bianchi identities appear to be identically satisfied [19]. A higher dimensional spacetime which admits a null vector  $\ell$  which is geodesic, non-expanding, shear-free and non-twisting, will be denoted as a higher-dimensional Kundt spacetime.

Let us assume that the spacetime admits a geodesic, non-twisting, non-expanding, shear-free null vector  $\ell$ . It can be then shown that the existence of a twist-free and geodesic null vector implies that there exists a local coordinate system  $(u, v, x^i)$  such that

$$(9) \quad ds^2 = 2du (dv + Hdu + W_i dx^i) + \tilde{g}_{ij}(v, u, x^k) dx^i dx^j.$$

In essence, the twist-free condition implies that the vector field is 'surface forming' so that there exists locally an exact null one-form  $du$ . Using a coordinate transformation and a null-rotation we can now bring the metric into the above form. In this coordinate system,  $\ell = \frac{\partial}{\partial v}$ . In addition, requiring that this vector field is non-expanding and shear-free implies that  $\tilde{g}_{ij,v} = 0$ .

We will therefore consider metrics of the form

$$(10) \quad ds^2 = 2du (dv + Hdu + W_i dx^i) + \tilde{g}_{ij}(u, x^k) dx^i dx^j,$$

where  $H = H(v, u, x^k)$  and  $W_i = W_i(v, u, x^k)$ . The metric (10) possesses a null vector field  $\ell$  obeying

$$(11) \quad \ell^A \ell_{B;A} = \ell^A_{;A} = \ell^{A;B} \ell_{(A;B)} = \ell^{A;B} \ell_{[A;B]} = 0;$$

i.e., it is geodesic, non-expanding, shear-free and non-twisting. We will refer to the metrics (10) as higher-dimensional *Kundt metrics* (or simply Kundt metrics), since they generalise the 4-dimensional Kundt metrics.

**2.5. Lorentzian CSI.** All examples to the authors knowledge of Lorentzian *CSI* spacetimes are of the following two forms:

- (1) Homogeneous spaces
- (2) A subclass of the Kundt spacetimes

For the homogeneous spaces, their decomposition can be of the most general type,

$$\nabla^{(k)} R = \sum_{b=-(2+k)}^{2+k} \left( \nabla^{(k)} R \right)_b,$$

since a frame can be chosen such that all components are constants. However, for all known examples of non-homogeneous *CSI* metrics, a frame can be found such

that we have the decomposition

$$\nabla^{(k)} R = \sum_{b=-(2+k)}^0 \left( \nabla^{(k)} R \right)_b,$$

with constant boost-weight zero components.

Since a homogeneous space is automatically a *CSI* space, we will in this paper consider metrics which are not necessarily homogeneous. Since all known examples of non-homogeneous metrics are of Kundt form, we will henceforth only consider Kundt metrics.

### 3. WARPED PRODUCT *CSI*

It is known that  $CSI_R$  spacetimes can be constructed in all dimensions via warped products. Let us first determine necessary and sufficient conditions so that the warped product of a homogeneous space and a *VSI* is *CSI*.

We consider the warped product metric  $g_{AB} = g_{ab} \oplus e^{2\tau} g_{\alpha\beta}^2$  where  $g^1$  is a Riemannian homogenous space and  $g^2$  is a  $k$ -dimensional *VSI* Lorentzian manifold, the warping function  $\tau$  only depends on points of the homogenous space. Using the left-invariant frame of the homogeneous space<sup>3</sup>,

$$(12) \quad \mathbf{e}_{\hat{a}} = \left\{ \mathbf{m}_{\hat{k}}, \dots, \widehat{\mathbf{m}_{\hat{N}-1}} \right\},$$

we complete it to a basis of the warped product by appending the null frame,

$$(13) \quad \mathbf{e}_{\hat{\alpha}} = \left\{ \ell = e^{-\tau} \ell', \mathbf{n} = e^{-\tau} \mathbf{n}', \mathbf{m}_{\hat{2}} = e^{-\tau} \mathbf{m}'_{\hat{2}}, \dots, \widehat{\mathbf{m}_{\hat{k}-1}} = e^{-\tau} \mathbf{m}'_{\hat{k}-1} \right\},$$

where the primed vectors represent the canonical *VSI* frame in which  $\ell'$  (and hence  $\ell$ ) is the aligned geodesic null congruence that is expansion, shear, and twist free. The non-vanishing inner products are such that

$$(14) \quad g_{AB} = 2\ell_{(A} n_{B)} + \delta_{\hat{\mu}\hat{\nu}} m^{\hat{\mu}}_A m^{\hat{\nu}}_B + \delta_{\hat{m}\hat{n}} m^{\hat{m}}_A m^{\hat{n}}_B.$$

Defining  $T_{\hat{m}\hat{p}} = \tau_{,\hat{m};\hat{p}} + \tau_{,\hat{m}} \tau_{,\hat{p}}$ , we shall show that if  $\tau_{,m} \tau^{,m}$  and  $T_{\hat{m}\hat{p}}$  are constant then this implies *CSI*. These two conditions imply that  $\tau^{,\hat{m}}$  is an eigenvector of  $T_{\hat{m}\hat{p}}$  with constant eigenvalue  $\tau_{,m} \tau^{,m}$ , and thus  $\tau_{,\hat{b}}$  must be constant.

Following [15], we perform a decomposition of the Riemann tensor for the warped product to obtain

$$(15) \quad \begin{aligned} R_{ABCD} = & 4R_{\hat{0}\hat{1}\hat{0}\hat{1}} n_{\{A} \ell_B n_C \ell_{D\}} + 8R_{\hat{0}\hat{\mu}\hat{1}\hat{\lambda}} n_{\{A} m^{\hat{\mu}}_B \ell_C m^{\hat{\lambda}}_{D\}} + 8R_{\hat{0}\hat{a}\hat{1}\hat{b}} n_{\{A} m^{\hat{a}}_B \ell_C m^{\hat{b}}_{D\}} \\ & + R_{\hat{\mu}\hat{\nu}\hat{\lambda}\hat{\sigma}} m^{\hat{\mu}}_{\{A} m^{\hat{\nu}}_B m^{\hat{\lambda}}_C m^{\hat{\sigma}}_{D\}} + 4R_{\hat{a}\hat{\mu}\hat{b}\hat{\lambda}} m^{\hat{a}}_{\{A} m^{\hat{\mu}}_B m^{\hat{b}}_C m^{\hat{\lambda}}_{D\}} \\ & + R_{\hat{a}\hat{b}\hat{c}\hat{d}} m^{\hat{a}}_{\{A} m^{\hat{b}}_B m^{\hat{c}}_C m^{\hat{d}}_{D\}} + \text{negative boost weight terms} \dots \end{aligned}$$

Evidently the Riemann tensor is of boost order zero with boost weight zero components:

<sup>3</sup>Hatted indices are the preferred null-frame indices.

$$(16) \quad R_{\hat{0}\hat{1}\hat{0}\hat{1}} = \tau_{,m}\tau^{,m}$$

$$(17) \quad R_{\hat{0}\hat{\mu}\hat{1}\hat{\lambda}} = -\tau_{,m}\tau^{,m} \delta_{\hat{\mu}\hat{\lambda}}^2$$

$$(18) \quad R_{\hat{0}\hat{a}\hat{1}\hat{b}} = -T_{\hat{a}\hat{b}}$$

$$(19) \quad R_{\hat{\mu}\hat{\nu}\hat{\lambda}\hat{\sigma}} = \tau_{,m}\tau^{,m} \left[ \delta_{\hat{\nu}\hat{\lambda}}^2 \delta_{\hat{\mu}\hat{\sigma}}^2 - \delta_{\hat{\mu}\hat{\lambda}}^2 \delta_{\hat{\nu}\hat{\sigma}}^2 \right]$$

$$(20) \quad R_{\hat{a}\hat{\mu}\hat{b}\hat{\lambda}} = -T_{\hat{a}\hat{b}} \delta_{\hat{\mu}\hat{\lambda}}^2$$

$$(21) \quad R_{\hat{a}\hat{b}\hat{c}\hat{d}} = R_{\hat{a}\hat{b}\hat{c}\hat{d}}^1.$$

The negative boost weight components, arising from the *VSI* spacetime, are given by

$$(22) \quad R_{\hat{1}\hat{0}\hat{1}\hat{\mu}} = e^{-2\tau} R_{\hat{1}\hat{0}\hat{1}\hat{\mu}}^2$$

$$(23) \quad R_{\hat{1}\hat{\mu}\hat{\nu}\hat{\lambda}} = e^{-2\tau} R_{\hat{1}\hat{\mu}\hat{\nu}\hat{\lambda}}^2$$

$$(24) \quad R_{\hat{1}\hat{\mu}\hat{1}\hat{\nu}} = e^{-2\tau} R_{\hat{1}\hat{\mu}\hat{1}\hat{\nu}}^2.$$

The condition that  $\tau_{,\hat{b}}$  is constant (and hence  $\tau_{,m}\tau^{,m}$  and  $T_{\hat{m}\hat{p}}$  are constant) implies that the boost weight zero components of the Riemann tensor are constant; the absence of positive boost weight components then gives *CSI*<sub>0</sub>.

The covariant derivative of the null frame has the form,

$$(25) \quad \ell_{K;E} = -\tau_{,\hat{b}} m_{\hat{K}}^{\hat{b}} \ell_E + \gamma_{\hat{\lambda}\hat{0}\hat{1}} m_{\hat{K}}^{\hat{\lambda}} \ell_E + \gamma_{\hat{1}\hat{0}\hat{\lambda}} \ell_K m_{\hat{E}}^{\hat{\lambda}} + \dots$$

$$(26) \quad n_{K;E} = -\tau_{,\hat{b}} m_{\hat{K}}^{\hat{b}} n_E - \gamma_{\hat{1}\hat{0}\hat{\lambda}} n_K m_{\hat{E}}^{\hat{\lambda}} + \dots$$

$$(27) \quad m_{\hat{\lambda}K;E} = -\tau_{,\hat{b}} m_{\hat{K}}^{\hat{b}} m_{\hat{\lambda}E} + \gamma_{\hat{\mu}\hat{\lambda}\hat{\nu}} m_{\hat{K}}^{\hat{\mu}} m_{\hat{E}}^{\hat{\nu}} - \gamma_{\hat{\lambda}\hat{0}\hat{1}} n_K \ell_E + \dots$$

$$(28) \quad m_{\hat{a}K;E} = \tau_{,\hat{a}} \ell_K n_E + \tau_{,\hat{a}} n_K \ell_E + \tau_{,\hat{a}} \delta_{\hat{\mu}\hat{\lambda}}^{\hat{\mu}} m_{\hat{K}}^{\hat{\mu}} m_{\hat{E}}^{\hat{\lambda}} + \gamma_{\hat{b}\hat{a}\hat{c}}^1 m_{\hat{K}}^{\hat{b}} m_{\hat{E}}^{\hat{c}}$$

where  $\dots$  denotes terms of lower order boost weight with respect to previous terms in an expression, and the unspecified  $\gamma$ 's correspond to *VSI* Ricci-rotation coefficients [15] [for example,  $\gamma_{\hat{1}\hat{0}\hat{\lambda}} = e^{-\tau} L_{\hat{1}\hat{\lambda}}$ ], except for the  $\gamma^1$  which are the rotation coefficients associated with the homogeneous space. Equations (25)-(28) have leading order boost weight terms of +1, -1, 0 and 0, respectively, which is the same boost weight as the corresponding null frame vector before taking a covariant derivative. It follows that the covariant derivative of the  $\{\cdot\}$  quantities appearing in (15) will produce terms of equal or lesser boost weight. Furthermore, in [15] it was proven that the negative boost weight curvature components appearing in (15) will remain negative upon covariant differentiation. Therefore, if the boost weight zero components of the Riemann tensor and its derivatives are constant then any number of covariant derivatives of the Riemann tensor will be of boost order zero, which implies that the warped product will be *CSI*.

Supposing that the boost weight zero components of  $\nabla^{(n)}R$  are constant, then for  $\nabla^{(n+1)}R$  the boost weight zero components will have contributions from  $\tau_{,\hat{a}}$ , which is constant, and the non-constant  $\gamma$ 's arising from the *VSI* spacetime, however these boost weight zero *VSI*  $\gamma$ 's do not occur as a result of Theorem 5.1. Therefore we have that if  $\tau_{,\hat{b}}$  is constant, then the warped product is *CSI*.



For reference, we include the covariant derivative of the Riemann tensor. By using the Ricci identity we have,

$$(29) \quad \tau^{\hat{a}} \overset{1}{R}_{\hat{a}\hat{b}\hat{c}\hat{d}} + 2T_{\hat{b}[\hat{d}\hat{c}\hat{e}]} = T_{\hat{b}\hat{c};\hat{d}} - T_{\hat{b}\hat{d};\hat{c}},$$

which can be used to express the non-vanishing boost weight zero components of the covariant derivative of the Riemann tensor as

$$(30) \quad R_{\hat{1}\hat{b}\hat{c}\hat{d};\hat{0}} = T_{\hat{b}\hat{c};\hat{d}} - T_{\hat{b}\hat{d};\hat{c}} = R_{\hat{0}\hat{b}\hat{c}\hat{d};\hat{1}}$$

$$(31) \quad R_{\hat{\mu}\hat{b}\hat{c}\hat{d};\hat{\lambda}} = \left[ T_{\hat{b}\hat{c};\hat{d}} - T_{\hat{b}\hat{d};\hat{c}} \right] \delta_{\hat{\mu}\hat{\lambda}}$$

$$(32) \quad R_{\hat{0}\hat{b}\hat{1}\hat{c};\hat{e}} = -T_{\hat{b}\hat{c};\hat{e}}$$

$$(33) \quad R_{\hat{b}\hat{\mu}\hat{c}\hat{\lambda};\hat{e}} = -T_{\hat{b}\hat{c};\hat{e}} \delta_{\hat{\mu}\hat{\lambda}}.$$

As expected all components in (30)-(33) are constant (as a result of requiring  $\tau_{\hat{b}}$  to be constant).

We shall now show the converse; that is, if the Ricci invariants are constant then  $\tau_{\hat{b}}$  is constant. In local coordinates, the Ricci tensor and Ricci scalar are

$$(34) \quad R_{\mu}^{\rho} = e^{-2\tau} \overset{2}{R}_{\mu}^{\rho} - [T + (k-1)\tau_{,m}\tau^{,m}] \delta_{\mu}^{\rho}$$

$$(35) \quad R_m^p = \overset{1}{R}_m^p - kT_m^p$$

$$(36) \quad R = \overset{1}{R} - k[2T + (k-1)\tau_{,m}\tau^{,m}],$$

where in (36) we have used that  $\overset{2}{R} = 0$  and set  $T = \text{Tr}(T_m^p)$ . Choosing new coordinates  $\tilde{x}^1 = \tau(x^m)$ ,  $\tilde{x}^2 = x^2, \dots$  so that  $\tau_{,m}\tau^{,m} = \tilde{g}^{11}$ , then

$$(37) \quad \tilde{T}_{mn} = -\tilde{\Gamma}_{mn}^1 + \delta_m^1 \delta_n^1.$$

Now, with normal coordinates along  $\tilde{x}^1$  we have  $\tilde{g}^{11} = 1$ . Therefore from (36), since  $\overset{1}{R}$  is constant we have that  $\tilde{T}$ , and hence  $T$ , is constant. In this coordinate system the constraint  $\tilde{T}$  constant implies that  $\tilde{g}^{mn}\tilde{g}_{mn,1}$  is constant. Constant Ricci scalar implies that  $T$  and  $\tau_{,m}\tau^{,m}$  are constant. Further conditions are then provided by the constancy of higher degree Ricci invariants.

From (34) and (35), the Ricci tensor of a warped product is  $R_M^N = R_m^n \oplus R_{\mu}^{\nu}$ , thus a degree  $d$  Ricci invariant always has the form

$$(38) \quad R_{N_1}^{N_2} R_{N_2}^{N_3} \dots R_{N_d}^{N_1} = R_{n_1}^{n_2} R_{n_2}^{n_3} \dots R_{n_d}^{n_1} + R_{\nu_1}^{\nu_2} R_{\nu_2}^{\nu_3} \dots R_{\nu_d}^{\nu_1}.$$

The second term of (38) is constant since  $T$  and  $\tau_{,m}\tau^{,m}$  are constant; hence invariants constructed from (35) must be constant. Assuming the determinant of  $R_m^p$  or  $\overset{1}{R}_m^p$  is nonzero, then there exists a frame in which  $R_m^p$  and  $\overset{1}{R}_m^p$  can simultaneously be expressed in block diagonal form with constant eigenvalues [18]. Therefore, (35) implies that in this frame  $T_m^p$  must also have constant components and thus constant invariants. In this case, the constancy of the Ricci invariants completely determines the conditions on  $T_m^p$  and specifies a frame in which the components of each term in (35) are constant. The frame in which  $\overset{1}{R}_m^p$  and  $T_m^p$  are constant is determined by the left-invariant 1-forms. As before, the conditions

that  $T_{\hat{a}\hat{b}}$  and  $\tau_m\tau^m$  are constant imply  $\tau_{,\hat{a}}$  is constant, and consequently constant Ricci invariants implies *CSI*.

Many examples of *CSI* spacetimes can be found among the cases where the homogeneous space is a solvable Lie group<sup>4</sup>:

**Theorem 3.1.** *Assume that  $\mathcal{M}_H$  is a solvable Lie group equipped with a left-invariant metric. Assume also that  $\mathcal{M}_H$  is connected and simply connected (i.e.,  $H^1(\mathcal{M}_H) = 0$ ). Then there exists a non-constant  $\tau$  such that the warped product of  $\mathcal{M}_H$  and a *VSI*, as constructed above, is a *CSI*.*

*Proof.* We identify  $\mathcal{M}_H = G$ , where  $G$  is a solvable Lie group, with the Lie algebra  $\mathfrak{g} = T_e G$ . Since  $\mathfrak{g}$  is solvable the set  $[\mathfrak{g}, \mathfrak{g}]$  will be a proper vector subspace of  $\mathfrak{g}$ . Thus for a solvable Lie algebra, there exists a non-zero  $\mathbf{X} \in \mathfrak{g} \cap [\mathfrak{g}, \mathfrak{g}]^\perp$  where  $[\mathfrak{g}, \mathfrak{g}]^\perp$  is the complement of the derived algebra  $[\mathfrak{g}, \mathfrak{g}]$ . This further implies that there is a left-invariant one-form  $\omega$  such that  $d\omega = 0$ . Since  $H^1(\mathcal{M}_H) = 0$ , there exists a non-constant function  $\phi$  such that  $\omega = d\phi$ . We can now choose  $\tau = p\phi$  for a constant  $p$ .  $\square$

In 2-dimensions flat space is the only *VSI* metric, and in the appendix we list all of the 3-dimensional *VSI* metrics. Examples of *CSI* warped products where the homogeneous space is either  $\mathbb{E}^n$  or  $\mathbb{H}^n$  yield a linear function for  $\tau$  (since both of these can be considered as solvmanifolds); however, for  $\mathbb{S}^n$ , since the coordinates are cyclic, we find that  $\tau$  is a constant. Another example, in 5-dimensions, is where the homogeneous space is a Bianchi type VIII Lie group ( $\cong SL(2, \mathbb{R})$ ) warped with a 2-dimensional *VSI*, with coordinates  $u, v, x, y, z$  the metric is

$$(39) \quad ds^2 = e^{2\tau} [2du(dv + Hdu)] + \frac{a^2}{y^2} (\cos z dx + \sin z dy)^2 + \frac{b^2}{y^2} (\cos z dy - \sin z dx)^2 + c^2 \left( dz + \frac{dx}{y} \right)^2.$$

In this case  $H(u, v)$  must be linear in  $v$  to give *CSI*, and it turns out that for the warped product to be *CSI* we have to consider two separate cases:

- (1) The maximal isometry group of the homogeneous space is 3-dimensional ( $a \neq b$ ):  $\tau$  is a constant.
- (2) The maximal isometry group of the homogeneous space is 4-dimensional ( $a = b$ ):  $\tau$  can be non-constant,  $\tau = \alpha \ln y + \beta z$ .

The reason why we have to distinguish these two cases is that in the first case, the isometry group is the semi-simple group,  $SL(2, \mathbb{R})$ . However, in the second case, the isometry group allows for a transitive group which is solvable (Bianchi type III).

#### 4. KUNDT *CSI* METRICS

Let us next consider the higher-dimensional Kundt metrics (10) in the form

$$(40) \quad ds^2 = 2du \left( dv + Hdu + W_i \mathbf{m}^i \right) + \delta_{\hat{i}\hat{j}} \mathbf{m}^{\hat{i}} \mathbf{m}^{\hat{j}}.$$

---

<sup>4</sup>Recall that a Lie group can be equipped with a left-invariant metric (i.e., invariant under the left action) in the standard way, see e.g. [20].

It is convenient to introduce the null frame

$$(41) \quad \ell = du,$$

$$(42) \quad \mathbf{n} = dv + Hdu + W_{\hat{i}} \mathbf{m}^{\hat{i}},$$

$$(43) \quad \mathbf{m}^{\hat{i}} = \mathbf{e}^{\hat{i}}_{\hat{j}}(u; x^k) dx^{\hat{j}},$$

$$(44) \quad (\eta_{AB}) = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \delta_{\hat{i}\hat{j}} \end{bmatrix}$$

such that

$$(45) \quad dS_H^2 \equiv \delta_{\hat{i}\hat{j}} \mathbf{m}^{\hat{i}} \mathbf{m}^{\hat{j}} = \tilde{g}_{ij} dx^i dx^j, \quad \tilde{g}_{ij,v} = 0.$$

There is a class of coordinate transformations that preserve the form of the metric (40). In particular, we can define new variables,

$$(46) \quad (v', u', x'^i) = (v, u, f^i(u; x^k)).$$

For our purposes we can always use this coordinate transformation to simplify the spatial metric  $\tilde{g}_{ij}$ :

**Theorem 4.1.** *Consider the metric (10) and assume that there exists a null frame  $\{\ell_A, n_A, m^{\hat{i}}_A\}$  such that all scalars  $R_{\hat{i}\hat{j}\hat{k}\hat{l}} \equiv R_{ABDC} m^{\hat{i}}_A m^{\hat{j}}_B m^{\hat{k}}_C m^{\hat{l}}_D$  and  $R_{\hat{i}\dots\hat{j}_1\dots\hat{j}_n} \equiv R_{A\dots C_1\dots C_n} m^{\hat{i}}_A \dots m^{\hat{j}_1}_{C_1} \dots m^{\hat{j}_n}_{C_n}$  are constants. Then there exists (locally) a coordinate transformation  $(v', u', x'^i) = (v, u, f^i(u; x^k))$  such that*

$$(47) \quad \tilde{g}_{ij} \equiv \tilde{g}'_{kl} \frac{\partial f^k}{\partial x^i} \frac{\partial f^l}{\partial x^j}, \quad \tilde{g}'_{ij,u'} = 0.$$

Moreover,  $dS_H^2 = \tilde{g}'_{ij} dx'^i dx'^j$  is a locally homogeneous space.

*Proof.* By calculating the curvature tensor of the metric (10), and its covariant derivatives, we note that

$$(48) \quad R_{\hat{i}\hat{j}\hat{k}\hat{l}} = \tilde{R}_{\hat{i}\hat{j}\hat{k}\hat{l}}, \quad R_{\hat{i}\dots\hat{j}_1\dots\hat{j}_n} = \tilde{R}_{\hat{i}\dots\hat{j}_1\dots\hat{j}_n},$$

where the tilde refers to curvature tensors with respect to the spatial metric  $\tilde{g}_{ij}(u, x^k)$ . Since this metric is Riemannian with all its components constant we can use the results of [1] which state that the metrics  $\tilde{g}_{ij}(u, x^k)$ , for different  $u$ , are equivalent up to isometries. Thus, given  $u_0$ , there exists for every  $u$  sufficiently close to  $u_0$  an isometry  $\psi_u(x^k)$  such that  $(\psi_u^* \tilde{g})(u) = \tilde{g}(u_0)$ . Since, we assume that  $\tilde{g}$  is sufficiently smooth in a neighbourhood of  $u_0$ , we can find a sufficiently smooth  $\psi_u$  in  $u$ . The map  $\psi_u(x^k)$  provides us with the functions  $f^i$  in the theorem by composition with the coordinate charts. Hence, we have  $\tilde{g}' = \tilde{g}|_{u=u_0}$ . Moreover,  $\tilde{g}'$  is a locally homogeneous space [1].  $\square$

Henceforth, we shall assume that we consider solutions in the set  $CSI_F \cap CSI_K$ . Consequently, from Theorem 4.1, there is no loss of generality in assuming that the metric (10) has  $\tilde{g}_{ij,u} = 0$ . The remaining coordinate freedom preserving the Kundt form is then:

$$(1) \quad (v', u', x'^i) = (v, u, f^i(x^k)) \text{ and } J^i_{\hat{j}} \equiv \frac{\partial f^i}{\partial x^{\hat{j}}}.$$

$$H' = H, \quad W'_i = W_j (J^{-1})^j_i, \quad \tilde{g}'_{ij} = \tilde{g}_{kl} (J^{-1})^k_i (J^{-1})^l_j.$$

$$(2) \quad (v', u', x^i) = (v + h(u, x^k), u, x^i)$$

$$H' = H - h_{,u}, \quad W'_i = W_i - h_{,i}, \quad \tilde{g}'_{ij} = \tilde{g}_{ij}.$$

$$(3) \quad (v, u, x^i) = (v/g_{,u}(u), g(u), x^i)$$

$$H' = \frac{1}{g_{,u}^2} \left( H + v \frac{g_{,uu}}{g_u} \right), \quad W'_i = \frac{1}{g_{,u}} W_i, \quad \tilde{g}'_{ij} = \tilde{g}_{ij}.$$

4.1. *CSI*<sub>0</sub> spacetimes. The linearly independent components of the Riemann tensor with boost weight 1 and 0 are:

$$(49) \quad R_{\hat{0}\hat{1}\hat{0}\hat{i}} = -\frac{1}{2} W_{\hat{i},vv},$$

$$(50) \quad R_{\hat{0}\hat{1}\hat{0}\hat{1}} = -H_{,vv} + \frac{1}{4} (W_{\hat{i},v}) (W^{\hat{i},v}),$$

$$(51) \quad R_{\hat{0}\hat{1}\hat{i}\hat{j}} = W_{[\hat{i}W_{\hat{j}],vv} + W_{[\hat{i};\hat{j}],v},$$

$$(52) \quad R_{\hat{0}\hat{i}\hat{1}\hat{j}} = \frac{1}{2} \left[ -W_{\hat{j}} W_{\hat{i},vv} + W_{\hat{i};\hat{j},v} - \frac{1}{2} (W_{\hat{i},v}) (W_{\hat{j},v}) \right],$$

$$(53) \quad R_{\hat{i}\hat{j}\hat{n}\hat{m}} = \tilde{R}_{\hat{i}\hat{j}\hat{n}\hat{m}}.$$

Hence, the spacetime is a *CSI*<sub>0</sub> spacetime if there exists a frame  $\{\ell, \mathbf{n}, \mathbf{m}^{\hat{i}}\}$ , a constant  $\sigma$ , anti-symmetric matrix  $\mathbf{a}_{\hat{i}\hat{j}}$ , and symmetric matrix  $\mathbf{s}_{\hat{i}\hat{j}}$  such that

$$(54) \quad W_{\hat{i},vv} = 0,$$

$$(55) \quad H_{,vv} - \frac{1}{4} (W_{\hat{i},v}) (W^{\hat{i},v}) = \sigma,$$

$$(56) \quad W_{[\hat{i};\hat{j}],v} = \mathbf{a}_{\hat{i}\hat{j}},$$

$$(57) \quad W_{(\hat{i};\hat{j}),v} - \frac{1}{2} (W_{\hat{i},v}) (W_{\hat{j},v}) = \mathbf{s}_{\hat{i}\hat{j}},$$

and the components  $\tilde{R}_{\hat{i}\hat{j}\hat{m}\hat{n}}$  are all constants (i.e.,  $dS_H^2$  is curvature homogeneous).

We note that the first equation implies that

$$(58) \quad W_{\hat{i}}(v, u, x^k) = v W_{\hat{i}}^{(1)}(u, x^k) + W_{\hat{i}}^{(0)}(u, x^k),$$

while the second implies

$$(59) \quad H(v, u, x^k) = \frac{v^2}{8} \left[ 4\sigma + (W_{\hat{i}}^{(1)})(W^{(1)\hat{i}}) \right] + v H^{(1)}(u, x^k) + H^{(0)}(u, x^k).$$

If  $dS_H^2 = \delta_{\hat{i}\hat{j}} \mathbf{m}^{\hat{i}} \mathbf{m}^{\hat{j}}$  is a (locally) homogeneous Riemannian space, then there exists a frame where  $\tilde{R}_{\hat{i}\hat{j}\hat{m}\hat{n}}$  are all constants. In general, curvature homogeneous does not imply homogeneous; however, since we are mostly interested in the *CSI* case, in light of Theorem 4.1, we will henceforth take  $dS_H^2$  as a (locally) homogeneous metric.

4.2. *CSI*<sub>1</sub> spacetimes. For a *CSI*<sub>0</sub> spacetime, we have

$$(60) \quad R_{\hat{0}\hat{1}\hat{0}\hat{i};\hat{1}} = -\frac{1}{2} \left[ \sigma W_{\hat{i},v} - \frac{1}{2} (\mathbf{s}_{\hat{j}\hat{i}} + \mathbf{a}_{\hat{j}\hat{i}}) W^{\hat{j},v} \right],$$

$$(61) \quad R_{\hat{0}\hat{i}\hat{j}\hat{k};\hat{1}} = -\frac{1}{2} \left[ W^{\hat{n},v} \tilde{R}_{\hat{n}\hat{i}\hat{j}\hat{k}} - W_{\hat{i},v} \mathbf{a}_{\hat{j}\hat{k}} + (\mathbf{s}_{\hat{i}[\hat{j}} + \mathbf{a}_{\hat{i}[\hat{j}}) W_{\hat{k}],v} \right].$$

For the spacetime to be  $CSI_1$ , it is sufficient to require that the above components are constants; i.e.

$$(62) \quad \sigma W_{\hat{i},v} - \frac{1}{2}(\mathfrak{s}_{\hat{j}\hat{i}} + \mathfrak{a}_{\hat{j}\hat{i}})W^{\hat{j},v} = \alpha_{\hat{i}},$$

$$(63) \quad \left(\mathfrak{s}_{\hat{i}\hat{j}} + \mathfrak{a}_{\hat{i}\hat{j}}\right)_{;\hat{k}} - \left(\mathfrak{s}_{\hat{i}\hat{k}} + \mathfrak{a}_{\hat{i}\hat{k}}\right)_{;\hat{j}} = \beta_{\hat{i}\hat{j}\hat{k}},$$

where the Ricci identity has been used to rewrite the latter condition.

## 5. EXAMPLES OF $CSI$ SPACETIMES OF KUNDT FORM

Various classes of  $CSI$  spacetimes that arise as members of  $CSI_F \cap CSI_K$  can now be found. The solutions come in classes according to the properties of  $W_{\hat{i}}^{(1)}(u, x^k)$ . In the following,  $\mathcal{M}_H$  is a locally homogeneous Riemannian space.

5.1.  $W_{\hat{i}}^{(1)}(u, x^k) = 0$ . Assuming  $W_{\hat{i}}^{(1)}(u, x^k) = 0$  immediately makes the metric (40) a  $CSI$  space.

A special subcase of this class which is worth mentioning is the Brinkmann metrics for which  $H$  and  $W_{\hat{i}}$  are all independent of  $v$ ; thus  $\sigma = \mathfrak{s}_{ij} = \mathfrak{a}_{ij} = 0$ . In this case, the expressions for the curvature tensors simplify drastically. In particular, if  $F_{\hat{j}\hat{i}} \equiv 2W_{[\hat{i};\hat{j}]}$ , the Ricci tensor is

$$(64) \quad R_{\hat{1}\hat{1}} = \tilde{\square}H - \frac{1}{4}\tilde{F}^2, \quad R_{\hat{1}\hat{i}} = \tilde{\nabla}^{\hat{j}}F_{\hat{j}\hat{i}}, \quad R_{\hat{i}\hat{j}} = \tilde{R}_{\hat{i}\hat{j}}.$$

5.2.  $W_{\hat{i}}^{(1)}(u, x^k) = \text{constant}$ . This case is  $CSI$  if and only if the simpler metric,  $\widetilde{\mathcal{M}}$ :

$$(65) \quad \widetilde{ds}^2 = 2du \left( dv + \frac{v^2}{2}\tilde{\sigma}du + vW_{\hat{i}}^{(1)}\mathbf{m}^{\hat{i}} \right) + \delta_{\hat{i}\hat{j}}\mathbf{m}^{\hat{i}}\mathbf{m}^{\hat{j}},$$

is a homogeneous space. The curvature invariants of the Kundt metrics in this class will have the same invariants as  $\widetilde{\mathcal{M}}$ ; i.e.  $\mathcal{I} = \mathcal{I}(\widetilde{\mathcal{M}})$ .

The homogeneous space  $\mathcal{M}_H$  can be considered as a quotient space  $\mathcal{M}_H = G/H$ , where  $G$  is the isometry group and  $H$  is the isotropy group. The corresponding Lie algebra decomposition is  $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$ , and we let this correspond to the algebra of left-invariant vectors. We note that if there exists a subalgebra  $\mathfrak{k} \subset \mathfrak{m}$  such that  $[\mathfrak{k}^\perp, \mathfrak{k}] = 0$ , where  $\mathfrak{k}^\perp$  is the complement of  $\mathfrak{k}$  in  $\mathfrak{g}$ , then we can choose an  $\text{Ad}(H)$ -invariant  $\mathbf{W}$ :

$$\mathbf{W} \in \mathfrak{k}.$$

By letting  $W_{\hat{i}}^{(1)}\mathbf{m}^{\hat{i}}$  be its corresponding left-invariant one-form, the associated Kundt spacetime will be a  $CSI$ .

Note that some homogeneous spaces may be represented by inequivalent Lie algebra decompositions. For example, hyperbolic space,  $\mathbb{H}^n$ , can both be considered as the quotient  $SO(1, n)/SO(n)$  and as a solvable Lie group  $G$  [7]. In the latter case the decomposition gives  $\mathfrak{g} = \mathfrak{m}$  and  $\mathfrak{h} = 0$ ; hence,  $[\mathfrak{h}, \mathfrak{m}] = 0$ .

5.3.  $W_{\hat{i}}^{(1)}(u, x^k) = \phi_{,\hat{i}} = \text{non-constant}$ .

5.3.1.  $\mathcal{I} = \mathcal{I}(\text{dS} \times \mathcal{M}_H)$ .  $\sigma > 0$ :

$$(66) \quad \begin{aligned} \text{d}s^2 &= \cos^2(\sqrt{\sigma}x) \left[ 2\text{d}u \left( \text{d}v + \tilde{H}\text{d}u + \tilde{W}_i \mathbf{m}^i \right) \right] \\ &+ \text{d}x^2 + \frac{1}{\sigma} \sin^2(\sqrt{\sigma}x) \text{d}S_{S^n}^2 + \text{d}S_H^2, \end{aligned}$$

where

$$(67) \quad \begin{aligned} W_{\hat{x}}^{(1)} &= 2\sqrt{\sigma} \tan(\sqrt{\sigma}x), \\ \tilde{W}_i &= \tilde{W}_i^{(0)}(u, x^j), \quad \tilde{H} = \frac{v^2}{2}\sigma + v\tilde{H}^{(1)}(u, x^j) + \tilde{H}^{(0)}(u, x^j). \end{aligned}$$

The Riemann tensor is of type:

$$R = (R)_0 + (R)_{-1} + (R)_{-2}.$$

Special cases:

- (1) If  $\tilde{W}_i^{(0)} = \frac{1}{\sigma} \tilde{H}_{,i}^{(1)}$  then  $(R)_{-1} = 0$ .
- (2) If  $\tilde{W}_i^{(0)} = \tilde{H}^{(1)} = \tilde{H}^{(0)} = 0$  then  $(R)_{-1} = (R)_{-2} = 0$  and  $\text{dS} \times \mathcal{M}_H$ .

5.3.2.  $\mathcal{I} = \mathcal{I}(\text{AdS} \times \mathcal{M}_H)$ .  $\sigma < 0$ :

$$(68) \quad \begin{aligned} \text{d}s_I^2 &= \cosh^2(\sqrt{|\sigma|x}) \left[ 2\text{d}u \left( \text{d}v + \tilde{H}\text{d}u + \tilde{W}_i \mathbf{m}^i \right) \right] \\ &+ \text{d}x^2 + \frac{1}{|\sigma|} \sinh^2(\sqrt{|\sigma|x}) \text{d}S_{S^n}^2 + \text{d}S_H^2, \end{aligned}$$

where

$$(69) \quad \begin{aligned} W_{\hat{x}}^{(1)} &= -2\sqrt{|\sigma|} \tanh(\sqrt{|\sigma|x}), \\ \tilde{W}_i &= \tilde{W}_i^{(0)}(u, x^j), \quad \tilde{H} = -\frac{v^2}{2}|\sigma| + v\tilde{H}^{(1)}(u, x^j) + \tilde{H}^{(0)}(u, x^j). \end{aligned}$$

The Riemann tensor is of type:

$$R = (R)_0 + (R)_{-1} + (R)_{-2}.$$

Special cases:

- (1) If  $\tilde{W}_i^{(0)} = -\frac{1}{|\sigma|} \tilde{H}_{,i}^{(1)}$  then  $(R)_{-1} = 0$ .
- (2) If  $\tilde{W}_i^{(0)} = \tilde{H}^{(1)} = \tilde{H}^{(0)} = 0$  then  $(R)_{-1} = (R)_{-2} = 0$  and  $\text{AdS} \times \mathcal{M}_H$ .

Let us present another example:

$$(70) \quad \begin{aligned} \text{d}s_{II}^2 &= \sinh^2(\sqrt{|\sigma|x}) \left[ 2\text{d}u \left( \text{d}v + \tilde{H}\text{d}u + \tilde{W}_i \mathbf{m}^i \right) \right] \\ &+ \text{d}x^2 + \frac{1}{|\sigma|} \cosh^2(\sqrt{|\sigma|x}) \text{d}S_{\mathbb{H}^n}^2 + \text{d}S_H^2, \end{aligned}$$

where

$$(71) \quad \begin{aligned} W_{\hat{x}}^{(1)} &= 2\sqrt{|\sigma|} \coth(\sqrt{|\sigma|x}), \\ \tilde{W}_i &= \tilde{W}_i^{(0)}(u, x^j), \quad \tilde{H} = \frac{v^2}{2}|\sigma| + v\tilde{H}^{(1)}(u, x^j) + \tilde{H}^{(0)}(u, x^j). \end{aligned}$$

The Riemann tensor is of type:

$$R = (R)_0 + (R)_{-1} + (R)_{-2}.$$

Special cases:

- (1) If  $\tilde{W}_i^{(0)} = \frac{1}{|\sigma|} \tilde{H}_{,i}^{(1)}$  then  $(R)_{-1} = 0$ .

(2) If  $\widetilde{W}_i^{(0)} = \widetilde{H}^{(1)} = \widetilde{H}^{(0)} = 0$  then  $(R)_{-1} = (R)_{-2} = 0$ .

There are more *CSI* spacetimes with  $\mathcal{I} = \mathcal{I}((A)dS \times \mathcal{M}_H)$ . They are constructed similarly to the ones above. They are all related to the fact that de Sitter space or Anti-de Sitter space can be written in many ways as fibered spaces over the spheres,  $S^n$ , or the hyperbolic space,  $\mathbb{H}^n$ , respectively (see Appendix B for the 4D case).

5.4.  $W_{\hat{a}}^{(1)}(u, x^k) = \text{non-constant}$ ,  $W_{\hat{\alpha}}^{(1)}(u, x^k) = \text{constant}$ . This is a generalisation of the Warped *CSI* spacetimes considered earlier. The metric can be written

$$(72) \quad ds^2 = e^{-2\phi(x^\alpha)} \left[ 2du \left( dv + \widetilde{H}du + \widetilde{W}_a dx^a + \widetilde{W}_\alpha \omega^\alpha \right) + \delta_{ab} dx^a dx^b \right] + \delta_{\alpha\beta} \omega^\alpha \omega^\beta,$$

where

$$(73) \quad \widetilde{W}_a = v\widetilde{W}_a^{(1)}(u, x^a) + \widetilde{W}_a^{(0)}(u, x^a, x^\alpha),$$

$$(74) \quad \widetilde{W}_\alpha = \widetilde{W}_\alpha^{(0)}(u, x^a, x^\alpha),$$

$$(75) \quad \phi = \phi(x^\alpha), \quad \phi_{,\alpha} = \text{constant},$$

$$(76) \quad \widetilde{H} = \frac{v^2}{8}(\widetilde{W}_a^{(1)})(\widetilde{W}^{(1)a}) + v\widetilde{H}^{(1)}(u, x^a, x^\alpha) + \widetilde{H}^{(0)}(u, x^a, x^\alpha).$$

Furthermore, we require that  $\delta_{\alpha\beta} \omega^\alpha \omega^\beta = \widetilde{g}_{\alpha\beta}(x^\sigma) dx^\alpha dx^\beta$  is a Riemannian homogeneous space,  $\mathcal{M}_H$ . The allowed forms of the function  $\phi$  depend on the homogeneous space  $\mathcal{M}_H$ . Topologically, this space is a fibered manifold with base manifold  $\mathcal{M}_H$  and we will require that the fiber is a *VSI* space; in particular, this means that  $\widetilde{W}_a$  satisfy the *VSI* equations (54)-(57) with  $\sigma = a_{ij} = s_{ij} = 0$ . Special cases:

(1) If  $\widetilde{W}_i^{(0)} = \Psi_{,i}$ ,  $\widetilde{W}_i^{(1)} = \widetilde{H}^{(1)} = 0$  then  $(R)_{-1} = 0$ .

(2) If  $\widetilde{W}_i^{(0)} = \widetilde{H}^{(1)} = \widetilde{H}^{(0)} = 0$  then  $(R)_{-1} = (R)_{-2} = 0$ .

The connection  $\nabla$  for the above metric can be decomposed as

$$(77) \quad \nabla = \widetilde{\nabla} - \tau,$$

where  $\tau$  is a  $(2,1)$  tensor (or operator acting in the obvious way) with only non-positive boost-weight components, while  $\widetilde{\nabla}$  has the following property: *There exists a null frame such that the connection coefficients of  $\widetilde{\nabla}$  has the following properties:*

(1) Positive boost weight connection coefficients are all zero.

(2) Zero boost weight connection coefficients are either all constants or are connection coefficients of  $\mathcal{M}_H$ .

Henceforth we will assume that this null frame is chosen. Moreover,  $\tau$  and  $\widetilde{\nabla}$  can be chosen such that

$$(78) \quad \tau_{e_\alpha} X = \tau_X e_\alpha = 0, \quad \widetilde{\nabla}_X g = 0, \quad g(\tau_X Y, Z) + g(Y, \tau_X Z) = 0, \quad \forall X, Y, Z \in TM.$$

Note that the torsion of  $\widetilde{\nabla}_X$  is  $S_X Y = \tau_Y X - \tau_X Y$ .

Let  $R$  and  $\widetilde{R}$  be the curvature tensors to the connections  $\nabla$  and  $\widetilde{\nabla}$  respectively. Then

$$(79) \quad R = \widehat{R} + \widetilde{R},$$

where  $(\widehat{R})_b = 0$  for  $b \geq 0$  while  $(\widetilde{R})_b = 0$  for  $b > 0$ . Moreover, all boost weight 0 components are constants<sup>5</sup>. We also have

**Theorem 5.1.** *For all  $k$ ,  $\nabla^{(k)}R$  has maximal boost order 0 where all boost weight 0 components are constants.*

*Proof.* We will prove the Theorem by induction. Let us in the following denote the projection of a tensor  $T$  onto the vector space spanned by the components of boost weight  $b$  by  $(T)_b$ . The key observation is that the  $(R)_0 = (\widetilde{R})_0$  is of constant curvature over the  $VSI$  fibers. As a result,  $\tau(R)_0 = 0$ . Thus the theorem is true for  $\nabla(R)_0$ . Moreover, by applying the Bianchi identity for  $\nabla R$ , we can show that it is true for  $\nabla(R)_{-1}$ . Hence, the theorem is true for  $k = 1$ .

Assume the theorem is true for  $k$ . Then consider  $\nabla\nabla^{(k)}R$ . Immediately, we have that it is true for  $\widetilde{\nabla}(\nabla^{(k)}R)_0$ . Moreover, the curvature tensor is of constant curvature over the  $VSI$  fibers, so  $\tau(\nabla^{(k)}R)_0 = 0$ ; hence, the theorem is true for  $\nabla(\nabla^{(k)}R)_0$ .

For  $\nabla(\nabla^{(k)}R)_{-1}$  the critical part is the one that raises the boost weight by one. However, by using the identity

$$(80) \quad [\nabla_A, \nabla_B]T_{C_1 \dots C_i \dots C_p} = \sum_{i=1}^p R^E_{C_i AB} T_{C_1 \dots E \dots C_p},$$

recursively, and also the Bianchi identity, we see that it is also true for  $\nabla(\nabla^{(k)}R)_{-1}$ . Hence, by induction, the Theorem follows.  $\square$

These results consequently imply that the 'fibered  $VSI$ ' spacetime given by eq.(72) is a  $CSI$  spacetime. A crucial part of the proof is that  $(R)_0$  is of constant curvature over the  $VSI$  fibers, which implies that  $\tau(\nabla^{(k)}R)_0 = 0$ .

**Theorem 5.2.** *Consider the metric (72). If for all  $k$   $\nabla^{(k)}R$  has maximal boost order 0, where all boost weight 0 components are constants, then there exists a homogenous space  $(\widetilde{\mathcal{M}}, \widetilde{g})$  with the same curvature invariants as the metric (72).*

*Proof.* The proof essentially follows along the same lines as above by noticing that the functions  $\widetilde{W}_i$  and  $\widetilde{H}$  do not contribute to the curvature invariants. The homogeneous space can be obtained by setting all of these to zero.  $\square$

## 6. DISCUSSION

We have obtained a number of general results. We have studied warped product  $CSI$  spacetimes (Theorem 3.1). We have presented the canonical form for the Kundt  $CSI$  metric (Theorem 4.1), and studied higher dimensional Kundt spacetimes in some detail. In sections 3 & 5 we have constructed higher dimensional examples of  $CSI$  spacetimes that arise as warped products and that belong to the Kundt class. In the appendices we shall present all of the 3-dimensional  $VSI$  metrics, explicitly construct the metrics in the 4-dimensional  $CSI$  spacetimes, and we shall establish the canonical higher dimensional  $VSI$  metric.

Motivated by our results and examples we propose the following conjectures:

**Conjecture 6.1** ( $CSI_F$  conjecture). *A spacetime is  $CSI$  iff there exists a null frame in which the Riemann tensor and its derivatives can be brought into one of the following forms: either*

<sup>5</sup>This can be seen by explicit calculation of the curvature tensors.



- (1) *the Riemann tensor and its derivatives are constant, in which case we have a locally homogeneous space, or*
- (2) *the Riemann tensor and its derivatives are of boost order zero with constant boost weight zero components at each order. This implies that Riemann tensor is of type II or less.*

Whereas the proof of the ‘if’ direction is trivial, the ‘only if’ part is significantly more difficult and would require considering second order curvature invariants. We also point out that if we relax  $CSI$  to  $CSI_1$  then in (1) we must now include all 1-curvature homogeneous spacetimes as well, since these naturally imply  $CSI_1$  and are not necessarily homogeneous spaces.

Assuming that the above conjecture is correct and that there exists such a preferred null frame, then

**Conjecture 6.2** ( $CSI_K$  conjecture). *If a spacetime is  $CSI$ , then the spacetime is either locally homogeneous or belongs to the higher dimensional Kundt  $CSI$  class.*

Finally, we have that

**Conjecture 6.3** ( $CSI_R$  conjecture). *If a spacetime is  $CSI$ , then it can be constructed from locally homogeneous spaces and  $VSI$  spacetimes.*

This construction can be done by means of fibering, warping and tensor sums.

From the results above and these conjectures, it is plausible that for  $CSI$  spacetimes that are not locally homogeneous, the Weyl type is  $II$ ,  $III$ ,  $N$  or  $O$ , and that all boost weight zero terms are constant.

We intend to study the general validity of these conjectures in future work. The relationship to curvature homogeneous spacetimes is addressed in [13]. Support for these conjectures in the 4D case is discussed in the next section.

## 7. FOUR-DIMENSIONAL $CSI$

Let us now consider the 4D  $CSI$  spacetimes in more detail. We assume that a  $CSI$  spacetime is *either* of Petrov (P)-type I or Plebanski-Petrov (PP)-type I, *or* of P-type II and PP-type II (branch A or B of Table 1, respectively). In branch A, we explicitly assume that the spacetime is not of P-type II and PP-type II, otherwise we would be in branch B. It is plausible that in branch A the spacetimes must be of P-type I *and* PP-type I (but we have not established this here). Since the spacetime is not of P-type II and PP-type II, it is necessarily completely backsolvable (CB) [11]. Since all of the zeroth order scalar curvature invariants are constant, there exists a frame in which the components of the Riemann tensor are all constant, and the spacetime is curvature homogeneous  $CH_0$  (i.e.,  $CSI_0 \equiv CH_0$ ). If the spacetime is also  $CH_1$ , then it is necessarily locally homogeneous  $H$  [13]. Therefore, if it is not locally homogeneous, it cannot be  $CH_1$ . Finally, by considering differential scalar invariants, we can obtain information on the spin coefficients in the  $CSI$  spacetime. Although a comprehensive analysis of the differential invariants is necessary, a preliminary investigation indicates that  $\tau$ ,  $\rho$  and  $\sigma$  must all be constant and are likely zero. Therefore, in branch A, there are severe constraints on those spacetimes that are not locally homogeneous, and it is plausible that there are no such spacetimes.

In branch B of Table 1, we have that the spacetime is necessarily of P-type II and PP-type II. It then follows from Theorem 7.1 below, that all boost weight zero terms are necessarily constant (i.e., the spacetime is  $CSI_{F,0}$ ). The spacetime is




TABLE 1. 4D  $CSI$ . All terms are defined in the text. A dashed line indicates a conjectured result. Footnotes: (1) this indicates that the spacetime is at least of type  $II$  (i.e., of type  $II$ ,  $III$ ,  $N$  or  $0$ ), (2) spacetimes in this set are described in the text, (3) all of the 4D  $CSI_K$  spacetimes are displayed explicitly in the text.

then either CB, in which case all of the results in branch A apply, or they are NCB and a number of further conditions apply (these conditions are very severe [11]). In either case, there are a number of different classes characterized by their P-type and PP-type (in each case at least of type II), and in each class there are a number of further restrictions. By investigating the differential scalar invariants we then find conditions on the spin coefficients. By considering each class separately, we can then establish that  $\tau = \rho = \sigma = 0$ , and that the spacetime is necessarily  $CSI_K$ . All of these spacetimes are constructed in Appendix B.

The results are summarized in Table 1. In branch A of Table 1, we have established that if the  $CSI$  spacetime is not locally homogeneous, then necessarily it is of P-type I or PP-type I (and does *not* belong to  $CSI_K$ ,  $CSI_F$ , or  $CSI_R$ ), it belongs

to  $CSI_0 \equiv CH_0$ , but not  $CH_1$  and there are a number of further constraints arising from the non-CB conditions and the differential constraints (i.e., constraints on the spin coefficients). This exceptional set is very sparse, and mostly likely empty; but in order to demonstrate this the conjectures in 4D must be proven, utilising a number of 4D invariants.

We have explicitly constructed the 4D  $CSI_F$ ,  $CSI_K$ , and  $CSI_R$ , and established the relationship between these  $CSI$  subsets themselves and with  $CSI \setminus H$ , thereby lending support to the conjectures in the previous section. The metrics of these  $CSI$  spacetimes are described in Appendix B. From this calculation of all of the members of the set  $CSI_F \cap CSI_K$ , we see that they all belong to  $CSI_R$ . That is, we have established the uniqueness of  $CSI_R$  within this particular context. In particular, we have found all  $CSI$  in branch B of Table 1.

Finally, let us consider Petrov (P)-type II and Plebanski-Petrov (PP)-type II Lorentzian  $CSI$  spacetimes. We shall show that if all zeroth order curvature invariants are constant then the boost weight zero curvature scalars,  $\Psi_2$ ,  $\Phi_{11}$  and  $\Phi_{02}$  are constant. For these algebraically special spacetimes the converse follows straightforwardly.

**Theorem 7.1.** *If all zeroth order curvature invariants are constant in Petrov-type II and Plebanski-Petrov-type II Lorentzian  $CSI$  spacetimes, then the boost weight zero curvature scalars,  $\Psi_2$ ,  $\Phi_{11}$  and  $\Phi_{02}$  are constant.*

*Proof.* We make use of the following curvature invariants

(81)

$$w_1 = \Psi_{ABCD}\Psi^{ABCD}, \quad r_1 = \Phi_{AB\dot{A}\dot{B}}\Phi^{AB\dot{A}\dot{B}}, \quad r_2 = \Phi_{AB\dot{A}\dot{B}}\Phi^B{}_{\dot{C}}{}^{\dot{B}}{}_{\dot{C}}\Phi^{CA\dot{C}\dot{A}}.$$

It is worth noting that in general there is an additional independent degree 4 Ricci invariant  $r_3$ ; however, in PP-types II or less there always exists a syzygy for  $r_3$  in terms of  $r_1$  and  $r_2$  (or else it vanishes). This is analogous to the well known syzygy for the Weyl invariants,  $I^3 = 27J^2$ , signifying an algebraically special spacetime.

Assuming first that the Weyl and Ricci canonical frames are aligned, then the non-vanishing curvature scalars are  $\Psi_2$ ,  $\Psi_4 = 1$ ,  $\Phi_{11}$ ,  $\Phi_{20} = \Phi_{02}$ ,  $\Phi_{22} = 1$  and we have  $w_1 = 6\Psi_2^2$ ,  $r_1 = 2\Phi_{02}^2 + 4\Phi_{11}^2$ ,  $r_2 = -6\Phi_{02}^2\Phi_{11}$ . Clearly  $w_1$  constant implies  $\Psi_2$  constant. If  $\Phi_{02}(2\Phi_{11} - \Phi_{02})(2\Phi_{11} + \Phi_{02}) \neq 0$  then  $\Phi_{02}$  and  $\Phi_{11}$  can be expressed in terms of  $r_1$  and  $r_2$ ; therefore, constant  $r_1$  and  $r_2$  gives constant  $\Phi_{11}$  and  $\Phi_{02}$ . If  $\Phi_{02} = 0$  or  $\Phi_{02} = \pm 2\Phi_{11}$  then  $r_1$  can be used to give the same result. In this aligned case we find that constant  $R$ ,  $w_1$ ,  $r_1$  and  $r_2$  implies that all curvature scalars are constant (including the boost weight 0 scalars); therefore, we have a curvature homogeneous spacetime.

In the non-aligned case we refer to Carminati and Zakhary [11] where it was shown that for this PP-type, complete back-solving (CB) of the Carminati-Zakhary (CZ) invariants can be achieved and all curvature scalars can be expressed in terms of zeroth order invariants; thus constant CZ invariants implies constant curvature scalars. In the exceptional case in which  $\Psi_0 = \Psi_1 = 0$  in the Ricci canonical frame, complete back-solvability is not possible since  $\Psi_3$  and  $\Psi_4$  cannot be determined from any zeroth order curvature invariant [11]. However, this exceptional case occurs when the Weyl and Ricci canonical frames are aligned, for P-type II  $\Psi_3 = 0$  and  $\Psi_4 = 1$  and the remaining curvature scalars were shown above to be constant (this is a particular instance of a not completely back-solvable (NCB) case becoming completely back-solvable; see comment in Ref. 3 of [11]).  $\square$

We have consequently also shown that for P-type II and PP-type II spacetimes, if the zeroth order invariants are constant then all curvature scalars are constant. These results support the higher dimensional conjectures discussed in the previous section.

#### ACKNOWLEDGEMENTS

We would like to thank R. Milson, V. Pravda and A. Pravdova for helpful comments. This work was supported by NSERC (AC and NP) and the Killam Foundation (SH).

$\gamma_{312}$  Constraints on  $W_1(u, v, x)$ ,  $W_2(u, v, x)$  and  $H(u, v, x)$

---

= 0 A1) (P-III ;  $\tau = 0$  ; PP-N)

$$W_1 = W_{01}(u, x), W_2 = W_{02}(u, x), H = vh_1(u, x) + h_0(u, x)$$

$\neq 0$  B1) (P-III ;  $\tau \neq 0$  ; PP-N)

$$W_1 = -\frac{2v}{x} + W_{01}(u, x), W_2 = W_{02}(u, x), H = -\frac{v^2}{2x^2} + vh_1(u, x) + h_0(u, x)$$

TABLE 2. Segre type  $\{3\}$  i.e.  $R_{22} \neq 0$  and  $R_{23} \neq 0$ .

$\gamma_{312}$  Constraints on  $W_1(u, v, x)$ ,  $W_2(u, v, x)$  and  $H(u, v, x)$

---

= 0 D1) (P-N ;  $\tau = 0$  ; PP-O) special case of A1)

$$W_1 = xf_1(u) + f_0(u), W_2 = xg_1(u) + g_0(u), H = h_0(u, x)$$

$\neq 0$  F1) (P-III ;  $\tau \neq 0$  ; PP-O) special case of B1)

$$W_1 = -\frac{2v}{x} + W_{01}(u), W_2 = W_{02}(u), H = -\frac{v^2}{2x^2} + \frac{v}{x}W_{01}(u) + h_0(u, x)$$

TABLE 3. Segre type  $\{(21)\}$  i.e.  $R_{22} \neq 0$  and  $R_{23} = 0$ .

#### APPENDIX A. 3-DIMENSIONAL *VSI* METRICS

By analogy with the four dimensional *VSI* spacetimes, we begin by considering a real null frame  $e_i = \{l, n, m\}$  such that the only non-vanishing inner products are  $l^a n_a = 1 = -m^a m_a$ . In three dimensions, the Riemann tensor is equivalent to the Ricci tensor, therefore all zeroth order invariants vanish if

$$(82) \quad R_{ab} = R_{22}l_a l_b + 2R_{23}l_{(a} m_{b)}.$$

We identify  $R_{22} \sim \phi_{22}$  and  $R_{23} \sim \phi_{12} + \phi_{21}$  and we use the n-dimensional version of the *VSI* theorem to conclude that the analogues of  $\kappa$ ,  $\sigma$  and  $\rho$  must vanish. In three dimensions we find that that  $\kappa \sim \gamma_{311}$ ,  $\sigma = \rho \sim \gamma_{313}$ , and  $\tau \sim \gamma_{312}$ , where  $\gamma_{ijk}$  are the Ricci-rotation coefficients. Therefore, we have that a three dimensional spacetime is *VSI* if and only if  $\gamma_{311} = \gamma_{313} = 0$  and has Segre type  $\{3\}$ ,  $\{(21)\}$  or  $\{(111)\}$  with the possibility that the (non-)vanishing of  $\gamma_{312}$  may lead to distinct subclasses.

The method we use to explicitly obtain the three dimensional *VSI* spacetimes involves writing the four dimensional *VSI* spacetimes in terms of real coordinates  $(u, v, x, y)$  then considering the resulting frame when either  $x = \text{const.}$  or  $y = \text{const.}$  We set  $\zeta = (x + iy)/\sqrt{2}$  and  $W = (W_1 + iW_2)/\sqrt{2}$  throughout, and in every case the restriction to  $y = \text{const.}$  yields nothing new, hence the relevant null frame, in coordinates  $(u, v, x)$ , is

$$(83) \quad l = \partial_v, \quad n = \partial_u - [H + \frac{1}{2}(W_1^2 + W_2^2)]\partial_v + W_1\partial_x, \quad m = \partial_x$$

Upon restriction to  $x = \text{const.}$  most of the four dimensional *VSI* spacetimes become either flat or a special case of a null frame given in the following tables. In addition, whenever  $\gamma_{312}$  is nonzero then it is always equal to  $-1/x$ .

---

 $\gamma_{312}$  Constraints on  $W_1(u, v, x)$ ,  $W_2(u, v, x)$  and  $H(u, v, x)$ 


---

$= 0$	I1) (P-III ; $\tau = 0$ ; vacuum) $W_1 = W_{01}(u)$ , $W_2 = W_{02}(u)$ , $H = xh_{01}(u) + h_{02}(u)$
$\neq 0$	L1) (P-N ; $\tau \neq 0$ ; vacuum) $W_1 = -\frac{2v}{x}$ , $W_2 = 0$ , $H = -\frac{v^2}{2x^2} + \sqrt{2}x[xh_{01}(u) + h_{02}(u)]$

TABLE 4. Segre type  $\{(111)\}$  i.e.  $R_{ij} = 0$ .

Tables 2–4 provide a list of the three-dimensional *VSI* spacetimes. Here, we have chosen to characterise the 3D cases according to Segre type. In this regard, the classification in [22] of three dimensional spacetimes is worth noting. In 4D the Segre types are related to the PP types [17]. We have also indicated the corresponding class of four dimensional *VSI* spacetimes (in brackets) from which the 3D solutions were obtained. It is also worth noting that some of the spacetimes given here may be equivalent and so these tables may reduce further. For example, in Table 4 the vanishing of the Ricci tensor implies that all spacetimes listed there are flat; therefore, using the full 3 parameter Lorentz group,  $SO(1, 2)$ , it should be possible to set  $\gamma_{312} = 0$  (i.e., the flat metric should appear in this table). In Tables 2 and 3 it is necessary to consider the frame freedom left after the Ricci tensor has been brought to its canonical form. This can then be used to determine if  $\gamma_{312}$  can be made to vanish; alternatively, it may be an invariant of this left over frame freedom and the metrics would be inequivalent. If  $\gamma_{312}$  can be set to zero, then it still remains to be shown if a coordinate transformation can be found relating the two metrics.

APPENDIX B. 4D *CSI*

We know that (Lorentzian) homogeneous spaces and *VSI* spacetimes are necessarily *CSI* spacetimes. Let us display all 4D spacetimes that are *CSI<sub>F</sub>* and *CSI<sub>K</sub>*. We shall find that all such spacetimes are necessarily in *CSI<sub>R</sub>*.

The 4D spacetimes in  $CSI_F \cap CSI_K$  are as follows.

B.1.  $\mathcal{I}(\text{AdS}_4)$ 

$$(84) \quad ds^2 = e^{-2py} \left[ 2du \left( dv + \tilde{H}du + \tilde{W}_x dx + \tilde{W}_y dy \right) + dx^2 \right] + dy^2,$$

where

$$\begin{aligned} \tilde{W}_x &= v\tilde{W}_x^{(1)}(u, x) + \tilde{W}_x^{(0)}(u, x, y), \\ \tilde{W}_y &= \tilde{W}_y^{(0)}(u, x, y), \\ \tilde{H} &= \frac{v^2}{8}(\tilde{W}_x^{(1)})^2 + v\tilde{H}^{(1)}(u, x, y) + \tilde{H}^{(0)}(u, x, y). \end{aligned}$$

and  $\tilde{W}_x^{(1)}$  satisfy the 3D *VSI* equations (see Appendix A).

Since  $\tilde{W}_x^{(0)}(u, x, y)$  depends on  $y$ , in addition to  $u$  and  $x$ , this metric is fibered. Due to the terms  $e^{-2py}$  and  $dy^2$  in the metric, the spacetime is a warped product. By redefining  $\tilde{W}_x$  in the metric above by omitting the term  $v\tilde{W}_x^{(1)}(u, x)$  (the ‘ $vW$ ’

term) and in  $\tilde{H}$  omitting the term  $\frac{v^2}{8}(\tilde{W}_x^{(1)})^2$  (the ' $v^2H$ ' term), we can rewrite the metric as the tensor sum of the redefined metric (84) and the metric

$$(85) \quad ds^2_+ = e^{-2py} \left[ 2du \left( dv + \frac{v^2}{8}(\tilde{W}_x^{(1)})^2 du + v\tilde{W}_x^{(1)} dx \right) + dx^2 \right] + dy^2,$$

Clearly the metric can be written in terms of the combined operations of fibering, warping and tensor summing, and hence belongs to  $CSI_R$ .

### B.2. $\mathcal{I}(\text{Sol})$

$$(86) \quad ds^2 = e^{-2py} \left[ 2du \left( dv + \tilde{H} du + \tilde{W}_x dx + \tilde{W}_y dy \right) \right] + e^{-2qy} dx^2 + dy^2,$$

where

$$\begin{aligned} \tilde{W}_x &= \tilde{W}_x^{(0)}(u, x, y), & \tilde{W}_y &= \tilde{W}_y^{(0)}(u, x, y), \\ \tilde{H} &= v\tilde{H}^{(1)}(u, x, y) + \tilde{H}^{(0)}(u, x, y). \end{aligned}$$

This metric is fibered and warped.

### B.3. $\mathcal{I}(M_1 \times M_2)$ :

$$(87) \quad ds^2 = 2du \left( dv + H du + W_i \mathbf{m}^i \right) + \delta_{ij} \mathbf{m}^i \mathbf{m}^j,$$

where

$$\begin{aligned} W_i &= vW_i^{(1)} + \tilde{W}_i^{(0)}(u, x, y), \\ H &= v^2H^{(2)} + v\tilde{H}^{(1)}(u, x, y) + \tilde{H}^{(0)}(u, x, y). \end{aligned}$$

and  $W_i^{(1)}$  and  $H^{(2)}$  are given in the table below. This metric is fibered and warped, and due to the ' $vW$ ' and ' $v^2H$ ' terms, can be written as a tensor sum.

$\delta_{ij} \mathbf{m}^i \mathbf{m}^j$	$dx^2 + dy^2$	$a^2(dx^2 + \sin^2 x dy^2)$	$dx^2 + e^{-2qx} dy^2$
$W_i^{(1)} \mathbf{m}^i$	$\alpha dx + \beta dy$	0	$\alpha dx + \beta e^{-qx} dy$
$H^{(2)}$	$\frac{1}{8}(4\sigma + \alpha^2 + \beta^2)$	$\frac{\sigma}{2}$	$\frac{1}{8}(4\sigma + \alpha^2 + \beta^2)$

B.4.  $\mathcal{I}((A)dS_3 \times \mathbb{R})$ : Metrics of the form (66), (68) and (70). These metrics are fibered, warped and tensor summed.

B.5.  $\mathcal{I}((A)dS_4)$ : Metrics of the form (66), (68) and (70). In addition, for  $\mathcal{I}(dS_4)$

$$(88) \quad ds^2 = \cos^2(\sqrt{\sigma}y) \cos^2(\sqrt{\sigma}x) \left[ 2du \left( dv + \tilde{H} du + \tilde{W}_i \mathbf{m}^i \right) \right] + \cos^2(\sqrt{\sigma}y) dx^2 + dy^2,$$

where

$$(89) \quad \tilde{W}_i = \tilde{W}_i^{(0)}(u, x, y), \quad \tilde{H} = \frac{v^2}{2}\sigma + v\tilde{H}^{(1)}(u, x, y) + \tilde{H}^{(0)}(u, x, y).$$

and in the case of  $\mathcal{I}(AdS_4)$ , similar versions of the metrics (68) and (70) are obtained. These metrics are fibered, warped and tensor summed.

All of the metrics in sections B.1-B.5 can be written in terms of the combined operations of fibering, warping and tensor summing. Therefore, all of the spacetimes constructed belong to  $CSI_R$ . We consequently have the result that in 4D  $CSI_F$ ,  $CSI_K$  and  $CSI_R$  are equivalent. It is plausible that a similar result applies in higher dimensions.

Finally, let us consider the tensor sum in a little more detail. Let us consider the  $CSI_R$  metric (66). Redefining the metric by omitting the ' $v^2H$ ' term, it can be rewritten as the 'tensor sum' of the redefined metric and

$$(90) \quad ds^2_+ = \left[ 2du \left( dv + \frac{v^2}{2}\sigma du + 2\sqrt{\sigma} \tan(\sqrt{\sigma}x) dx \right) \right] + dx^2.$$

The term  $\left[ 2du \left( dv + \frac{v^2}{2}\sigma du \right) \right]$  can be rewritten as  $\left[ \left( 1 + \frac{\sigma}{2}UV \right)^{-2} dU dV \right]$ , which is a metric of constant Ricci scalar curvature.

It is plausible that we have constructed all of the 4D  $CSI$  spacetimes here. To prove this, we must prove the conjectures outlined above. This is perhaps best done by considering higher order differential scalar invariants in 4D [2], which we hope to do in a future paper.

### APPENDIX C. VSI METRICS

We can now write down the metric for higher dimensional VSI spacetimes in a canonical form. An immediate corollary of Theorem 4.1 is:

**Corollary C.1.** *Any VSI metric can be written*

$$(91) \quad ds^2 = 2du \left[ dv + H(v, u, x^k) du + W_i(v, u, x^k) dx^i \right] + \delta_{ij} dx^i dx^j.$$

*Proof.* In [15] it was proven that all VSI metrics have  $L_{ij} = 0$  and  $R_{\hat{i}\hat{j}\hat{k}\hat{l}} = 0$ . The first condition implies that the metric is Kundt, hence of the form (10), while the second implies that the spatial metric,  $\tilde{g}_{ij}$ , is flat. Using Theorem 4.1 the corollary now follows.  $\square$

We are now in a position to determine the explicit metric forms for higher dimensional VSI spacetimes [21], which we shall return to in future work.



## REFERENCES

- [1] F. Prüfer, F. Tricerri, L. Vanhecke, 1996, *Trans. Am. Math. Soc.* **348**, 4643.
- [2] V. Pravda, A. Pravdová, A. Coley and R. Milson,, 2002, *Class. Q. Grav.* **19**, 6213 [gr-qc/0209024].
- [3] J. Bicak and J. Podolsky, 1999, *J. Math. Phys.* **40**, 4495; J. Podolsky and M. Ortaggio, 2003, *Class. Q. Grav.* **20**, 1685 & J. B. Griffiths, P. Docherty and J. Podolsky, 2004, *Class. Q. Grav.* **21**, 207.
- [4] Y. N. Obukhov, 2004, *Phys. Rev. D* **69**, 024013.
- [5] S.T.C. Siklos, 1985, *Lobatchevski plane wave gravitational waves, in Galaxies, Axisymmetric systems and Relativity*, ed. M. MacCallum (Cambridge University Press, Cambridge).
- [6] H. -J. Schmidt, 1998, *Int. J. Th. Phys.* **37**, 1691.
- [7] S. Hervik, 2004, *Class. Q. Grav.* **21**, 4273.
- [8] N. Pelavas, A. Coley, R. Milson, V. Pravda and A. Pravdova, 2005, *J. Math. Phys.* **46** 063501 [gr-qc/0503040].
- [9] A. Coley, R. Milson, V. Pravda and A. Pravdova, 2004, *Class. Q. Grav.* **21**, L35 [gr-qc/0401008].
- [10] R. Milson, A. Coley, V. Pravda and A. Pravdova, 2005, *Int. J. Geom. Meth. Mod. Phys.* **2**, 41 [gr-qc/0401010].
- [11] J. Carminati and E. Zakhary, 2002, *J. Math. Phys.* **43**, 4020.
- [12] G.S. Hall, 2003, *Symmetries and Curvature Structure in General Relativity* (World Scientific, Singapore).
- [13] R. Milson and N. Pelavas, 2006; preprint.
- [14] I. M. Singer, 1960, *Comm. Pure and App. Math.* **13**, 685.
- [15] A. Coley, R. Milson, V. Pravda and A. Pravdova, 2004, *Class. Q. Grav.* **21**, 5519.
- [16] F. Podesta and A. Spiro, 1995, preprint *Introduzione ai gruppi transformation*.
- [17] H. Stephani, D. Kramer, M. MacCallum and E. Herlt, 2003, *Exact Solutions to Einstein's Field Equations*, (Second Edition, Cambridge Univ. Press).
- [18] A. Z. Petrov, 1969, *Einstein Spaces* (Pergamon Press).
- [19] V. Pravda, A. Pravdová, A. Coley and R. Milson, 2004, *Class. Quant. Grav.* **21**, 2873 [gr-qc/0401013].
- [20] J. Milnor, 1976, *Adv. Math.* **21** 293
- [21] A. Coley, R. Milson, V. Pravda, A. Pravdova, N. Pelavas and R. Zaleletdinov 2003, *Phys. Rev. D.* **67**, 104020 [gr-qc/0212063].
- [22] G.S. Hall and S. Capocci, 1999, *J. Math. Phys.* **40**, 1466

DEPARTMENT OF MATHEMATICS AND STATISTICS, DALHOUSIE UNIVERSITY, HALIFAX, NOVA SCOTIA, CANADA B3H 3J5

*E-mail address:* aac@mathstat.dal.ca, herviks@mathstat.dal.ca, pelavas@mathstat.dal.ca