ON THE RELATION BETWEEN ADM AND BONDI ENERGY-MOMENTA III – PERTURBED RADIATIVE SPATIAL INFINITY

WEN-LING HUANG AND XIAO ZHANG

ABSTRACT. In a vacuum spacetime equipped with the Bondi's radiating metric which is asymptotically flat at spatial infinity including gravitational radiation (**Condition D**), we establish the relation between the ADM total energy-momentum and the Bondi energy-momentum for perturbed radiative spatial infinity. The perturbation is given by defining the "real" time the sum of the retarded time, the Euclidean distance and certain function f.

1. INTRODUCTION

It is a fundamental problem in gravitational radiation what the relation is between the ADM energy-momentum and the Bondi energy-momentum. Under certain asymptotic flatness conditions at spatial infinity, it was shown the ADM energy-momentum at spatial infinity is the past limit of the Bondi energy-momentum in certain vacuum spacetimes [2, 7, 10, 11, 4, 13]. However, it is presumably believed the assumed asymptotic flatness at spatial infinity in all above works precludes gravitational radiation, at least near spatial infinity.

In [13], the second author introduced a weaker assumption of asymptotic flatness at spatial infinity in Bondi's radiating spacetimes (**Condition D**). This condition can be viewed as Sommerfeld's radiation condition at spatial infinity [13]. And it should not preclude gravitational radiation. Under this condition, it is found that the ADM total energy of a *t*-slice is no longer the past limit of the Bondi mass, and they differ by a quantity related to the *news functions*. In [8], the authors established the relation between the ADM total linear momentum and the past limit of the Bondi "momentum" under **Condition D**. It is surprising that in this case the second fundamental form h on the *t*-slice falls off as $O(\frac{1}{r})$, however, some nice cancellation occurs and the ADM total linear momentum is still finite. The ADM total linear momentum at spatial infinity is no longer the past limit of the Bondi "momentum", and the difference is calculated.

The definition of the "real" time t is essential in Bondi's radiating spacetimes and the spatial infinity is defined as a t slice. In the works of [13, 8],

Key words and phrases. Gravitational radiation, ADM energy-momentum, Bondi energy-momentum .

the "real" time t is defined as the sum of the retarded coordinate u and the Euclidean distance r. However, this definition is rather restricted, it does not hold true even in the Schwarzschild spacetime. In this paper, we will perturb the "real" time t by adding certain function f. The spatial infinity is also defined in terms of this new t-slice. We establish the relation between the ADM energy-momentum and the Bondi energy-momentum under **Condition D**.

We do not consider the polyhomogeneous Bondi expansions in the present paper, as studied in [6]. In [6] (Appendix D), the logarithmic singularities at null infinity can be removed in the axisymmetric case if the free function $\gamma_2(u, x^{\alpha})$ is chosen to be zero and $\gamma_{3,1}(u_0, x^{\alpha})$ is chosen to be zero for some u_0 . We believe this is a generic property that under certain class of Bondi's radiating metrics which make the ADM total energy-momentum and the Bondi energy-momentum well-defined, the standard Bondi expansions without any logarithmic singularity at null infinity can always be achieved.

The paper is organized as follows: In Section 2, we state some well-known results of Bondi, van der Burg, Metzner and Sachs. In Section 3, we define the "real" time t via adding a perturbation f. We study the asymptotic behaviors of the induced metric and the the second fundamental form of a tslice. In Section 4, we derive the difference between the ADM total energy and the past limit of the Bondi mass. Under certain conditions, we can find a good perturbation such that the ADM total energy equals the past limit of the Bondi mass. In Section 5, we derive the difference between the ADM total linear momentum and the past limit of the Bondi "momentum". We also find certain condition ensuring the existence of good perturbation such that the ADM total linear momentum equals the past limit of the Bondi "momentum". However, we find, in general, it is impossible to perturb spatial infinity so that both the ADM total energy and the ADM total linear momentum are the past limit of the Bondi energy-momentum.

2. Bondi's radiating vacuum spacetimes

The Bondi's radiating vacuum spacetime $(L^{3,1}, \tilde{g})$ is a vacuum spacetime equipped with the following metric $\tilde{g} = \tilde{g}_{ij} dx^i dx^j$

$$\tilde{g} = \left(\frac{V}{r}e^{2\beta} + r^2e^{2\gamma}U^2\cosh 2\delta + r^2e^{-2\gamma}W^2\cosh 2\delta + 2r^2UW\sinh 2\delta\right)du^2 - 2e^{2\beta}dudr \\ -2r^2\left(e^{2\gamma}U\cosh 2\delta + W\sinh 2\delta\right)dud\theta \\ -2r^2\left(e^{-2\gamma}W\cosh 2\delta + U\sinh 2\delta\right)\sin\theta dud\psi \\ +r^2\left(e^{2\gamma}\cosh 2\delta d\theta^2 + e^{-2\gamma}\cosh 2\delta\sin^2\theta d\psi^2 + 2\sinh 2\delta\sin\theta d\theta d\psi\right),$$
(2.1)

where $\beta, \gamma, \delta, U, V, W$ are functions of

$$x^0 = u, \ x^1 = r, \ x^2 = \theta, \ x^3 = \psi,$$

u is a retarded coordinate, *r* is Euclidean distance, θ and ψ are spherical coordinates, $0 \le \theta \le \pi$, $0 \le \psi \le 2\pi$. We assume that \tilde{g} satisfies the outgoing radiation condition.

The metric (2.1) was studied by Bondi, van der Burg, Metzner and Sachs in the theory of gravitational waves in general relativity [3, 9, 12]. They proved that the following asymptotic behavior holds for r sufficiently large if the spacetime satisfies the outgoing radiation condition [12]

$$\begin{split} \gamma &= \frac{c(u,\theta,\psi)}{r} + \frac{C(u,\theta,\psi) - \frac{1}{6}c^3 - \frac{3}{2}cd^2}{r^3} + O\Big(\frac{1}{r^4}\Big), \\ \delta &= \frac{d(u,\theta,\psi)}{r} + \frac{H(u,\theta,\psi) + \frac{1}{2}c^2d - \frac{1}{6}d^3}{r^3} + O\Big(\frac{1}{r^4}\Big), \\ \beta &= -\frac{c^2 + d^2}{4r^2} + O\Big(\frac{1}{r^4}\Big), \\ U &= -\frac{l(u,\theta,\psi)}{r^2} + \frac{p(u,\theta,\psi)}{r^3} + O\Big(\frac{1}{r^4}\Big), \\ W &= -\frac{\bar{l}(u,\theta,\psi)}{r^2} + \frac{\bar{p}(u,\theta,\psi)}{r^3} + O\Big(\frac{1}{r^4}\Big), \\ V &= -r + 2M(u,\theta,\psi) + \frac{\bar{M}(u,\theta,\psi)}{r} + O\Big(\frac{1}{r^2}\Big), \end{split}$$

where

$$\begin{split} l &= c_{,2} + 2c\cot\theta + d_{,3}\csc\theta, \\ \bar{l} &= d_{,2} + 2d\cot\theta - c_{,3}\csc\theta, \\ p &= 2N + 3(cc_{,2} + dd_{,2}) + 4(c^2 + d^2)\cot\theta \\ &- 2(c_{,3}d - cd_{,3})\csc\theta, \\ \bar{p} &= 2P + 2(c_{,2}d - cd_{,2}) + 3(cc_{,3} + dd_{,3})\csc\theta, \\ \bar{M} &= N_{,2} + \cot\theta + P_{,3}\csc\theta - \frac{c^2 + d^2}{2} \\ &- [(c_{,2})^2 + (d_{,2})^2] - 4(cc_{,2} + dd_{,2})\cot\theta \\ &- 4(c^2 + d^2)\cot^2\theta - [(c_{,3})^2 + (d_{,3})^2]\csc^2\theta \\ &+ 4(c_{,3}d - cd_{,3})\csc\theta\cot\theta + 2(c_{,3}d_{,2} - c_{,2}d_{,3})\csc\theta. \end{split}$$

(We denote $f_{,\nu} = \frac{\partial f}{\partial x^{\nu}}$ for $\nu = 0, 1, 2, 3$ throughout the paper.) M is the mass aspect and $c_{,0}, d_{,0}$ are the news functions and they satisfy the following equation [12]:

$$M_{,0} = -\left[(c_{,0})^2 + (d_{,0})^2 \right] + \frac{1}{2} \left(l_{,2} + l \cot \theta + \bar{l}_{,3} \csc \theta \right)_{,0}.$$
 (2.2)

Let N_{u_0} be a null hypersurface which is given by $u = u_0$ at null infinity. The Bondi energy-momentum of N_{u_0} is defined by [3]:

$$m_{\nu}(u_0) = \frac{1}{4\pi} \int_{S^2} M(u_0, \theta, \psi) n^{\nu} dS$$

where $\nu = 0, 1, 2, 3, S^2$ is the unit sphere,

$$n^0 = 1$$
, $n^1 = \sin \theta \cos \psi$, $n^2 = \sin \theta \sin \psi$, $n^3 = \cos \theta$.

And m_0 is the Bondi mass, m_i is the Bondi momentum.

Denote by $\{\check{e}^i\}$ the coframe of the standard flat metric g_0 on \mathbb{R}^3 in polar coordinates,

$$\breve{e}^1 = dr, \ \breve{e}^2 = r d\theta, \ \breve{e}^3 = r \sin \theta d\psi.$$

Denote by $\{\check{e}_i\}$ the dual frame (i = 1, 2, 3). The connection 1-form $\{\check{\omega}_{ij}\}$ is defined by

$$d\breve{e}^i = -\breve{\omega}_{ij} \wedge \breve{e}^j.$$

It is easy to find that

$$\breve{\omega}_{12} = -\frac{1}{r}\breve{e}^2, \ \ \breve{\omega}_{13} = -\frac{1}{r}\breve{e}^3, \ \ \breve{\omega}_{23} = -\frac{\cot\theta}{r}\breve{e}^3.$$

The Levi-Civita connection $\breve{\nabla}$ of g_0 is given by

$$\breve{\nabla}\breve{e}_i = -\breve{\omega}_{ij}\otimes\breve{e}_j.$$

We denote $\breve{\nabla}_i \equiv \breve{\nabla}_{\breve{e}_i}$ for i = 1, 2, 3 throughout the paper.

Define $C_{\{a_1,a_2,a_3\}}$ the space of smooth functions in the spacetime which satisfies the following asymptotic behavior at spatial infinity

$$\mathcal{C}_{\{a_1,a_2,a_3\}} = \begin{cases} \lim_{r \to \infty} \lim_{u \to -\infty} r^{a_1} f = O(1), \\ f: \quad \lim_{r \to \infty} \lim_{u \to -\infty} r^{a_2} \breve{\nabla}_i f = O(1), \\ \lim_{r \to \infty} \lim_{u \to -\infty} r^{a_3} \breve{\nabla}_i \breve{\nabla}_j f = O(1) \end{cases} \right\}.$$
(2.3)

In [13], the following four conditions were introduced:

Condition A: Each of the six functions β , γ , δ , U, V, W together with its derivatives up to the second orders are equal at $\psi = 0$ and 2π .

Condition B: For all u,

$$\begin{split} & \int_{0}^{2\pi} c(u,0,\psi)d\psi = 0, \ \int_{0}^{2\pi} c(u,\pi,\psi)d\psi = 0.\\ \text{Condition C:} \quad \gamma \in \mathcal{C}_{\{1,2,3\}}, \quad \delta \in \mathcal{C}_{\{1,2,3\}}, \quad \beta \in \mathcal{C}_{\{2,3,4\}}, \quad U \in \mathcal{C}_{\{2,3,4\}}, \quad W \in \mathcal{C}_{\{2,3,4\}}, \quad V + r \in \mathcal{C}_{\{0,1,2\}}.\\ \text{Condition D:} \quad \gamma \in \mathcal{C}_{\{1,1,1\}}, \quad \delta \in \mathcal{C}_{\{1,1,1\}}, \quad \beta \in \mathcal{C}_{\{2,2,2\}}, \quad U \in \mathcal{C}_{\{1,2,3\}}, \quad U \in \mathcal{C}_{\{1,3,3\}}, \quad U \in \mathcal{C}$$

Condition D:
$$\gamma \in C_{\{1,1,1\}}, \quad \delta \in C_{\{1,1,1\}}, \quad \beta \in C_{\{2,2,2\}}, \quad U \in C_{\{2,2,2\}}, \quad W \in C_{\{2,2,2\}}, \quad V + r \in C_{\{0,0,0\}}.$$

Condition A and Condition B ensure that the metric (2.1) is regular, also ensure the following Bondi mass loss formula

$$\frac{d}{du}m_{\nu} = -\frac{1}{4\pi}\int_{S^2} \left[(c_{,0})^2 + (d_{,0})^2 \right] n^{\nu} dS.$$

Condition C ensures the Schoen-Yau's positive mass theorem at spatial infinity. However, it precludes gravitational radiation. **Condition D** should include gravitational radiation. It indicates that, for r sufficiently large,

$$\lim_{u \to -\infty} \left\{ M, c, d, M_{,0}, c_{,0}, d_{,0}, M_{,A}, c_{,A}, d_{,A} \right\} = O(1)$$

where $2 \leq A \leq 3$. We refer to [13] for some physical interpretation of **Condition D**.

3. Initial data sets

From now on, we assume the "real" time t is defined as

$$t = u + r + f(r, \theta, \psi) \tag{3.1}$$

for r sufficiently large, where f is smooth function which has the following asymptotic behavior

$$f = a_1 \ln r + a_2(\theta, \psi) + a_3(r, \theta, \psi)$$
(3.2)

for r sufficiently large, where a_1 is constant, a_2 , a_3 are smooth functions which satisfy

$$a_i|_{\psi=0} = a_i|_{\psi=2\pi}, \ a_{i,A}|_{\psi=0} = a_{i,A}|_{\psi=2\pi}, \ a_{i,AB}|_{\psi=0} = a_{i,AB}|_{\psi=2\pi}$$

for i = 2, 3, A, B = 2, 3. Moreover,

$$a_3 = O\left(\frac{1}{r}\right), \quad \breve{\nabla}_k a_3 = O\left(\frac{1}{r^2}\right), \quad \breve{\nabla}_l \breve{\nabla}_k a_3 = O\left(\frac{1}{r^3}\right).$$

for r sufficiently large.

Substituting (3.1) into (2.1), we obtain the spacetime metric

$$\tilde{g} = \tilde{g}_{tt}dt^2 + 2\tilde{g}_{ti}dtdx^i + \tilde{g}_{ij}dx^i dx^j$$

 $(1 \leq i, j \leq 3)$. An initial data set (N_{t_0}, g, h) is a spacelike hypersurface in $L^{3,1}$ which is given by $\{t = t_0\}$. Here g is the induced metric of \tilde{g} and h is the second fundamental form. The lapse \mathcal{N} and the shift X_i (i = 1, 2, 3) of the spacelike hypersurface N_{t_0} are

$$\mathcal{N} = \left(-\tilde{g}^{tt}\right)^{-\frac{1}{2}} = \left(-\tilde{g}_{tt} + \tilde{g}_{ti}\tilde{g}_{tj}g^{ij}\right)^{\frac{1}{2}},$$

$$X_i = \tilde{g}_{ti}.$$

The second fundamental form is then given by

$$h_{ij} = \frac{1}{2\mathcal{N}} \Big(\nabla_i X_j + \nabla_j X_i - \partial_t \tilde{g}_{ij} \Big)_{t=t_0}$$

With the help of asymptotic behavior of β , γ , δ , U, V, W, f and Mathematica 5.0, we obtain the asymptotic expansion of g_{ij} and h_{ij} :

$$\begin{array}{rcl} g_{11} &=& 1+\frac{2M}{r}+\frac{1}{r^2}\Big[-\frac{c^2}{2}-\frac{d^2}{2}+l^2+\bar{l}^2+\bar{M}-a_1^2-4Ma_1\Big]\\ &+O\Big(\frac{1}{r^3}\Big), \\ g_{22} &=& r^2+2rc+2c^2+2d^2+2la_{2,2}-a_{2,2}^2+O\Big(\frac{1}{r}\Big), \\ g_{33} &=& \Big(r^2-2rc+2c^2+2d^2\Big)\sin^2\theta+2\bar{l}\sin\theta a_{2,3}-a_{2,3}^2+O\Big(\frac{1}{r}\Big), \\ g_{12} &=& -l+\frac{1}{r}\Big(-2cl-2d\bar{l}+p+a_1l-a_1a_{2,2}-2Ma_{2,2}\Big)+O\Big(\frac{1}{r^2}\Big), \\ g_{13} &=& -\bar{l}\sin\theta+\frac{1}{r}\Big[\Big(2c\bar{l}-2dl+\bar{p}\Big)\sin\theta+\bar{l}\sin\theta a_1-a_1a_{2,3}-2Ma_{2,3}\Big] \\ &+O\Big(\frac{1}{r^2}\Big), \\ g_{23} &=& 2rd\sin\theta+la_{2,3}+\bar{l}\sin\theta a_{2,2}-a_{2,2}a_{2,3}+O\Big(\frac{1}{r}\Big), \\ h_{11} &=& \frac{M_0}{r}+\frac{1}{r^2}\Big(2M-MM_0+\frac{cc_0}{2}+\frac{dd_0}{2}+ll_0+\frac{\bar{M}_0}{2} \\ &+a_1-3M_0a_1+\bar{l}_0\csc\theta a_{2,3}\Big)+O\Big(\frac{1}{r^3}\Big), \\ h_{22} &=& -rc_0-2M+Mc_0-2cc_0-2dd_0+l_{,2}-a_1-a_{2,22}+c_0a_1 \\ &+O\Big(\frac{1}{r}\Big), \\ h_{33} &=& \Big(rc_0-2M-Mc_0-2cc_0-2dd_0+l\cot\theta+\bar{l}_3\csc\theta\Big)\sin^2\theta \\ &-\sin^2\theta a_1-a_{2,33}-\sin\theta\cos\theta a_{2,2}-c_0\sin^2\theta a_1+O\Big(\frac{1}{r}\Big), \\ h_{12} &=& \frac{1}{r}\Big[-M_{,2}-l+c_0l+d_0\bar{l}-(c_0+M_0-1)a_{2,2}-d_0\csc\theta a_{2,3}\Big] \\ &+O\Big(\frac{1}{r^2}\Big), \\ h_{13} &=& \frac{1}{r}\Big[\Big(-M_3\csc\theta-\bar{l}-c_0\bar{l}+d_0l\Big)\sin\theta+(c_0-M_0+1)a_{2,3} \\ &-d_0\sin\theta a_{2,2}\Big]+O\Big(\frac{1}{r^2}\Big), \\ h_{23} &=& \Big[-rd_0+Md_0+\frac{1}{2}(\bar{l}_2-\bar{l}\cot\theta+l_3\csc\theta)\Big]\sin\theta-a_{2,23} \\ &+d_0\sin\theta a_1+\cot\theta a_{2,3}+O\Big(\frac{1}{r}\Big). \\ \end{array}$$

6

The trace of the second fundamental form is

$$tr_g(h) = \frac{M_{,0}}{r} + \frac{1}{r^2} \Big[-2M - 3MM_{,0} + l_{,2} + l\cot\theta + \bar{l}_{,3}\csc\theta \\ + \frac{cc_{,0}}{2} + \frac{dd_{,0}}{2} + ll_{,0} + \frac{\bar{M}_{,0}}{2} - (a_1 + 3M_{,0}a_1 + a_{2,22} \\ + \csc^2\theta a_{2,33} + \cot\theta a_{2,2}) + \bar{l}_{,0}\csc\theta a_{2,3} \Big] + O\Big(\frac{1}{r^3}\Big).$$

4. ADM and Bondi total energy

In [13, 8], the authors derived the relation between the ADM total energy, the total linear momentum and the Bondi energy-momentum under **Condition A**, **Condition B** and **Condition D** and under the assumption t = u + r. In this section, we study the relation between them under these conditions and under (3.1) with condition (3.2). Let Euclidean coordinates

$$y^1 = r \sin \theta \cos \psi, \quad y^2 = r \sin \theta \sin \psi, \quad y^3 = r \cos \theta.$$

In polar coordinates, the ADM total energy \mathbb{E} and the ADM total linear momentum \mathbb{P}_k are [1, 13, 8]

$$\mathbb{E} = \frac{1}{16\pi} \lim_{r \to \infty} \int_{S_r} \left[\breve{\nabla}^j g(\breve{e}_1, \breve{e}_j) - \breve{\nabla}_1 tr_{g_0}(g) \right] \breve{e}^2 \wedge \breve{e}^3,$$

$$\mathbb{P}_k = \frac{1}{8\pi} \lim_{r \to \infty} \int_{S_r} \left[h\left(\frac{\partial}{\partial y^k}, \frac{\partial}{\partial r}\right) - g\left(\frac{\partial}{\partial y^k}, \frac{\partial}{\partial r}\right) tr_g(h) \right] \breve{e}^2 \wedge \breve{e}^3.$$

Theorem 4.1. Let $\mathbb{E}(t_0)$ be the ADM total energy of the initial data set (N_{t_0}, g, h) where t is given by (3.1) with condition (3.2). Under Condition A, Condition B and Condition D, we have

$$\mathbb{E}(t_0) = m_0(-\infty) + \frac{1}{2\pi} \lim_{u \to -\infty} \int_0^{\pi} \int_0^{2\pi} (cc_{,0} + dd_{,0}) \sin\theta d\psi d\theta + \frac{1}{16\pi} \lim_{u \to -\infty} \int_0^{\pi} \int_0^{2\pi} (a_{2,2} \sin\theta l_{,0} + a_{2,3}\bar{l}_{,0}) d\psi d\theta.$$

Proof : Under Condition A, Condition B and Condition D, we have, as $r \to \infty$, (or $u \to -\infty$),

$$\begin{split} \breve{\nabla}^{j}g(\breve{e}_{1},\breve{e}_{j}) - \breve{\nabla}_{1}tr_{g_{0}}(g) &= \breve{e}_{j}\Big(g(\breve{e}_{1},\breve{e}_{j})\Big) - \breve{e}_{1}tr_{g_{0}}(g) \\ &-g(\breve{e}_{j},\breve{e}_{i})\breve{\omega}_{i1}(\breve{e}_{j}) - g(\breve{e}_{1},\breve{e}_{i})\breve{\omega}_{ij}(\breve{e}_{j}) \\ &= \frac{4M}{r^{2}} - \frac{l_{,2}}{r^{2}} - \frac{l\cot\theta}{r^{2}} - \frac{\bar{l}_{,3}}{r^{2}\sin\theta} - 4\breve{e}_{1}\Big(\frac{c^{2}+d^{2}}{r^{2}}\Big) \\ &+ \frac{a_{2,2}\,l_{,0} + a_{2,3}\csc\theta\bar{l}_{,0}}{r^{2}} + O\Big(\frac{1}{r^{3}}\Big) \\ &= \frac{4M}{r^{2}} - \frac{l_{,2}}{r^{2}} - \frac{l\cot\theta}{r^{2}} - \frac{\bar{l}_{,3}}{r^{2}\sin\theta} + \frac{4(c^{2}+d^{2})_{,0}}{r^{2}} \\ &+ \frac{a_{2,2}\,l_{,0} + a_{2,3}\csc\theta\bar{l}_{,0}}{r^{2}} + O\Big(\frac{1}{r^{3}}\Big). \end{split}$$

Therefore the theorem is a direct consequence of integrating it over S_r and using that for fixed $t = t_0, r \to \infty$ is equivalent to $u \to -\infty$. Q.E.D.

Remark 4.1. If a_2 is chosen to be a constant, the difference $\mathbb{E}(t_0) - m_0(-\infty)$ is independent on the choice of f, which is invariant in the perturbed class (3.2) that a_1 , a_2 are constant.

The following theorem indicates that, under certain conditions, we can choose suitable f such that, at spatial infinity defined by (3.2), the ADM total energy is the past limit of the Bondi mass.

Theorem 4.2. Suppose that Condition A, Condition B and Condition D hold in the Bondi's radiating spacetime (2.1). If

$$\lim_{u \to -\infty} c_{,0} \big|_{\theta=0} = \lim_{u \to -\infty} c_{,0} \big|_{\theta=\pi} = 0,$$
(4.1)

$$\lim_{u \to -\infty} d_{,30} \big|_{\theta=0} = \lim_{u \to -\infty} d_{,30} \big|_{\theta=\pi} = 0,$$
(4.2)

$$\lim_{u \to -\infty} \left[M_{,0} + (c_{,0})^2 + (d_{,0})^2 \right] \neq 0,$$
(4.3)

then there exists f_0 such that

$$\mathbb{E}_{f_0}(t_0) = m_0(-\infty)$$

where $\mathbb{E}_{f_0}(t_0)$ is the ADM total energy of the initial data set (N_{t_0}, g, h) where t is given by (3.1) and $f = f_0$ given by (3.2) with

$$a_2 = \lim_{u \to -\infty} \frac{4cc_{,0} + 4dd_{,0}}{M_{,0} + (c_{,0})^2 + (d_{,0})^2}.$$
(4.4)

Proof : Note that

$$\begin{split} \lim_{u \to -\infty} \int_{0}^{\pi} a_{2,2} \sin \theta l_{,0} d\theta &= \lim_{u \to -\infty} a_{2} \big(c_{,20} \sin \theta + 2c_{,0} \cos \theta + d_{,30} \big) \Big|_{0}^{\pi} \\ &- \lim_{u \to -\infty} \int_{0}^{\pi} a_{2} \big(\sin \theta l_{,0} \big)_{,2} d\psi d\theta \\ &= -\lim_{u \to -\infty} \int_{0}^{\pi} a_{2} \big(\cos \theta l_{,0} + \sin \theta l_{,02} \big) d\theta, \\ \lim_{u \to -\infty} \int_{0}^{2\pi} a_{2,3} \bar{l}_{,0} d\psi &= \lim_{u \to -\infty} a_{2} \big(d_{,20} + 2d_{,0} \cot \theta - c_{,30} \csc \theta \big) \Big|_{0}^{2\pi} \\ &- \lim_{u \to -\infty} \int_{0}^{2\pi} a_{2} \bar{l}_{,03} d\psi \\ &= -\lim_{u \to -\infty} \int_{0}^{2\pi} a_{2} \bar{l}_{,03} d\psi. \end{split}$$

We obtain

$$\lim_{u \to -\infty} \int_0^{\pi} \int_0^{2\pi} \left(a_{2,2} \sin \theta \, l_{,0} + a_{2,3} \bar{l}_{,0} \right) d\psi d\theta$$

= $-\lim_{u \to -\infty} \int_0^{\pi} \int_0^{2\pi} a_2 \left(l_{,2} + l \cot \theta + \bar{l}_{,3} \csc \theta \right)_{,0} \sin \theta d\psi d\theta$
= $-2 \lim_{u \to -\infty} \int_0^{\pi} \int_0^{2\pi} a_2 \left[M_{,0} + (c_{,0})^2 + (d_{,0})^2 \right] \sin \theta d\psi d\theta.$

Therefore the theorem is a direct consequence of Theorem 4.1. Q.E.D.

5. ADM and Bondi total momenta

In this section, we study the relation between the ADM total linear momentum and the Bondi momentum.

Theorem 5.1. Let $\mathbb{P}_k(t_0)$ be the ADM total linear momentum of the initial data set (N_{t_0}, g, h) where t is given by (3.1) with condition (3.2). Under Condition A, Condition B and Condition D, we have

$$\mathbb{P}_k(t_0) = m_k(-\infty) + \frac{1}{8\pi} \lim_{u \to -\infty} \int_0^{\pi} \int_0^{2\pi} \overline{\mathcal{P}}_k d\psi d\theta$$

for k = 1, 2, 3, where where

$$\overline{\mathcal{P}}_{1} = \left[(c_{,0} + M_{,0}) \cos \theta \cos \psi - d_{,0} \sin \psi \right] (l - a_{2,2}) \sin \theta \\ + \left[(c_{,0} - M_{,0}) \sin \psi + d_{,0} \cos \theta \cos \psi \right] (\bar{l} \sin \theta - a_{2,3}), \\ \overline{\mathcal{P}}_{2} = \left[(c_{,0} + M_{,0}) \cos \theta \sin \psi + d_{,0} \cos \psi \right] (l - a_{2,2}) \sin \theta \\ - \left[(c_{,0} - M_{,0}) \cos \psi - d_{,0} \cos \theta \sin \psi \right] (\bar{l} \sin \theta - a_{2,3}), \\ \overline{\mathcal{P}}_{3} = -(c_{,0} + M_{,0}) (l - a_{2,2}) \sin^{2} \theta - d_{,0} (\bar{l} \sin \theta - a_{2,3}) \sin \theta.$$

Proof: Denote $\mathbb{K}_k = h\left(\frac{\partial}{\partial y^k}, \frac{\partial}{\partial r}\right) - g\left(\frac{\partial}{\partial y^k}, \frac{\partial}{\partial r}\right) tr_g(h)$. Using the asymptotic expansions of g_{ij} and h_{ij} , we obtain

$$\begin{split} \mathbb{K}_{1} &= \frac{1}{r^{2}} \Big\{ 4M \sin \theta \cos \psi - M_{,2} \cos \theta \cos \psi + M_{,3} \csc \theta \sin \psi \\ &\quad -l_{,2} \sin \theta \cos \psi - \bar{l}_{,3} \cos \psi \\ &\quad + \left[(c_{,0} + M_{,0} - 2) \cos \theta \cos \psi - d_{,0} \sin \psi \right] (l - a_{2,2}) \\ &\quad + \left[(c_{,0} - M_{,0} + 1) \sin \psi + d_{,0} \cos \theta \cos \psi \right] (\bar{l} - \csc \theta a_{2,3}) \\ &\quad + 2 \sin \theta \cos \psi a_{1} + \csc \theta \cos \psi a_{2,33} + \sin \theta \cos \psi a_{2,22} \Big\} + O\Big(\frac{1}{r^{3}}\Big), \\ \mathbb{K}_{2} &= \frac{1}{r^{2}} \Big\{ 4M \sin \theta \sin \psi - M_{,2} \cos \theta \sin \psi - M_{,3} \csc \theta \cos \psi \\ &\quad - l_{,2} \sin \theta \sin \psi - \bar{l}_{,3} \sin \psi \\ &\quad + \left[(c_{,0} + M_{,0} - 2) \cos \theta \sin \psi + d_{,0} \cos \psi \right] (l - a_{2,2}) \\ &\quad - \left[(c_{,0} - M_{,0} + 1) \cos \psi - d_{,0} \cos \theta \sin \psi \right] (\bar{l} - \csc \theta a_{2,3}) \\ &\quad + 2a_{1} \sin \theta \sin \psi + \csc \theta \sin \psi a_{2,33} + \sin \theta \sin \psi a_{2,22} \Big\} + O\Big(\frac{1}{r^{3}}\Big). \\ \mathbb{K}_{3} &= \frac{1}{r^{2}} \Big\{ 4M \cos \theta - M_{,2} \sin \theta - l_{,2} \cos \theta - \bar{l}_{,3} \cot \theta \\ &\quad - \left[(c_{,0} + M_{,0} - 2) \sin \theta + \csc \theta \right] l - d_{,0} (\bar{l} \sin \theta - a_{2,3}) \\ &\quad + 2a_{1} \cos \theta + \cos \theta \cot \theta a_{2,22} + \cot \theta \csc \theta a_{2,33} \\ &\quad + (c_{,0} + M_{,0} - 1) \sin \theta a_{2,2} \Big\} + O\Big(\frac{1}{r^{3}}\Big). \end{split}$$

Since a_1 is constant,

$$\int_{S^2} a_1 n^i dS = 0$$

for i = 1, 2, 3, then the theorem is a direct consequence of integrating \mathbb{K}_k over S_r and using that for fixed $t = t_0, r \to \infty$ is equivalent to $u \to -\infty$. Q.E.D. **Remark 5.1.** If a_2 is chosen to be a constant, then the difference $\mathbb{P}_k(t_0) - m_k(-\infty)$ are independent on the choice of f, which are invariant in the perturbed class (3.2) that a_1 , a_2 are constant.

Theorem 5.2. Suppose that Condition A, Condition B and Condition D hold in the Bondi's radiating spacetime (2.1). If

$$\lim_{u \to -\infty} \left(\bar{l}_{,2} + \bar{l} \cot \theta - l_{,3} \csc \theta \right) = 0, \tag{5.1}$$

then there exists f_1 such that

$$\mathbb{P}_{k,f_1}(t_0) = m_k(-\infty)$$

where $\mathbb{P}_{k,f_1}(t_0)$ is the ADM total linear momentum of the initial data set (N_{t_0}, g, h) where t is given by (3.1) and $f = f_1$ given by (3.2) with a_2 satisfying

$$a_{2,2} = l, \quad a_{2,3} = \bar{l}\sin\theta.$$
 (5.2)

Proof: The condition (5.1) ensures the existence of (5.2). Therefore the theorem follows. Q.E.D.

Remark 5.2. From Theorem 4.2 and Theorem 5.2, we know that, in general, it is impossible to perturb spatial infinity via the function f so that both the ADM total energy and the ADM total linear momentum are the past limit of the Bondi energy-momentum.

Acknowledgements. Xiao Zhang is partially supported by National Natural Science Foundation of China under grants 10231050, 10421001 and the Innovation Project of Chinese Academy of Sciences. This work was partially done when Wen-ling Huang visited the Morningside Center of Mathematics, Chinese Academy of Sciences, and she would like to thank the center for its hospitality.

References

- S. Arnowitt, S. Deser, C. Misner, Coordinate invariance and energy expressions in general relativity, Phys. Rev. 122(1961), 997-1006.
- [2] A. Ashtekar, A. Magnon-Ashtekar, *Energy-momentum in general relativity*, Phys. Rev. Lett. 43(1979), 181-184.
- [3] H. Bondi, M. van der Burg, A. Metzner, Gravitational waves in general relativity VII. Waves from axi-symmetric isolated systems, Proc. Roy. Soc. London A 269(1962), 21-52.
- [4] D. Christodoulou, S. Klainerman, The global nonlinear stablity of Minkowski space, Princeton Math. Series 41, Princeton Univ. Press, Princeton, 1993.
- [5] P. Chruściel, J. Jezierski, J. Kijowski, Hamiltonian field theory in the radiating regime. Lecture Notes in Physics. Monographs, 70. Springer-Verlag, Berlin, 2002.
- [6] P. Chruściel, M. MacCallum, D. Singleton, Gravitational waves in general relativity XIV. Bondi expansions and the "polyhomogeneity" of Scri, Phil. Trans. Roy. Soc. A 350(1995), 113-141.
- [7] S. Hayward, Spatial and null infinity via advanced and retarded conformal factors, Phys. Rev. D68(2003), 104015.

- [8] W.-l. Huang, X. Zhang, On the relation between ADM and Bondi energy-momenta II radiative spatial infinity, gr-qc/0511037.
- R. Sachs, Gravitational waves in general relativity VIII. Waves in asymptotically flat space-time, Proc. Roy. Soc. London, A 270(1962), 103-126.
- [10] J. Valiente Kroon, Early radiative properties of the developments of time-symmetric conformally flat initial data, Class. Quantum Grav. 20(2003)L53-L59.
- J. Valiente Kroon, Does asymptotic simplicity allow for radiation near spatial infinity, Commun. Math. Phys. 251(2004), 211-234.
- [12] M. van der Burg, Gravitational waves in general relativity IX. Conserved quantities, Proc. Roy. Soc. London A 294(1966), 112-122.
- [13] X. Zhang, On the relation between ADM and Bondi energy-momenta, Adv. Theor. Math. Phys., 10(2006), 261-282.

(Wen-ling Huang) DEPARTMENT MATHEMATIK, SCHWERPUNKT GD, UNIVERSITÄT HAMBURG, BUNDESSTR. 55, D-20146 HAMBURG, GERMANY

E-mail address: huang@math.uni-hamburg.de

(Xiao Zhang) Institute of Mathematics, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100080, China

E-mail address: xzhang@amss.ac.cn