CLASSICAL ASPECTS OF LIGHTLIKE DIMENSIONAL REDUCTION

E. MINGUZZI

ABSTRACT. Some aspects of lightlike dimensional reduction in flat spacetime are studied with emphasis to classical applications. Among them the Galilean transformation of shadows induced by inertial frame changes is studied in detail by proving that, (i) the shadow of an object has the same shape in every orthogonal-to-light screen, (ii) if two shadows are simultaneous in an orthogonal-to-light screen then they are simultaneous in any such screen. In particular, the Galilean group in 2+1 dimensions is recognized as an exact symmetry of Nature which acts on the shadows of the events instead that on the events themselves. The group theoretical approach to lightlike dimensional reduction is used to solve the reconstruction problem of a trajectory starting from its acceleration history or from its projected (shadow) trajectory. The possibility of obtaining a Galilean projected physics starting from a Poincaré invariant physics is stressed through the example of relativistic collisions. In particular, it is shown that the projection of a relativistic collision between massless particles gives a non-relativistic collision in which the kinetic energy is conserved.

Department of Applied Mathematics, Florence University, Via S. Marta 3, I-50139 Florence, Italy. E-mail: ettore.minguzzi@unifi.it.

[...] they see only their own shadows, or the shadows of one another, which the fire throws on the opposite wall of the cave [...]. To them, I said, the truth would be literally nothing but the shadows of the images. Plato, The allegory of the cave, Book VII of the Republic

1. INTRODUCTION

In the famous book *The Republic* Plato, talking to one of his followers, Glaucon, introduced a powerful image, *The allegory of the cave*, in order to explain how much the knowledge can make the man free.

Plato imagined some "observers" chained since their childhood and constrained to look in front of them at a wall in the deep of a cave. Light came behind them projecting their own shadows and the shadows of other objects and people on the wall. According to Plato the chained observers would identify themselves and the other objects with their respective shadows, eventually loosing their ability to perceive the third space dimension.

To the physical minded reader Plato's allegory represents the first historical example of a dimensional reduction process. Indeed, Plato's thought experiment represents a particular type of dimensional reduction, that is, a *lightlike* dimensional reduction. It differs from that considered much later by Kaluza and Nordström [27] (spacelike dimensional reduction) or from that used in the hydrodynamic formalism of general relativity or in the study of stationary metrics (timelike dimensional reduction). In this work we shall study how the physics laws on the full spacetime reduce to the quotient spacetime. A general feature will be that relativistic physics projects to non-relativistic physics. In Plato's allegory terms: the chained observers would not be able to observe any relativistic effect, independently of the speed reached by the shadows, that is, at any degree of accuracy.

Our notations are as follows. Let $d \in \mathbb{N}$. The indexes i, j, k, take the values $1, \ldots, d + 1$, the indexes a, b, c, take the values $1, \ldots, d$. Vectors in *d*-dimensional vector spaces are denoted in boldface, so that v^a reads \mathbf{v} . The Greek indexes α, β, μ, ν , take the values $0, 1, \ldots, d + 1$, and the indexes A, B, C, take the values $0, 1, \ldots, d + 1$, and the indexes A, B, C, take the values $0, 1, \ldots, d + 1$, and the indexes A, B, C, take the values $0, 1, \ldots, d$. For compactness the transpose of a vector is denoted with the same letter. On Minkowski spacetime M of dimension (d + 1) + 1, we use coordinates $\{x^{\mu}\}, \mu = 0, 1, \ldots, d + 1$, the spacelike convention $\eta_{00} = -1$, and units such that c = 1. In the list of commutation relations defining a Lie algebra we shall omit the vanishing ones.

Mathematically the study of lightlike dimensional reductions began in a 1929 work by Eisenhart [10] who showed that the trajectories of a Lagrangian system with d degrees of freedom

(1)
$$L(q^{a}, \dot{q}^{a}, t) = \frac{1}{2}a_{ab}(q)\dot{q}^{a}\dot{q}^{b} + b_{c}(q)\dot{q}^{c} - U(q),$$

may be obtained as the projection of geodesics of a d + 2-dimensional Lorentzian¹ spacetime of metric [10, 23]²

(2)
$$ds^{2} = a_{ab}dq^{a}dq^{b} + 2b_{c}dq^{c}dq^{0} - (2U+1)(dq^{0})^{2} - 2dq^{0}dq^{d+1},$$

where $q^0 = t$ and q^{d+1} is an auxiliary variable.

In particular given a solution of the dynamical system $q^{a}(t)$, set

(3)
$$q^{d+1}(t) = C - t + \int_0^t L(q^a(t), \dot{q}^a(t)) \mathrm{d}t,$$

where C is an arbitrary constant. The trajectory $\{q^0, q^a(q^0), q^{d+1}(q^0)\}$ can be regarded as an affine parametrization of a geodesic of the Eisenhart metric (2) with respect to a natural parameter [23], $s = t = q^0$. Moreover, every solution of the Lagrangian system can be regarded in this way. The Eisenhart metric takes its simplest and most symmetric form in the case of a free classical particle $a_{bc} =$ $\delta_{bc}, b_c = 0, U = 0$. Remarkably in this case the Eisenhart metric becomes the Minkowski metric as can be seen introducing alternative coordinates $x^i = q^i, x^0 =$ $q^0 + q^{d+1}$.

This result was the first signal of an interesting correspondence between Poincaré invariance and Galilei invariance in a lightlike reduced spacetime with one dimension less. Nevertheless, it took half a century to fully realize the correspondence. Some progress was made after the impulse of Dirac's work [6] that led to the so called front wave (or light front or infinite momentum frame) dynamics [5, 20, 13, 19]. Later, a work by Bargmann [2] inspired a series of works where this correspondence was fully appreciated and led to the Bargmann structure approach to the Newton-Cartan theory [17, 7, 8, 3, 9]. The algebraic conditions at its foundations were slightly weakened in subsequent works [15]. In both lines of research the authors studied an alternative formulation of quantum field theory by using null Cauchy surfaces. In the latter case the authors considered the (d+1)+1 spacetime mainly as a tool for expressing in a simple way the symmetries of Galilei invariant quantum mechanical systems in d+1 spacetime dimensions.

In this work we focus on classical applications of lightlike dimensional reduction. Indeed, we feel that already at the classical level this structure may have interesting applications. For instance, the correspondence between Poincaré invariance and Galilei invariance in a lightlike reduced spacetime may be grasped particularly well in its simplest application: the study of the transformation properties of shadows under inertial frame changes. In order to keep the work at a reasonable size we shall limit ourselves to the flat spacetime case as it has some peculiarities which are worth studying in their own right. Indeed, in Minkowski spacetime the Poincaré group is left unbroken and the group theoretical approach becomes particularly advantageous. On the contrary, in the curved spacetime case differential geometric tools should be preferred, and new concepts such as the mentioned Bargmann structures or Eisenhart's spacetime should be introduced.

This work is organized as follows.

¹The Lorentzianity of the metric follows immediately by introducing the base of 1-forms $\omega^0 = dq^0$, $\omega^a = dq^a$, $\omega^{d+1} = dq^{d+1} + (U + 1/2)dq^0 - b_c dq^c$ and by taking into account the positive definiteness of a_{bc} .

²In our convention the roles of q^0 and q^{d+1} are inverted with respect to [23, Book II, Sect. 11] and there is also a different choice of sign.

- In section 2 we model our problem by considering a distant source of light and some sets of inertial observers suitably oriented with respect to the direction of light (subsection 2.1). Then the groups acting transitively on each set are studied (subsection 2.2), and it is shown that suitably accelerating and rotating observers can have comoving inertial frames which keep staying, time by time, in the same original set (subsection 2.2). In subsection 2.4 it is shown that a subgroup of the Poincaré group is in fact, a central extension of the Galilei group. Thus, although the Galilei group does not act on events, its action is well defined on the quotient space Q of their shadows. The application to the transformation properties of shadows is studied in detail (subsections 2.5 and 2.6), and several conclusions are finally drawn (subsection 2.7).
- Section 3 is devoted to various reconstruction problems. First, the problem of reconstructing the trajectory starting from the decomposition of the proper acceleration in acceleration along the direction of light and orthogonal to it is considered. A new particular (lightlike) parallel transport related to the presence of a preferential lightlike vector is introduced by means of group theoretical methods, and it is shown to give rise to an absolute transverse orientation (subsection 3.1). The new parallel transport is completely integrated and the relevance of this solution for the problem of autonomous spacetime navigation is emphasized (subsection 3.2). In subsection 3.3 the problem of reconstructing a trajectory starting from its shadow is considered, and the relation with the classical action is pointed out. In subsection 3.4 a formula for the time dilation given the acceleration history of a non-inertial observer is obtained. It is the first formula of this kind developed in three or more space dimensions, the analogous problem for a decomposition of the acceleration with respect a Fermi-Walker triad being still open. In the last subsection 3.5 we point out that the knowledge, by an inertial observer, of the sky position and longitudinal frequency of a signal emitted by a source does not allow to deduce the behavior of the distance. A position drift may always occur which is related to the classical action.
- Section 4 is devoted to an example in which the projection of a Poincaré symmetry into a Galilei symmetry becomes particularly clear, that is, that of a relativistic collision which projects into a non-relativistic collision. The reduced Lagrangian is calculated showing that it is of classical type and the kinetic and internal energies are identified. It is shown that any shadow worldline has an associated (inertial reference frame) invariant, i.e. the *shadow mass*, which plays the role of a non-relativistic mass. The conservation of shadow mass, momentum and energy is shown to follow from the conservation of relativistic momentum in the full spacetime. Although the kinetic energy is not necessarily conserved, it is shown that it is conserved in the projection of a relativistic collisions between massless particles. Finally, the inverse problem of finding the relativistic collision from which a non-relativistic collision comes from, is solved.
- In section 5 we give some conclusions.
- In Appendix A a differential geometric formulation of the lightlike parallel transport is given.

4

• In Appendix B the formal analogy between the problem of transforming shadows between different inertial frames, and the problem of transforming the photon polarization vector is pointed out. Some ambiguities regarding the role of the transverse Galilean boosts generators in this context are clarified.

2. Shadows and group aspects of lightlike dimensional reduction

In this section the reader is introduced to the geometry of lightlike dimensional reduction through the study of shadows and their transformation properties. However, we stress that the results of this section are not confined to the application to shadows, but are rather general properties of the geometry of lightlike dimensional reduction.

2.1. Subsets of inertial observers. let M be Minkowski spacetime. By observer we mean a timelike worldline $\gamma(\tau)$ parametrized with respect to proper time and an orthonormal tetrad field $\{e_0 = \partial_{\tau}, e_i\}$ over it. If the tetrad is parallely transported the observer is inertial. An orthonormal tetrad at $m \in M$, $\{e_0, e_i\}$ with e_0 timelike will be said to be an *inertial frame* since canonical coordinates $\{x^{\mu}\}$ with origin m can be introduced such that $e_{\mu} = \partial/\partial x^{\mu}$, $ds^2 = \eta_{\mu\nu} dx^{\mu} dx^{\nu}$. Sometimes an inertial frame will be denoted with the letter K. Thus every observer is a sequence of inertial frames $K(\tau)$. The inertial frames of an inertial observer are related by time translations.

Consider a point like source of monochromatic light Σ placed very far from a spacetime region $\mathcal{U} \subset M$ of Minkowski spacetime. For an inertial observer inside \mathcal{U} the light coming from the source is described by a congruence of parallel null geodesics, the photons coming from the source having a well defined momentum p^{μ} . The inertial observers inside \mathcal{U} , without changing their covariant velocity, can orient their own space axes $\{e_i\}$, so that the photons travel in direction e_{d+1} , $p^{\mu} = \omega(1, 0, \ldots, 0, 1)$ with a constant ω possibly depending on the observer. We are focusing here on the subset $S(\hat{p})$ of the inertial frames for which the null momentum p^{μ} has the same spatial components up to a multiplicative constant. The subset $S(\hat{p})$ splits into subsets $S_{\omega}(\hat{p})$ dependent on the value of the frequency ω measured by the observer, $p^i = \omega \hat{p}^i$.

Let $L(e_{d+1}) \subset SO(1, d+1)$ be the little group of the null vector $n^{\mu} = (1, 0, \ldots, 1)$ (by SO(1, d+1) we mean the connected component of the Lorentz group containing the identity), that is, the subgroup of SO(1, d+1) which leaves n^{μ} invariant, and analogously let $IL(e_{d+1}) \subset ISO(1, d+1)$ be the little group with the translations included, $L(e_{d+1}) \subset IL(e_{d+1})$. Moreover, let $G(e_{d+1})$ be the group obtained from $L(e_{d+1})$ by including the boosts in the direction e_{d+1} , i.e. the group of those Lorentz transformations that send n^{μ} to a vector proportional to it, and let $IG(e_{d+1})$ be the group obtained by including the boosts in direction e_{d+1} to $IL(e_{d+1})$, or the translations to $G(e_{d+1})$. It is known [33] that the group $IL(e_{d+1})$ acts freely and transitively on $S_{\omega}(e_{d+1})$ and that the group $IG(e_{d+1})$ acts freely and transitively on $S(e_{d+1})$. Under changes of n the groups remain the same up to isomorphisms.

The group $IG(e_{d+1})$ represents Poincaré transformations between inertial frames for which the light from Σ comes from the same direction $-e_{d+1}$. The group $IL(e_{d+1})$ represents Poincaré transformations between inertial frames for which the light from Σ comes from the same direction $-e_{d+1}$ with the same frequency.




FIGURE 1. Shadow of an object on the plane perpendicular to the last axis. The worldlines of the points at rest with the plane span on spacetime the hyperplane $x^{d+1} = 0$. Assume the plane is a semitransparent screen so that the shadow has different images, one for each orthogonal-to-light screen. How does the shadow image change under change of orthogonal-to-light screen?

The groups $G(e_{d+1})$ and $L(e_{d+1})$ add the condition that the related inertial frames share the same origin of coordinates.

We assume that the inertial observers set up semitransparent screens perpendicular to the direction of light, at their respective coordinate $x^{d+1} = 0$ in such a way that light comes from the region $x^{d+1} < 0$. In this way the shadow of an object projects on their respective screens leaving a dark image on each one. One should be careful because, although the screens of different observers are perpendicular to the same direction of light, the screens are in no sense parallel due to the aberration of light under boosts. Our aim is to show that the different images of the same shadow are related between the screens of observers in $S_{\omega}(e_{d+1})$ by a Galilean transformation and between the screens of observers in $S(e_{d+1})$ by a Galilean transformation plus a time scaling.

This problem is somewhat related to that of finding how the appearance of the night sky transforms under Lorentz transformations [28, 29, 26]. Whereas there one focuses on light reaching an event from all directions, here we focus on light of a given direction projecting on planes.

2.2. The little group of the massless particle. Let us briefly recall how to construct $L(e_{d+1})$ [33]. Let

(4)
$$x'^{\mu} = L^{\mu}_{\ \nu} x^{\nu} - b^{\mu},$$

be a transformation in $IL(e_{d+1})$. The invariance of the (d+2)-momentum ωn^{μ} reads $L^{\mu}_{\ \nu}n^{\nu} = n^{\mu}$. Let $t^{\mu} = (1, 0, \dots, 0)$, the vector $(Lt)^{\mu} = L^{\mu}_{\ \nu}t^{\nu}$ is a unit vector that satisfies $n_{\mu}(Lt)^{\mu} = (Ln) \cdot (Lt) = n \cdot t = -1$. As a consequence Lt can be written

(5)
$$(Lt)^{\mu} = (1+\zeta, -\alpha^1, \dots, -\alpha^d, \zeta),$$

for a suitable α^a and $\zeta = \boldsymbol{\alpha} \cdot \boldsymbol{\alpha}/2$. The Lorentz transformation

(6)
$$S(\boldsymbol{\alpha})^{\mu}_{\nu} = \begin{pmatrix} 1+\zeta & -\boldsymbol{\alpha} & -\zeta \\ -\boldsymbol{\alpha} & I & \boldsymbol{\alpha} \\ \zeta & -\boldsymbol{\alpha} & 1-\zeta \end{pmatrix},$$

where I_b^a is the identity matrix, leaves n^{ν} invariant and sends t to Lt. Thus the Lorentz transformation $L^{-1}S$ leaving invariant both t and n (and hence $e_{d+1}^{\mu} = n^{\mu} - t^{\mu}$) must be an element of SO(d), that is a rotation $(R^{-1})_{\nu}^{\mu}$ that leaves e_{d+1} invariant. The matrix R_{ν}^{μ} takes the form, with obvious notation,

(7)
$$R^{\mu}_{\ \nu} = \begin{pmatrix} 1 & \mathbf{0} & 0 \\ \mathbf{0} & \mathbf{R} & \mathbf{0} \\ 0 & \mathbf{0} & 1 \end{pmatrix},$$

where $\mathbf{R}^{a}_{\ b} = \mathbf{R}_{ab} \in SO(d)$. The generic element $L^{\mu}_{\ \nu}$ of the little group $L(e_{d+1})$ takes the form $L^{\mu}_{\ \nu} = S^{\mu}_{\ \beta}R^{\beta}_{\ \nu}$ or

(8)
$$L^{\mu}_{\ \nu} = \begin{pmatrix} 1+\zeta & -\alpha^{c}\mathbf{R}_{cb} & -\zeta \\ -\alpha^{a} & \mathbf{R}^{a}_{\ b} & \alpha^{a} \\ \zeta & -\alpha^{c}\mathbf{R}_{cb} & 1-\zeta \end{pmatrix}.$$

Note that the matrices of type (6) form an Abelian subgroup of dimension d, the so called 'translations' of the little group. From the expression (8) we can recover the Lie algebra $\mathcal{L}(e_{d+1})$ of $L(e_{d+1})$. The infinitesimal transformation has the form $(J^{ab} = J_{ab})$

(9)
$$L \simeq I + \frac{1}{2}\Omega_{ab}J^{ab} + \alpha^a W_a$$

where W_a are the generators of 'translations' and (here and throughout the work we omit vanishing commutation relations such as $[W_a, W_b] = 0$ while reporting the commutation relations of an entire Lie algebra)

(10)
$$[J_{ab}, J_{cd}] = \delta_{ad}J_{bc} + \delta_{bd}J_{ad} - \delta_{ac}J_{bd} - \delta_{bd}J_{ac},$$

(11)
$$[W_a, J_{bc}] = \delta_{ab} W_c - \delta_{ac} W_b,$$

thus $L(e_{d+1})$ is isomorphic to the group ISO(d). From (8) one can also find the expression of these generators in terms of the generators of the Lorentz group

(12)
$$[J^{\alpha\beta}, J^{\gamma\delta}] = \eta^{\alpha\delta} J^{\beta\gamma} + \eta^{\beta\gamma} J^{\alpha\delta} - \eta^{\alpha\gamma} J^{\beta\delta} - \eta^{\beta\delta} J^{\alpha\gamma}.$$

Indeed, the usual matrix representation of the group $x'^{\mu} = \Lambda^{\mu}_{\nu} x^{\nu}$ induces the Lie algebra representation $(J^{\alpha\beta})^{\mu}_{\ \nu} = \eta^{\alpha\mu} \delta^{\beta}_{\ \nu} - \eta^{\beta\mu} \delta^{\alpha}_{\ \nu}$. From the infinitesimal version of Eq. (8) we obtain

$$W_a = J^{0 a} - J^{d+1 a}$$

and, as the notation suggests, J_{ab} is nothing but $J_{\mu\nu}$ with the indexes restricted to the values $1, \ldots d$. Moreover, as expected, $(J_{ab})^{\mu}_{\ \nu}n^{\nu} = (W_a)^{\mu}_{\ \nu}n^{\nu} = 0$.

2.3. Poincaré group, inhomogeneous little group and non-inertial observers. The infinitesimal Poincaré transformation $x'^{\mu} = \Lambda^{\mu}_{\ \nu} x^{\nu} - b^{\mu}$ in a (d+1)+1 spacetime can be written

(13)
$$I + \frac{1}{2}\Omega_{\alpha\beta}J^{\alpha\beta} - b_{\gamma}P^{\gamma},$$

where the generators $J^{\alpha\beta}$, P^{γ} , satisfy the Lie algebra

(14)
$$[J^{\alpha\beta}, J^{\gamma\delta}] = \eta^{\alpha\delta} J^{\beta\gamma} + \eta^{\beta\gamma} J^{\alpha\delta} - \eta^{\alpha\gamma} J^{\beta\delta} - \eta^{\beta\delta} J^{\alpha\gamma},$$

(15)
$$[P^{\alpha}, J^{\beta\gamma}] = \eta^{\alpha\beta} P^{\gamma} - \eta^{\alpha\gamma} P^{\beta}.$$

We shall write $H = P^0$.

Any non-inertial observer passes through a sequence of inertial frames $K(\tau)$. The coordinate transformation between $K(\tau)$ and $K(\tau+d\tau)$ is an infinitesimal Poincaré transformation $I + \frac{1}{2}\tilde{\omega}_{\alpha\beta}J^{\alpha\beta}d\tau + Hd\tau$ where the space translations do not enter. Indeed, the presence of space translations would imply a violation of causality. Here the motion is regarded as a sequence of infinitesimal time translations, boosts and rotations. The infinite product of those infinitesimal transformations may generate space translations that, however, are not present in the infinitesimal transformation.

Let $x_{K(\tau)}^{\mu} = \Lambda_{\nu}^{\mu}(\tau)[x_{K(0)}^{\nu} - x^{\mu}(\tau)]$ be the Poincaré transformation from K(0)to $K(\tau)$. Here $x_{K(\tau)}^{\mu}$ are the coordinates of $K(\tau)$, and $x^{\mu}(\tau)$ represents the origin of $K(\tau)$ with respect to K(0). This general transformation arises from an infinite product of the said infinitesimal transformations. The covariant velocity of $K(\tau)$ in its own coordinates is δ_{0}^{μ} hence the covariant velocity of $K(\tau)$ in K(0)'s coordinates is $u^{\mu}(\tau) = (\Lambda^{-1})_{0}^{\mu}$. We have

(16)
$$\frac{\mathrm{d}}{\mathrm{d}\tau}\Lambda = \frac{1}{2}\tilde{\omega}_{\alpha\beta}(\tau)J^{\alpha\beta}\Lambda,$$

(17)
$$x^{\mu}(\tau) = \int_{0}^{\tau} \mathrm{d}\tau' (\Lambda^{-1}(\tau'))^{\mu}_{0}.$$

The motion of the non-inertial observer can be recovered from the knowledge of $\tilde{\omega}_{\alpha\beta}(\tau)$. Its physical meaning is the following. The acceleration of the non-inertial observer in K(0)'s coordinates is $\frac{\mathrm{d}}{\mathrm{d}\tau}(\Lambda^{-1}(\tau'))^{\mu}_{0}$ hence the acceleration of the non-inertial observer in its own coordinates is

$$\tilde{a}^{\mu} = \Lambda^{\mu}_{\ \nu} \frac{\mathrm{d}}{\mathrm{d}\tau} (\Lambda^{-1})^{\nu}_{\ 0} = -\tilde{\omega}^{\mu}_{\ 0}.$$

The tilde reminds us that \tilde{a}^{μ} is not only the covariant acceleration a^{μ} , which makes sense in arbitrary frames, but the covariant acceleration in the coordinates of the accelerating frame (hence $\tilde{a}^0 = 0$). The components of the acceleration in the non-inertial observer's axes are $\tilde{\omega}^{\mu 0} = \tilde{\omega}_{0\mu}$. Analogously $\tilde{\omega}^{ij} = \tilde{\omega}_{ij}$ represents the tensorial angular velocity of the non-inertial frame in its own coordinates.

Now, let us consider a non-inertial observer such that $K(\tau)$ passes through inertial frames belonging to $S_{\omega}(e_{d+1})$. Let $x_{K(\tau)}^{\mu} = L^{\mu}_{\nu}(\tau)[x_{K(0)}^{\nu} - x^{\mu}(\tau)]$ be the inhomogeneous little group transformation from K(0) to $K(\tau)$. This time we have that $L(\tau + d\tau) = (I + \frac{1}{2}\tilde{\omega}_{ab}d\tau J^{ab} + \tilde{\alpha}^a d\tau W_a)L$. Using the expression for W_a , we obtain the constraints

(18)
$$\tilde{a}^{d+1} = \tilde{\omega}^{d+10} = 0,$$

(19)
$$\tilde{a}^a = \tilde{\omega}_{0a} = -\tilde{\omega}_{d+1a} = \tilde{\alpha}_a.$$

In other words the non-inertial observer may keep staying time by time into $S_{\omega}(e_{d+1})$ despite its non-inertial motion provided (i) its acceleration along the direction e_{d+1} vanishes and (ii) (this interpretation holds only for d = 2 in which case an angular velocity vector can be defined), the projection of the angular velocity vector on the plane perpendicular to e_{d+1} and the acceleration are perpendicular to each other

and of the same magnitude, moreover, their vector product equals e_{d+1} up to a non-negative factor.

Due to the different dimensionality of acceleration and angular velocity we find, restoring c, that the projected angular velocity is indeed very small. It is required in order to correct for the aberration of light due to the acceleration which would change the night sky position of the source Σ from the direction $-e_{d+1}$.

2.4. The Galilei quotient group. Let

(20)
$$\mathcal{M} = H - P^{d+1},$$

in such a way that the infinitesimal transformation in $\mathcal{IL}(e_{d+1})$ reads

$$I + \frac{1}{2}\Omega_{ab}J^{ab} + \alpha^{a}W_{a} - b^{b}P_{b} + (b^{0} - b^{d+1})H + b^{d+1}\mathcal{M}.$$

The Lie algebra $\mathcal{IL}(e_{d+1})$ in terms of the generators J_{ab} , W_a , P_a , H and \mathcal{M} is obtained by adding to Eqs. (10)-(11) the commutations relations that follow from Eq. (15)

(21)
$$[P_a, J_{bc}] = \delta_{ab} P_c - \delta_{ac} P_b,$$

$$(22) [W_a, H] = P_a,$$

$$[W_a, P_b] = \delta_{ab}\mathcal{M}$$

The generators J_{ab} , W_a , P_a and H would form a Galilei subalgebra $\mathcal{G}al(d)$ given by the only non-vanishing commutation relations (10), (11), (21) and (22) if the right hand side of (23) would vanish. Since \mathcal{M} commutes with the other generators, \mathcal{IL} is a central extension of the Galilei algebra $\mathcal{G}al(d)$ in d + 1 spacetime

(24)
$$0 \to \{\mathcal{M}\} \to \mathcal{IL}(e_{d+1}) \to \mathcal{Gal}(d) \to 0,$$

where $\{\mathcal{M}\}$ is the 1-dimensional Lie algebra spanned by \mathcal{M} .

The inequivalent central extensions of the Galilei algebra have been classified by Bargmann [2] for the case $d \ge 3$ and by Bose [4] for the case d = 2. The vector space of inequivalent extensions is one dimensional in the former case and three dimensional in the latter case. Thus, the central extension considered here is the only one available for d > 3, and in particular it is the one that makes sense physically. Indeed, in quantum mechanics the unitary ray representations of a group G are induced by the unitary representations of the central extensions of the universal covering of G, G^* . Bargmann has shown that the projective unitary representation of the Galilei group that makes sense physically is the one obtained from the central extension considered above. The operator \mathcal{M} , in that quantum mechanical context, is known as the mass operator. Its presence implies the mass superselection rule which forbids the superpositions of states with different mass [22, 33]. We shall see below that it has the same meaning in our classical context with the difference that by mass here we shall mean the one that makes sense in a suitable d+1 Galilean spacetime Q. In the application to shadows, for instance, it will be called the *shadow mass*, to distinguish it from the mass of the particle that projects the shadow.

Let N be the 1-dimensional normal subgroup generated by \mathcal{M} , then $IL(e_{d+1})/N \sim Gal(d)$ has the correct Lie algebra. The central extension at the group level is

(25)
$$1 \to N(\sim T_1) \to IL(e_{d+1}) \to Gal(d) \to 1.$$




FIGURE 2. The decomposition of the event of coordinates x^{μ} into transversal (shadow) position **x**, hitting time t, and longitudinal coordinate x^{d+1} . The interpretation of **x** as the position of the shadow makes sense only if $x^{d+1} \leq 0$.

The group Gal(d) does not act on events of coordinates x^{μ} but rather on the worldlines of the form Nx. Since $\mathcal{M} = H - P^{d+1} = -P^{\gamma}n_{\gamma}$, the effect of an element of N on x^{μ} is that of adding a vector proportional to n^{μ} , i.e. the elements Nx are the light rays emitted from Σ . The reduced spacetime Q is the space of null geodesics of M of direction n^{μ} . The dimensional reduction of M onto Q along a lightlike direction n sends the little group $IL(e_{d+1})$ into the Galilei group. Thus, at least for the case d = 2 we can regard the Galilean symmetry in 2+1 spacetime as an exact symmetry of Nature rather than as an approximate symmetry for small velocities. In order to reveal this symmetry one has to focus on the shadows of the objects in place of the material objects themselves. On the contrary, the Galilean symmetry of the full 3+1 spacetime remains an approximate symmetry valid only for small velocities.

2.5. The transformation of shadows. Chosen an inertial frame in $S_{\omega}(e_{d+1})$ of coordinates $\{x^{\mu}\}$ every event x^{μ} admits the unique decomposition $(t = x^0 - x^{d+1})$

(26)
$$x^{\mu} = \begin{pmatrix} x^{d+1} + t \\ \mathbf{x} \\ x^{d+1} \end{pmatrix} = x^{d+1} n^{\mu} + \begin{pmatrix} t \\ \mathbf{x} \\ 0 \end{pmatrix}.$$

The parameter t represents the time at which the light beam from Σ passing through x^{μ} hits the surface (screen) $x^{d+1} = 0$. The vector **x** gives the hitting point on the screen (see figure 2). By shadow of event x we mean the null geodesic of direction n^{μ} passing through x or its representative (t, \mathbf{x}) . Note that physically the shadow exists only if the object of worldline $x^{\mu}(\tau)$ that projects the shadow passes between the source and the screen, $x^{d+1}(\tau) < 0$. We shall not impose this condition because it is not restrictive (the frame origin can be translated) and because we are more interested on the mathematical definition of shadow.

The shadow of a (C^1) particle worldline $x^{\mu}(\lambda)$ is that shadow worldline $\mathbf{x}(t) = \mathbf{x}(\lambda(t))$ composed of the shadows of the events of the original worldline. We shall

assume that $x^{\mu}(\lambda)$ is causal, future pointing and that $dx^{\mu}/d\lambda$ is not proportional to n^{μ} , that is $\frac{dx^{\mu}}{d\lambda}n_{\mu} < 0 \Rightarrow dt/d\lambda > 0$, then $\mathbf{x}(t)$ is future pointing in the sense that $t(\lambda)$ is increasing. In this work we shall consider massive and massless particles on M. One should be careful with the massless particles having a four momentum proportional to n^{μ} , since they have a special role. They do not project on worldlines $\mathbf{x}(t)$ but rather on isolated events (t, \mathbf{x}) . Thus such particles represent the events of our reduced Galilean spacetime Q rather than the particles moving on it.

Since the light ray passing though x is determined by the parameters t and **x** one would expect that under a transformation in $IL(e_{d+1})$ these parameters transform according to the Galilei quotient group. Indeed, this is the case. If x' is related to x by Eq. (4) with L^{μ}_{ν} given by (8) then

(27)
$$x'^{\mu} = \begin{pmatrix} {x'}^{d+1} + t' \\ \mathbf{x}' \\ {x'}^{d+1} \end{pmatrix} = {x'}^{d+1} n^{\mu} + \begin{pmatrix} t' \\ \mathbf{x}' \\ 0 \end{pmatrix}$$

with

(28)
$$x'^{d+1} = x^{d+1} + t\zeta - \alpha^a \mathbf{R}_{ab} x^b - b^{d+1},$$

and (Galilei transformation)

(29)
$$t' = t - (b^0 - b^{d+1}),$$

(30)
$$x'^{b} = \mathbf{R}^{b}_{c}x^{c} - t\alpha^{b} - b^{b}.$$

The transformation (28)-(30) makes it clear that the generators W_a generate Galilean boosts on the d + 1 quotient spacetime. Their commutativity expresses the commutativity of Galilean boosts.

In the coordinates x^{d+1} , t and x^a the Minkowski metric reads

(31)
$$ds^{2} = -dt^{2} - 2dt \, dx^{d+1} + dx^{a} dx_{a}.$$

It is invariant under the transformation (28)-(30).

2.6. The inclusion of boosts of direction e_{d+1} . In the previous sections we studied the group $IL(e_{d+1})$ that acts freely and transitively on $S_{\omega}(e_{d+1})$. The boosts in direction e_{d+1} do not belong to this group since they would change the observed frequency ω . Thus the inertial frames belonging to $S_{\omega}(e_{d+1})$, not only are suitably oriented but also their covariant velocity satisfies a constraint and is not completely general. The situation is quite different with the set $S(e_{d+1})$: up to a reorientation of the axes every inertial frame belongs to this set. It is important to generalize our conclusion to the group $IG(e_{d+1})$ that acts freely and transitively on $S(e_{d+1})$ as the results would hold for any observer as long as we identify an observer with its worldline.

The subgroup $G(e_{d+1})$ of the Lorentz group sends the null vector $n^{\mu} = (1, 0, 0, 1)$ to a vector proportional to it.³ Let $G^{\mu}_{\nu} \in G(e_{d+1})$, we have $G^{\mu}_{\nu}n^{\nu} = e^{-r(G)}n^{\mu}$ for a suitable constant r(G) (by Lorentz group we mean the connected component which contains the identity, thus, since it sends the forward light cone into the

³The group $G(e_{d+1})$ sends the null plane $n^{\mu}x_{\mu} = 0$ into itself. This fact explains the close relation between this work and studies on *front wave dynamics*.

forward light cone, the proportionality constant is positive). The ratio between the frequencies in the two frames is $\omega'/\omega = e^{-r}$. Note that the boost

(32)
$$B(r, e_{d+1}) = \begin{pmatrix} \cosh r & \mathbf{0} & -\sinh r \\ \mathbf{0} & \delta^a_b & \mathbf{0} \\ -\sinh r & \mathbf{0} & \cosh r \end{pmatrix}$$

satisfies $B^{\mu}_{\ \nu}(r, e_{d+1})n^{\nu} = e^{-r}n^{\mu}$. Thus $B^{-1}{}^{\mu}_{\ \nu}G^{\nu}{}_{\alpha}n^{\mu} = n^{\mu}$, that is $L^{\mu}_{\ \alpha} = B^{-1}{}^{\mu}_{\ \nu}G^{\nu}{}_{\alpha}$ belongs to the little group $L(e_{d+1})$. The parametric form of $G^{\mu}_{\ \nu} = B^{\mu}{}_{\beta}L^{\beta}{}_{\alpha}$ is

(33)
$$G^{\mu}_{\nu} = \begin{pmatrix} \cosh r + e^{-r}\zeta & -e^{-r}\alpha^{c}\mathbf{R}_{cb} & -\sinh r - e^{-r}\zeta \\ -\alpha^{a} & \mathbf{R}^{a}_{\ b} & \alpha^{a} \\ -\sinh r + e^{-r}\zeta & -e^{-r}\alpha^{c}\mathbf{R}_{cb} & \cosh r - e^{-r}\zeta \end{pmatrix}.$$

Using the decomposition (26) the generic $IG(e_{d+1})$ transformation, $x'^{\mu} = G^{\mu}_{\ \nu} x^{\nu} - b^{\mu}$, takes the form

$$x'^{d+1} = e^{-r}x^{d+1} + t(e^{-r}\zeta - \sinh r) - e^{-r}\alpha^a \mathbf{R}_{ab}x^b - b^{d+1},$$

(34)
$$t' = e^r t - (b^0 - b^{d+1})$$

(35) $x'^{b} = \mathbf{R}^{b}_{c}x^{c} - t\alpha^{b} - b^{b}.$

Let $K^{d+1} = J^{0 d+1}$. The Lie algebra of $IG(e_{d+1})$ is spanned by J_{ab} , W_a , P^b , H, \mathcal{M} and K^{d+1} and has non-vanishing commutations relations given by Eqs. (10), (11), (21), (22), (23) and

$$[K^{d+1},H] = H - \mathcal{M},$$

$$[K^{d+1}, \mathcal{M}] = -\mathcal{M},$$

$$[K^{d+1}, W_a] = -W_a$$

Fortunately $\mathcal{I} = \{\mathcal{M}\}$ is still an ideal for the enlarged Lie algebra, and $N = \exp \mathcal{I}$ is a normal subgroup. The quotient group $IG(e_{d+1})/N$ has a well defined action on the space Q of light rays Nx. The Lie algebra of $IG(e_{d+1})/N$ is spanned by $\tilde{J}_{ab} = J_{ab} + \mathcal{I}, \tilde{W}_a = W_a + \mathcal{I}, \tilde{P}^b = P^b + \mathcal{I}, \tilde{H} = H + \mathcal{I}, \text{ and } \tilde{K}^{d+1} = K^{d+1} + \mathcal{I}$ and satisfies the non-vanishing commutation relations

(39)
$$[\tilde{J}_{ab}, \tilde{J}_{cd}] = \delta_{ad}\tilde{J}_{bc} + \delta_{bd}\tilde{J}_{ad} - \delta_{ac}\tilde{J}_{bd} - \delta_{bd}\tilde{J}_{ac},$$

(40)
$$[\tilde{W}_a, \tilde{J}_{bc}] = \delta_{ab}\tilde{W}_c - \delta_{ac}\tilde{W}_b,$$

(41)
$$[P_a, J_{bc}] = \delta_{ab} P_c - \delta_{ac} P_b,$$

(42)
$$[\tilde{W}_a, \tilde{H}] = \tilde{P}_a,$$

(43)
$$[\tilde{K}^{d+1}, \tilde{H}] = \tilde{H},$$

(44)
$$[\tilde{K}^{d+1}, \tilde{W}_a] = -\tilde{W}_a.$$

This is the Lie algebra of the Galilei group plus time dilations whose action on Q is given by Eqs. (34) and (35). The boost K^{d+1} on directions e_{d+1} generates time dilations in the quotient spacetime.

The study of section 2.3 on the non-inertial observers can be generalized. The result is that condition (i) of that section can be dropped. In order to stay time by time in $S(e_{d+1})$ an observer must satisfy the following condition. The projections of the angular velocity vector and acceleration on the plane perpendicular to e_{d+1} must be perpendicular to each other and of the same magnitude, while their vector product must equal e_{d+1} up to a non-negative factor. Thus, contrary to what




FIGURE 3. The observer O' has velocity $\mathbf{v} = v\mathbf{e}_2$ with respect to O and is slightly rotated in such a way that for both frames the light from Σ has direction along the third axis. Consider two light beams from Σ . If they hit the first screen at the same time according to O's time then they hit the second screen at the same time according to O''s. The Minkowskian distance d between the two pairs of events is the same which, due to the previous result, is another way to say that the distance between the spots on the 'screen' $x^3 = 0$ as measured by O is the same of the distance between the spots on the screen $x'^3 = 0$ as measured by O'. The moving screens in this figure are obtained from the system $x'^3 = 0$ and $x^0 = C$, by varying C.

happens for the $S_{\omega}(e_{d+1})$ subset, here the acceleration is not constrained, since the angular velocity (orientation) can always be chosen such that the observer belongs time by time to $S(e_{d+1})$.

As a consequence of Eq. (34), if two light spots are simultaneous in a frame belonging to $S(e_{d+1})$ then they are simultaneous in every frame in the same set (universality of Galilean simultaneity). This property gives a way of splitting the space of null geodesics Q of direction n into subsets of 'simultaneous' null geodesics S_t where t is the time of arrival of the light beam on the screen of a representative observer in $S(e_{d+1})$. These subsets are the planes $n \cdot x = cnst$. The label t is required in order to distinguish between the different sets but its actual value is not important.

The Eq. (35) implies that the distance between two simultaneous light spots does not change under frame changes, that is, the scalar $x^a x_a$ is an invariant. Thus, this scalar gives a well defined Euclidean metric on the d-dimensional space of simultaneous null geodesics of a given direction S_t . Figure 3 summarizes these results.

2.7. The shape invariance of shadows. Shadows are produced by the absence of a light. Where there is a dark spot we can think that a light beam has been screened by the object that produces the shadow. Shadows propagate as light does and thus we have found a transformation rule for shadows on perpendicular-to-light screens under inertial frame changes. The shadow of a pointlike particle worldline is a pointlike shadow worldline which we can think as composed of a continuous succession of dark spots.

Some observations are in order:

- : (i) Shadows do not deform their shape under the change of frame in $S(e_{d+1})$. This result holds for non-inertial observers as well as long as they are oriented in such a way that the light from Σ propagates in the direction of their axis e_{d+1} (i.e. $K(\tau)$ belongs to $S(e_{d+1})$). Indeed at any non-inertial observer proper time τ , the statement holds for the image of the shadow on the screen of $K(\tau)$ which coincides with that of the non-inertial observer. Thus, the shape of a shadow can not be deformed by the arbitrary motion of a orthogonal-to-light screen.
- : (ii) If a shadow changes in time in an orthogonal-to-light screen of an inertial observer, due for instance to the motion, rotation or deformation of the object that projects the shadow, then, up to a Galilean transformation, exactly the same projected movement is seen by a different inertial observer with possibly a different time rate if the respective frames are not related by an inhomogeneous little group transformation.
- : (iii) Since the Galilean group does not impose an upper bound on the velocities we recover the known fact [31, Chap. 6] that shadows can move faster than light or, in more suggestive terms that "darkness is faster than light".
- : (iv) A dark (or light) spot can be at rest in the screen of a suitable inertial frame as it happens if the object that projects the shadow is at rest in that same frame (or as it happens if a screen is opaque and has a hole). Reasoning in the rest frame one can easily conclude that the shadow of an object in uniform motion has the shape of the shadow of the object at rest taken with respect to a suitable direction. In particular the shadow of a sphere is always a disk (as it happens in the spherical picture case [28]).
- : (v) As it is well known the attempt at defining what is a "rigid motion" in special relativity leads to several problems. Rigid motions are particularly restrictive (for instance, Born's rigidity does not allow to put into rotation a disk which was initially at rest) and ultimately, all the difficulties arise from the finite speed of light which makes any kind of "rigidity" constraint quite unnatural. Due to the same reason the "rigid body" does not exist in special relativity. Nevertheless, given a system of particles and a preferential lightlike direction n, it makes perfect sense to define its motion as *shadow rigid* if the shadows of the particles preserve their Euclidean distance in time. Due to the Galilean nature of the transformations, this definition is clearly independent of the orthogonal-to-light screen chosen.

We recall that the little group of the massless particle $L(e_{d+1})$ has been investigated in connection with the classification of elementary particles [33]. There the generators W_a were essentially removed from the little group algebra on the ground that the two degrees of freedom to which they give rise are not observed. Weinberg [32] concludes that the ISO(d) structure of the little group has no clear physical significance in particle physics and that the generators W_a must therefore vanish in the unitary representation. In our physical problem the generators W_a acquire the clear role of Galilean boosts for the transformation of shadows between inertial frames. The role of W_a as 'translations' of the little group ISO(d) expresses the commutativity of Galilean boosts.

3. The reconstruction problem

In this section we consider a non-inertial observer whose comoving inertial frame $K(\tau)$ belongs to $S(e_{d+1})$. Our aim is to recover the motion of the observer starting from the (observable) acceleration $\tilde{a}^{d+1}(\tau)$ along the longitudinal direction e_{d+1} , the acceleration $\tilde{a}^a(\tau)$ along the transverse directions e_a and the angular velocity $\tilde{\omega}_{ab}(\tau)$ in the plane perpendicular to e_{d+1} . Remarkably, the analogous problem in which the proper acceleration is decomposed with respect to a Fermi-Walker transported tetrad has not yet been solved. Nevertheless, as we shall see, it can be solved in this case where, however, the orthonormal basis $\{u, e_a, e_{d+1}\}$ is not Fermi-Walker transported because of the correction needed to reorient the axis e_{d+1} after an infinitesimal boost. In other words, whereas the Fermi-Walker transport is a consequence of the operator $e^{a_i K^i d\tau + H d\tau}$, in our case the infinitesimal boost (plus time translation) has the form $e^{(a_a W^a + a_{d+1}K^{d+1})d\tau + H d\tau}$ and therefore includes an infinitesimal rotation. The transport induced by the operator $e^{(a_a W^a + a_{d+1}K^{d+1})d\tau + H d\tau}$ will be called *lightlike parallel transport* (relative to the null direction n^{μ}). A differential geometric characterization of this parallel transport will be given in appendix A.

Let $x_{K(\tau)}^{\mu} = G^{\mu}_{\ \nu}[x_{K(0)}^{\nu} - x^{\mu}(\tau)]$ be the transformation from K(0) to $K(\tau)$. Taking into account the decomposition $G^{\mu}_{\ \nu} = B^{\mu}_{\ \alpha}S^{\alpha}_{\ \beta}R^{\beta}_{\ \nu}$, the operator that sends $x_{K(0)}^{\mu}$ to $x_{K(\tau)}^{\mu}$ can be written

$$e^{r(\tau)K^{d+1}}e^{\alpha_a(\tau)W^a}e^{\frac{1}{2}\Omega_{ab}(\tau)J^{ab}}e^{t(\tau)H-x^a(\tau)P_a+x^{d+1}(\tau)\mathcal{M}}.$$

where we used the decomposition of x^{μ} , (26). We have to find an expression for $r(\tau)$, $\alpha_a(\tau)$, $\Omega_{ab}(\tau)$ (or alternatively $\mathbf{R}^a_{\ b}(\tau)$, see (7)), $t(\tau)$, $x^a(\tau)$ and $x^{d+1}(\tau)$ given $\tilde{a}^{d+1}(\tau)$, $\tilde{a}^a(\tau)$ and $\tilde{\omega}_{ab}(\tau)$. The infinitesimal transformation of the observer can be modelled as a (proper) time translation followed by a transformation in $L(e_{d+1})$

$$e^{\tilde{a}^{d+1}(\tau)K^{d+1}\mathrm{d}\tau}e^{\tilde{a}^{a}(\tau)W_{a}\mathrm{d}\tau}e^{\frac{1}{2}\tilde{\omega}_{ab}(\tau)J^{ab}\mathrm{d}\tau}e^{H\mathrm{d}\tau}.$$

Note that the space translations do not enter at the infinitesimal level, as they would imply a violation of causality.

The dependence of $r(\tau)$, $\alpha_a(\tau)$, $\Omega_{ab}(\tau)$, $t(\tau)$, $x^a(\tau)$ and $x^{d+1}(\tau)$ on time is recovered from the constraint

$$\begin{split} & [e^{\tilde{a}^{d+1}(\tau)K^{d+1}\mathrm{d}\tau}e^{\tilde{a}^{a}(\tau)W_{a}\mathrm{d}\tau}e^{\frac{1}{2}\tilde{\omega}_{ab}(\tau)J^{ab}\mathrm{d}\tau}e^{H\mathrm{d}\tau}][e^{r(\tau)K^{d+1}}\\ & e^{\alpha_{a}(\tau)W^{a}}e^{\frac{1}{2}\Omega_{ab}(\tau)J^{ab}}e^{t(\tau)H-x^{a}(\tau)P_{a}+x^{d+1}(\tau)\mathcal{M}}]\\ & = e^{r(\tau+\mathrm{d}\tau)K^{d+1}}e^{\alpha_{a}(\tau+\mathrm{d}\tau)W^{a}}e^{\frac{1}{2}\Omega_{ab}(\tau+\mathrm{d}\tau)J^{ab}}\\ & e^{t(\tau+\mathrm{d}\tau)H-x^{a}(\tau+\mathrm{d}\tau)P_{a}+x^{d+1}(\tau+\mathrm{d}\tau)\mathcal{M}}, \end{split}$$

In order to find $r(\tau + d\tau)$, $\alpha_a(\tau + d\tau)$, $\Omega_{ab}(\tau + d\tau)$, $t(\tau + d\tau)$, $x^a(\tau + d\tau)$ and $x^{d+1}(\tau + d\tau)$, we have to use the commutation rules of $\mathcal{IG}(e_{d+1})$ several times. The following observation will be particularly useful. The Campbell-Baker-Hausdorff formula

(45)
$$e^{A} e^{B} = e^{A+B+\frac{1}{2}[A,B]+\frac{1}{12}[A,[A,B]]},$$

is correct at any order if [B, [B, A]] = 0 and [A, [A, B]] commutes with A, B and [A, B]. This fact implies the identity

(46)
$$e^B e^A = e^A e^B e^{-[A,B]} e^{\frac{1}{2}[A,[A,B]]},$$

which is very useful in order to move $e^{Hd\tau}$ on the right hand side of $e^{\alpha_a(\tau)W^a}$. Also from the expression (33) or (32) it can be easily checked that

$$e^{H\mathrm{d}\tau}e^{r(\tau)K^{d+1}} = e^{r(\tau)K^{d+1}}e^{e^{-r(\tau)}H\mathrm{d}\tau + \sinh r(\tau)\mathcal{M}\mathrm{d}\tau},$$

and

$$e^{\tilde{a}^{a}(\tau)W_{a}\mathrm{d}\tau}e^{r(\tau)K^{d+1}} = e^{r(\tau)K^{d+1}}e^{e^{r(\tau)}\tilde{a}^{a}(\tau)W_{a}\mathrm{d}\tau}.$$

In the end we arrive at the following set of differential equations

1.0.0

(47)
$$\frac{\mathrm{d}\mathbf{R}^a_{\ b}}{\mathrm{d}\tau} = \tilde{\omega}^a_{\ c}\mathbf{R}^c_{\ b}$$

(48)
$$\frac{\mathrm{d}r}{\mathrm{d}\tau} = \tilde{a}^{d+1},$$

(49)
$$\frac{\mathrm{d}t}{\mathrm{d}\tau} = e^{-r},$$

(50)
$$\frac{\mathrm{d}\alpha^a}{\mathrm{d}\tau} = e^r \tilde{a}^a + \tilde{\omega}^a_{\ b} \alpha^b,$$

(51)
$$\frac{\mathrm{d}x^a}{\mathrm{d}\tau} = e^{-r}\alpha_b \mathrm{R}^{ba},$$

(52)
$$\frac{\mathrm{d}x^{d+1}}{\mathrm{d}\tau} = \sinh r + \frac{e^{-r}}{2}\alpha^a\alpha_a.$$

The lightlike parallel transport is obtained for $\tilde{\omega}_c^a = 0$. Note that each one of the first three equations is independent of the others. It can be easily checked that

(53)
$$u^{\mu} = \frac{\mathrm{d}x^{\mu}}{\mathrm{d}\tau} = \begin{pmatrix} \cosh r + \frac{e^{-r}}{2} \alpha^{a} \alpha_{a} \\ e^{-r} \alpha_{b} \mathrm{R}^{ba} \\ \sinh r + \frac{e^{-r}}{2} \alpha^{a} \alpha_{a} \end{pmatrix},$$

and $u^{\mu}u_{\mu} = -1$. Thus $x^{\mu}(\tau)$ is timelike as expected.

An interesting consequence of Eq. (49) is that if, for any τ , $K(\tau) \in S_{\omega}(e_{d+1})$ for a suitable ω , then r = 0 and $t = \tau$, that is the proper time of $K(\tau)$'s origin coincides with the proper time of its shadow on K(0)'s screen.

3.1. Absolute transverse orientation. The equation (47), although with a different interpretation, has been studied in many references [21]. Fortunately, it can be completely integrated in the 4-dimensional spacetime case, since d = 2. Let $\tilde{\omega}^{\mu}$ be the angular velocity vector

$$\mathbf{R}^{a}_{\ b}(\tau) = \begin{pmatrix} \cos\theta(\tau) & \sin\theta(\tau) \\ -\sin\theta(\tau) & \cos\theta(\tau) \end{pmatrix}, \quad \omega^{a}_{\ b}(\tau) = \tilde{\omega}^{d+1}(\tau) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

then $\theta(\tau) = \int_0^{\tau} \tilde{\omega}^{d+1}(\tau) d\tau$. More generally the fact that Eq. (47) is independent of the other equations implies that $\mathbf{R}^a_{\ b}(\tau)$ (and hence $R^{\mu}_{\ \nu}(\tau)$) is independent of the acceleration history of the frame. This feature is due to the existence of an absolute transverse orientation induced by the lightlike parallel transport. In other words assume that $\tilde{\omega}^a_{\ c} = 0$, then if the frame K(0) can be moved staying time by time into $S(e_{d+1})$ to $K(\tau)$ it can not be moved to any other frame $K'(\tau)$ which differs from $K(\tau)$ only for a rotation which keeps e_{d+1} fixed. The unique orientation of $K(\tau)$, dependent on the choice of initial orientation of K(0), and the arbitrariness in the origin of coordinates of $K(\tau)$, implies an absolute transverse orientation on spacetime. To see this consider the following argument. On the set $S(e_{d+1})$ (resp. $S_{\omega}(e_{d+1})$) acts freely the subgroup $IGP(e_{d+1})$ generated by K^{d+1} , W_a , H, \mathcal{M} , P_a (resp. $ILP(e_{d+1})$ generated by the same set without K^{d+1}). The quotient space is isomorphic to the group SO(d). Thus the set $S(e_{d+1})$ (resp. $S_{\omega}(e_{d+1})$) splits into classes $S_{\mathbf{R}}(e_{d+1})$ (resp. $S_{\mathbf{R},\omega}(e_{d+1})$) where the abstract symbol R belongs to the quotient space isomorphic to SO(d) and denotes the absolute orientation of the frame induced by the lightlike parallel transport. An observer belonging to $S_{\mathbf{R}}(e_{d+1})$ can not move, at a later proper time, to a frame belonging $S_{\mathbf{R}'}(e_{d+1})$, $R' \neq R$, if over it does not act an operator which does not belong to $IGP(e_{d+1})$ (resp. $ILP(e_{d+1})$), i.e. if in the mean time $\tilde{\omega}_{ab} = 0$.

The following fact can be easily checked using Eq.(47)-(52). Let

$$\{{\mathbf{R'}}^a_b, r', t', {\alpha'}^a, {x'}^a, {x'}^{d+1}\}(\tau)$$

be a solution of the above equations given $\{\tilde{\omega}'_{ab}, \tilde{a}'^{a}, \tilde{a}'^{d+1}\}(\tau)$, then

$$\{\mathbf{R}^{a}_{\ b} = \delta^{a}_{b}, r = r', t = t', \alpha^{a} = \alpha'_{b} \mathbf{R'}^{ba}, {x'}^{a} = x^{a}, x^{d+1} = {x'}^{d+1}\}(\tau)$$

is a solution of the above equations given $\{\tilde{\omega}_{ab} = 0, \tilde{a}^a = \tilde{a}'_b \mathbf{R}'^{ba}, \tilde{a}^{d+1} = \tilde{a}'^{d+1}\}(\tau)$. It follows that the integration of the system reduces to equation (47), that can be completely solved if d = 2, and to the system (48)-(52) with $\omega_{ab} = 0$, $R^a_{\ b} = \delta^a_b$ which describes a lightlike transported frame.

3.2. Integration of the lightlike parallel transport and spacetime navigation. We integrate the system in the case of a lightlike transported frame, i.e. $\omega_{ab} = 0$, $R^a_{\ b} = \delta^a_b$. Since for $\tau = 0$ the frame of the non-inertial observer in $S(e_{d+1})$ coincides with K(0) we have, to begin with,

(54)
$$r(\tau) = \int_0^\tau \tilde{a}^{d+1}(\tau') \mathrm{d}\tau',$$

(55)
$$t(\tau) = \int_{0}^{\infty} e^{-r(\tau')} \mathrm{d}\tau',$$

(56)
$$\alpha^a(\tau) = \int_0^\tau e^{r(\tau')} \tilde{a}^a(\tau') \mathrm{d}\tau'.$$

Eqs. (49) and (51) imply

(57)
$$\frac{\mathrm{d}x^a}{\mathrm{d}t} = \alpha^a,$$

that is, the shadow of $K(\tau)$'s origin has a velocity on K(0)'s screen which equals the group parameter α^a . Moreover, if $K(\tau) \in S_{\omega}(e_{d+1})$ for any τ then r = 0 and using Eqs. (49) and (50)

(58)
$$\frac{\mathrm{d}\alpha^a}{\mathrm{d}t} = \tilde{a}^a,$$

which means that the acceleration of the shadow on the screen equals the actual transverse acceleration measured by the non-inertial observer using comoving accelerometers.

In general any timelike worldline $x^{\mu}(\tau)$ starting at the origin of K(0) can be interpreted as the motion of the origin of a lightlike transported non-inertial frame. Given $x^{\mu}(\tau)$ one can calculate $u^{\mu}(\tau)$, hence $r(\tau) = -\ln(u^0 - u^{d+1})$, $\alpha_a(\tau) = e^r u_a$ and finally the longitudinal and transversal acceleration measured by the noninertial observer through Eqs. (48) and (50). The inverse problem of obtaining the trajectory from the measured acceleration is the reconstruction problem.

The worldline $x^{\mu}(\tau)$ can be obtained without difficulties from Eqs. (53), (54) and (56). The provided analytical solution of the reconstruction problem could be applied in futurable spacetime navigation. Far from the massive sources of gravity the spacetime is almost Minkowskian. A method that gives to the non-inertial observer a way to recover its own inertial coordinates would be welcome as in this way a direct communication of the observer with an inertial observer K(0) would be avoided. Indeed, as the distance between the observers increases, a communication between them becomes unlikely. In any case, such communication would introduce a delay that would practically forbid an autonomous spacetime navigation, and hence the possibility of correcting the trajectory in short decisional times. The formulas given here allow an autonomous navigation of the non-inertial observer (spaceship).

The comoving laboratory should be provided with an accelerometer and three orthogonal gyroscopes that time by time correct their own orientation so that the last one (the direction e_{d+1}) points always towards the opposite direction of a given star in the night sky sphere (the light from the star determines the null direction n). The correction must consist in infinitesimal rotations along axes perpendicular to e_{d+1} . If not corrected the gyroscopes would give a Fermi-Walker transported frame. In other words a suitable onboard instrumentation can reproduce without difficulties a lightlike transported frame (coincident up to a reorientation to the comoving Fermi-Walker transported triad). The measured acceleration can therefore be projected on the gyroscopic directions to obtain $\tilde{a}^{d+1}(\tau)$ and $\tilde{a}^b(\tau)$. Given the acceleration history, the non-iniertial observer can recover the coordinates $x^{\mu}(\tau)$ by integration using the above formulas. As far as we know, the analytical solution to the reconstruction problem provided here represents the only one available for the case d > 0.

3.3. Motion reconstruction from projection and meaning of the action. In this subsection we consider another kind of reconstruction problem. This time we assume to have given the shadow motion of $K(\tau)$'s origin on K(0)'s screen and r(t) (or alternatively the frequency $\frac{\omega'(t)}{\omega(0)} = e^{-r(t)}$). Then, $K(\tau(t))$'s origin in K(0)'s coordinates is (with a dot we denote the differentiation with respect to t)

(59)
$$x^{\mu}(t) = \begin{pmatrix} \int_{0}^{t} \frac{e^{2r(t')} + 1}{2} dt' + \int_{0}^{t} \frac{1}{2} \dot{x}^{a} \dot{x}_{a} dt' \\ x^{a}(t) \\ \int_{0}^{t} \frac{e^{2r(t')} - 1}{2} dt' + \int_{0}^{t} \frac{1}{2} \dot{x}^{a} \dot{x}_{a} dt' \end{pmatrix}.$$

This equation gives a meaning to the classical action as it shows that $\int_0^t \frac{1}{2} \frac{dx^a}{dt} \frac{dx^a}{dt} dt'$ represents the coordinate $x^{d+1}(t)$ of $K(\tau)$'s origin if $K(\tau) \in S_{\omega}(e_{d+1})$ (hence r = 0), given the shadow worldline $x^a(t)$ of $K(\tau)$'s origin.

3.4. Differential aging from acceleration. In this section we generalize to arbitrary spacetime dimensions (d + 1) + 1, a formula for the differential aging in terms of the acceleration history given by the author in [25] for the d = 0 case. By differential aging Δ we mean the difference between the proper time T needed by an inertial observer to go from $x^{\mu}(0)$ to $x^{\mu}(\tau)$ and the proper time τ of the non-inertial

observer that follows the trajectory $x^{\mu}(\tau)$. It is well known that $T(\tau) \geq \tau$ where the equality holds iff $x^{\mu}(\tau)$ is a straight line, however, an explicit formula for T or $\Delta = T - \tau$, in terms of the acceleration history is in general difficult to obtain. The above results lead to

$$T^{2} - \tau^{2} = \sum_{a=1}^{d} \{ [\int_{0}^{\tau} e^{-r(\tau')} d\tau'] [\int_{0}^{\tau} e^{-r(\tau')} \alpha^{a} \alpha^{a} d\tau'] - [\int_{0}^{\tau} e^{-r(\tau')} \alpha^{a} d\tau']^{2} \}$$

(60)
$$+ \{ [\int_{0}^{\tau} e^{-r(\tau')} d\tau'] [\int_{0}^{\tau} e^{r(\tau')} d\tau'] - \tau^{2} \},$$

where r and α are given by Eqs. (54) and (56). We did not use the sum-overrepeated-indexes convention to point out that the *i*-th term between braces on the right-hand side is, thank to the Cauchy-Schwarz inequality, greater than zero and vanishing iff $\tilde{a}^i(\tau') = 0$ for every $\tau' \in [0, \tau]$. This observation leads to the expected inequality $T \geq \tau$, where the equality holds iff the acceleration vanishes. If d = 0 the right hand side reduces to the last term and the formula reduces the the differential aging formula given in [25].

The expression for T given the shadow worldline $x^{a}(t)$ and r(t) reads instead

(61)
$$T^{2} = t \int_{0}^{t} \dot{x}^{a} \dot{x}_{a} dt' - x^{a} x_{a} + t \int_{0}^{t} e^{2r(t')} dt'.$$

3.5. Position and frequency drifts. The last component of the non-inertial observer worldline $x^{\mu}(\tau)$ is

(62)
$$\begin{aligned} x^{d+1}(t) &= x^{d+1}(0) + \frac{1}{2} \int_0^t [e^{2r(t')} - 1 + \dot{x}^a \dot{x}_a] dt' \\ &= x^{d+1}(0) + \frac{1}{2} \int_0^t [(\frac{\omega}{\omega'(t)})^2 - 1 + \dot{x}^a \dot{x}_a] dt', \end{aligned}$$

where ω is the frequency of light going in direction e_{d+1} as measured in the inertial frame K and $\omega'(t)$ is the frequency measured by the non-inertial observer. The relevance of this equation comes from its unexpected consequences. For instance, assume that the non-inertial observer stays very close to the last axis of K, so that $0 \simeq |x^a| \ll |x^{d+1}|$, and the distance is given by $|x^{d+1}|$ with a small error. Moreover, assume that $\omega' = \omega$. If $x^{d+1} < 0$ the signal of frequency ω could be thought as emitted by the non-inertial observer. The inertial observer measures the same frequency and in practice knows quite accurately the position of the non-inertial observer up to the value of coordinate x^{d+1} . From the fact that the frequency does not change her could be tempted to infer that x^{d+1} is constant in time as a velocity component along the axis would imply a Doppler effect. However, the previous formula gives

$$x^{d+1}(t) = x^{d+1}(0) + \frac{1}{2} \int_0^t \dot{x}^a \dot{x}_a \mathrm{d}t',$$

which means that, due to the transversal motion, the position may drift along the last axis without any frequency change. In particular the drift is such that the coordinate x^{d+1} can only increase. Thus the knowledge of the transversal position of the non-inertial observer does not place a bound on the transversal velocities which are at the origin of the effect. Remarkably, transversal velocities are difficult to measure and hence the effect may be present in practical applications.

A related effect takes place if the second non-inertial observer is positioned near the last axis of K, i.e. $0 \simeq |x^a| \ll |x^{d+1}|$ at a fixed value of the coordinate x^{d+1} . Assume that $x^{d+1} > 0$, so that ω can be identified with the frequency of a wave emitted by K and received by the non-inertial observer. It could be naively expected that the measured frequency ω' is constant since the non-inertial observer is almost at rest, but differentiation of Eq. (62) gives

$$\omega'(t) = \frac{\omega}{\sqrt{1 - \dot{x}^a \dot{x}_a(t)}}$$

The transversal velocity can change the frequency without affecting the average position.

4. FROM POINCARÉ TO GALILEI INVARIANCE: PARTICLE COLLISIONS

If a Poincaré invariant physical phenomenon in (d+1)+1-Minkwoski spacetime is projectable on Q then the projected phenomenon is Galilei invariant [11]. Indeed, the original physical phenomenon is invariant under the Poincaré subgroup $IL(\mathbf{e}_{d+1})$ which means that the projection is invariant under the Galilei group Gal(d). The aim of these last sections is to show explicitly the Galilei invariance of the shadow of a relativistic collision.

We choose an arbitrary inertial observer in $S_{\omega}(e_{d+1})$ of coordinates $\{x^{\mu}\}$ and consider a particle of worldline $x^{\mu}(\lambda)$ and momentum $p^{\mu} = dx^{\mu}/d\lambda$. We are interested on the lightlike projection $x^{a}(t)$ of the curve $x^{\mu}(\lambda)$ on the screen $x^{d+1} = 0$, where the projected curve is parametrized with the Galilean time t. The velocity of the shadow on the screen is

(63)
$$\dot{x}^a = \frac{\mathrm{d}x^0}{\mathrm{d}t}v^a = \frac{v^a}{1 - v^{d+1}} = \frac{p^a}{p^0 - p^{d+1}}.$$

Let us temporarily focus on a massive particle on M and then generalize the results to the massless case. The equation of motion for the free particle of mass m is obtained from Hamilton's principle in configuration space (see Eq. (31))

(64)
$$0 = \delta \int m \mathrm{d}\tau = \delta \int_{t_0}^{t_1} \sqrt{2\dot{x}^{d+1} + 1 - \dot{\mathbf{x}}^2} \, \mathrm{d}t.$$

The momentum conjugated to x^{d+1} is the shadow mass

(65)
$$\tilde{m} = -n^{\mu}p_{\mu} = p^0 - p^{d+1}$$

The last expression makes sense also for massless particles on M and it is invariant under inertial frame changes generated by $IL(\mathbf{e}_{d+1})$. The cyclic variable x^{d+1} can be removed by using Routh's reduction [24, Sect. 8.9]. On the reduced spacetime Q Hamilton's action principle holds where the new Lagrangian is replaced by the Routhian, i.e. a suitable Legendre transform of the original Lagrangian in which the conjugated momentum \tilde{m} is regarded as a constant. In our case the Routhian is

$$\frac{\tilde{m}}{2}\dot{\mathbf{x}}^2 - \frac{1}{2}[\frac{m^2}{\tilde{m}} + \tilde{m}]$$

From this expression we deduce the shadow kinetic energy T, which has the same form as in classical mechanics, and the shadow internal energy I

(66)
$$T = \frac{\tilde{m}}{2}\dot{\mathbf{x}}^2$$

(67)
$$I = \frac{1}{2} \left[\frac{m^2}{\tilde{m}} + \tilde{m} \right]$$

With minor differences these identifications can also be found in [5]. The internal energy depends on a parameter m which has no (shadow) kinematical interpretation. Indeed, in classical non-relativistic mechanics the internal energy of a particle can not be expressed in terms of the mass alone as in special relativity. For instance, a body A composed by bodies B, C and a compressed spring has a total mass given by $m_A = m_B + m_C$. Nevertheless, if the particle A separates into Band C, the potential energy of the spring must be taken into account in the energy balance. It is therefore natural that a non-kinematical parameter m enters in the expression of the shadow internal energy. The fact that the above identification of kinetic and internal energy is correct will be confirmed in a moment when we shall study the conservation of energy.

Note that in any case the potential energy is a constant and therefore it is irrelevant in the variational principle. We conclude that the reduced action coincides with the classical action with the *shadow masses* provided by Eq. (65).

The shadow mass $\tilde{m} = p^0 - p^{d+1}$ is positive unless the particle on M is massless with momentum of direction n^{μ} , in which case it vanishes. This fact is coherent with the previous observation that those massless particles on M do not represent particles on Q but, rather, events. Analogously, the definitions of shadow kinetic energy and shadow internal energy generalize to massless particles on M. Indeed they can be directly expressed in terms of the covariant momentum p^{μ}

(68)
$$T = \frac{\tilde{m}}{2}\dot{\mathbf{x}}^2 = \frac{p^a p_a}{2(p^0 - p^{d+1})},$$

(69)
$$I = \frac{1}{2} \left[\frac{m^2}{\tilde{m}} + \tilde{m} \right] = \frac{1}{2} \left[\frac{m^2}{p^0 - p^{d+1}} + p^0 - p^{d+1} \right].$$

The shadow momentum can be obtained from Eqs. (65) and (63)

(70)
$$\tilde{m}\dot{x}^a = p^a$$

The energy reads

(71)
$$E = T + I = \frac{1}{2} \frac{p^a p_a}{p^0 - p^{d+1}} + \frac{1}{2} [\frac{m^2}{p^0 - p^{d+1}} + p^0 - p^{d+1})] = p^0.$$

4.1. Conservation of shadow mass, momentum and energy. Let us now consider a collision of N particles of momentum $p_{(i)}^{\mu}$, i = 1, ..., N. From the collision N' final particles of momentum $p_{(j)}^{\prime \mu}$, j = 1, ..., N', emerge. The conservation of momentum in Minkowski spacetime reads

(72)
$$\sum_{i=1}^{N} p_{(i)}^{\mu} = \sum_{i=1}^{N'} p_{(i)}^{\prime \mu}$$

The particle worldlines are geodesics that project into geodesics of Q: the shadow worldlines. These straight lines in Q collide. Subtracting Eq. (72) for $\mu = 0$ and

 $\mu = d + 1$ we obtain

(73)
$$\sum_{i=1}^{N} \tilde{m}_{(i)} = \sum_{i=1}^{N'} \tilde{m}'_{(i)},$$

that is, the total shadow mass is conserved in the shadow collision. From Eq. (72) with $\mu = 1, \ldots, d$ we obtain

(74)
$$\sum_{i=1}^{N} \tilde{m}_{(i)} \dot{\mathbf{x}}_{(i)} = \sum_{i=1}^{N'} \tilde{m}'_{(i)} \dot{\mathbf{x}}'_{(i)},$$

that is, the shadow momentum is conserved. The found conservation principles are characteristic of a non-relativistic system. Indeed they are consequence of the underlying Galilean symmetry.

Let us come to the conservation of energy. Since the relativistic energy is conserved, from Eq. (71) we deduce that the shadow energy is conserved

(75)
$$\sum_{i=1}^{N} E_{(i)} = \sum_{i=1}^{N'} E'_{(i)}.$$

In order to prove the conservation of shadow mass, momentum and energy we used four independent momentum conservation equations on M. Thus the obtained shadow conservation principles are equivalent to the original ones given by Eq. (72). However, there is an important difference. The internal energy in the classical case is not a kinematical observable contrary to what happens in the relativistic case where the energy of the particle can be inferred from the mass and the covariant velocity. In some interesting cases this unknown internal energy plays no role in the shadow collision.

4.2. Conservation of shadow kinetic energy and massless particles. We give some definitions. An *elastic* relativistic collision in M is a collision (i.e. Eq. (72) holds) in which the particle species are conserved, N = N' and $m_{(i)} = m'_{(i)}$, $i = 1, \ldots, N$. An *elastic classical* collision in Q is a collision (i.e. Eqs. (73) and (74) hold) in which the kinetic energy is conserved

(76)
$$\sum_{i=1}^{N} \frac{\tilde{m}_{(i)}}{2} \dot{\mathbf{x}}_{(i)}^2 = \sum_{i=1}^{N'} \frac{\tilde{m}_{(i)}'}{2} \dot{\mathbf{x}}_{(i)}'^2.$$

We shall say that a classical elastic collision is *proper* if the particle species are conserved N = N', $\tilde{m}_{(i)} = \tilde{m}'_{(i)}$, i = 1, ..., N and the total kinetic energy is conserved

From the expression of the shadow internal energy we obtain that the shadow of an elastic relativistic collision is a proper elastic classical collision provided the shadow masses are preserved in the collision, $\tilde{m}_{(i)} = \tilde{m}'_{(i)}$, i = 1, ..., N.

Finally, let us consider the shadow of a collision in which the initial and final particles are massless. Since the sum of two non-collinear null vectors is a timelike vector, a collision of this form must involve at least two initial and two final particles. From the expression of the internal energy, $I = \tilde{m}/2$, we conclude that the shadow of a collision that involves only massless particles is a classical elastic collision.

22

4.3. The inverse problem. Suppose we are given the data of a collision on d+1 Galilean spacetime. In this section we want to construct a collision in (d+1)+1 Minkowski spacetime such that the original collision can be regarded as its shadow. Thus assume that we are given numbers N, N', of initial and final particles, their (positive) masses $\tilde{m}_{(i)}$ and $\tilde{m}'_{(i)}$ and their velocities $\dot{\mathbf{x}}_{(i)}, \dot{\mathbf{x}}'_{(i)}$. Assume the conservation of total mass

(77)
$$\sum_{i=1}^{N} \tilde{m}_{(i)} = \sum_{i=1}^{N'} \tilde{m}'_{(i)},$$

and momentum

(78)
$$\sum_{i=1}^{N} \tilde{m}_{(i)} \dot{\mathbf{x}}_{(i)} = \sum_{i=1}^{N'} \tilde{m}'_{(i)} \dot{\mathbf{x}}'_{(i)}.$$

We want to lift the collision to M in a way invariant under Galilei transformations, i.e. the lift to M must depend only on the collision on Q and not on the choice of Galilean frame on Q. We known that the kinetic energy $T_{(i)}$ of particle (i) and the total kinetic energy $T = \sum_{i}^{N} T_{(i)}$, are not invariant under Galilean transformation. Nevertheless, the difference between final and initial kinetic energies is invariant

(79)
$$\Delta T = T' - T = \sum_{i=1}^{N'} \frac{1}{2} \tilde{m}'_{(i)} \dot{\mathbf{x}}'^2_{(i)} - \sum_{i=1}^{N} \frac{1}{2} \tilde{m}_{(i)} \dot{\mathbf{x}}^2_{(i)}.$$

This fact can be easily checked using the conservation of Galilean mass and momentum assumed above. One should be careful because $\Delta T_{(i)}$ is not Galilei invariant. The quantity ΔT and the masses $\tilde{m}_{(i)}$, $\tilde{m}'_{(i)}$, do not depend on the Galilean frame chosen. Assume that constants $m_{(i)}$ and $m'_{(i)}$ are given such that

(80)
$$2\Delta T = \sum_{i}^{N} \frac{m_{(i)}^{2}}{\tilde{m}_{(i)}} - \sum_{i}^{N'} \frac{m_{(i)}^{\prime 2}}{\tilde{m}_{(i)}^{\prime}}.$$

By giving these numbers one fixes the (Galilei invariant) internal energy of the particles involved in the collision on Q according to the formula (67). Clearly with this definition of $I_{(i)}$ the total energy is conserved

(81)
$$\sum_{(i)}^{N} E_{(i)} = \sum_{(i)}^{\bar{N}} \bar{E}_{(i)}$$

Then the collision on Q is the shadow of a collision in (d+1)+1 Minkowski spacetime constructed as follows. The particles have masses $m_{(i)}$ $(m'_{(i)})$ and covariant momentum

$$p_{(i)}^{\mu} = \begin{pmatrix} E \\ \tilde{m}\dot{\mathbf{x}} \\ E - \tilde{m} \end{pmatrix}_{(i)} = (E - \tilde{m})_{(i)}n^{\mu} + \begin{pmatrix} \tilde{m} \\ \tilde{m}\dot{\mathbf{x}} \\ 0 \end{pmatrix}_{(i)} = \frac{\tilde{m}}{2} \begin{pmatrix} \dot{\mathbf{x}}^2 + \frac{m^2}{\tilde{m}^2} + 1 \\ 2\dot{\mathbf{x}} \\ \dot{\mathbf{x}}^2 + \frac{m^2}{\tilde{m}^2} - 1 \end{pmatrix}_{(i)},$$

such that $p_{(i)}^{\mu}p_{\mu(i)} = -m_{(i)}^2$. With these definitions the total momentum is conserved. Now, we have to give some more data in order to fix the motion of the particles in M. Since we have fixed the momentum of the particles we have only to give the event of collision. It must be chosen in the null worldline that projects on the event of collision in Q. This gives one more real parameter. Once fixed the motion of the particles is determined both forward and backward in time with respect to the instant of collision.

Particularly interesting are the elastic collisions on Q, i.e. $\Delta T = 0$. In this case, up to translations generated by \mathcal{M} , there is a canonical $\frac{1}{\mu}$ -lift from Q to M, $\frac{1}{\mu} \in [0, +\infty)$ obtained by setting $m_{(i)} = \frac{1}{\mu}\tilde{m}_{(i)}$ and analogously for $m'_{(i)}$. Indeed with this choice, due to the conservation of (shadow) mass, the constraint (80) is satisfied. The internal energy becomes $I_{(i)} = \frac{1+1/\mu^2}{2}\tilde{m}_{(i)}$, and (restoring c) we find that for $\mu = 1$ it takes the usual relativistic form, $\tilde{m}_{(i)}c^2$.

There are two natural choices for $1/\mu$

Lightlike 0-lift, $\frac{1}{\mu} = m_{(i)} = m'_{(i)} = 0$: . The elastic collision on Q can be regarded as the shadow of a collision on M in which only massless particles of momentum

(82)
$$p_{(i)}^{\mu} = \frac{\tilde{m}_{(i)}}{2} \begin{pmatrix} \dot{\mathbf{x}}^2 + 1\\ 2\dot{\mathbf{x}}\\ \dot{\mathbf{x}}^2 - 1 \end{pmatrix}_{(i)}$$

are involved.

Timelike 1-lift, $\frac{1}{\mu} = 1$, $m_{(i)} = \tilde{m}_{(i)}$, $m'_{(i)} = \tilde{m}'_{(i)}$: The elastic collision on Q can be regarded as the shadow of a collision between particles of momentum

(83)
$$p_{(i)}^{\mu} = \frac{\tilde{m}_{(i)}'}{2} \begin{pmatrix} \dot{\mathbf{x}}^2 + 2\\ 2\dot{\mathbf{x}}\\ \dot{\mathbf{x}}^2 \end{pmatrix}_{(i)},$$

for which the shadow mass coincides with the mass.

5. Conclusions

Some classical aspects of lightlike dimensional reduction have been studied. The Galilean transformation property of shadows represents surely the most intuitive way to grasp the underlying mathematics. As an interesting consequence it gives to the Galilei group in 2+1 dimensions the status of exact physical symmetry once we agree that it should be applied to the transformation of shadows rather than to events (at least in absence of curvature otherwise even the Poincaré group is broken).

Also, the emphasis made on the role of observers on the full spacetime M allowed us to recognize the usefulness of lightlike dimensional reduction for autonomous spacetime navigation. An important role was played by the concept of lightlike parallel transport which we introduced twice, using a group theoretical definition or an equivalent differential geometric definition (in the appendix).

In the last section we studied in detail the shadow of a relativistic collision. Apart from results that could have been expected in view of the existence of a Galilei quotient subgroup inside the Poincaré group, a further interesting result was obtained which relates the masslessness of the particles involved in the collision to the conservation of kinetic energy in the projected shadow collision.

Acknowledgments

I thank C. Tejero Prieto for suggesting the analogy between this work and Plato's allegory of the cave, and M. Santander and M. Modugno for pointing out some works

on, respectively, front wave dynamics and Newton-Cartan theory. I also would like to thank M. Sánchez for its accurate reading of the manuscript and for the many suggestions. This work has been partially supported by INFN, Grant No. 9503/02.

APPENDIX A. LIGHTLIKE PARALLEL TRANSPORT

In this appendix we give a differential geometric characterization of the lightlike parallel transport already introduced in section 3 by group theoretical means. The results of this section hold as well in a curved spacetime having a covariantly constant null vector field n, $n_{\mu;\nu} = 0$ (the direction of light).

Consider a non-inertial observer on M, that is a timelike worldline $\gamma(\tau)$ parametrized with respect to proper time, and an orthonormal frame $\{u = \partial_{\tau}, e_i\}$ along it. The Fermi-Walker derivative ∇_u^{FW} is a minimal modification of the covariant derivative ∇_u , such that $\nabla_u^{FW} u = 0$. A vector field $v^{\mu}(\tau)$ along γ is Fermi-Walker transported, $\nabla_u^{FW} v = 0$ iff its components with respect to the Fermi-Walker transported tetrad $\{u, e_i^{FW}\}, \nabla_u^{FW} e_i^{FW} = 0$ do not change. As it is well know the condition of preserving the orthonormality, gives, for a generic v,

(84)
$$\nabla_u v_\mu - \nabla_u^{FW} v_\mu = \Omega_{\mu\nu}^{FW} v^\nu,$$

where $\Omega_{\mu\nu}^{FW}$ is a 2-form on γ . The minimal modification which leads to $\nabla_u^{FW} u = 0$ is then the choice

(85)
$$\Omega_{\mu\nu}^{FW} = u_{\mu}a_{\nu} - u_{\nu}a_{\mu}.$$

Analogously we are looking for a lightlike covariant derivative ∇_u^L along γ which measures the variation of the components of a vector field v with respect to the tetrad $\{u, e_i^L\}$ of a lightlike transported frame. Here the lightlike transported frame is such that the null vector field n has always the same components up to a factor that may change in time, i.e. the null vector field n has always the same direction with respect to a lightlike transported frame. Then, with respect to the lightlike transported frame the vector $n^{\mu}/(n^{\beta}u_{\beta})$ has constant components and hence the further condition which defines the lightlike transported frame is f $\nabla_u^L[n^{\mu}/(n^{\beta}u_{\beta})] =$ 0. Let us write

(86)
$$\nabla_u v_\mu - \nabla^L_u v_\mu = \Omega^L_{\mu\nu} v^\nu$$

it is easily seen that the minimal modification of the covariant derivative which satisfies the said properties is obtained by defining

(87)
$$\Omega^{L}_{\mu\nu} = \frac{1}{u^{\beta}n_{\beta}}[a_{\mu}n_{\nu} - a_{\nu}n_{\mu}]$$

(88)
$$= [u_{\mu}a_{\nu} - u_{\nu}a_{\mu}] + \{a_{\mu}[u_{\nu} + \frac{n_{\nu}}{u^{\beta}n_{\beta}}] - a_{\nu}[u_{\mu} + \frac{n_{\mu}}{u^{\beta}n_{\beta}}]\}.$$

The last term represents the angular velocity required to preserve the direction of light n with respect to the comoving tetrad.

APPENDIX B. TRANSFORMATION OF PHOTON POLARIZATION VECTORS

In this appendix we clarify the relation between this work and previous works on the transformation properties of the photon polarization vector. In particular we investigate the relation between gauge transformations and transformations generated by \mathcal{M} . The electromagnetic field $F_{\mu\nu} = 2\partial_{[\mu}A_{\nu]}$ in vacuum satisfies the Maxwell equation $F^{\mu\nu}_{;\nu} = 0$, that, in terms of the potential A_{μ} , reads

(89)
$$\Box A_{\mu} = \partial_{\mu} (\partial \cdot A).$$

By construction this equation is invariant under gauge transformations $A'_{\mu} = A_{\mu} + \partial_{\mu}\alpha(x)$. In this section we are interested in solutions of the form (plane waves)

(90)
$$A_{\mu}(x) = \operatorname{Re}[\epsilon_{\mu}e^{ik_{\nu}x^{\nu}}]$$

where k^{μ} is a null vector and ϵ_{μ} is a complex vector known as *polarization* (for a more general treatment with a complete spectral decomposition of the general solution and a momentum dependent polarization see [30]). This plane wave satisfies Eq. (89) iff $\epsilon^{\nu}k_{\nu} = 0$ which is the equivalent to the Lorentz gauge $\partial \cdot A = 0$. The Lorentz gauge is invariant under the restricted gauge transformations such that $\alpha(x)$ satisfies $\Box \alpha = 0$. In particular $\alpha(x) = \text{Im}[Ce^{ik_{\mu}x^{\mu}}], C \in \mathbb{C}$, satisfies this condition and hence the transformation $\epsilon'_{\mu} = \epsilon_{\mu} + Ck_{\mu}$ comes from a restricted gauge transformation. In order to fix the ideas let us take $k^{\mu} = \omega n^{\mu} = \omega(1, 0, \dots, 0, 1)$.

Choose an inertial frame. For suitable constants $\lambda, \tau \in \mathbb{C}$ (τ is the analog of t), the polarization can be uniquely decomposed in the form

(91)
$$\epsilon^{\mu} = \begin{pmatrix} \lambda + \tau \\ \epsilon \\ \lambda \end{pmatrix} = \lambda n^{\mu} + \begin{pmatrix} \tau \\ \epsilon \\ 0 \end{pmatrix}$$

where $\boldsymbol{\epsilon}$ is the transverse polarization vector. We want to find out how the transverse polarization component $\boldsymbol{\epsilon}$, changes under Lorentz transformations $G(e_{d+1})$ that preserve the direction of n^{μ} .

Formally, the problem considered here is very similar to the one considered previously. However, there are some minor differences: (i) the group considered does not include the translations, indeed here we are are considering the subgroup $G(e_{d+1})$ of the Lorentz group, (ii) despite the fact that there are no translations the gauge transformations play the role of the transformations generated by \mathcal{M} , in that they add to ϵ^{μ} (the analog of x^{μ}) an arbitrary quantity proportional to the null vector, (iii) here there is the additional Lorentz gauge condition $n^{\mu}\epsilon_{\mu} = 0$ which implies $\tau = 0$.

Let us come to the solution of the problem. The transformation G^{μ}_{ν} has the form given by Eq. (33). Thus, if $x'^{\mu} = G^{\mu}_{\ \nu} x^{\nu}$

(92)
$$A'^{\mu} = G^{\mu}_{\ \nu} A^{\nu} = \operatorname{Re}[\epsilon'^{\mu} e^{ik'_{\beta}x'^{\beta}}]$$

where $k'_{\beta} = e^{-r}k_{\beta}$ and $\epsilon'^{\mu} = G^{\mu}_{\nu}\epsilon^{\nu}$. From Eqs. (33) we obtain the transformation of the polarization under the Lorentz transformation G^{μ}_{ν} followed by a restricted gauge transformation of phase $\alpha(x) = \text{Im}[Ce^{ik_{\mu}x^{\mu}}]$

(93)
$$\lambda' = e^{-r}\lambda + \tau(e^{-r}\zeta - \sinh r) - e^{-r}\alpha^a \mathbf{R}_{ab}\epsilon^b + C\omega,$$

(94)
$$\tau' =$$

(95)
$$\epsilon'^{b} = R^{b}_{\ c}\epsilon^{c} - \tau \alpha^{b}, \qquad b = 1, 2$$

 τe^r

We see clearly that if $\lambda = 0$ then in general $\lambda' \neq 0$, but λ can be sent to zero with a suitable gauge transformation (choice of C). The condition $\tau = 0$ implies $\tau' = 0$. In other words the existence of a invariant simultaneity in Galilean relativity $(t' = e^r t \text{ implies that the splitting of spacetime in 'simultaneity' slices <math>t = cnst$. is independent of the frame) is related to the Lorentz invariance of the Lorentz

gauge condition ($\tau = 0 \Rightarrow \tau' = 0$). We conclude that the splitting (91) is invariant even with $\tau = \lambda = 0$ provided the Lorentz transformation in $G(e_{d+1})$ is followed by a suitable gauge transformation [18, 12]. The effect of the transformation on ϵ amounts to a rotation given by Eq. (95) with $\tau = 0$. Note that if $\tau = 0$ and $R^a_{\ b} = I^a_{\ b}$, the parameter λ changes while τ and ϵ do not. This fact led some authors [12, 16, 1, 30] to the conclusion that the generators W_a generate gauge transformations instead of Galilean boosts.

In the study of the polarization and its transformation properties the group N of the previous sections is replaced by a subgroup of the restricted gauge transformations. The invariance of Galilean simultaneity, t = cnst, is replaced by the invariance of the Lorentz condition and, due to this same condition ($\tau = 0$), the Galilean group is not fully appreciated since it reduces to the group of rotations in a *d*-dimensional Euclidean space. Indeed, the invariance of ϵ under Galilean boosts (6) was noticed already in [14] but, as we have shown, the role of the Galilei group could not emerge from studies of the polarization.

References

- Banerjee, R. and Chakraborty, B.: The translation groups as generators of gauge transformation in B ∧ F theory. Phys. Lett. B 502, 291–299 (2001)
- [2] Bargmann, V.: On unitary ray representations of continuous groups. Ann. of Math. 59, 1–46 (1954)
- Bernal, A. N. and Sánchez, M.: Leibnizian, Galilean and Newtonian structures of spacetime. J. Math. Phys. 44, 1129–1149 (2003)
- Bose, S. K.: The Galilean group in 2 + 1 space-times and its central extension. Commun. Math. Phys. 169, 385–395 (1995)
- [5] Bouchiat, C., Fayet, P., and Meyer, P.: Galilean invariance in the infinite momentum frame and the parton model. Nucl. Phys. B 34, 157–176 (1971)
- [6] Dirac, P. A. M.: Forms of relativistic dynamics. Rev. Mod. Phys. 21, 392-399 (1949)
- [7] Duval, C., Burdet, G., Künzle, H. P., and Perrin, M.: Bargmann structures and Newton-Cartan theory. Phys. Rev. D 31, 1841–1853 (1985)
- [8] Duval, C., Gibbons, G., and Horváthy, P.: Celestial mechanics, conformal structures, and gravitational waves. Phys. Rev. D 31, 1841–1853 (1985)
- [9] Ehlers, J.: Examples of Newtonian limits of relativistic spacetimes. Class. Quantum Grav. 14, A119–A126 (1997)
- [10] Eisenhart, L. P.: Dynamical trajectories and geodesics. Ann. Math. (Ser 2) 30, 591–606 (1929)
- [11] Elizade, E. and Gomis, J.: The groups of Poincaré and Galilei in arbitrary dimensional spaces. J. Math. Phys. 19, 1790–1792 (1978)
- [12] Han, D. and Kim, Y. S.: Little group for photons and gauge transformations. Am. J. Phys. 49, 348–351 (1981)
- [13] Harindranath, A.: An Introduction to Light-Front Dynamics for Pedestrians, Ames: International Institute of Theoretical and Applied Physics, vol. Light-Front Quantization and Non-Perturbative QCD (1996)
- [14] Janner, A. and Janssen, T.: Electromagnetic compensating gauge transformations. Physica A 53, 1–27 (1971)
- [15] Julia, B. and Nicolai, H.: Null-Killing vector dimensional reduction and Galilean geometrodynamics. Nucl. Phys. B 439, 291–323 (1994)
- [16] Kim, Y. S.: Internal space-time symmetries of massive and massless particles and their unification. Nucl. Phys. B 102/103, 369–376 (2001)
- [17] Künzle, H. P.: Galilei and Lorentz structures on space-time: comparison of the correspondig geometry and physics. Ann. Inst. H. Poincaré Phys. Theor. 17, 337–362 (1972)
- [18] Kupersztych, J.: Is there a link between gauge invariance, relativistic invariance and electron spin? Il Nuovo Cimento 31, 1–11 (1976)
- [19] Kutach, D.: Non-relativistic quantum mechanics on a Kaluza-Klein manifold (2004)

- [20] Leutwyler, H. and Stern, F.: Relativistic dynamics on a null plane. Ann. Phys. 112, 94–164 (1978)
- [21] Levi, M.: Composition of rotations and parallel transport. Nonlinearity 9, 413–419 (1996)
- [22] Lévy-Leblond, J. M.: Galilei group and nonrelativistic quantum mechanics. J. Math. Phys. 4, 776–788 (1963)
- [23] Lichnerowicz, A.: Théories relativistes de la gravitation et de l'électromagnetisme, Relativité Générale et théories unitaires. Paris: Masson (1955)
- [24] Marsden, J. E. and Ratiu, T. S.: Introduction to Mechanics and Symmetry. New York: Springer (1999)
- [25] Minguzzi, E.: Differential aging from acceleration: An explicit formula. Am. J. Phys. 73, 876–880 (2005)
- [26] Naber, G. L.: The geometry of Minkowski spacetime. New York: Springer-Verlag (1992)
- [27] O'Raifeartaigh, L. and Straumann, N.: Gauge theory: Historical origins and some modern developments. Rev. Mod. Phys. 72, 1–23 (2000)
- [28] Penrose, R.: The apparent shape of a relativistically moving sphere. Proc. Camb. Phil. Soc. 55, 137–139 (1959)
- [29] Penrose, R. and Rindler, W.: Spinors and space-time. Vol. 1. Cambridge Monographs on Mathematical Physics. Cambridge: Cambridge University Press (1984)
- [30] Scaria, T.: Translational groups as generators of gauge transformations. Phys. Rev. D 68, 105013 (2003)
- [31] Schiller, C.: Motion mountain the physics textbook (2003). Available at www.motionmountain.net
- [32] Weinberg, S.: Feynman rules for any spin. II. Massless particles. Phys. Rep. 134, B882–B896 (1964)
- [33] Weinberg, S.: *The Quantum Theory of Fields*, vol. I. Cambridge: Cambridge University Press (1995)

28