Irrotational and Incompressible Ellipsoids in the First Post-Newtonian Approximation of General Relativity

Keisuke Taniguchi, *) Hideki Asada **) and Masaru Shibata ***)

*Department of Physics, Kyoto University, Kyoto 606-8502, Japan

** Faculty of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan

*** Department of Earth and Space Science, Graduate School of Science

Osaka University, Toyonaka, Osaka 560-0043, Japan

First post-Newtonian (1PN) hydrostatic equations for an irrotational fluid which have been recently derived are solved for an incompressible star. The 1PN configurations are expressed as a deformation of the Newtonian irrotational Riemann ellipsoid using Lagrangian displacement vectors introduced by Chandrasekhar. For the 1PN solutions, we also calculate the luminosity of gravitational waves in the 1PN approximation using the Blanchet-Damour formalism. It is found that the solutions of the 1PN equations exhibit singularities at points where the axial ratios of semi-axes are 1: 0.5244: 0.6579 and 1: 0.2374: 0.2963, and the singularities seem to show that at the points, the irrotational Riemann ellipsoid is unstable to the deformation induced by the effect of general relativity. For stable cases $(a_2/a_1 > 0.5244, where a_1 and a_2$ are the semi-major and minor axes, respectively) we find that when increasing the 1PN correction, the angular velocity and total angular momentum increase, while the total energy and luminosity of gravitational waves decrease. These 1PN solutions will be useful when examining the accuracy of numerical code for obtaining relativistic irrotational stars.

We also investigate the validity of an ellipsoidal approximation, in which a 1PN solution is obtained assuming an ellipsoidal figure and neglecting the deformation. It is found that for $a_2/a_1 > 0.7$, the ellipsoidal approximation gives a fairly accurate result for the energy, angular momentum, and angular velocity, although in the approximation we cannot find the singularities.

§1. Introduction

Preparation of reliable theoretical models on the late inspiraling stage of binary neutron stars is one of the most important issues for gravitational wave astronomy. This is because they represent one promising source for gravitational wave detectors such as LIGO, ¹⁾ VIRGO, ²⁾ GEO600³⁾ and TAMA.⁴⁾ From their signals, we will obtain a wide variety of physical information on neutron stars such as their mass and spin, if we have a theoretical template of them.⁵⁾ In particular, a signal from the very late inspiraling stage just prior to merging may contain physically important information on neutron stars such as their radius,⁵⁾ which will be utilized for determining the equation of state of neutron stars.⁶⁾

Binary neutron stars evolve due to the radiation reaction of gravitational waves, so that they never settle down to equilibrium states. However, the emission time scale will always be longer than the orbital period outside their innermost stable

typeset using $\mathcal{PTPTEX}.sty < ver.0.8 >$

^{*)} E-mail: taniguci@tap.scphys.kyoto-u.ac.jp,

^{**)} E-mail: asada@phys.hirosaki-u.ac.jp

^{***)} E-mail: shibata@vega.ess.sci.osaka-u.ac.jp

circular orbit (ISCO), so that we may consider them to be in quasiequilibrium states in their inspiraling phase even near the ISCO.

Until now, all the reliable relativistic works devoted to obtaining a quasiequilibrium state have been undertaken assuming a corotational velocity field, $^{7),8)}$ since there was no formalism to compute non-corotational solutions. As pointed out previously, $^{9),10)}$ however, corotation is not an adequate assumption for the velocity field of realistic binary neutron stars, because the effect of viscosity is negligible for the evolution of neutron stars in a binary system and, as a result, their velocity fields are expected to be irrotational (or nearly irrotational). Formalism for the computation of realistic quasiequilibrium states of coalescing binary neutron stars just prior to merging has recently been developed by several authors in general relativity. $^{11), 12), 13), 14), 15)$ In this formalism, we have only to solve two hydrostatic equations as for the fluid equations. One of them is the integrated form of the Euler equation and the other is the Poisson equation for the velocity potential. Thus, the formalism seems to be very tractable for computing equilibrium configurations of relativistic irrotational bodies.

In this paper, we apply the formalism for solving an incompressible, irrotational single star in the first post-Newtonian (1PN) approximation as a first step. Incompressible rotating stars were studied extensively at the 1PN order by Chandrasekhar about 30 years ago. He derived many equilibrium configurations of the Maclaurin, Jacobi, Dedekind ellipsoids, ¹⁶, ¹⁷, ¹⁸, ¹⁹) developing original methods (see also Ref. 20)), and he found many interesting features of rotating stars deformed by relativistic effects.

The purpose of this paper is twofold. One is to develop a formalism to obtain an equilibrium state of an irrotational, incompressible fluid in the 1PN approximation. As mentioned above, Chandrasekhar studied 1PN equilibrium stars extensively, but he did not do so for irrotational stars. We develop the method in this paper which can be applied even for irrotational binary systems. The other comes from a demand in performing numerical computation: Our final aim is to study general relativistic irrotational binary neutron stars with compressible equations of state. Thus, numerical computation is necessary. Although some promising numerical methods are proposed, ^{21), 11)} the computation, it is necessary to check its validity by comparing with an analytic solution. Thus, preparation of exact solutions is required.

This paper is organized as follows. In §2, we describe the basic equations to calculate equilibrium configurations of the 1PN irrotational star. The deformation of the star from an ellipsoid and the angular velocity in the 1PN approximation are calculated in §3. In §4, we give the boundary conditions to determine the velocity field and the deformation in the 1PN approximation, and the total energy and angular momentum are obtained in §5. We present formalism for computation of gravitational radiation from a rotating star with arbitrary internal motion in §6. In §7, we present the numerical results obtained by solving the equations derived in previous sections. We also compare the results with those in the ellipsoidal approximation.²² Section 8 is devoted to summary and discussion.

Throughout this paper, c denotes the light velocity and we use the units in which

G = 1. Latin indices i, j, k, \cdots take values 1 to 3, and δ_{ij} denotes the Kronecker delta.

§2. Formulation

Non-axisymmetric equilibrium configurations with non-uniform velocity fields are obtained by solving the Euler, continuity, and Poisson equations, consistently. Since we consider an incompressible fluid, all the calculations are carried out analytically even in the 1PN case. The procedure is as follows.

(1) We calculate a Newtonian equilibrium state, i.e., the irrotational Riemann ellipsoid, 20 as a non-perturbed state. For simplicity, we consider only the case in which the directions of the vorticity vector and the angular velocity vector lie along x_3 -axis.

(2) 1PN corrections for the velocity potential and gravitational potentials are obtained from the 1PN Poisson equations for them.

(3) We calculate the deformation from the Newtonian ellipsoid induced by 1PN gravity using Lagrangian displacement vectors introduced by Chandrasekhar.²⁰⁾ Then, corrections of the Newtonian quantities due to the deformation of the star are estimated.

(4) We substitute all the 1PN corrections obtained in (2) and (3) into the 1PN Euler and continuity equations. Then, coefficients of the Lagrangian displacement vectors and 1PN velocity potential are determined from the boundary conditions on the stellar surface.

In this section, we calculate the Newtonian and 1PN terms which we need in the above procedures. In the following, we assume that the center of mass of the star is located at the origin of the coordinate system, and the direction of the coordinate axes are parallel to the principal axes of the star, whose lengths are defined as a_1, a_2 and a_3 .

2.1. Hydrostatic and Poisson equations for 1PN irrotational stars

For an irrotational fluid, the relativistic Euler equation can be integrated, and in the 1PN case, it is written as $^{13)}$

$$\operatorname{const} = \frac{P}{\rho} - U + \frac{1}{2} \sum_{k} (\partial_{k} \phi_{\mathrm{N}})^{2} - \sum_{k} \ell^{k} \partial_{k} \phi_{\mathrm{N}}$$
$$+ \frac{1}{c^{2}} \left[-\frac{P}{\rho} U + \frac{1}{2} U^{2} + X - \frac{1}{2} \left(\frac{P}{\rho} + 3U \right) \sum_{k} (\partial_{k} \phi_{\mathrm{N}})^{2} - \frac{1}{8} \left(\sum_{k} (\partial_{k} \phi_{\mathrm{N}})^{2} \right)^{2} + \sum_{k} (\partial_{k} \phi_{\mathrm{N}}) (\partial_{k} \phi_{\mathrm{PN}}) - \sum_{k} \ell^{k} \partial_{k} \phi_{\mathrm{PN}} - \sum_{k} \hat{\beta}^{k} \partial_{k} \phi_{\mathrm{N}} \right], \qquad (2.1)$$

where we assume that fluid is incompressible, i.e., $\rho = \text{const.}$ In Eq. (2·1), P, ρ , ℓ^k , ϕ_N , ϕ_{PN} , U, X, and $\hat{\beta}^k$, respectively, denote the pressure, the density, the velocity field of the figure rotation (spatial component of the Killing vector), the Newtonian and 1PN velocity potentials, and the last three terms are the Newtonian and 1PN

potentials, which are derived by solving Poisson equations as

$$\Delta U = -4\pi\rho, \tag{2.2}$$

$$\Delta X = 4\pi \rho \left[2U + \frac{3P}{\rho} + 2\sum_{k} (\partial_k \phi_N)^2 \right], \qquad (2.3)$$

$$\Delta P_k = -4\pi\rho\partial_k\phi_{\rm N},\tag{2.4}$$

$$\Delta \chi = 4\pi \rho \sum_{k} (\partial_k \phi_{\rm N}) x^k.$$
(2.5)

Here $\hat{\beta}_k$ is expressed as

$$\hat{\beta}_k = -\frac{7}{2}P_k + \frac{1}{2} \Big(\partial_k \chi + \sum_l x^l \partial_k P_l \Big).$$
(2.6)

We note that using the gravitational potentials, the spacetime line element to 1PN order can be written as

$$ds^{2} = -\alpha^{2}c^{2}dt^{2} + \frac{2}{c^{2}}\sum_{i}\hat{\beta}_{i}dx^{i}dt + \left(1 + \frac{2U}{c^{2}}\right)\sum_{i}dx^{i}dx^{i}, \qquad (2.7)$$

where

$$\alpha = 1 - \frac{U}{c^2} + \frac{1}{c^4} \left(\frac{U^2}{2} + X \right) + O(c^{-6}).$$
(2.8)

A characteristic feature of the irrotational fluid is that the continuity equation reduces to a Poisson type equation for a velocity potential ϕ , ^{13), 14)} and in the 1PN incompressible case, it is

$$\sum_{i} \rho \partial_i C_i = 0, \qquad (2.9)$$

where

$$C_i = -\ell^i + \partial_i \phi_{\rm N} + \frac{1}{c^2} \left[-\ell^i \left(\frac{1}{2} \sum_k (\partial_k \phi_{\rm N})^2 + 3U \right) - \frac{P}{\rho} \partial_i \phi_{\rm N} - \hat{\beta}_i + \partial_i \phi_{\rm PN} \right].$$
(2·10)

In the following subsections, we obtain terms appearing in Eq. $(2 \cdot 1)$ by separately solving equations for each.

2.2. Newtonian terms

The solution of the Newtonian potential for the ellipsoidal star is

$$U = \pi \rho \Big(A_0 - \sum_k A_k x_k^2 \Big), \qquad (2.11)$$

where $A_{ij\dots}$ are index symbols introduced by Chandrasekhar, ²⁰⁾ and A_0 is calculated from ²⁰⁾

$$A_{0} = a_{1}a_{2}a_{3} \int_{0}^{\infty} \frac{du}{\sqrt{(a_{1}^{2} + u)(a_{2}^{2} + u)(a_{3}^{2} + u)}}$$
$$= a_{1}^{2}\alpha_{2}\alpha_{3} \int_{0}^{\infty} \frac{dt}{\sqrt{(1 + t)(\alpha_{2}^{2} + t)(\alpha_{3}^{2} + t)}},$$
(2.12)

where $\alpha_2 = a_2/a_1$ and $\alpha_3 = a_3/a_1$. Also, the pressure at Newtonian order is written as

$$P = P_0 \left(1 - \sum_k \frac{x_k^2}{a_k^2} \right), \tag{2.13}$$

where $P_0 = \pi \rho^2 a_3^2 A_3$ denotes the pressure at the center of the star.

The Newtonian velocity field in the inertial frame $v_i \equiv \partial_i \phi_N$ is written as

$$v_{1} = \ell^{1} + u_{1} = -\left(\frac{a_{1}^{2}f_{R}}{a_{1}^{2} + a_{2}^{2}} + 1\right)\Omega x_{2},$$

$$v_{2} = \ell^{2} + u_{2} = \left(\frac{a_{2}^{2}f_{R}}{a_{1}^{2} + a_{2}^{2}} + 1\right)\Omega x_{1},$$

$$v_{3} = 0,$$
(2.14)

where u_i is the velocity field in the corotating frame and Ω denotes the angular velocity of the figure. ℓ^i and u_i are given by

$$\ell^{i} = (-\Omega x_{2}, \ \Omega x_{1}, \ 0), \qquad (2.15)$$
$$u_{i} = \left(\frac{a_{1}}{a_{2}}\Lambda x_{2}, \ -\frac{a_{2}}{a_{1}}\Lambda x_{1}, \ 0\right)$$
$$= \left(-\frac{a_{1}^{2}f_{R}}{a_{1}^{2} + a_{2}^{2}}\Omega x_{2}, \ \frac{a_{2}^{2}f_{R}}{a_{1}^{2} + a_{2}^{2}}\Omega x_{1}, \ 0\right), \qquad (2.16)$$

where
$$\Lambda$$
 is the angular velocity of the internal motion. f_R is defined as ζ/Ω , where $\zeta \equiv (\text{rot}\mathbf{u})_3$ denotes the vorticity in the corotating frame. For the irrotational ellipsoid, f_R becomes -2 . Thus, $\phi_N = F_a \Omega x_1 x_2$ and

$$v_i = (F_a \Omega x_2, F_a \Omega x_1, 0), \qquad (2.17)$$

where

ζ

$$F_a \equiv \frac{a_1^2 - a_2^2}{a_1^2 + a_2^2}.$$
 (2.18)

2.3. 1PN terms

2.3.1. X

For the convenience of the calculation, we separate X into two parts as

$$X = X_0 + X_v, \tag{2.19}$$

where X_0 and X_v are derived from Poisson-like equations as

$$\Delta X_0 = 4\pi \rho \left(2U + \frac{3P}{\rho} \right), \tag{2.20}$$

$$\Delta X_v = 8\pi \rho \sum_k (\partial_k \phi_N)^2.$$
(2.21)

Then, the equation for X_0 becomes

$$\Delta X_0 = 4\pi \rho \left[\left(2\pi \rho A_0 + \frac{3P_0}{\rho} \right) - \sum_k \left(2\pi \rho A_k + \frac{3P_0}{\rho a_k^2} \right) x_k^2 \right],$$
(2.22)

and the solution is

$$X_0 = -\alpha_0 U + \sum_k \eta_k D_{kk}, \qquad (2.23)$$

where

$$\alpha_0 = 2\pi\rho A_0 + \frac{3P_0}{\rho},\tag{2.24}$$

$$\eta_k = 2\pi\rho A_k + \frac{3P_0}{\rho a_k^2},$$
(2.25)

$$D_{kk} = \pi \rho \bigg[a_k^4 \Big(A_{kk} - \sum_m A_{kkm} x_m^2 \Big) x_k^2 + \frac{1}{4} a_k^2 \Big(B_k - 2 \sum_m B_{km} x_m^2 + \sum_m \sum_n B_{kmn} x_m^2 x_n^2 \Big) \bigg], \quad (2.26)$$

and $B_{ij\cdots}$ are index symbols introduced by Chandrasekhar.²⁰⁾ $D_{ij\cdots}$ is calculated from a Poisson equation as ²⁰⁾

$$\Delta D_{ij\dots} = -4\pi\rho x_i x_j \cdots . \tag{2.27}$$

The equation for X_v is

4

$$\Delta X_v = 8\pi\rho F_a^2 \Omega^2 (x_1^2 + x_2^2), \qquad (2.28)$$

and hence, the solution is

$$X_v = -2F_a^2 \Omega^2 (D_{11} + D_{22}). (2.29)$$

2.3.2. $\hat{\beta}_k$

Substituting v_i of Newtonian order into the equations for P_i and χ , we immediately find the solutions as

$$P_1 = F_a \Omega D_2, \tag{2.30}$$

$$P_2 = F_a \Omega D_1, \tag{2.31}$$

$$P_3 = 0, \tag{2.32}$$

$$\chi = -2F_a\Omega D_{12},\tag{2.33}$$

where $^{20)}$

$$D_i = \pi \rho a_i^2 \left(A_i - \sum_k A_{ik} x_k^2 \right) x_i, \qquad (2.34)$$

$$D_{12} = \pi \rho a_1^2 a_2^2 \Big(A_{12} - \sum_k A_{12k} x_k^2 \Big) x_1 x_2.$$
(2.35)

Using the above solutions of P_k and χ , we obtain $\hat{\beta}_k$:

$$\hat{\beta}_{1} = \pi \rho F_{a} \Omega \left[\left(\frac{1}{2} a_{1}^{2} A_{1} - \frac{7}{2} a_{2}^{2} A_{2} - a_{1}^{2} a_{2}^{2} A_{12} \right) x_{2} + \left(-\frac{3}{2} a_{1}^{2} A_{11} + \frac{5}{2} a_{2}^{2} A_{12} + 3 a_{1}^{2} a_{2}^{2} A_{112} \right) x_{1}^{2} x_{2} + \left(-\frac{1}{2} a_{1}^{2} A_{12} + \frac{7}{2} a_{2}^{2} A_{22} + a_{1}^{2} a_{2}^{2} A_{122} \right) x_{1}^{3} + \left(-\frac{1}{2} a_{1}^{2} A_{12} + \frac{7}{2} a_{2}^{2} A_{23} + a_{1}^{2} a_{2}^{2} A_{123} \right) x_{2} x_{3}^{2} \right], \qquad (2.36)$$

$$\hat{\beta}_{2} = \pi \rho F_{a} \Omega \left[\left(\frac{1}{2} a_{2}^{2} A_{2} - \frac{7}{2} a_{1}^{2} A_{1} - a_{1}^{2} a_{2}^{2} A_{12} \right) x_{1} + \left(-\frac{1}{2} a_{2}^{2} A_{12} + \frac{7}{2} a_{1}^{2} A_{11} + a_{1}^{2} a_{2}^{2} A_{112} \right) x_{1}^{3} + \left(-\frac{3}{2} a_{2}^{2} A_{22} + \frac{5}{2} a_{1}^{2} A_{12} + 3 a_{1}^{2} a_{2}^{2} A_{122} \right) x_{1} x_{2}^{2} + \left(-\frac{1}{2} a_{2}^{2} A_{23} + \frac{7}{2} a_{1}^{2} A_{13} + a_{1}^{2} a_{2}^{2} A_{123} \right) x_{1} x_{3}^{2} \right], \qquad (2.37)$$

$$\hat{\beta}_{*} = \pi \rho F_{a} \Omega \left[\left(2 a_{2}^{2} a_{2}^{2} A_{12} - a_{2}^{2} A_{23} - a_{2}^{2} A_{23} - a_{2}^{2} A_{23} \right) x_{2} x_{3}^{2} \right], \qquad (2.37)$$

$$\hat{\beta}_3 = \pi \rho F_a \Omega \Big(2a_1^2 a_2^2 A_{123} - a_2^2 A_{23} - a_1^2 A_{13} \Big) x_1 x_2 x_3.$$
(2.38)

Then, the divergence of $\hat{\beta}_k$ is

$$\sum_{k} \partial_k \hat{\beta}_k = 6\pi \rho F_a \Omega(a_1^2 + a_2^2) A_{12} x_1 x_2, \qquad (2.39)$$

where we have used some relations among index symbols. $^{20)}\,$

2.3.3. $\phi_{\rm PN}$

An explicit form of the Poisson-type equation for $\phi_{\rm PN}$ can be written as

$$\Delta\phi_{\rm PN} = \left[6\pi\rho \Omega (A_1 - A_2) - 2F_a \Omega \left(\frac{1}{a_1^2} + \frac{1}{a_2^2} \right) \frac{P_0}{\rho} \right] x_1 x_2 + \sum_i \partial_i \hat{\beta}_i$$
$$= -2\Omega \frac{(a_1^2 - a_2^2)}{a_1^2 a_2^2} \frac{P_0}{\rho} x_1 x_2. \tag{2.40}$$

The solution of this equation can be written up to biquadratic terms in x_i as^{*)}

$$\phi_{\rm PN} = (p + qx_1^2 + rx_2^2 + sx_3^2)x_1x_2 + \text{const},$$
 (2.41)

where q, r and s satisfy the condition

$$3q + 3r + s = -\Omega \frac{(a_1^2 - a_2^2)}{a_1^2 a_2^2} \frac{P_0}{\rho}.$$
 (2.42)

^{*)} Although we can add higher order terms which satisfy $\Delta \phi_{\rm PN} = 0$, we negelet them for simplicity because we consider only biquadratic deformation of ellipsoids in this paper.

2.4. Collection

Substituting the terms derived above into Eq. (2.1), we obtain

$$\mathcal{G} \equiv \frac{P}{\rho} - U - \delta U + \frac{1}{2} F_a^2 \left(\Omega_R^2 + \frac{1}{c^2} \delta \Omega^2 \right) (x_1^2 + x_2^2) + F_a \left(\Omega_R^2 + \frac{1}{c^2} \delta \Omega^2 \right) (x_2^2 - x_1^2) + \frac{1}{c^2} \left(\gamma_0 + \sum_l \gamma_l x_l^2 + \sum_{l \le m} \gamma_{lm} x_l^2 x_m^2 \right) = \text{const},$$
(2.43)

where \varOmega_R denotes the angular velocity of an irrotational Riemann ellipsoid and the γ_{ij} are expressed as

$$\begin{split} \gamma_{0} &= \pi \rho \left[-A_{0} \left(\frac{4P_{0}}{\rho} + \frac{3}{2} \pi \rho A_{0} \right) + \frac{1}{4} \sum_{l} \left(2\pi \rho a_{l}^{2} A_{l} + \frac{3P_{0}}{\rho} \right) B_{l} \\ &\quad -\frac{1}{2} F_{a}^{2} \Omega_{R}^{2} (a_{1}^{2} B_{1} + a_{2}^{2} B_{2}) \right], \end{split} \tag{2.44} \\ \gamma_{1} &= \pi \rho \left[4A_{1} \frac{P_{0}}{\rho} + \pi \rho A_{0} A_{1} + A_{0} \frac{P_{0}}{\rho a_{1}^{2}} \\ &\quad + \left(2\pi \rho a_{1}^{2} A_{1} + \frac{3P_{0}}{\rho} \right) \left(a_{1}^{2} A_{11} - \frac{1}{2} B_{11} \right) - \frac{1}{2} \left(2\pi \rho a_{2}^{2} A_{2} + \frac{3P_{0}}{\rho} \right) B_{12} \\ &\quad -\frac{1}{2} \left(2\pi \rho a_{3}^{2} A_{3} + \frac{3P_{0}}{\rho} \right) B_{13} \\ &\quad + F_{a}^{2} \Omega_{R}^{2} (3a_{1}^{2} A_{1} - a_{2}^{2} A_{2} - 2a_{3}^{2} A_{3} - 3a_{1}^{4} A_{11}) - \frac{2a_{2}^{2}}{a_{1}^{2} + a_{2}^{2}} \frac{\Omega_{R}}{\pi \rho} \rho \right], \qquad (2.45) \\ \gamma_{2} &= \pi \rho \left[4A_{2} \frac{P_{0}}{\rho} + \pi \rho A_{0} A_{2} + A_{0} \frac{P_{0}}{\rho a_{2}^{2}} + \left(2\pi \rho a_{2}^{2} A_{2} + \frac{3P_{0}}{\rho} \right) \left(a_{2}^{2} A_{22} - \frac{1}{2} B_{22} \right) \\ &\quad -\frac{1}{2} \left(2\pi \rho a_{1}^{2} A_{1} + \frac{3P_{0}}{\rho} \right) B_{12} - \frac{1}{2} \left(2\pi \rho a_{3}^{2} A_{3} + \frac{3P_{0}}{\rho} \right) B_{23} \\ &\quad + F_{a}^{2} \Omega_{R}^{2} (3a_{2}^{2} A_{2} - a_{1}^{2} A_{1} - 2a_{3}^{2} A_{3} - 3a_{2}^{4} A_{22}) + \frac{2a_{1}^{2}}{a_{1}^{2} + a_{2}^{2}} \frac{\Omega_{R}}{\pi \rho} \rho \right], \qquad (2.46) \\ \gamma_{3} &= \pi \rho \left[4A_{3} \frac{P_{0}}{\rho} + \pi \rho A_{0} A_{3} + A_{0} \frac{P_{0}}{\rho a_{3}^{2}} + \left(2\pi \rho a_{3}^{2} A_{3} + \frac{3P_{0}}{\rho} \right) \left(a_{3}^{2} A_{33} - \frac{1}{2} B_{33} \right) \\ &\quad -\frac{1}{2} \left(2\pi \rho a_{1}^{2} A_{1} + \frac{3P_{0}}{\rho} \right) B_{13} - \frac{1}{2} \left(2\pi \rho a_{2}^{2} A_{2} + \frac{3P_{0}}{\rho} \right) B_{23} \\ &\quad + F_{a}^{2} \Omega_{R}^{2} (a_{1}^{2} B_{13} + a_{2}^{2} B_{23}) \right], \qquad (2.47) \\ \gamma_{11} &= \pi \rho \left[-A_{1} \frac{P_{0}}{\rho a_{1}^{2}} + \frac{1}{2} \pi \rho A_{1}^{2} + \left(2\pi \rho a_{1}^{2} A_{1} + \frac{3P_{0}}{\rho} \right) \left(-a_{1}^{2} A_{111} + \frac{1}{4} B_{111} \right) \\ \end{cases} \end{cases}$$

$$\begin{split} \gamma_{12} &= \pi \rho \left[-\frac{P_0}{\rho} \left(\frac{A_1}{a_2^2} + \frac{A_2}{a_1^2} \right) + \pi \rho A_1 A_2 + \left(2\pi \rho a_1^2 A_1 + \frac{3P_0}{\rho} \right) \left(-a_1^2 A_{112} + \frac{1}{2} B_{112} \right) \right. \\ &+ \left(2\pi \rho a_2^2 A_2 + \frac{3P_0}{\rho} \right) \left(-a_2^2 A_{122} + \frac{1}{2} B_{122} \right) + \frac{1}{2} \left(2\pi \rho a_3^2 A_3 + \frac{3P_0}{\rho} \right) B_{123} \\ &+ \frac{F_a^2 \Omega_R^2}{2} \left(3A_1 + 3A_2 + \frac{a_3^2 (a_1^2 + a_2^2)}{a_1^2 a_2^2} A_3 \right) \\ &- 4(a_1^2 + a_2^2) A_{12} + 3a_1^2 (a_1^2 - a_2^2) A_{112} - 3a_2^2 (a_1^2 - a_2^2) A_{122} \right) \\ &- \frac{F_a^4 \Omega_R^4}{4\pi \rho} + \frac{6a_1^2}{a_1^2 + a_2^2} \frac{\Omega_R}{\pi \rho} q - \frac{6a_2^2}{a_1^2 + a_2^2} \frac{\Omega_R}{\pi \rho} r \right], \end{split}$$

$$\gamma_{13} = \pi \rho \left[-\frac{P_0}{\rho} \left(\frac{A_1}{a_3^2} + \frac{A_3}{a_1^2} \right) + \pi \rho A_1 A_3 + \left(2\pi \rho a_1^2 A_1 + \frac{3P_0}{\rho} \right) \left(-a_1^2 A_{113} + \frac{1}{2} B_{113} \right) \right. \\ \left. + \frac{1}{2} \left(2\pi \rho a_2^2 A_2 + \frac{3P_0}{\rho} \right) B_{123} + \left(2\pi \rho a_3^2 A_3 + \frac{3P_0}{\rho} \right) \left(-a_3^2 A_{133} + \frac{1}{2} B_{133} \right) \right. \\ \left. + \frac{F_a^2 \Omega_R^2}{2} (4A_3 - a_2^2 A_{23} - 9a_1^2 A_{13} + 6a_1^4 A_{113}) - \frac{2a_2^2}{a_1^2 + a_2^2} \frac{\Omega_R}{\pi \rho} s \right], \quad (2.50)$$

$$\gamma_{22} = \pi \rho \left[-A_2 \frac{P_0}{\rho a_2^2} + \frac{1}{2} \pi \rho A_2^2 + \left(2\pi \rho a_2^2 A_2 + \frac{3P_0}{\rho} \right) \left(-a_2^2 A_{222} + \frac{1}{4} B_{222} \right) \right. \\ \left. + \frac{1}{4} \left(2\pi \rho a_1^2 A_1 + \frac{3P_0}{\rho} \right) B_{122} + \frac{1}{4} \left(2\pi \rho a_3^2 A_3 + \frac{3P_0}{\rho} \right) B_{223} \right. \\ \left. + \frac{F_a^2 \Omega_R^2}{2a_2^2} (3a_2^2 A_2 + a_3^2 A_3 - 8a_2^4 A_{22} - a_1^2 a_2^4 A_{122} + 5a_2^6 A_{222}) \right. \\ \left. - \frac{F_a^4 \Omega_R^4}{8\pi\rho} + \frac{2a_1^2}{a_1^2 + a_2^2} \frac{\Omega_R}{\pi\rho} r \right],$$

$$(2.51)$$

$$\gamma_{23} = \pi \rho \left[-\frac{P_0}{\rho} \left(\frac{A_2}{a_3^2} + \frac{A_3}{a_2^2} \right) + \pi \rho A_2 A_3 + \left(2\pi \rho a_2^2 A_2 + \frac{3P_0}{\rho} \right) \left(-a_2^2 A_{223} + \frac{1}{2} B_{223} \right) + \frac{1}{2} \left(2\pi \rho a_1^2 A_1 + \frac{3P_0}{\rho} \right) B_{123} + \left(2\pi \rho a_3^2 A_3 + \frac{3P_0}{\rho} \right) \left(-a_3^2 A_{233} + \frac{1}{2} B_{233} \right) + \frac{F_a^2 \Omega_R^2}{2} (4A_3 - a_1^2 A_{13} - 9a_2^2 A_{23} + 6a_2^4 A_{223}) + \frac{2a_1^2}{a_1^2 + a_2^2} \frac{\Omega_R}{\pi \rho} s \right], \quad (2.52)$$

$$\gamma_{33} = \pi \rho \left[-A_3 \frac{P_0}{\rho a_3^2} + \frac{1}{2} \pi \rho A_3^2 + \left(2 \pi \rho a_3^2 A_3 + \frac{3P_0}{\rho} \right) \left(-a_3^2 A_{333} + \frac{1}{4} B_{333} \right) \right. \\ \left. + \frac{1}{4} \left(2 \pi \rho a_1^2 A_1 + \frac{3P_0}{\rho} \right) B_{133} + \frac{1}{4} \left(2 \pi \rho a_2^2 A_2 + \frac{3P_0}{\rho} \right) B_{233} \right. \\ \left. - \frac{F_a^2 \Omega_R^2}{2} (a_1^2 B_{133} + a_2^2 B_{233}) \right].$$

$$(2.53)$$

 δU denotes the gravitational potential induced by the deformation of the figure. The explicit form of δU is given in the §4.

§3. Second tensor virial equation

The angular velocity is determined from the second tensor virial equation, $^{18)}$ which is written as

$$\int d^3x \rho x_j \frac{\partial}{\partial x_i} \mathcal{G} = 0. \tag{3.1}$$

Substituting Eq. (2.43) into the above equation, we obtain

$$0 = -\delta_{ij}(\Pi + \delta\Pi) - W_{ij} - \delta W_{ij} + F_a(F_a - 2)(\Omega_R^2 + \delta\Omega^2)\delta_{1i}(I_{1j} + \delta I_{1j}) + F_a(F_a + 2)(\Omega_R^2 + \delta\Omega^2)\delta_{2i}(I_{2j} + \delta I_{2j}) + \frac{1}{c^2} \Big[2\gamma_i I_{ij} + 4\gamma_{11}I_{111j}\delta_{1i} + 2\gamma_{12}(I_{122j}\delta_{1i} + I_{112j}\delta_{2i}) + 2\gamma_{13}(I_{133j}\delta_{i1} + I_{113j}\delta_{3i}) + 4\gamma_{22}I_{222j}\delta_{2i} + 2\gamma_{23}(I_{233j}\delta_{2i} + I_{223j}\delta_{3i}) + 4\gamma_{33}I_{333j}\delta_{3i} \Big],$$
(3.2)

where

$$\Pi = \int d^3 x P, \tag{3.3}$$

$$W_{ij} = \int d^3x \rho x_i \frac{\partial U}{\partial x_j} = -2\pi\rho A_i I_{ij}, \qquad (3.4)$$

$$I_{ij} = \int d^3x \rho x_i x_j = \frac{M}{5} a_i^2 \delta_{ij}, \qquad (3.5)$$

$$I_{iijj} = \int d^3x \rho x_i^2 x_j^2 = \frac{M}{35} a_i^2 a_j^2 (1 + 2\delta_{ij}), \qquad (3.6)$$

$$\delta \Pi = \delta \int d^3 x P, \tag{3.7}$$

$$\delta W_{ij} = \delta \int d^3 x \rho x_i \frac{\partial U}{\partial x_j},\tag{3.8}$$

$$\delta I_{ij} = \delta \int d^3x \rho x_i x_j. \tag{3.9}$$

 ${\cal M}$ denotes the Newtonian mass:

$$M \equiv \int d^3x \rho. \tag{3.10}$$

10

 $\delta \Pi$, δW_{ij} , and δI_{ij} must be taken into account because the ellipsoidal figure deforms due to the 1PN effect. The explicit forms are given later.

3.1. Second virial equations at Newtonian order

First, we derive equations for the angular velocity and axial ratios of the ellipsoid at Newtonian order. The second tensor virial equation at Newtonian order is written

$$0 = -\delta_{ij}\Pi - W_{ij} + F_a(F_a - 2)\Omega_R^2 \delta_{1i} I_{1j} + F_a(F_a + 2)\Omega_R^2 \delta_{2i} I_{2j}.$$
 (3.11)

The non-vanishing components are

$$\Pi = -W_{11} + F_a(F_a - 2)\Omega_R^2 I_{11}, \qquad (3.12)$$

$$\Pi = -W_{22} + F_a(F_a + 2)\Omega_R^2 I_{22}, \qquad (3.13)$$

$$\Pi = -W_{33}.$$
 (3.14)

Then, we obtain

$$\Omega_R^2 \left[1 + \left(\frac{2a_1 a_2}{a_1^2 + a_2^2} \right)^2 \right] = 2\pi\rho B_{12}, \qquad (3.15)$$

$$-\frac{2a_1^2a_2^2}{a_1^2+a_2^2}\Omega_R^2 = \pi\rho(a_1^2a_2^2A_{12}-a_3^2A_3).$$
(3.16)

From these equations, we can determine the axial ratios a_2/a_1 and a_3/a_1 , and the angular velocity.

3.2. Deformation of the figure

The deformation of an incompressible ellipsoid due to the 1PN gravity is written by using the Lagrangian displacement vectors $\xi_k^{(ij)}$ of $\sum_k \partial_k \xi_k^{(ij)} = 0^{18}$ as

$$\xi_k = \frac{1}{c^2} \sum_{ij} S_{ij} \xi_k^{(ij)}, \qquad (3.17)$$

where

$$\xi_k^{(11)} = (x_1, \ 0, \ -x_3), \tag{3.18}$$

$$\xi_k^{(12)} = (0, \ x_2, \ -x_3), \tag{3.19}$$

$$\xi_k^{(31)} = \left(\frac{1}{3}x_1^3, \ -x_1^2x_2, \ 0\right),\tag{3.20}$$

$$\xi_k^{(32)} = \left(0, \ \frac{1}{3}x_2^3, \ -x_2^2x_3\right),\tag{3.21}$$

$$\xi_k^{(33)} = \left(-x_3^2 x_1, \ 0, \ \frac{1}{3} x_3^3\right). \tag{3.22}$$

Here, we consider only up to the cubic deformation and also assume triplane symmetries, so that the Lagrangian displacement vectors of higher order functions in x_i and of even functions can be neglected.

Using the Lagrangian displacement vectors, we can calculate Eqs. (3.7) \sim (3.9). We find $^{20)}$

$$\delta \Pi = 0, \tag{3.23}$$

$$\delta I_{ij} = \int d^3x \rho \sum_l \xi_l \frac{\partial}{\partial x_l} (x_i x_j), \qquad (3.24)$$

$$\delta I_{11} = \frac{2}{c^2} \left(S_{11} I_{11} + \frac{1}{3} S_{31} I_{1111} - S_{33} I_{1133} \right), \tag{3.25}$$

$$\delta I_{22} = \frac{2}{c^2} \left(S_{12} I_{22} - S_{31} I_{1122} + \frac{1}{3} S_{32} I_{2222} \right), \tag{3.26}$$

$$\delta I_{33} = \frac{2}{c^2} \left(-S_{11}I_{33} - S_{12}I_{33} - S_{32}I_{2233} + \frac{1}{3}S_{33}I_{3333} \right), \tag{3.27}$$

$$\delta W_{ij} = \pi \rho \Big(-2B_{ij} \delta I_{ij} + a_i^2 \delta_{ij} \sum_l A_{il} \delta I_{ll} \Big), \tag{3.28}$$

$$\delta W_{11} = \frac{2\pi\rho}{c^2} \left[S_{11} \left\{ (a_1^2 A_{11} - 2B_{11})I_{11} - a_1^2 A_{13}I_{33} \right\} + S_{12}(a_1^2 A_{12}I_{22} - a_1^2 A_{13}I_{33}) + S_{31} \left\{ \frac{1}{3}(a_1^2 A_{11} - 2B_{11})I_{1111} - a_1^2 A_{12}I_{1122} \right\} + S_{32} \left(\frac{1}{3}a_1^2 A_{12}I_{2222} - a_1^2 A_{13}I_{2233} \right) + S_{33} \left\{ -(a_1^2 A_{11} - 2B_{11})I_{1133} + \frac{1}{3}a_1^2 A_{13}I_{3333} \right\} \right], \quad (3.29)$$

$$\begin{split} \delta W_{22} &= \frac{2\pi\rho}{c^2} \left[S_{11} (a_2^2 A_{12} I_{11} - a_2^2 A_{23} I_{33}) \\ &+ S_{12} \left\{ (a_2^2 A_{22} - 2B_{22}) I_{22} - a_2^2 A_{23} I_{33} \right\} \\ &+ S_{31} \left\{ \frac{1}{3} a_2^2 A_{12} I_{1111} - (a_2^2 A_{22} - 2B_{22}) I_{1122} \right\} \\ &+ S_{32} \left\{ \frac{1}{3} (a_2^2 A_{22} - 2B_{22}) I_{2222} - a_2^2 A_{23} I_{2233} \right\} \\ &+ S_{33} \left(-a_2^2 A_{12} I_{1133} + \frac{1}{3} a_2^2 A_{23} I_{3333} \right) \right], \end{split}$$
(3.30)
$$\delta W_{33} &= \frac{2\pi\rho}{c^2} \left[S_{11} \left\{ a_3^2 A_{13} I_{11} - (a_3^2 A_{33} - 2B_{33}) I_{33} \right\} \\ &+ S_{12} \left\{ a_3^2 A_{23} I_{22} - (a_3^2 A_{33} - 2B_{33}) I_{33} \right\} \\ &+ S_{31} \left(\frac{1}{3} a_3^2 A_{13} I_{111} - a_3^2 A_{23} I_{1122} \right) \\ &+ S_{32} \left\{ \frac{1}{3} a_3^2 A_{23} I_{2222} - (a_3^2 A_{33} - 2B_{33}) I_{2233} \right\} \end{split}$$

$$+S_{33}\bigg\{-a_3^2A_{13}I_{1133}+\frac{1}{3}(a_3^2A_{33}-2B_{33})I_{3333}\bigg\}\bigg]. (3.31)$$

3.3. Second virial equations in the 1PN approximation

Substituting the 1PN terms into Eq. (3.2), we have the second virial equations in the 1PN approximation. Their components are given as follows:

1. i = j = 1:

$$0 = 2\gamma_{1}a_{1}^{2} + \frac{12}{7}\gamma_{11}a_{1}^{4} + \frac{2}{7}\gamma_{12}a_{1}^{2}a_{2}^{2} + \frac{2}{7}\gamma_{13}a_{1}^{2}a_{3}^{2} + F_{a}(F_{a}-2)a_{1}^{2}\delta\Omega^{2} + 2F_{a}(F_{a}-2)\Omega_{R}^{2}\left(S_{11}a_{1}^{2} + \frac{1}{7}S_{31}a_{1}^{4} - \frac{1}{7}S_{33}a_{1}^{2}a_{3}^{2}\right) - 2\pi\rho\left[S_{11}(a_{1}^{4}A_{11}-2a_{1}^{2}B_{11}-a_{1}^{2}a_{3}^{2}A_{13}) + S_{12}(a_{1}^{2}a_{2}^{2}A_{12}-a_{1}^{2}a_{3}^{2}A_{13}) + \frac{S_{31}}{7}(a_{1}^{6}A_{11}-2a_{1}^{4}B_{11}-a_{1}^{4}a_{2}^{2}A_{12}) + \frac{S_{32}}{7}(a_{1}^{2}a_{2}^{4}A_{12}-a_{1}^{2}a_{2}^{2}a_{3}^{2}A_{13}) + \frac{S_{33}}{7}(-a_{1}^{4}a_{3}^{2}A_{11}+2a_{1}^{2}a_{3}^{2}B_{11}+a_{1}^{2}a_{3}^{4}A_{13})\right],$$
(3.32)

2. i = j = 2:

$$0 = 2\gamma_{2}a_{2}^{2} + \frac{2}{7}\gamma_{12}a_{1}^{2}a_{2}^{2} + \frac{12}{7}\gamma_{22}a_{2}^{4} + \frac{2}{7}\gamma_{23}a_{2}^{2}a_{3}^{2} + F_{a}(F_{a}+2)a_{2}^{2}\delta\Omega^{2} + 2F_{a}(F_{a}+2)\Omega_{R}^{2}\left(S_{12}a_{2}^{2} - \frac{1}{7}S_{31}a_{1}^{2}a_{2}^{2} + \frac{1}{7}S_{32}a_{2}^{4}\right) - 2\pi\rho\left[S_{11}(a_{1}^{2}a_{2}^{2}A_{12} - a_{2}^{2}a_{3}^{2}A_{23}) + S_{12}(a_{2}^{4}A_{22} - 2a_{2}^{2}B_{22} - a_{2}^{2}a_{3}^{2}A_{23}) + \frac{S_{32}}{7}(a_{2}^{6}A_{22} - 2a_{2}^{4}B_{22} - a_{2}^{4}a_{3}^{2}A_{23}) + \frac{S_{33}}{7}(-a_{1}^{2}a_{2}^{2}a_{3}^{2}A_{12} + a_{2}^{2}a_{3}^{4}A_{23}) + \frac{S_{31}}{7}(a_{1}^{4}a_{2}^{2}A_{12} - a_{1}^{2}a_{2}^{4}A_{22} + 2a_{1}^{2}a_{2}^{2}B_{22})\right],$$
(3.33)

3. i = j = 3:

$$0 = 2\gamma_{3}a_{3}^{2} + \frac{2}{7}\gamma_{13}a_{1}^{2}a_{3}^{2} + \frac{2}{7}\gamma_{23}a_{2}^{2}a_{3}^{2} + \frac{12}{7}\gamma_{33}a_{3}^{4}$$

$$-2\pi\rho \Big[S_{11}(a_{1}^{2}a_{3}^{2}A_{13} - a_{3}^{4}A_{33} + 2a_{3}^{2}B_{33}) + S_{12}(a_{2}^{2}a_{3}^{2}A_{23} - a_{3}^{4}A_{33} + 2a_{3}^{2}B_{33})$$

$$+ \frac{S_{33}}{7}(a_{3}^{6}A_{33} - 2a_{3}^{4}B_{33} - a_{1}^{2}a_{3}^{4}A_{13}) + \frac{S_{31}}{7}(a_{1}^{4}a_{3}^{2}A_{13} - a_{1}^{2}a_{2}^{2}a_{3}^{2}A_{23})$$

$$+ \frac{S_{32}}{7}(a_{2}^{4}a_{3}^{2}A_{23} - a_{2}^{2}a_{3}^{4}A_{33} + 2a_{2}^{2}a_{3}^{2}B_{33}) \Big].$$
(3.34)

Subtracting the (2,2) component of the 1PN second virial equation from the (1,1) component, we have the equation for $\delta \Omega^2$ as

$$0 = 2\gamma_1 a_1^2 - 2\gamma_2 a_2^2 + \frac{2}{7} (6\gamma_{11}a_1^4 + \gamma_{13}a_1^2a_3^2 - 6\gamma_{22}a_2^4 - \gamma_{23}a_2^2a_3^2)$$

$$-(a_1^2 - a_2^2) \left[1 + \left(\frac{2a_1 a_2}{a_1^2 + a_2^2} \right)^2 \right] \delta \Omega^2 -2\pi \rho \left[S_{11} \tau_1 + S_{12} \tau_2 + \frac{a_1^2}{7} S_{31} (\tau_1 - \tau_2) + \frac{a_2^2}{7} S_{32} \tau_2 - \frac{a_3^2}{7} S_{33} \tau_1 \right], \quad (3.35)$$

where we use the equilibrium equations of Newtonian order and define new variables as

$$\tau_1 \equiv 3a_1^4 A_{11} - a_1^2 a_2^2 A_{12} - a_1^2 a_3^2 A_{13} + a_2^2 a_3^2 A_{23} - 2a_3^2 A_3, \qquad (3.36)$$

$$\tau_2 \equiv a_1^2 a_2^2 A_{12} - 3a_2^4 A_{22} - a_1^2 a_3^2 A_{13} + a_2^2 a_3^2 A_{23} + 2a_3^2 A_3.$$
(3.37)

Then, $\delta \Omega^2$ is expressed as

$$\delta\Omega^{2} = \frac{\Omega_{R}^{2}}{\pi\rho(a_{1}^{2}-a_{2}^{2})B_{12}} \bigg[\gamma_{1}a_{1}^{2} - \gamma_{2}a_{2}^{2} + \frac{1}{7}(6\gamma_{11}a_{1}^{4} + \gamma_{13}a_{1}^{2}a_{3}^{2} - 6\gamma_{22}a_{2}^{4} - \gamma_{23}a_{2}^{2}a_{3}^{2}) \\ -\pi\rho\bigg(S_{11}\tau_{1} + S_{12}\tau_{2} + \frac{a_{1}^{2}}{7}S_{31}(\tau_{1}-\tau_{2}) + \frac{a_{2}^{2}}{7}S_{32}\tau_{2} - \frac{a_{3}^{2}}{7}S_{33}\tau_{1}\bigg) \bigg]. \quad (3.38)$$

By calculating (1,1) - (3,3) and (2,2) - (3,3), we can find other expressions of $\delta \Omega^2$ as

$$0 = 2\gamma_1 a_1^2 - 2\gamma_3 a_3^2 + \frac{2}{7} (6\gamma_{11} a_1^4 + \gamma_{12} a_1^2 a_2^2 - \gamma_{23} a_2^2 a_3^2 - 6\gamma_{33} a_3^4) + F_a (F_a - 2) a_1^2 \delta \Omega^2 -2\pi \rho \Big[S_{11} (\tau_1 - \tau_3) - S_{12} \tau_3 + \frac{a_1^2}{7} S_{31} \tau_1 - \frac{a_2^2}{7} S_{32} \tau_3 - \frac{a_3^2}{7} S_{33} (\tau_1 - \tau_3) \Big], \quad (3.39)$$
$$0 = 2\gamma_1 a_1^2 - 2\gamma_2 a_2^2 + \frac{2}{7} (\gamma_1 - \gamma_2^2 a_2^2 + 6\gamma_2 a_3^4 - \gamma_2 a_2^2 a_2^2 - 6\gamma_3 a_3^4) + F_a (F_a - 2) a_1^2 \delta \Omega^2$$

$$0 = 2\gamma_2 a_2^2 - 2\gamma_3 a_3^2 + \frac{2}{7} (\gamma_{12} a_1^2 a_2^2 + 6\gamma_{22} a_2^4 - \gamma_{13} a_1^2 a_3^2 - 6\gamma_{33} a_3^4) + F_a (F_a + 2) a_2^2 \delta \Omega^2 - 2\pi \rho \Big[-S_{11} \tau_3 - S_{12} (\tau_2 + \tau_3) + \frac{a_1^2}{7} S_{31} \tau_2 - \frac{a_2^2}{7} S_{32} (\tau_2 + \tau_3) + \frac{a_3^2}{7} S_{33} \tau_3 \Big], (3.40)$$

where

$$\tau_3 \equiv a_1^2 a_3^2 A_{13} - 3a_3^4 A_{33} - a_1^2 a_2^2 A_{12} + a_2^2 a_3^2 A_{23} + 2a_3^2 A_3.$$
(3.41)

Eliminating $\delta \Omega^2$ from Eq. (3.39) by using Eq. (3.40), we obtain the identity

$$0 = (3a_1^2 + a_2^2)a_2^2 \left[\gamma_1 a_1^2 - \gamma_3 a_3^2 + \frac{1}{7} (6\gamma_{11}a_1^4 + \gamma_{12}a_1^2a_2^2 - \gamma_{23}a_2^2a_3^2 - 6\gamma_{33}a_3^4) \right] \\ + (a_1^2 + 3a_2^2)a_1^2 \left[\gamma_2 a_2^2 - \gamma_3 a_3^2 + \frac{1}{7} (\gamma_{12}a_1^2a_2^2 + 6\gamma_{22}a_2^4 - \gamma_{13}a_1^2a_3^2 - 6\gamma_{33}a_3^4) \right] \\ - \pi \rho (3a_1^2 + a_2^2)a_2^2 \left[S_{11}(\tau_1 - \tau_3) - S_{12}\tau_3 + \frac{a_1^2}{7}S_{31}\tau_1 \\ - \frac{a_2^2}{7}S_{32}\tau_3 - \frac{a_3^2}{7}S_{33}(\tau_1 - \tau_3) \right] \\ - \pi \rho (a_1^2 + 3a_2^2)a_1^2 \left[-S_{11}\tau_3 - S_{12}(\tau_2 + \tau_3) + \frac{a_1^2}{7}S_{31}\tau_2 \\ - \frac{a_2^2}{7}S_{32}(\tau_2 + \tau_3) + \frac{a_3^2}{7}S_{33}\tau_3 \right].$$

$$(3.42)$$

Equation (3.42) is one of the equations to determine p, q, r, s, and S_{ij} .

§4. Boundary conditions

In this section we derive equations to determine the coefficients of the 1PN velocity potential and the deformation of the figure, p, q, r, s, and S_{ij} , from the boundary conditions on the stellar surface. The stellar surfaces of the Riemann ellipsoid and its deformed figure are expressed as $S_R(x) = 0$ and S(x) = 0, respectively, where

$$S_{R}(x) = \sum_{l} \frac{x_{l}^{2}}{a_{l}^{2}} - 1, \qquad (4.1)$$

$$S(x) = \sum_{l} \frac{x_{l}^{2}}{a_{l}^{2}} - 1 - \sum_{j} \xi_{j} \frac{\partial S_{R}}{\partial x_{j}}$$

$$= S_{R}(x) - \frac{2}{c^{2}} \left[S_{11} \left(\frac{x_{1}^{2}}{a_{1}^{2}} - \frac{x_{3}^{2}}{a_{3}^{2}} \right) + S_{12} \left(\frac{x_{2}^{2}}{a_{2}^{2}} - \frac{x_{3}^{2}}{a_{3}^{2}} \right) + S_{31} \left(\frac{x_{1}^{4}}{3a_{1}^{2}} - \frac{x_{1}^{2}x_{2}^{2}}{a_{2}^{2}} \right) + S_{32} \left(\frac{x_{2}^{4}}{3a_{2}^{2}} - \frac{x_{2}^{2}x_{3}^{2}}{a_{3}^{2}} \right) + S_{33} \left(\frac{x_{3}^{4}}{3a_{3}^{2}} - \frac{x_{3}^{2}x_{1}^{2}}{a_{1}^{2}} \right) \right]. \qquad (4.2)$$

The boundary conditions for the continuity equation and integrated Euler equation are, respectively,

(A)
$$\sum_{j} C_{j} \frac{\partial S}{\partial x_{j}} = 0$$
 on $S(x) = 0$, (4.3)

(B)
$$\frac{P}{\rho} = 0$$
 on $S(x) = 0.$ (4.4)

The condition (A) comes from the constraint that the normal component of the velocity on the surface vanishes. The condition (B) determines the stellar surface. Eqs. (4.3) and (4.4) must hold to order of c^{-2} .

4.1. Condition (A)

Substituting C_i given by Eq. (2.10) into Eq. (4.3), we obtain an equation of 1PN order as

$$0 = x_1 x_2 (c_0 + c_1 x_1^2 + c_2 x_2^2 + c_3 x_3^2), \qquad (4.5)$$

where c_k (k = 0, 1, 2, 3) are functions of a_k , p, q, r, s, and S_{ij} . Equation (4.5) must be satisfied on S_R . Then, we obtain the following three equations for p, q, r, s and S_{ij} :

$$\begin{split} C_1^{(A)} &\equiv \frac{a_1^2 + a_2^2}{a_1^4 a_2^2} p + \frac{a_1^2 + 3a_2^2}{a_1^2 a_2^2} q + \frac{4\Omega}{a_1^2 + a_2^2} \bigg[\frac{1}{a_1^2} (-S_{11} + S_{12}) - \frac{5}{3} S_{31} \bigg] \\ &\quad -3\pi\rho \Omega \bigg(\frac{A_0}{a_1^2} - A_1 \bigg) \frac{a_1^2 - a_2^2}{a_1^2 a_2^2} - \frac{1}{2} F_a^2 \Omega^3 \frac{a_1^2 - a_2^2}{a_1^2 a_2^2} \\ &\quad + \frac{\pi\rho F_a \Omega}{2a_1^2} \bigg[\bigg(7\frac{a_1^2}{a_2^2} - 1 \bigg) B_{11} + \bigg(7\frac{a_2^2}{a_1^2} - 1 \bigg) B_{12} + 4(a_1^2 + a_2^2) B_{112} \bigg] = 0, (4\cdot6) \\ C_2^{(A)} &\equiv \frac{a_1^2 + a_2^2}{a_1^2 a_2^4} p + \frac{3a_1^2 + a_2^2}{a_1^2 a_2^2} r + \frac{4\Omega}{a_1^2 + a_2^2} \bigg[\frac{1}{a_2^2} (-S_{11} + S_{12}) + \frac{a_1^2}{a_2^2} S_{31} + \frac{2}{3} S_{32} \bigg] \end{split}$$

$$\begin{split} & -3\pi\rho\Omega\Big(\frac{A_0}{a_2^2} - A_2\Big)\frac{a_1^2 - a_2^2}{a_1^2 a_2^2} - \frac{1}{2}F_a^2\Omega^3\frac{a_1^2 - a_2^2}{a_1^2 a_2^2} \\ & +\frac{\pi\rho F_a\Omega}{2a_2^2}\Big[\Big(7\frac{a_1^2}{a_2^2} - 1\Big)B_{12} + \Big(7\frac{a_2^2}{a_1^2} - 1\Big)B_{22} + 4(a_1^2 + a_2^2)B_{122}\Big] = 0, (4\cdot7) \\ C_3^{(A)} &\equiv \frac{a_1^2 + a_2^2}{a_1^2 a_2^2 a_3^2}p + \Big(\frac{1}{a_1^2} + \frac{1}{a_2^2} + \frac{2}{a_3^2}\Big)s + \frac{4\Omega}{a_1^2 + a_2^2}\Big[\frac{1}{a_3^2}(-S_{11} + S_{12}) - \frac{a_2^2}{a_3^2}S_{32} + S_{33}\Big] \\ & -3\pi\rho\Omega\Big(\frac{A_0}{a_3^2} - A_3\Big)\frac{a_1^2 - a_2^2}{a_1^2 a_2^2} \\ & +\frac{\pi\rho F_a\Omega}{2a_3^2}\Big[\Big(7\frac{a_1^2}{a_2^2} - 1\Big)B_{13} + \Big(7\frac{a_2^2}{a_1^2} - 1\Big)B_{23} + 4(a_1^2 + a_2^2)B_{123}\Big] = 0.(4\cdot8) \end{split}$$

We can make two equations which do not contain S_{11} , S_{12} and p as

$$C_1^{(A)}a_1^2 - C_2^{(A)}a_2^2 = 0, (4.9)$$

$$C_1^{(A)}a_1^2 - C_3^{(A)}a_3^2 = 0. (4.10)$$

Thus, Eqs. (4.9) and (4.10) depend only on the coefficients q, r, s, S_{31} , S_{32} and S_{33} ; i.e., the coefficients for biquadratic deformation. This is an important feature in determining these coefficients.

4.2. Condition (B)

To obtain \mathcal{G} consistently, we must calculate the correction of the self-gravity potential due to the deformation of the figure δU , which is given by ¹⁸

$$\delta U = \frac{1}{c^2} \sum_{ij} S_{ij} \delta U^{(ij)}, \qquad (4.11)$$

where

$$\delta U^{(ij)} = -\sum_k \frac{\partial}{\partial x_k} \int \frac{\rho(x')\xi_k^{(ij)}(x')}{|x-x'|} d^3x'.$$
(4.12)

Substituting the Lagrangian displacement vectors into Eq. (4.12), we have

$$\delta U^{(11)} = -D_{1,1} + D_{3,3}, \tag{4.13}$$

$$\delta U^{(12)} = -D_{2,2} + D_{3,3}, \tag{4.14}$$

$$\delta U^{(31)} = -\frac{1}{3}D_{111,1} + D_{112,2}, \qquad (4.15)$$

$$\delta U^{(32)} = -\frac{1}{3}D_{222,2} + D_{223,3}, \qquad (4.16)$$

$$\delta U^{(33)} = -\frac{1}{3}D_{333,3} + D_{133,1}, \qquad (4.17)$$

where $^{20)}$ for $i \neq j$,

$$D_{iii} = \pi \rho \bigg[a_i^6 \Big(A_{iii} - \sum_l A_{iiil} x_l^2 \Big) x_i^3 \bigg]$$

$$+\frac{3}{4}a_{i}^{4}\left(B_{ii}-2\sum_{l}B_{iil}x_{l}^{2}+\sum_{l}\sum_{m}B_{iilm}x_{l}^{2}x_{m}^{2}\right)x_{i}\right],\qquad(4.18)$$

$$D_{iij} = \pi \rho \bigg[a_i^4 a_j^2 \Big(A_{iij} - \sum_l A_{iijl} x_l^2 \Big) x_i^2 x_j + \frac{1}{4} a_i^2 a_j^2 \Big(B_{ij} - 2 \sum_l B_{ijl} x_l^2 + \sum_l \sum_m B_{ijlm} x_l^2 x_m^2 \Big) x_j \bigg]. \quad (4.19)$$

Substituting δU and the equation S = 0 into Eq. (2.43), we obtain

$$\begin{split} \left(\frac{P}{\rho}\right)_{S} &= -\frac{2P_{0}}{\rho c^{2}} \bigg[S_{11} \bigg(\frac{x_{1}^{2}}{a_{1}^{2}} - \frac{x_{3}^{2}}{a_{3}^{2}} \bigg) + S_{12} \bigg(\frac{x_{2}^{2}}{a_{2}^{2}} - \frac{x_{3}^{2}}{a_{3}^{2}} \bigg) + S_{31} \bigg(\frac{x_{1}^{4}}{3a_{1}^{2}} - \frac{x_{1}^{2}x_{2}^{2}}{a_{2}^{2}} \bigg) \\ &\quad + S_{32} \bigg(\frac{x_{2}^{4}}{3a_{2}^{2}} - \frac{x_{2}^{2}x_{3}^{2}}{a_{2}^{2}} \bigg) + S_{33} \bigg(\frac{x_{3}^{4}}{3a_{3}^{2}} - \frac{x_{1}^{2}x_{3}^{2}}{a_{1}^{2}} \bigg) \bigg] \\ &\quad - \frac{1}{c^{2}} \bigg[S_{11} (D_{1,1} - D_{3,3}) + S_{12} (D_{2,2} - D_{3,3}) + S_{31} \bigg(\frac{1}{3} D_{111,1} - D_{112,2} \bigg) \\ &\quad + S_{32} \bigg(\frac{1}{3} D_{222,2} - D_{223,3} \bigg) + S_{33} \bigg(-D_{133,1} + \frac{1}{3} D_{333,3} \bigg) \bigg] \\ &\quad - \frac{1}{c^{2}} \delta \Omega^{2} \bigg[\frac{F_{a}^{2}}{2} (x_{1}^{2} + x_{2}^{2}) + F_{a} (x_{2}^{2} - x_{1}^{2}) \bigg] \\ &\quad - \frac{1}{c^{2}} \bigg(\gamma_{0} + \sum_{l} \gamma_{l} x_{l}^{2} + \sum_{l \leq m} \gamma_{lm} x_{l}^{2} x_{m}^{2} \bigg) + \text{const} \\ &\equiv \frac{1}{c^{2}} \bigg[Q_{1} x_{1}^{2} + Q_{2} x_{2}^{2} + Q_{3} x_{3}^{2} + Q_{11} x_{1}^{4} + Q_{22} x_{2}^{4} + Q_{33} x_{3}^{4} \\ &\quad + Q_{12} x_{1}^{2} x_{2}^{2} + Q_{13} x_{1}^{2} x_{3}^{2} + Q_{23} x_{2}^{2} x_{3}^{2} + \text{const} \bigg], \tag{4.20}$$

where

$$\begin{aligned} Q_{1} &= -\gamma_{1} - \frac{F_{a}}{2} (F_{a} - 2) \delta \Omega^{2} - S_{11} \pi \rho \left(\frac{2a_{3}^{2}}{a_{1}^{2}} A_{3} - 3a_{1}^{2} A_{11} + a_{3}^{2} A_{13} \right) \\ &+ S_{12} \pi \rho (a_{2}^{2} A_{12} - a_{3}^{2} A_{13}) \\ &- S_{31} \pi \rho a_{1}^{2} \left(a_{1}^{4} A_{111} - a_{1}^{2} a_{2}^{2} A_{112} - \frac{3}{2} a_{1}^{2} B_{111} + \frac{1}{2} a_{2}^{2} B_{112} \right) \\ &+ S_{32} \pi \rho \frac{a_{2}^{2}}{2} (a_{2}^{2} B_{122} - a_{3}^{2} B_{123}) + S_{33} \pi \rho \frac{a_{3}^{2}}{2} (a_{3}^{2} B_{133} - 3a_{1}^{2} B_{113}), \end{aligned}$$
(4·21)
$$Q_{2} &= -\gamma_{2} - \frac{F_{a}}{2} (F_{a} + 2) \delta \Omega^{2} + S_{11} \pi \rho (a_{1}^{2} A_{12} - a_{3}^{2} A_{23}) \\ &- S_{12} \pi \rho \left(\frac{2a_{3}^{2}}{a_{2}^{2}} A_{3} - 3a_{2}^{2} A_{22} + a_{3}^{2} A_{23} \right) + S_{31} \pi \rho \frac{a_{1}^{2}}{2} (a_{1}^{2} B_{112} - 3a_{2}^{2} B_{122}) \\ &- S_{32} \pi \rho a_{2}^{2} \left(a_{2}^{4} A_{222} - a_{2}^{2} a_{3}^{2} A_{223} - \frac{3}{2} a_{2}^{2} B_{222} + \frac{1}{2} a_{3}^{2} B_{223} \right) \end{aligned}$$

$$+S_{33}\pi\rho \frac{a_3^2}{2}(a_3^2B_{233} - a_1^2B_{123}), \qquad (4.22)$$

$$Q_3 = -\gamma_3 + S_{11}\pi\rho(2A_3 + a_1^2A_{13} - 3a_3^2A_{33}) + S_{12}\pi\rho(2A_3 + a_2^2A_{23} - 3a_3^2A_{33}) + S_{31}\pi\rho \frac{a_1^2}{2}(a_1^2B_{113} - a_2^2B_{123}) + S_{32}\pi\rho \frac{a_2^2}{2}(a_2^2B_{223} - 3a_3^2B_{233}) - S_{33}\pi\rho a_3^2 \left(a_3^4A_{333} - a_1^2a_3^2A_{133} - \frac{3}{2}a_3^2B_{333} + \frac{1}{2}a_1^2B_{133}\right), \qquad (4.23)$$

$$Q_{11} = -\gamma_{11} - S_{31}\pi\rho a_1^2 \left(\frac{2a_3^2}{3a_1^4}A_3 - \frac{5}{3}a_1^4A_{1111} + a_1^2a_2^2A_{1112} + \frac{5}{4}a_1^2B_{1111} - \frac{1}{4}a_2^2B_{1112}\right) -S_{32}\pi\rho \frac{a_2^2}{4}(a_2^2B_{1122} - a_3^2B_{1123}) - S_{33}\pi\rho \frac{a_3^2}{4}(a_3^2B_{1133} - 5a_1^2B_{1113}), \quad (4.24)$$

$$Q_{22} = -\gamma_{22} - S_{31}\pi\rho \frac{a_1^2}{4} (a_1^2 B_{1122} - 5a_2^2 B_{1222}) -S_{32}\pi\rho a_2^2 \left(\frac{2a_3^2}{3a_2^4} A_3 - \frac{5}{3}a_2^4 A_{2222} + a_2^2 a_3^2 A_{2223} + \frac{5}{4}a_2^2 B_{2222} - \frac{1}{4}a_3^2 B_{2223}\right) -S_{33}\pi\rho \frac{a_3^2}{4} (a_3^2 B_{2233} - a_1^2 B_{1223}),$$

$$(4.25)$$

$$Q_{33} = -\gamma_{33} - S_{31}\pi\rho \frac{a_1^2}{4} (a_1^2 B_{1133} - a_2^2 B_{1233}) - S_{32}\pi\rho \frac{a_2^2}{4} (a_2^2 B_{2233} - 5a_3^2 B_{2333}) -S_{33}\pi\rho a_3^2 \left(\frac{2}{3a_3^2} A_3 - \frac{5}{3}a_3^4 A_{3333} + a_1^2 a_3^2 A_{1333} + \frac{5}{4}a_3^2 B_{3333} - \frac{1}{4}a_1^2 B_{1333}\right),$$

$$(4.26)$$

$$Q_{12} = -\gamma_{12} + S_{31}\pi\rho a_1^2 \left(\frac{2a_3^2}{a_1^2 a_2^2} A_3 + a_1^4 A_{1112} - 3a_1^2 a_2^2 A_{1122} - \frac{3}{2} a_1^2 B_{1112} + \frac{3}{2} a_2^2 B_{1122} \right) + S_{32}\pi\rho a_2^2 \left(a_2^4 A_{1222} - a_2^2 a_3^2 A_{1223} - \frac{3}{2} a_2^2 B_{1222} + \frac{1}{2} a_3^2 B_{1223} \right) - S_{33}\pi\rho \frac{a_3^2}{2} \left(a_3^2 B_{1233} - 3a_1^2 B_{1123} \right),$$

$$Q_{13} = -\gamma_{13} + S_{31}\pi\rho a_1^2 \left(a_1^4 A_{1113} - a_1^2 a_2^2 A_{1123} - \frac{3}{2} a_1^2 B_{1113} + \frac{1}{2} a_2^2 B_{1123} \right)$$

$$(4.27)$$

$$-S_{32}\pi\rho \frac{a_2^2}{2} \left(a_2^2 B_{1223} - 3a_3^2 B_{1233} \right) +S_{33}\pi\rho a_3^2 \left(\frac{2}{a_1^2} A_3 + a_3^4 A_{1333} - 3a_1^2 a_3^2 A_{1133} - \frac{3}{2} a_3^2 B_{1333} + \frac{3}{2} a_1^2 B_{1133} \right), (4.28)$$
$$Q_{23} = -\gamma_{23} - S_{31}\pi\rho \frac{a_1^2}{2} \left(a_1^2 B_{1123} - 3a_2^2 B_{1223} \right) +S_{32}\pi\rho a_2^2 \left(\frac{2}{a_2^2} A_3 + a_2^4 A_{2223} - 3a_2^2 a_3^2 A_{2233} - \frac{3}{2} a_2^2 B_{2223} + \frac{3}{2} a_3^2 B_{2233} \right) +S_{33}\pi\rho a_3^2 \left(a_3^4 A_{2333} - a_1^2 a_3^2 A_{1233} - \frac{3}{2} a_3^2 B_{2333} + \frac{1}{2} a_1^2 B_{1233} \right).$$
(4.29)

Since $(P/\rho)_S$ must vanish on the surface, we have five conditions,

$$a_1^4 Q_{11} + a_2^4 Q_{22} - a_1^2 a_2^2 Q_{12} = 0, (4.30)$$

$$a_2^4 Q_{22} + a_3^4 Q_{33} - a_2^2 a_3^2 Q_{23} = 0, (4.31)$$

$$a_{2}^{4}Q_{22} + a_{3}^{4}Q_{33} - a_{2}^{2}a_{3}^{2}Q_{23} = 0, \qquad (4\cdot31)$$

$$a_{3}^{*}Q_{33} + a_{1}^{*}Q_{11} - a_{3}^{*}a_{1}^{*}Q_{13} = 0, \qquad (4.32)$$

$$a_1^* Q_{11} - a_2^* Q_{22} + a_1^2 Q_1 - a_2^2 Q_2 = 0, (4.33)$$

$$a_3^4 Q_{33} - a_1^4 Q_{11} + a_3^2 Q_3 - a_1^2 Q_1 = 0. (4.34)$$

When we substitute $\delta \Omega^2$ into Eqs. (4.33) and (4.34), we have two identical equations:

$$\begin{aligned} &(a_3^4Q_{33} + a_1^4Q_{11} - a_1^2a_3^2Q_{13}) - (a_2^4Q_{22} + a_3^4Q_{33} - a_2^2a_3^2Q_{23}) \\ &= 7(a_1^4Q_{11} - a_2^4Q_{22} + a_1^2Q_1 - a_2^2Q_2), \\ &(a_2^4Q_{22} + a_3^4Q_{33} - a_2^2a_3^2Q_{23}) - (a_1^4Q_{11} + a_2^4Q_{22} - a_1^2a_2^2Q_{12}) \\ &= 7(a_3^4Q_{33} - a_1^4Q_{11} + a_3^2Q_3 - a_1^2Q_1). \end{aligned}$$
(4.35)

This implies that Eqs. (4·33) and (4·34) can be derived from Eqs. (4·30) ~ (4·32), and only three of five conditions (4·30) ~ (4·34) are available. Here, we choose Eqs. (4·30) ~ (4·32) as the equations to be solved. Then, the three equations do not contain p, S_{11} and S_{12} , so that they depend only on q, r, s, S_{31} , S_{32} and S_{33} . This means that if we use Eqs. (4·30) ~ (4·32) together with Eqs. (2·42), (4·9) and (4·10), we can determine the six coefficients.

The question arises when we determine the rest three coefficients p, S_{11} and S_{12} because we have only *two* equations (3.42) and (4.6) (or (4.7) or (4.8)) for the *three* variables. There is one degree of freedom. This is because there is no unique definition of the 1PN solution as the counterpart of a Newtonian solution, as pointed out by Chandrasekhar^{17), 18)} and Bardeen.²³⁾ (Or, we may say that we have a degree to fix the PN correction of the angular velocity ($\delta\Omega$) or axial ratio (S_{11} or S_{12}) in obtaining a PN equilibrium solution.) There are many possible ways to compare a 1PN solution with a Newtonian solution, and to fix the 1PN counterpart to a Newtonian solution, we must impose an additional condition. (For example, Bardeen has proposed that we should compare two solutions fixing the angular momentum and the baryon number.²³)

Since the method for connecting a 1PN solution to a Newtonian solution can be arbitrarily chosen, in this paper, we simply give the condition as

$$S_{11} = S_{12}, \tag{4.37}$$

i.e., we fix the PN correction of the axial ratio. The reason why we choose this condition is simply that S_{11} and S_{12} in Eqs. (4.6) ~ (4.8) are erased, and manipulations are simplified. Then, we have three equations (3.42), (4.6), and (4.37) for solving p, S_{11} , and S_{12} . $\delta\Omega$ is determined from Eq. (3.38) if we obtain these coefficients.

§5. The total energy and angular momentum

5.1. Total energy

The total energy is written as

$$E = \int d^3x \rho \left[\frac{v^2}{2} - \frac{1}{2}U + \frac{1}{c^2} \left(\frac{5}{8} v^4 + \frac{5}{2} v^2 U + \frac{1}{2} \hat{\beta}_i v^i + \frac{P}{\rho} v^2 - \frac{5}{2} U^2 \right) + O(c^{-4}) \right]$$

= $E_{\rm N} + \frac{1}{c^2} E_{\rm PN} + O(c^{-4}),$ (5.1)

where

$$E_{\rm N} = \frac{M_*}{10} \bigg[F_a^2 \Omega_R^2 (a_{1*}^2 + a_{2*}^2) - 4\pi \rho A_{0*} \bigg], \tag{5.2}$$

$$E_{\rm PN} = E_{\rm N \to PN} + E_{v^4} + E_{v^2U} + E_{\beta_i v^i} + E_{Pv^2/\rho} + E_{U^2}, \qquad (5.3)$$

$$\begin{split} E_{\mathrm{N}\to\mathrm{PN}} &= -E_{\mathrm{N}} \left[\frac{1}{6} F_{a}^{2} \Omega_{R}^{2} (a_{1*}^{2} + a_{2*}^{2}) + 4\pi\rho A_{0*} \right] \\ &+ \frac{M_{*}}{5} F_{a}^{2} \Omega_{R}^{2} \left[a_{1*}^{2} S_{11} + a_{2*}^{2} S_{12} + \frac{1}{7} a_{1*}^{2} (a_{1*}^{2} - a_{2*}^{2}) S_{31} + \frac{1}{7} a_{2*}^{4} S_{32} \right. \\ &- \frac{1}{7} a_{1*}^{2} a_{3*}^{2} S_{33} \right] \\ &+ \frac{M_{*}}{10} F_{a}^{2} \delta \Omega^{2} (a_{1*}^{2} + a_{2*}^{2}) \\ &+ \frac{M_{*}}{10} F_{a}^{2} \delta \Omega^{2} (a_{1*}^{2} + a_{2*}^{2}) \left(p + \frac{3}{7} q a_{1*}^{2} + \frac{3}{7} r a_{2*}^{2} + \frac{1}{7} s a_{3*}^{2} \right) \\ &+ \frac{2M_{*}}{5} F_{a} \Omega_{R} (a_{1*}^{2} + a_{2*}^{2}) \left(p + \frac{3}{7} q a_{1*}^{2} + \frac{3}{7} r a_{2*}^{2} + \frac{1}{7} s a_{3*}^{2} \right) \\ &+ \frac{2M_{*}}{5} \pi \rho \left[S_{11} (a_{1*}^{2} A_{1*} - a_{3*}^{2} A_{3*}) + S_{12} (a_{2*}^{2} A_{2*} - a_{3*}^{2} A_{3*}) \right. \\ &+ \frac{1}{7} a_{1*}^{2} S_{31} (a_{1*}^{2} A_{1*} - a_{2*}^{2} A_{2*}) + \frac{1}{7} a_{2*}^{2} S_{32} (a_{2*}^{2} A_{2*} - a_{3*}^{2} A_{3*}) \\ &- \frac{1}{7} a_{3*}^{2} S_{33} (a_{1*}^{2} A_{1*} - a_{3*}^{2} A_{3*}) \right], \end{split}$$

$$E_{v^4} = \frac{M_*}{56} F_a^4 \Omega_R^4 (3a_{1*}^4 + 2a_{1*}^2 a_{2*}^2 + 3a_{2*}^4), \tag{5.5}$$

$$E_{v^{2}U} = \frac{M_{*}}{7} \pi \rho F_{a}^{2} \Omega_{R}^{2} \bigg[3A_{0*}(a_{1*}^{2} + a_{2*}^{2}) - (a_{1*}^{4}A_{1*} + a_{2*}^{4}A_{2*}) \bigg], \qquad (5.6)$$

$$E_{\beta_i v^i} = \frac{M_*}{35} \pi \rho F_a^2 \Omega_R^2 \bigg[a_{1*}^2 (a_{2*}^2 - 7a_{1*}^2) A_{1*} + a_{2*}^2 (a_{1*}^2 - 7a_{2*}^2) A_{2*} \\ -2a_{1*}^2 a_{2*}^2 (a_{1*}^2 + a_{2*}^2) A_{12*} \bigg],$$
(5.7)

$$E_{Pv^2/\rho} = \frac{2M_*}{35} \frac{P_0}{\rho} F_a^2 \Omega_R^2 (a_{1*}^2 + a_{2*}^2), \tag{5.8}$$

$$E_{U^2} = -\frac{M_*}{7} (\pi \rho)^2 (11A_{0*}^2 + a_{1*}^4 A_{1*}^2 + a_{2*}^4 A_{2*}^2 + a_{3*}^4 A_{3*}^2).$$
(5.9)

In the above equations, we have used Eq. (3.38) for $\delta \Omega^2$ and the conserved mass

$$M_* \equiv \int d^3x \rho \left[1 + \frac{1}{c^2} \left(\frac{v^2}{2} + 3U \right) \right],$$

20

$$= M \left[1 + \frac{1}{c^2} \left\{ \frac{1}{10} F_a^2 \Omega_R^2 (a_1^2 + a_2^2) + \frac{12}{5} \pi \rho A_0 \right\} \right], \tag{5.10}$$

instead of the Newtonian mass M. Also we substitute the mean radius of the star defined by the conserved mass as

$$a_* \equiv \left(\frac{M_*}{4\pi\rho/3}\right)^{1/3} = a_1(\alpha_2\alpha_3)^{1/3} \left[1 + \frac{1}{c^2} \left\{\frac{1}{30} F_a^2 \Omega_R^2(a_1^2 + a_2^2) + \frac{4}{5}\pi\rho A_0\right\}\right], \quad (5.11)$$

and a_{i*} defined by $a_{1*}a_{2*}a_{3*} = a_*^3$, $a_{2*}/a_{1*} = a_2/a_1$ and $a_{3*}/a_{1*} = a_3/a_1$. $A_{ijk\cdots*}$ and $B_{ijk\cdots*}$ are computed by using a_{2*}/a_{1*} and a_{3*}/a_{1*} . In Eq. (5.4) we have used some relations among index symbols.²⁰

5.2. Total angular momentum

The total angular momentum is written as

$$J = \int d^3x \rho \left[v_{\varphi} \left\{ 1 + \frac{1}{c^2} \left(v^2 + 6U + \frac{P}{\rho} \right) \right\} + \frac{\hat{\beta}_{\varphi}}{c^2} + O(c^{-4}) \right]$$

= $J_{\rm N} + \frac{1}{c^2} J_{\rm PN},$ (5.12)

where

$$v_{\varphi} = -x_2 v_1 + x_1 v_2, \tag{5.13}$$

$$\hat{\beta}_{\varphi} = -x_2 \hat{\beta}_1 + x_1 \hat{\beta}_2, \tag{5.14}$$

$$J_{\rm N} = \frac{M_*}{5} F_a \Omega_R (a_{1*}^2 - a_{2*}^2), \tag{5.15}$$

$$J_{\rm PN} = J_{\rm N \to PN} + J_{v_{\varphi}v^2} + J_{v_{\varphi}U} + J_{v_{\varphi}P/\rho} + J_{\beta_{\varphi}}, \qquad (5.16)$$

$$J_{\mathrm{N}\to\mathrm{PN}} = -J_{\mathrm{N}} \left[\frac{1}{6} F_{a}^{2} \Omega_{R}^{2} (a_{1*}^{2} + a_{2*}^{2}) + 4\pi\rho A_{0*} \right] + \frac{M_{*}}{5} (a_{1*}^{2} - a_{2*}^{2}) \left(p + \frac{3}{7} q a_{1*}^{2} + \frac{3}{7} r a_{2*}^{2} + \frac{1}{7} s a_{3*}^{2} \right) + \frac{M_{*}}{5} F_{a} \left[2\Omega_{R} \left\{ a_{1*}^{2} S_{11} - a_{2*}^{2} S_{12} + \frac{1}{7} a_{1*}^{2} S_{31} (a_{1*}^{2} + a_{2*}^{2}) - \frac{1}{7} a_{2*}^{4} S_{32} \right. \left. - \frac{1}{7} a_{1*}^{2} a_{3*}^{2} S_{33} \right\} + \delta \Omega (a_{1*}^{2} - a_{2*}^{2}) \right],$$

$$(5.17)$$

$$J_{\nu_{\varphi}\nu^{2}} = \frac{3M_{*}}{35} F_{a}^{3} \Omega_{R}^{3} (a_{1*}^{4} - a_{2*}^{4}), \tag{5.18}$$

$$J_{\nu_{\varphi}U} = \frac{12M_*}{35} \pi \rho F_a \Omega_R \bigg[3A_{0*}(a_{1*}^2 - a_{2*}^2) - a_{1*}^4 A_{1*} + a_{2*}^4 A_{2*} \bigg], \qquad (5.19)$$

$$J_{\nu_{\varphi}P/\rho} = \frac{2M_*}{35} \frac{P_0}{\rho} F_a \Omega_R (a_{1*}^2 - a_{2*}^2),$$

$$J_{\beta_{\varphi}} = \frac{2M_*}{35} \pi \rho F_a \Omega_R (-a_{1*}^2 a_{2*}^2 A_{1*} - 7a_{1*}^4 A_{1*} + a_{1*}^2 a_{2*}^2 A_{2*})$$
(5.20)

$$+7a_{2*}^4A_{2*} - 2a_{1*}^4a_{2*}^2A_{12*} + 2a_{1*}^2a_{2*}^4A_{12*}), (5.21)$$

and

$$\delta \Omega = \frac{\delta \Omega^2}{2\Omega_R}.\tag{5.22}$$

§6. Gravitational radiation

In this section, we derive an equation for the luminosity of the gravitational radiation from a rotating star.

6.1. Frames

In previous sections, we used the coordinate system $\{x_i\}$ associated with the principal axis of the ellipsoid, which is considered as the rotating frame. In these coordinates, the estimation of multipole moments is quite simple. On the other hand, it is convenient to use the inertial frame $\{X_i\}$ for the estimation of the luminosity of gravitational waves. These coordinate systems are related to each other as

$$X_1 = x_1 \cos \Omega t - x_2 \sin \Omega t,$$

$$X_2 = x_1 \sin \Omega t + x_2 \cos \Omega t,$$

$$X_3 = x_3.$$
(6.1)

In the inertial frame, the Newtonian internal velocity is

$$v_i = (-\Omega F_1 X_2, \ \Omega F_2 X_1, \ 0),$$
 (6.2)

where we define F_i for i = 1, 2 as

$$F_i \equiv 1 + \frac{a_i^2}{a_1^2 + a_2^2} f_R.$$
 (6.3)

Note that this expression of the Newtonian velocity is different from Eq. (2.17) because we do not restrict our attention to the irrotational case in this section. If one takes -2 for f_R , Eq. (6.2) reduces to Eq. (2.17).

6.2. Multipole moments

In the 1PN approximation, the energy loss due to gravitational radiation is written as $^{24)}$

$$\frac{dE}{dt} = -\frac{1}{5c^5} \left[\sum_{i,j} M_{ij}^{(3)} M_{ij}^{(3)} + \frac{1}{c^2} \left(\frac{5}{189} \sum_{i,j,k} M_{ijk}^{(4)} M_{ijk}^{(4)} + \frac{16}{9} \sum_{i,j} S_{ij}^{(3)} S_{ij}^{(3)} \right) \right], \quad (6.4)$$

where (n) denotes that n time derivatives are taken, and M_{ij} , M_{ijk} and S_{ij} are written as

$$M_{ij} \equiv K_{ij} + V_{ij} + U_{ij} + P_{ij} + Y_{ij} + R_{ij}, \tag{6.5}$$

$$K_{ij} \equiv \int d^3 X \rho \hat{X}_{ij}, \tag{6.6}$$

$$V_{ij} \equiv 2 \int d^3 X \rho v^2 \hat{X}_{ij}, \qquad (6.7)$$

$$U_{ij} \equiv 2 \int d^3 X \rho U \hat{X}_{ij}, \tag{6.8}$$

$$P_{ij} \equiv 3 \int d^3 X P \hat{X}_{ij}, \tag{6.9}$$

$$Y_{ij} \equiv \frac{1}{14} \frac{d^2}{dt^2} \int d^3 X \rho X^2 \hat{X}_{ij},$$
 (6.10)

$$R_{ij} \equiv -\frac{20}{21} \frac{d}{dt} \int d^3 X \rho \sum_k v_k \hat{X}_{ijk}, \qquad (6.11)$$

$$M_{ijk} \equiv \int d^3 X \rho \hat{X}_{ijk}, \qquad (6.12)$$

$$S_{ij} \equiv \int d^3 X \rho \sum_{k,l} \epsilon_{kl < i} \hat{X}_{j > k} v_l, \qquad (6.13)$$

where ϵ_{kli} denotes the spatial Levi-Civita symbol, and both 'hat' and <> denote the symmetric traceless part; that is,

$$\hat{X}_{ij} = X_i X_j - \frac{1}{3} \delta_{ij} X^2, \tag{6.14}$$

$$\hat{X}_{ijk} = X_i X_j X_k - \frac{1}{5} X^2 (\delta_{ij} X_k + \delta_{jk} X_i + \delta_{ki} X_j), \qquad (6.15)$$

$$A_{\langle ij \rangle} = \frac{1}{2} (A_{ij} + A_{ji}) - \frac{1}{3} \delta_{ij} \sum_{k} A_{kk}.$$
 (6.16)

First, we consider the mass multipole moments associated with the principal axis of the ellipsoid. We obtain mass quadrupole moments to 1PN order as

$$I_{11} \equiv \int_{V} d^{3}x \rho x_{1}^{2}$$

= $\frac{4\pi}{15} \rho a_{1}^{3} a_{2} a_{3} \left[1 + \frac{2}{c^{2}} \left(S_{11} + \frac{a_{1}^{2}}{7} S_{31} - \frac{a_{3}^{2}}{7} S_{33} \right) \right] + O(c^{-4}), \quad (6.17)$

$$I_{22} \equiv \int_{V} d^{3}x \rho x_{2}^{2}$$

= $\frac{4\pi}{15} \rho a_{1} a_{2}^{3} a_{3} \left[1 + \frac{2}{c^{2}} \left(S_{12} - \frac{a_{1}^{2}}{7} S_{31} + \frac{a_{2}^{2}}{7} S_{32} \right) \right] + O(c^{-4}), \quad (6.18)$

$$I_{12} \equiv \int_{V} d^{3}x \rho x_{1} x_{2} = 0.$$
 (6.19)

These moments are related to the reduced moments in the inertial frame as

$$K_{11} = \frac{1}{2}(I_{11} - I_{22})\cos 2\Omega t + \text{const}, \qquad (6.20)$$

$$K_{22} = -K_{11} + \text{const}, \tag{6.21}$$

$$K_{12} = \frac{1}{2}(I_{11} - I_{22})\sin 2\Omega t + \text{const.}$$
 (6.22)

6.3. Energy loss

The quadrupole moments relevant to the energy loss are obtained as

$$V_{11} - V_{22} = \Omega^2 \left[-\frac{1}{2} (I_{1111} + I_{2222} - 6I_{1122}) (F_1^2 - F_2^2) \cos 4\Omega t + (I_{1111} - I_{2222}) (F_1^2 + F_2^2) \cos 2\Omega t \right] + \text{const}, \quad (6.23)$$

$$V_{12} = \Omega^2 \left[-\frac{1}{4} (I_{1111} + I_{2222} - 6I_{1122}) (F_1^2 - F_2^2) \sin 4\Omega t + \frac{1}{2} (I_{1111} - I_{2222}) (F_1^2 + F_2^2) \sin 2\Omega t \right] + \text{const}, \qquad (6.24)$$

$$U_{11} - U_{22} = 2\pi\rho \left[A_0(I_{11} - I_{22}) - \sum_l A_l(I_{11ll} - I_{22ll}) \right] \cos 2\Omega t, \qquad (6.25)$$

$$U_{12} = \pi \rho \left[A_0(I_{11} - I_{22}) - \sum_l A_l(I_{11ll} - I_{22ll}) \right] \sin 2\Omega t, \qquad (6.26)$$

$$P_{11} - P_{22} = \frac{3P_0}{\rho} \left[(I_{11} - I_{22}) - \sum_l \frac{1}{a_l^2} (I_{11ll} - I_{22ll}) \right] \cos 2\Omega t, \tag{6.27}$$

$$P_{12} = \frac{3P_0}{2\rho} \left[(I_{11} - I_{22}) - \sum_l \frac{1}{a_l^2} (I_{11ll} - I_{22ll}) \right] \sin 2\Omega t, \tag{6.28}$$

$$Y_{11} - Y_{22} = -\frac{2}{7}\Omega^2 (I_{1111} - I_{2222} + I_{1133} - I_{2233})\cos 2\Omega t, \qquad (6.29)$$

$$Y_{12} = -\frac{1}{7}\Omega^2 (I_{1111} - I_{2222} + I_{1133} - I_{2233})\sin 2\Omega t, \qquad (6.30)$$

$$R_{11} - R_{22} = \Omega^2 \left[\frac{20}{21} (F_1 - F_2) (I_{1111} + I_{2222} - 6I_{1122}) \cos 4\Omega t - \frac{8}{21} (F_1 + F_2) (I_{1111} - I_{2222} + I_{1133} - I_{2233}) \cos 2\Omega t \right], (6.31)$$

$$R_{12} = \Omega^2 \left[\frac{10}{21} (F_1 - F_2) (I_{1111} + I_{2222} - 6I_{1122}) \sin 4\Omega t - \frac{4}{21} (F_1 + F_2) (I_{1111} - I_{2222} + I_{1133} - I_{2233}) \sin 2\Omega t \right]. (6.32)$$

The higher multipole moments relevant to gravitational radiation in the 1PN approximation vanish due to the symmetry, i.e.,

$$M_{ijk} = O(c^{-2}), (6.33)$$

$$S_{ij} = O(c^{-2}).$$
 (6.34)

In total, we obtain

$$\frac{dE}{dt} = -\frac{32}{5c^5}(I_{11} - I_{22})\Omega^6 \left[(I_{11} - I_{22}) + \frac{1}{c^2} \left\{ 2(I_{1111} - I_{2222})(F_1^2 + F_2^2)\Omega^2 + 4\pi\rho \left(A_0(I_{11} - I_{22}) - \sum_l A_l(I_{11ll} - I_{22ll}) \right) \right\}$$

$$+ \frac{6P_0}{\rho} \left(I_{11} - I_{22} - \sum_l \frac{1}{a_l^2} (I_{11ll} - I_{22ll}) \right) - \left(\frac{4}{7} + \frac{16}{21} (F_1 + F_2) \right) (I_{1111} - I_{2222} + I_{1133} - I_{2233}) \Omega^2 \bigg\} \bigg] + O(c^{-8}),$$
 (6.35)

where temporal averaging has been done.

Using some algebraic relations among I_{ii} and I_{iiii} , the formula for the energy loss can be expressed in terms of $(\rho, \Omega, a_1, a_2, a_3, S_{11}, \dots, S_{33}, A_0, f_R)$ as

$$\begin{aligned} \frac{dE}{dt} &= -\frac{32}{5c^5} \left(\frac{4\pi}{15}\right)^2 (\rho a_1 a_2 a_3)^2 (a_1^2 - a_2^2) \Omega^6 \\ &\times \left[(a_1^2 - a_2^2) + \frac{2}{c^2} \left\{ 2a_1^2 \left(S_{11} + \frac{a_1^2}{7} S_{31} - \frac{a_3^2}{7} S_{33} \right) \right. \\ &\left. -2a_2^2 \left(S_{12} - \frac{a_1^2}{7} S_{31} + \frac{a_2^2}{7} S_{32} \right) \right. \\ &\left. + (a_1^2 - a_2^2) \left(\frac{38}{21} \pi \rho A_0 + \frac{1}{147} \frac{\Omega_R^2}{a_1^2 + a_2^2} \mathcal{F}(f_R, a_1, a_2, a_3) \right) \right\} \right], \quad (6.36) \end{aligned}$$

where we define $\mathcal{F}(f_R, a_1, a_2, a_3)$ as

$$\mathcal{F}(f_R, a_1, a_2, a_3) \equiv (63f_R^2 + 102f_R + 11)(a_1^4 + a_2^4) -(49f_R^2 - 92f_R - 22)a_1^2a_2^2 - (8f_R + 22)a_3^2(a_1^2 + a_2^2).$$
(6.37)

6.4. The Energy loss rate for irrotational Riemann ellipsoids

For a 1PN irrotational ellipsoid $(f_R = -2)$, ²⁰⁾ we obtain

$$\begin{aligned} \frac{dE}{dt} &= -\frac{32}{5c^5} \left(\frac{4\pi}{15}\right)^2 (\rho a_1 a_2 a_3)^2 (a_1^2 - a_2^2) \Omega^6 \\ &\times \left[(a_1^2 - a_2^2) + \frac{2}{c^2} \left[2a_1^2 \left(S_{11} + \frac{a_1^2}{7} S_{31} - \frac{a_3^2}{7} S_{33} \right) - 2a_2^2 \left(S_{12} - \frac{a_1^2}{7} S_{31} + \frac{a_2^2}{7} S_{32} \right) \right. \\ &+ (a_1^2 - a_2^2) \left\{ \frac{38}{21} \pi \rho A_0 \right. \\ &\left. + \frac{1}{147} \left(59(a_1^4 + a_2^4) - 358a_1^2 a_2^2 - 6a_3^2(a_1^2 + a_2^2) \right) \frac{\Omega_R^2}{a_1^2 + a_2^2} \right\} \right] \right].$$
(6·38)

We can rewrite Eq. (6.38) by using the conserved mass as

$$\frac{dE}{dt} = \left(\frac{dE}{dt}\right)_{\rm N} + \frac{1}{c^2} \left(\frac{dE}{dt}\right)_{\rm PN},\tag{6.39}$$

where

$$\left(\frac{dE}{dt}\right)_{\rm N} = -\frac{32M_*^2}{125c^5}(a_{1*}^2 - a_{2*}^2)^2 \Omega_R^6,\tag{6.40}$$

$$\begin{split} \left(\frac{dE}{dt}\right)_{\rm PN} &= -\frac{64M_*^2}{125c^5} (a_{1*}^2 - a_{2*}^2) \Omega_R^6 \\ &\times \left[2a_{1*}^2 \left(S_{11} + \frac{a_{1*}^2}{7} S_{31} - \frac{a_{3*}^2}{7} S_{33} \right) - 2a_{2*}^2 \left(S_{12} - \frac{a_{1*}^2}{7} S_{31} + \frac{a_{2*}^2}{7} S_{32} \right) \\ &+ (a_{1*}^2 - a_{2*}^2) \left\{ \frac{3\delta\Omega^2}{2\Omega_R^2} - \frac{46}{21} \pi \rho A_{0*} - \frac{1}{6} F_a^2 \Omega_R^2 (a_{1*}^2 + a_{2*}^2) \right. \\ &\left. + \frac{1}{147} \left(59(a_{1*}^4 + a_{2*}^4) - 358a_{1*}^2 a_{2*}^2 - 6a_{3*}^2 (a_{1*}^2 + a_{2*}^2) \right) \frac{\Omega_R^2}{a_{1*}^2 + a_{2*}^2} \right\} \right]. \end{split}$$

$$(6.41)$$

Note that in Eq. (6.39) there are no terms dependent on the 1PN velocity, since the quadrupole moment M_{ij} in Eq. (6.4) does not depend on the velocity.

§7. Numerical results

7.1. Normalization

First of all, we change to non-dimensional parameters according to

$$\tilde{p} \equiv \frac{p}{\pi\rho\Lambda a_1^2}, \quad \tilde{q} \equiv \frac{q}{\pi\rho\Lambda}, \quad \tilde{r} \equiv \frac{r}{\pi\rho\Lambda}, \quad \tilde{s} \equiv \frac{s}{\pi\rho\Lambda}, \\ \tilde{S}_{11} \equiv \frac{S_{11}}{\pi\rho a_1^2}, \quad \tilde{S}_{12} \equiv \frac{S_{12}}{\pi\rho a_1^2}, \quad \tilde{S}_{31} \equiv \frac{S_{31}}{\pi\rho}, \quad \tilde{S}_{32} \equiv \frac{S_{32}}{\pi\rho}, \quad \tilde{S}_{33} \equiv \frac{S_{33}}{\pi\rho}, \quad (7.1)$$

and also

$$\tilde{\Omega} \equiv \frac{\Omega}{\sqrt{\pi\rho}}, \quad \delta \tilde{\Omega}^2 \equiv \frac{\delta \Omega^2}{(\pi\rho a_1)^2}, \quad \delta \tilde{\Omega} \equiv \frac{\delta \Omega}{(\pi\rho)^{3/2} a_1^2}.$$
(7.2)

Using these parameters with the relation $S_{11} = S_{12}$, we can rewrite equations (2·42), (3·42), (4·6) ~ (4·8), and (4·30) ~ (4·32) into the non-dimension forms. From these eight equations, we can determine eight variables, \tilde{p} , \tilde{q} , \tilde{r} , \tilde{s} , \tilde{S}_{11} , \tilde{S}_{31} , \tilde{S}_{32} and \tilde{S}_{33} .

After determination of the variables, we can calculate the energy and angular momentum of the star from Eqs. $(5\cdot1)$ and $(5\cdot12)$ and also the luminosity of gravitational radiation from Eq. $(6\cdot38)$. In the numerical calculations these values must be normalized. The normalized energy, angular momentum and luminosity of gravitational radiation are written as

$$\tilde{E} \equiv \frac{E}{M_*^2/a_*} = \tilde{E}_{\rm N} + C_{\rm s}\tilde{E}_{\rm PN},\tag{7.3}$$

$$\tilde{J} \equiv \frac{J}{(M_*^3 a_*)^{1/2}} = \tilde{J}_{\rm N} + C_{\rm s} \tilde{J}_{\rm PN},$$
(7.4)

$$\frac{\tilde{dE}}{dt} \equiv \left(\frac{dE}{dt}\right) \left/ \left(\frac{M_*^5}{c^5 a_*^5}\right) = \left(\frac{\tilde{dE}}{dt}\right)_{\rm N} + C_{\rm s} \left(\frac{\tilde{dE}}{dt}\right)_{\rm PN},\tag{7.5}$$

where $C_{\rm s}$ is the compactness parameter defined as

$$C_{\rm s} \equiv \frac{M_*}{c^2 a_*}.\tag{7.6}$$

26

In the PN approximation, we assume $C_{\rm s} \ll 1$.

7.2. Ellipsoidal approximation

The ellipsoidal approximation, in which the equilibrium configuration is assumed to have an ellipsoidal shape, is useful in order to investigate features of stars or binary systems in the 1PN approximation, $^{25), 26), 27}$ because the whole calculation is done by setting $S_{ij} = 0$, and hence is very simplified. The ellipsoidal approximation gives an exact solution for the rotating incompressible star in the Newtonian theory. However, it gives only an approximate solution for the 1PN case. If the ellipsoidal approximation is really an excellent approximation, it becomes a robust method for the study of the 1PN effects. This is motivation for our investigation of the validity of the ellipsoidal approximation.

Table I. The Newtonian and 1PN angular velocity, energy, angular momentum, and luminosity of gravitational radiation along the equilibrium sequence of the irrotational Riemann S-type ellipsoid.

a_2/a_1	a_3/a_1	$ ilde{\Omega}_R^2$	$\tilde{E}_{\rm N}$	$\tilde{J}_{ m N}$	$\delta \tilde{\Omega}^2$	$\tilde{E}_{\rm PN}$	$\tilde{J}_{\rm PN}$	$(d\tilde{E}/dt)_{\rm N}$	$(d\tilde{E}/dt)_{\rm PN}$
0.99	0.9950	0.2667	-0.6000	1.807(-5)	8.050(-2)	-4.284(-2)	2.957(-5)	-8.275(-7)	5.153(-6)
0.95	0.9741	0.2667	-0.5998	4.707(-4)	7.837(-2)	-4.235(-2)	7.312(-4)	-2.160(-5)	1.341(-4)
0.90	0.9464	0.2669	-0.5991	1.986(-3)	7.776(-2)	-4.071(-2)	2.879(-3)	-9.184(-5)	5.660(-4)
0.85	0.9167	0.2673	-0.5979	4.728(-3)	7.936(-2)	-3.775(-2)	6.359(-3)	-2.214(-4)	1.348(-3)
0.80	0.8848	0.2677	-0.5960	8.917(-3)	8.301(-2)	-3.324(-2)	1.106(-2)	-4.258(-4)	2.546(-3)
0.75	0.8505	0.2683	-0.5934	1.483(-2)	8.841(-2)	-2.694(-2)	1.682(-2)	-7.269(-4)	4.244(-3)
0.70	0.8136	0.2688	-0.5898	2.282(-2)	9.515(-2)	-1.861(-2)	2.338(-2)	-1.156(-3)	6.544(-3)
0.65	0.7736	0.2692	-0.5851	3.332(-2)	0.1026	-8.211(-3)	3.006(-2)	-1.757(-3)	9.594(-3)
0.60	0.7304	0.2691	-0.5789	4.691(-2)	0.1102	3.220(-3)	3.425(-2)	-2.591(-3)	1.370(-2)
0.55	0.6835	0.2683	-0.5711	6.433(-2)	0.1188	1.989(-3)	1.035(-2)	-3.739(-3)	2.110(-2)
$\dagger 0.5244$	0.6579	0.2673	-0.5663	7.505(-2)	∞	∞	∞	-4.481(-3)	∞
0.50	0.6325	0.2660	-0.5612	8.660(-2)	0.1111	9.301(-2)	0.1499	-5.305(-3)	1.530(-2)
0.45	0.5772	0.2614	-0.5485	0.1151	0.1117	0.1021	0.1357	-7.403(-3)	2.337(-2)
0.40	0.5174	0.2534	-0.5324	0.1516	0.1025	0.1379	0.1699	-1.012(-2)	2.763(-2)
0.35	0.4531	0.2403	-0.5121	0.1988	8.362(-2)	0.1895	0.2423	-1.345(-2)	2.898(-2)
0.30	0.3850	0.2206	-0.4864	0.2606	5.094(-2)	0.2762	0.4161	-1.707(-2)	2.507(-2)
0.25	0.3143	0.1927	-0.4540	0.3428	-7.291(-2)	0.7824	1.632	-2.018(-2)	-3.537(-3)
$\ddagger 0.2374$	0.2963	0.1842	-0.4447	0.3678	∞	∞	∞	2.071(-2)	∞
0.20	0.2430	0.1559	-0.4135	0.4551	6.538(-2)	5.760(-3)	1.493(-3)	-2.127(-2)	2.552(-2)
0.15	0.1737	0.1117	-0.3630	0.6154	2.253(-2)	0.1183	0.8808	-1.861(-2)	1.067(-2)
0.10	0.1094	6.454(-2)	-0.2994	0.8666	5.964(-3)	0.1163	2.973	-1.172(-2)	2.893(-3)
0.05	5.167(-2)	2.252(-2)	-0.2137	1.370	5.969(-4)	6.554(-2)	14.44	-3.460(-3)	9.916(-5)

In the ellipsoidal approximation, a solution is obtained by setting $S_{ij} = 0$ in all the equations. After we set $S_{ij} = 0$, we can calculate the velocity field in the 1PN approximation from Eqs. (2·42) and (4·6) ~ (4·8). We note that in the ellipsoidal approximation, the boundary condition (4·4) is not satisfied.

7.3. Results

We give the results in Tables I, II and III. In Table I, the angular velocity, energy, angular momentum, and luminosity of gravitational radiation of Newtonian and 1PN

orders are presented along the Newtonian sequence. Coefficients of the 1PN velocity potential and the Lagrangian displacement vectors are shown in Table II. In these tables, † and ‡ denote singularities at which all the 1PN terms diverge. We believe that at those points, some instabilities which are concerned with the fourth order harmonics of the Riemann ellipsoid are induced by relativistic corrections.

Table II. Coefficients of the 1PN velocity potential and the Lagrangian displacement vectors shown along the equilibrium sequence of the irrotational Riemann S-type ellipsoid.

/	~	~	~	~	ã	ã	ã	ã
a_2/a_1	p	q	r	s	S_{11}	S_{31}	S_{32}	S_{33}
0.99	3.337(-2)	-1.893(-3)	-1.936(-3)	-1.914(-3)	-9.723(-6)	4.082(-5)	8.268(-6)	-2.560(-6)
0.95	0.1635	-9.213(-3)	-1.034(-2)	-9.786(-3)	-2.393(-4)	1.017(-3)	2.096(-4)	-5.569(-5)
0.90	0.3180	-1.781(-2)	-2.245(-2)	-2.021(-2)	-9.335(-4)	3.996(-3)	8.393(-4)	-1.655(-4)
0.85	0.4623	-2.593(-2)	-3.643(-2)	-3.147(-2)	-2.033(-3)	8.639(-3)	1.841(-3)	-1.926(-4)
0.80	0.5947	-3.403(-2)	-5.209(-2)	-4.389(-2)	-3.470(-3)	1.425(-2)	3.050(-3)	1.025(-4)
0.75	0.7137	-4.334(-2)	-6.855(-2)	-5.794(-2)	-5.159(-3)	1.929(-2)	4.050(-3)	1.133(-3)
0.70	0.8177	-5.704(-2)	-8.293(-2)	-7.450(-2)	-7.021(-3)	2.019(-2)	3.860(-3)	3.615(-3)
0.65	0.9051	-8.443(-2)	-8.605(-2)	-9.551(-2)	-9.071(-3)	7.270(-3)	-4.744(-5)	8.805(-3)
0.60	0.9768	-0.1620	-4.032(-2)	-0.1273	-1.192(-2)	-5.589(-2)	-1.615(-2)	1.885(-2)
0.55	1.060	-0.6220	0.4026	-0.2216	-2.342(-2)	-0.4951	-0.1118	3.530(-2)
$\dagger 0.5244$	∞	∞	∞	∞	∞	∞	∞	∞
0.50	0.9322	0.9441	-1.289	-1.317(-2)	1.656(-2)	1.043	0.1764	9.268(-2)
0.45	0.9589	0.4818	-0.8558	-0.1228	8.354(-3)	0.5938	4.776(-2)	0.1726
0.40	0.9099	0.4682	-0.9012	-0.1789	1.439(-2)	0.5780	-4.266(-2)	0.3572
0.35	0.8081	0.5694	-1.073	-0.2463	2.877(-2)	0.6705	-0.2372	0.8163
0.30	0.6232	0.8617	-1.440	-0.3617	6.274(-2)	0.9574	-0.8477	2.320
0.25	-0.2187	3.306	-3.834	-0.9421	0.3245	3.571	-7.001	18.19
$\ddagger 0.2374$	∞	∞	∞	∞	∞	∞	∞	∞
0.20	0.8031	-0.6934	-0.2521	-0.2627	-9.644(-2)	-0.8229	3.568	-9.625
0.15	0.4650	-8.768(-2)	-1.024	-0.6262	-2.999(-2)	-0.2098	2.168	-6.409
0.10	0.2583	7.056(-2)	-1.555	-1.133	-1.064(-2)	-6.378(-2)	1.877	-6.251
0.05	0.1006	0.1021	-2.758	-2.463	-2.574(-3)	-1.361(-2)	1.885	-7.055

As shown in Table II, the velocity field of 1PN order has an x_3 component which is written as

$$v_3 = \partial_3 \phi_{\rm PN} = 2sx_1 x_2 x_3, \tag{7.7}$$

where we use Eq. (2.41). This feature has also been found in the study of the Dedekind ellipsoid.¹⁹⁾ In the 1PN (and relativistic) case, we believe that the velocity field cannot be restricted in the planes orthogonal to the figure rotation axis in general.

In Table III, the angular velocity, energy, angular momentum, luminosity of gravitational radiation and velocity field of 1PN order in the ellipsoidal approximation are presented. There appear no singularities which are associated with instabilities because we do not take into account deformation of the ellipsoidal figure. From Table III, it is found that the velocity field of 1PN order has an x_3 component, as in the case of the exact calculation. This implies that it is necessary to include the x_3 component of the velocity field even in the ellipsoidal approximation.

In Fig. 1, we show the angular velocity of Newtonian and 1PN orders as a

Table III. The 1PN angular velocity, energy, angular momentum, luminosity of gravitational radiation and velocity field in the ellipsoidal approximation along the equilibrium sequence of the irrotational Riemann S-type ellipsoid. The subscript e denotes the case of the *ellipsoidal* approximation.

a_2/a_1	$\delta \tilde{\Omega}_{ m e}^2$	$\tilde{E}_{\rm PNe}$	\tilde{J}_{PNe}	$(d\tilde{E}/dt)_{\rm PNe}$	$ ilde{p}_{ ext{e}}$	\tilde{q}_{e}	\tilde{r}_{e}	\tilde{s}_{e}
0.99	8.050(-2)	-4.284(-2)	2.954(-5)	5.154(-6)	3.336(-2)	-1.921(-3)	-1.907(-3)	-1.914(-3)
0.95	7.826(-2)	-4.235(-2)	7.275(-4)	1.343(-4)	0.1633	-9.959(-3)	-9.595(-3)	-9.787(-3)
0.90	7.735(-2)	-4.073(-2)	2.849(-3)	5.676(-4)	0.3172	-2.088(-2)	-1.938(-2)	-2.021(-2)
0.85	7.844(-2)	-3.780(-2)	6.259(-3)	1.354(-3)	0.4608	-3.287(-2)	-2.948(-2)	-3.149(-2)
0.80	8.138(-2)	-3.336(-2)	1.084(-2)	2.562(-3)	0.5929	-4.601(-2)	-4.012(-2)	-4.388(-2)
0.75	8.592(-2)	-2.713(-2)	1.645(-2)	4.276(-3)	0.7120	-6.030(-2)	-5.165(-2)	-5.774(-2)
0.70	9.168(-2)	-1.884(-2)	2.296(-2)	6.596(-3)	0.8169	-7.563(-2)	-6.466(-2)	-7.353(-2)
0.65	9.812(-2)	-8.131(-3)	3.027(-2)	9.633(-3)	0.9056	-9.168(-2)	-8.001(-2)	-9.187(-2)
0.60	0.1045	5.362(-3)	3.835(-2)	1.350(-2)	0.9762	-0.1079	-9.899(-2)	-0.1136
0.55	0.1099	2.202(-2)	4.729(-2)	1.826(-2)	1.026	-0.1232	-0.1234	-0.1398
0.50	0.1130	4.218(-2)	5.749(-2)	2.388(-2)	1.054	-0.1360	-0.1560	-0.1720
0.45	0.1126	6.598(-2)	6.993(-2)	3.009(-2)	1.055	-0.1439	-0.2003	-0.2122
0.40	0.1071	9.315(-2)	8.678(-2)	3.621(-2)	1.026	-0.1438	-0.2611	-0.2633
0.35	9.567(-2)	0.1226	0.1130	4.096(-2)	0.9658	-0.1320	-0.3441	-0.3289
0.30	7.828(-2)	0.1519	0.1599	4.251(-2)	0.8710	-0.1049	-0.4562	-0.4147
0.25	5.664(-2)	0.1763	0.2559	3.900(-2)	0.7430	-6.093(-2)	-0.6049	-0.5289
0.20	3.434(-2)	0.1886	0.4789	2.994(-2)	0.5874	-3.273(-3)	-0.8009	-0.6868
0.15	1.596(-2)	0.1799	1.078	1.763(-2)	0.4161	5.746(-2)	-1.070	-0.9242
0.10	4.719(-3)	0.1427	3.088	6.679(-3)	0.2469	0.1024	-1.512	-1.357
0.05	4.973(-4)	7.597(-2)	14.52	9.189(-4)	9.926(-2)	0.1057	-2.718	-2.594

function of the axial ratio a_2/a_1 . Figures 1(a) and (b) are drawn for the Newtonian solutions, and (c) and (d) are for the 1PN solutions. The solid and dotted lines denote the results of the exact calculation and those of the ellipsoidal approximation, respectively. In the Newtonian case (or in the spherical case), the above two lines coincide. On the other hand, in the 1PN (and non-spherical) cases, the differences between the two lines appear, but are not large.

The energy and angular momentum of Newtonian and 1PN orders are shown in Fig. 2 as functions of the axial ratio a_2/a_1 . In Fig. 2, we present the results only for $1 \ge a_2/a_1 \ge 0.55$, i.e., for the stable branch. The solid and dotted lines denote the same quantities as those in Fig. 1. The luminosity of gravitational radiation is shown in Fig. 3. The solid and dotted lines are again the same as in Fig. 1. From the figures, it is found that with increasing the 1PN correction, the angular velocity and angular momentum increase, while the total energy and luminosity of gravitational waves decrease. These features are reasonable, because with increasing relativistic correction, the self-gravity of the star becomes strong (and hence the binding energy becomes large), and to support the self-gravity, angular velocity must be larger.

§8. Summary and discussion

We have calculated an equilibrium sequence and the luminosity of gravitational waves of an irrotational and incompressible star in the 1PN approximation. We have



Fig. 1. The angular velocity of Newtonian (a) and (b) and 1PN (c) and (d) orders as functions of the axial ratio a_2/a_1 . Figures 1(a) and (c) are magnifications of Figs. 1(b) and (d). The solid and dotted lines denote the results of the exact calculation and of the ellipsoidal approximation, respectively.

developed a scheme to solve the hydrostatic equations for an irrotational fluid in the 1PN approximation. By extending the present study, we will investigate irrotational binary stars in the 1PN approximation. Also, the 1PN solutions presented in this paper will be useful when examining the accuracy of numerical code for obtaining relativistic irrotational stars.

In addition to establishing the method, we have an interesting result. The 1PN velocity field has an x_3 component. This fact implies that internal motion never remains in the planes orthogonal to the figure rotation axis. (Note that such a waving motion closes even if it has an x_3 component.) We can find in the Chandrasekhar and



Fig. 2. The total energy of Newtonian (a) and 1PN (c) orders, and the total angular momentum of Newtonian (b) and 1PN (d) orders as a function of the axial ratio a_2/a_1 . The solid and dotted lines are the same as in Fig. 1.

Elbert paper¹⁹⁾ that there is also an x_3 component of the internal motion in the case of the 1PN Dedekind ellipsoid. Moreover, Uryū and Eriguchi²¹⁾ numerically showed that the internal flow of the compressible Dedekind-like star has an x_3 component even in the Newtonian case. Therefore, except for *incompressible*, *Newtonian* cases, non-axisymmetric stars with an internal motion will have an x_3 component of the velocity field in general.

Finally, we discuss the validity of the ellipsoidal approximation. One can see from Figs. 1 ~ 3 and Tables I ~ III that the error of the 1PN angular velocity, energy, and angular momentum in the ellipsoidal approximation are less than about ~ 5% in the range of $1 \ge a_2/a_1 \ge 0.7$. Although some of the coefficients of the



Fig. 3. The luminosity of gravitational radiation of the Newtonian (a) and 1PN (b) orders as functions of the axial ratio a_2/a_1 . The solid and dotted lines are the same as in Fig. 1.

1PN velocity potential, \tilde{q} and \tilde{r} , deviate from exact values by a large factor near $a_2/a_1 \simeq 0.55$, the error is not so large for larger a_2/a_1 . Therefore, we may conclude that the ellipsoidal approximation is a good one if the star is not highly nonspherical $(1 \ge a_2/a_1 \ge 0.7)$. This result is encouraging for study of binary stars in the ellipsoidal approximation because in the binary case, it is expected that a_2/a_1 and a_3/a_1 do not become very small.²⁵⁾

Acknowledgements

We thank Y. Eriguchi for helpful comments. K. T. would like to thank T. Nakamura and H. Sato for helpful comments and continuous encouragement. This work was partly supported by a Grant-in-Aid for Scientific Research Fellowship (No.9402) and a Grant-in-Aid (Nos. 08NP0801 and 09740336) of the Japanese Ministry of Education, Science, Sports and Culture.

References

- [1] A. Abramovici et al., Science **256** (1992), 325.
- [2] C. Bradaschia et al., Nucl. Instrum. and Methods A289 (1990), 518.
- [3] J. Hough, in Proceedings of the Sixth Marcel Grossmann Meeting, ed. H. Sato and T. Nakamura (World Scientific, Singapore, 1992), p. 192.
- K. Kuroda et al., in Proceedings of International Conference on Gravitational Waves: Sources and Detectors, ed. I. Ciufolini and F. Fidecaro (World Scientific, Singapore, 1997), p. 100.
- [5] For example, K. S. Thorne, in Proceeding of Snowmass 94 Summer Study on Particle and Nuclear Astrophysics and Cosmology, ed. E. W. Kolb and R. Peccei (World Scientific, Singapore, 1995) and references cited therein.
- [6] L. Lindblom, Astrophys. J. **398** (1992), 569.
- M. Shibata, Phys. Rev. D55 (1997), 6019; Prog. Theor. Phys. 96 (1996), 317.
 M. Shibata, K. Taniguchi, and T. Nakamura, Prog. Theor. Phys. Supple. 128 (1997), 295.
- [8] T. W. Baumgarte, G. B. Cook, M. A. Scheel, S. L. Shapiro and S. A. Teukolsky, Phys. Rev. Lett. 79 (1997), 1182; Phys. Rev. D57 (1998), 7299 and references cited therein.
- [9] C. S. Kochanek, Astrophys. J. 398 (1992), 234.
- [10] L. Bildsten and C. Cutler, Astrophys. J. 400 (1992), 175.
- [11] S. Bonazzola, E. Gourgoulhon, and J.-A. Marck, Phys. Rev. D56 (1997), 7740.
- [12] H. Asada, Phys. Rev. **D57** (1998), 7292.
- [13] M. Shibata, Phys. Rev. **D58** (1998), 024012.
- [14] S. A. Teukolsky, accepted to Astrophys. J. (1998), preprint gr-qc/9803082.
- [15] E. Gourgoulhon, preprint gr-qc/9804054.
- [16] S. Chandrasekhar, Astrophys. J. 142 (1965), 1488.
- [17] S. Chandrasekhar, Astrophys. J. 142 (1965), 1513; 147 (1967), 334; 167 (1971), 447. See also Ref. 18).
- [18] S. Chandrasekhar, Astrophys. J. 148 (1967), 621; 167, 455 (1971).
- [19] S. Chandrasekhar and D. D. Elbert, Astrophys. J. 192 (1974), 731; 220, 303 (1978).
- [20] S. Chandrasekhar, Ellipsoidal Figures of Equilibrium (Yale University Press, New Haven, CT, 1969).
- [21] K. Uryū and Y. Eriguchi, Mon. Not. R. Astron. Soc. 282 (1996), 653; 296 (1998), L1; preprint astro-ph/9808102, 9808118.
- [22] D. Lai, F. Rasio and S. L. Shapiro, Astrophys. J. Supple. 88 (1993), 205.
- [23] J. M. Bardeen, Astrophys. J. 167 (1971), 425.
- [24] L. Blanchet and T. Damour, Ann. Inst. H. Poincare (Phys. Theor.) 50 (1989), 377.
- [25] K. Taniguchi and M. Shibata, Phys. Rev. **D56** (1997), 798.
- M. Shibata and K. Taniguchi, **56** (1997), 811.
- [26] J. C. Lombardi, F. A. Rasio and S. L. Shaipro, Phys. Rev. D56 (1997), 3416.
- [27] S. L. Shapiro and S. Zane, submitted to Astrophys. J. (1998), preprint gr-qc/9711050.