

**Abelian chiral gauge theories on the lattice  
with exact gauge invariance**

Martin Lüscher

*Deutsches Elektronen-Synchrotron DESY  
Notkestrasse 85, D-22603 Hamburg, Germany  
E-mail: luscher@mail.desy.de*

**Abstract**

It is shown that  $U(1)$  chiral gauge theories with anomaly-free multiplets of Weyl fermions can be put on the lattice without breaking the gauge invariance or violating any other fundamental principle. The Ginsparg-Wilson relation plays a key rôle in this construction, which is non-perturbative and includes all topological sectors of the theory in finite volume. In particular, the cancellation of the gauge anomaly and the absence of global topological obstructions can be established on the basis of this relation and the lattice symmetries alone.

November 2024

## 1. Introduction

For well-known reasons the formulation of chiral gauge theories on the lattice proves to be difficult and no completely satisfactory solution of the problem has been found so far [1]. One of the propositions that have been made is to put the gauge-fixed theory on the lattice and to include a set of counterterms in the action with coefficients chosen in such a way that the BRS symmetry is restored in the continuum limit [2] (for a review and further references see refs. [3,4]). Using lattices with different lattice spacings for the gauge and the fermion fields is another idea which is being actively pursued [5–8]. The symmetry breaking terms can then be suppressed by choosing the lattice spacing in the fermion sector to be much smaller than the other lattice spacing.

A few years ago an entirely different approach was suggested by Kaplan [9], who noted that fermion modes which are bound to a four-dimensional defect in a five-dimensional lattice are chiral under certain conditions. Later this led to the so-called overlap formulation of chiral gauge theories [10,11], where the fermion partition function is written as a transition matrix element (the “overlap”) between the ground states of two auxiliary Hamilton operators. This development no doubt represents a big step forward, but as in the other cases the gauge symmetry is broken on the lattice. Moreover the locality properties of the theory are not transparent.

In this paper we consider U(1) gauge theories where the gauge field couples to  $N$  left-handed Weyl fermions with charges  $e_\alpha$  satisfying

$$\sum_{\alpha=1}^N e_\alpha^3 = 0. \tag{1.1}$$

This is the classical condition for anomaly cancellation and the continuum theory is thus well-defined to all orders of perturbation theory. In particular, the effective gauge field action generated by the fermions is uniquely determined up to finite renormalizations of the gauge coupling [12–14].

The main result obtained here is that these theories can be put on the lattice without breaking the gauge invariance or violating other basic principles such as the requirement of locality. The construction is non-perturbative and one has the right number and type of Weyl fermions from the beginning. Not surprisingly it is technically rather involved and perhaps not as explicit as one would like, particularly in finite volume, where the non-trivial topology of the space of gauge fields gives rise to additional complications. The present paper is hence mainly of theoretical interest, clarifying a question of principle, but it does not provide a formulation

of chiral gauge theories on the lattice which would be immediately usable for non-perturbative studies through numerical simulations. One should however note that chiral gauge theories are anyway a difficult case for numerical simulations, because the effective action has a non-zero imaginary part in general.

The starting point in this paper is the recent discovery that chiral symmetry can be preserved on the lattice without having to compromise in any other ways [15–18]. One achieves this by choosing a lattice Dirac operator  $D$  satisfying †

$$\gamma_5 D + D \gamma_5 = D \gamma_5 D. \tag{1.2}$$

This relation (which is originally due to Ginsparg and Wilson [19]) guarantees that the fermion action is invariant under a group of infinitesimal transformations which may be regarded as a lattice form of the usual chiral symmetries. Moreover the non-invariance of the fermion integration measure under flavour-singlet transformations straightforwardly leads to the expected axial anomaly [18,20].

Having an exact chiral symmetry of the action, it turns out to be relatively easy to introduce left- and right-handed fields [24]. Because of the anomaly the fermion integration measure however does not decompose in a unique way and one ends up with a gauge field dependent phase ambiguity. To fix the phase of the measure so that the gauge symmetry and the locality of the theory are preserved is the principal problem which has to be solved if one would like to set up chiral gauge theories along these lines.

All this will be explained in more detail in the next two sections. We then discuss the conditions which an ideal fermion integration measure should fulfil (sect. 4). Whether such measures exist is far from obvious and the rest of the paper is in fact entirely devoted to this question. For clarity the results are first presented in sect. 5 in a concise form, with all proofs and technical details being deferred to sections 6–11. A few concluding remarks are collected in sect. 12.

---

† For notational convenience the lattice spacing  $a$  is set to 1 so that all length scales are given in numbers of lattice spacings. In particular, the right-hand side of eq. (1.2) should be multiplied with  $a$  if physical units are employed

## 2. Lattice fields and functional integral

The lattice theories constructed in this paper are defined in the traditional manner, where one begins by specifying the space of fields and the lattice action. Expectation values of arbitrary products of the fields are then obtained as usual from the functional integral. The definition of the integration measure for Weyl fermions is non-trivial, however, and there are further technical details which need to be discussed carefully. A summary of notational conventions is included in appendix A.

### 2.1 Gauge fields

We choose lattice units and construct the theory on a finite lattice of size  $L$  with periodic boundary conditions.  $U(1)$  gauge fields on such a lattice may be represented through periodic link fields,

$$U(x, \mu) \in U(1), \quad x = (x_0, \dots, x_3) \in \mathbb{Z}^4, \quad (2.1)$$

$$U(x + L\hat{\nu}, \mu) = U(x, \mu) \quad \text{for all } \mu, \nu = 0, \dots, 3, \quad (2.2)$$

on the infinite lattice. The independent degrees of freedom are then the link variables at the points  $x$  in the block

$$\Gamma = \{x \in \mathbb{Z}^4 \mid 0 \leq x_\mu < L\} \quad (2.3)$$

( $L \geq 1$  is assumed to be an integer). Under gauge transformations

$$U(x, \mu) \rightarrow \Lambda(x)U(x, \mu)\Lambda(x + \hat{\mu})^{-1}, \quad (2.4)$$

the periodicity of the field will be preserved if  $\Lambda(x) \in U(1)$  is periodic. This is not the most general possibility, but the convention is here adopted that only periodic functions are referred to as gauge transformations.

For the gauge field action  $S_G$  we take a somewhat unusual expression which effectively imposes an upper bound on the lattice field tensor. The reasons for this will become clear later. As in the case of the standard Wilson action we write

$$S_G = \frac{1}{4g_0^2} \sum_{x \in \Gamma} \sum_{\mu, \nu} \mathcal{L}_{\mu\nu}(x) \quad (2.5)$$

with  $g_0$  being the bare coupling. The plaquette action is however taken to be of the more complicated form

$$\mathcal{L}_{\mu\nu}(x) = \begin{cases} [F_{\mu\nu}(x)]^2 \{1 - [F_{\mu\nu}(x)]^2 / \epsilon^2\}^{-1} & \text{if } |F_{\mu\nu}(x)| < \epsilon, \\ \infty & \text{otherwise,} \end{cases} \quad (2.6)$$

where  $\epsilon$  is a fixed number in the range  $0 < \epsilon < \frac{1}{3}\pi$  and the field tensor  $F_{\mu\nu}(x)$  is defined through

$$F_{\mu\nu}(x) = \frac{1}{i} \ln P(x, \mu, \nu), \quad -\pi < F_{\mu\nu}(x) \leq \pi, \quad (2.7)$$

$$P(x, \mu, \nu) = U(x, \mu)U(x + \hat{\mu}, \nu)U(x + \hat{\nu}, \mu)^{-1}U(x, \nu)^{-1}. \quad (2.8)$$

Note that the Boltzmann factor  $e^{-S_G}$  is a smooth function of the link variables with this choice of action. In particular, the functional integral can be set up in the usual way with the standard integration measure for U(1) lattice gauge fields.

This concludes the definition of the pure gauge part of the theory. There are a few remarks which should be added here.

(a) The Boltzmann factor is a product of local factors, one for each plaquette on the lattice. The locality of the theory is thus guaranteed. Moreover since it is differentiable, no special precautions are required when performing partial integrations in the functional integral (such as those needed when deriving the field equations).

(b) As already mentioned, our choice of action is such that the functional integral is effectively restricted to the space of fields satisfying

$$|F_{\mu\nu}(x)| < \epsilon \quad \text{for all } x, \mu, \nu. \quad (2.9)$$

Gauge fields of this type will be referred to as *admissible* in the following.

(c) When physical units are employed, the parameter  $\epsilon$  should be replaced by  $\epsilon/a^2$  where  $a$  denotes the lattice spacing. It is then immediately clear that the curly bracket in the definition (2.6) of the action and the bound (2.9) are irrelevant in the classical continuum limit. As far as the weak coupling phase is concerned, there is in fact little doubt that the lattice theory defined here is in the same universality class as the standard lattice theory. In particular, it is a valid lattice regularization of the free U(1) gauge theory.

## 2.2 Magnetic flux sectors

Before proceeding to the fermion fields, we briefly discuss the topology of the space of admissible fields. Proofs and further details will be given in sect. 7. One might expect that there is no interesting topological structure in this simple theory, but this is not so. The key observation is that the magnetic flux

$$\phi_{\mu\nu}(x) = \sum_{s,t=0}^{L-1} F_{\mu\nu}(x + s\hat{\mu} + t\hat{\nu}) \quad (2.10)$$

through the  $(\mu, \nu)$ -planes of the lattice is conserved and quantized. In other words, for any admissible field the associated flux satisfies

$$\phi_{\mu\nu}(x) = 2\pi m_{\mu\nu}, \quad (2.11)$$

where  $m_{\mu\nu} = -m_{\nu\mu}$  is an integer tensor independent of  $x$ .

Evidently the flux quantum numbers  $m_{\mu\nu}$  cannot change when the gauge field is continuously deformed. The field space is thus a disjoint union of the sectors of all admissible fields with a given magnetic flux. Moreover it can be shown that each of these sectors has the topology a multi-dimensional torus times a convex space.

## 2.3 Lattice Dirac operator

We first consider Dirac fermions and discuss the projection to the left-handed components in the next subsection. Dirac fields  $\psi(x)$  on the lattice carry a Dirac index and a flavour index  $\alpha = 1, \dots, N$ . As in the case of the gauge field it is convenient to assume that the fermion fields are defined on the infinite lattice. Periodic boundary conditions are then imposed through the requirement that

$$\psi(x + L\hat{\mu}) = \psi(x) \quad \text{for all } \mu = 0, \dots, 3. \quad (2.12)$$

Other types of periodic boundary conditions could be admitted here with little change in the following.

Under gauge transformations  $\Lambda(x)$  the fermion fields transform according to the representation

$$\psi(x) \rightarrow R[\Lambda(x)]\psi(x), \quad R[\Lambda(x)]_{\alpha\beta} = \delta_{\alpha\beta} \Lambda(x)^{e_\alpha}, \quad (2.13)$$

where  $e_\alpha \in \mathbb{Z}$  is the charge of the fermion with flavour  $\alpha$ . Throughout the paper we take it for granted that the condition for anomaly cancellation, eq. (1.1), is satisfied.

A simple example of an acceptable charge assignment is thus given by

$$e_1 = e_2 = \dots = e_8 = 1, \quad e_9 = -2. \quad (2.14)$$

Taking pairs of charges with opposite sign is another possibility, but in the present context this is a less interesting case, because one ends up with a chiral theory which is effectively vector-like.

The proper choice of the lattice Dirac operator  $D$  is of central importance in the following. Apart from being a solution of the Ginsparg-Wilson relation (1.2), the operator should fulfil a number of technical requirements. In particular, it should be local, gauge covariant and differentiable in the gauge field. The complete list of requirements is given in appendix B.

Gauge covariant solutions of the Ginsparg-Wilson relation are not easy to find. The “perfect” lattice Dirac operator of refs. [15,16] is one of them and another solution has been derived by Neuberger [17] from the overlap formalism. In this case all properties described in appendix B have been established rigorously [21]. Note that it suffices to define the Dirac operator for all admissible gauge fields since only these contribute to the functional integral. The relevant results of ref. [21] in fact apply for admissible fields only and if  $\epsilon$  is such that  $|e_\alpha|\epsilon < \frac{1}{30}$  for all  $\alpha$ .

In infinite volume the action of the Dirac operator is given by

$$D\psi(x) = \sum_{y \in \mathbb{Z}^4} D(x, y)\psi(y), \quad (2.15)$$

where the kernel  $D(x, y)$  is a matrix in Dirac and flavour space. For periodic fields eq. (2.15) may be rewritten in the form

$$D\psi(x) = \sum_{y \in \Gamma} D_L(x, y)\psi(y), \quad D_L(x, y) = \sum_{n \in \mathbb{Z}^4} D(x, y + Ln), \quad (2.16)$$

i.e. the finite-volume kernel  $D_L(x, y)$  is obtained from the kernel on the infinite lattice by applying the reflection principle. Evidently, since we are dealing with the same operator, the Ginsparg-Wilson relation holds in finite volume too.

From the properties listed in appendix B it follows that  $D_L(x, y)$  is periodic in  $x$  and  $y$  separately. Moreover it transforms in the same way as  $D(x, y)$  under the gauge and lattice symmetries and from the locality of the operator one infers that

$$D_L(x, y) = D(x, y) + O(e^{-L/\varrho}), \quad (2.17)$$

where  $\varrho$  is the localization range of  $D$ .

## 2.4 Weyl fermions

Using the Ginsparg-Wilson relation, the infinitesimal transformation

$$\delta\psi = \gamma_5(1 - D)\psi, \quad \delta\bar{\psi} = \bar{\psi}\gamma_5, \quad (2.18)$$

is easily shown to be a symmetry of the fermion action

$$S_F = \sum_{x \in \Gamma} \bar{\psi}(x) D\psi(x). \quad (2.19)$$

Eq. (2.18) is the chiral transformation of ref. [18] except that the fermion and the anti-fermion fields are here treated asymmetrically. The reason for this is that the present formulation allows one to decompose the fields into left- and right-handed components in a natural way. First note that the operator  $\hat{\gamma}_5 = \gamma_5(1 - D)$  satisfies

$$(\hat{\gamma}_5)^\dagger = \hat{\gamma}_5, \quad (\hat{\gamma}_5)^2 = 1. \quad (2.20)$$

So if we define the projectors

$$\hat{P}_\pm = \frac{1}{2}(1 \pm \hat{\gamma}_5), \quad P_\pm = \frac{1}{2}(1 \pm \gamma_5), \quad (2.21)$$

it is immediately clear that the left-handed fields  $\hat{P}_-\psi$  and  $\bar{\psi}P_+$  (and the complementary components) transform under lattice chiral rotations in the same way as the corresponding fields in the continuum theory. In particular, left- and right-handed fields decouple in the action (2.19) †.

We now eliminate the right-handed components by imposing the constraints

$$\hat{P}_-\psi = \psi, \quad \bar{\psi}P_+ = \bar{\psi}, \quad (2.22)$$

on the fermion fields. An important point to note here is that these conditions are local and gauge-invariant. The same is true for the action (2.19) and we thus have a completely satisfactory definition of the theory at the classical level.

## 2.5 Fermion integration measure

To set up the quantum theory we also need to specify an integration measure for left-handed fields. The basic difficulty which one has here is that the subspace of

---

† That the action can be split this way has independently been noted by Hasenfratz and Niedermayer [23,24]. A closely related observation has also been made by Narayanan in the context of the overlap formalism [25]



left-handed fermion fields depends on the gauge field. As a consequence there is a non-trivial phase ambiguity in the integration measure.

To make this explicit let us suppose that  $v_j(x)$ ,  $j = 1, 2, 3, \dots$ , is a basis of complex valued, periodic fermion fields such that

$$\hat{P}_- v_j = v_j, \quad (v_k, v_j) = \delta_{kj} \quad (2.23)$$

(the bracket denotes the obvious scalar product for fermion fields in finite volume). The quantum field may then be expanded according to

$$\psi(x) = \sum_j v_j(x) c_j, \quad (2.24)$$

where the coefficients  $c_j$  generate a Grassmann algebra. They represent the independent degrees of freedom of the field and an integration measure for left-handed fermion fields is thus given by

$$D[\psi] = \prod_j dc_j. \quad (2.25)$$

An important mathematical fact which should be kept in mind in the following is that the measure is independent of the particular basis that has been chosen up to a phase factor. One can quickly see this by noting that a change of basis

$$\tilde{v}_j(x) = \sum_l v_l(x) (\mathcal{Q}^{-1})_{lj}, \quad \tilde{c}_j = \sum_l \mathcal{Q}_{jl} c_l, \quad (2.26)$$

implies a change of the measure by the factor  $\det \mathcal{Q}$  which is a pure phase factor since  $\mathcal{Q}$  is unitary. On the other hand, the remark shows that one has a phase ambiguity which is cannot be ignored because the basis (and hence the phase of the measure) depends on the gauge field. One can try to fix the ambiguity in some ad hoc manner, but as will become clear in sect. 4 such prescriptions are likely to be unsatisfactory. For the time being we assume that some particular basis has been chosen and proceed with the definition of the theory.

In the case of the anti-fermion fields the subspace of left-handed fields is independent of the gauge field and one can take the same orthonormal basis  $\bar{v}_k(x)$  for all gauge fields. The ambiguity in the integration measure

$$D[\bar{\psi}] = \prod_k d\bar{c}_k, \quad \bar{\psi}(x) = \sum_k \bar{c}_k \bar{v}_k(x), \quad (2.27)$$

is then only a constant phase factor.

Fermion expectation values of any product  $\mathcal{O}$  of fields are now obtained as usual through the functional integral

$$\langle \mathcal{O} \rangle_{\text{F}} = w[m] \int \text{D}[\psi] \text{D}[\bar{\psi}] \mathcal{O} e^{-S_{\text{F}}}. \quad (2.28)$$

Note that this integral is completely well-defined. The integration variables are the coefficients  $c_j$  and  $\bar{c}_k$  in terms of which the action assumes the form

$$S_{\text{F}} = \sum_{k,j} \bar{c}_k M_{kj} c_j, \quad M_{kj} = \sum_{x \in \Gamma} \bar{v}_k(x) D v_j(x). \quad (2.29)$$

The fermion fields in the product  $\mathcal{O}$  should be expanded similarly and the integral can then be evaluated following the standard rules for Grassmann integration.

In the definition (2.28) a complex factor  $w[m]$  has been included, which allows one to adjust the relative phase and absolute weight of the topological sectors. The factor only depends on the magnetic flux quantum numbers  $m_{\mu\nu}$  and we are free to set  $w[0] = 1$ . One might be tempted to do the same in all other sectors as well, but this may not be the proper choice since the number of integration variables depends on the sector which is being considered (cf. subsect. 3.2).

Full normalized expectation values are finally given by

$$\langle \mathcal{O} \rangle = \frac{1}{\mathcal{Z}} \int \text{D}[U] e^{-S_{\text{G}}} \langle \mathcal{O} \rangle_{\text{F}}, \quad (2.30)$$

where the normalization factor  $\mathcal{Z}$  is defined through the requirement that  $\langle 1 \rangle = 1$  and  $\text{D}[U]$  denotes the usual integration measure for  $\text{U}(1)$  lattice gauge fields.

### 3. Correlation functions and effective action

Apart from the fact that we have not fixed the phase of the fermion integration measure, the theory is completely defined at this point and one can begin to study its properties. In the following paragraphs we work out a few quantities and address some of the basic questions which one may have in order to demonstrate the consistency of the approach.

### 3.1 Fermion propagator

If  $D$  has no zero-modes it is straightforward to show that

$$\langle \psi(x) \bar{\psi}(y) \rangle_{\text{F}} = \langle 1 \rangle_{\text{F}} \times \hat{P}_- S_L(x, y) P_+, \quad (3.1)$$

where the fermion propagator  $S_L(x, y)$  is a periodic function satisfying

$$\sum_{y \in \Gamma} D_L(x, y) S_L(y, z) = \delta_{xz} \quad \text{for all } x, z \in \Gamma. \quad (3.2)$$

In other words,  $S_L(x, y)$  is the kernel of the inverse of the Dirac operator in finite volume. Note that there is no dependence on the bases  $v_j(x)$  and  $\bar{v}_k(x)$  here since the phase ambiguity of the fermion integration measure cancels in eq. (3.1).

From the above and the definition of the chiral projectors it follows that

$$\hat{P}_- S_L(x, y) P_+ = P_- S_L(x, y) P_+ + \frac{1}{2} P_+ \delta_{xy} \quad (3.3)$$

for all points  $x, y$  in  $\Gamma$ . This expression makes it evident that the propagating fermion modes are chiral. The theory thus describes the right number and type of Weyl fermions and there is little doubt that one recovers the correct Feynman rules in the continuum limit for the propagator in an external field.

### 3.2 Fermion number violation

A characteristic feature of chiral gauge theories is that fermion number violating processes can take place. This is possible whenever the numbers of left- and right-handed zero-modes of the Dirac operator,  $n_-$  and  $n_+$ , are not the same.

We can now easily check this in the lattice theory. First note that the dimensions of the spaces of left-handed fermion and anti-fermion fields can be different. Since these spaces are the eigenspaces of the corresponding chiral projectors, the difference of their dimensions is given by †

$$\text{Tr}_L \{ \hat{P}_- \} - \text{Tr}_L \{ P_+ \} = \frac{1}{2} \text{Tr}_L \{ \gamma_5 D \} = n_- - n_+, \quad (3.4)$$

where the second equality follows from the index theorem [16,18]. The index  $n_+ - n_-$  is a topological invariant which assumes a fixed and in general non-zero value in each magnetic flux sector.

---

† Here and below the symbol “ $\text{Tr}_L$ ” implies a trace over the space of fermion fields in finite volume, “ $\text{Tr}$ ” the same in infinite volume and “ $\text{tr}$ ” a trace over Dirac and flavour indices only

In all sectors where the index does not vanish, the matrix  $M_{kj}$  which appears in the action (2.29) has thus a rectangular shape. So if we temporarily choose the basis vectors  $v_j(x)$  and  $\bar{v}_k(x)$  such that the first of them are the zero-modes, the action becomes

$$S_F = \sum_{k>n_+} \sum_{j>n_-} \bar{c}_k M_{kj} c_j, \quad (3.5)$$

which is a non-degenerate quadratic form in the integration variables  $c_j$  and  $\bar{c}_k$  associated with the other modes. The functional integral (2.28) hence vanishes unless  $\mathcal{O}$  is a product of  $n_-$  fermion and  $n_+$  anti-fermion fields times an arbitrary polynomial in pairs of these fields and the gauge field variables. In other words,  $\mathcal{O}$  has to have a net fermion number equal to  $n_- - n_+$  and the lattice theory thus complies with the expected selection rules for fermion number violating processes.

### 3.3 Effective action

In the vacuum sector the dimensions of the spaces of left-handed fermion and anti-fermion fields are the same and the fermion partition function is hence given by

$$\langle 1 \rangle_F = \det M. \quad (3.6)$$

Chiral determinants in the continuum theory are usually studied by computing their variation under infinitesimal deformations of the gauge field [12–14]. We can do the same here and it will soon become clear that this is a useful exercise.

So let us consider a variation

$$\delta_\eta U(x, \mu) = i\eta_\mu(x)U(x, \mu) \quad (3.7)$$

of the gauge field, where  $\eta_\mu(x)$  is any real periodic vector field. After some algebra the associated variation of the effective action is then found to be given by

$$\delta_\eta \ln \det M = \text{Tr}_L \{ \delta_\eta D \hat{P}_- D^{-1} P_+ \} - i\mathfrak{L}_\eta. \quad (3.8)$$

One might have expected to end up with the first term only, but since the basis vectors  $v_j(x)$  depend on the gauge field one has a second term,

$$\mathfrak{L}_\eta = i \sum_j (v_j, \delta_\eta v_j), \quad (3.9)$$

which may be regarded as a contribution of the fermion integration measure.

The current  $j_\mu(x)$  which is defined through

$$\mathfrak{L}_\eta = \sum_{x \in \Gamma} \eta_\mu(x) j_\mu(x) \quad (3.10)$$

is going to play an important rôle in the following. In particular, it will be shown later that the measure can be reconstructed from the current if certain conditions are fulfilled. Note that the measure term transforms according to

$$\tilde{\mathfrak{L}}_\eta = \mathfrak{L}_\eta - i \delta_\eta \ln \det \mathcal{Q} \quad (3.11)$$

under basis transformations (2.26) and  $\mathfrak{L}_\eta$  is hence unchanged if the transformation preserves the integration measure. As a consequence the current should be thought of as a quantity which is associated with the measure rather than the basis vectors  $v_j(x)$ . It is also immediately clear from this that any two measures with the same current are related to each other by a constant phase factor in each topological sector.

#### 3.4 Integrability condition

The significance of the measure term  $\mathfrak{L}_\eta$  may be further elucidated by computing the ‘‘curvature’’  $\delta_\eta \mathfrak{L}_\zeta - \delta_\zeta \mathfrak{L}_\eta$ . Starting from eq. (3.9), this is easily done and in a few lines one obtains

$$\delta_\eta \mathfrak{L}_\zeta - \delta_\zeta \mathfrak{L}_\eta = i \text{Tr}_L \{ \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \}. \quad (3.12)$$

As expected from the transformation law (3.11), the curvature does not depend on the choice of the basis vectors  $v_j(x)$ . In particular, if it is not equal to zero it cannot be made to vanish by adjusting the basis and in these cases the measure term is hence required to ensure the integrability of eq. (3.8).

It is interesting to note in this connection that essentially the same happens in Leutwyler’s construction of the chiral determinant in the continuum theory [12], where a local counterterm has to be added to restore the integrability after applying a finite-part prescription to the variation of the determinant. The analogy will be even more striking after the discussion in the next section, which will lead us to require that the current  $j_\mu(x)$  should be a local expression in the gauge field. The measure term then assumes the form of a local counterterm.

### 3.5 Gauge anomaly

Although the fermion action and the projection to the left-handed fields are gauge-invariant, the effective action tends to be non-invariant due to the anomaly and the fact that the fermion integration measure depends on the gauge field. To work this out, let us consider a gauge variation

$$\eta_\mu(x) = -\partial_\mu \omega(x), \quad (3.13)$$

where  $\omega(x)$  is any periodic gauge function and  $\partial_\mu$  the forward difference operator defined in appendix A. If we introduce the generator

$$T_{\alpha\alpha'} = \delta_{\alpha\alpha'} e_\alpha \quad (3.14)$$

of the fermion representation (2.13) of the gauge group, it is then obvious that

$$\delta_\eta D = i[\omega T, D] \quad (3.15)$$

and taking eq. (3.8) into account one obtains

$$\delta_\eta \ln \det M = i \sum_{x \in \Gamma} \omega(x) \{ \mathcal{A}_L(x) - \partial_\mu^* j_\mu(x) \}, \quad (3.16)$$

$$\mathcal{A}_L(x) = -\frac{1}{2} \text{tr} \{ \gamma_5 T D_L(x, x) \}, \quad (3.17)$$

for the gauge variation of the effective action. Note that  $\mathcal{A}_L(x)$  is equal to the sum of the axial anomalies associated with the  $N$  flavours of fermions in the theory, weighted with their charge [16,18]. In other words,  $\mathcal{A}_L(x)$  is the anomaly of the current which couples to the gauge field.

### 3.6 Vector-like theories

If the charges  $e_\alpha$  come in pairs with opposite sign, the continuum theory is formally equivalent to a vector-like theory, where the gauge field couples to  $\frac{1}{2}N$  Dirac fermions with positive charges. On the lattice we can choose a basis of left-handed fermion fields such that the basis vectors in the sectors with positive and negative charges are related to each other through

$$v_j^-(x) = \gamma_5 C^{-1} [v_j^+(x)]^*, \quad (3.18)$$

where  $C$  denotes the charge conjugation matrix. The associated fermion integration measure is the same for any such basis and it is also easy to show that the measure term  $\mathfrak{L}_\eta$  vanishes.

If the basis  $\bar{v}_k(x)$  of left-handed anti-fermion fields is taken to be of the same type, the partition function (3.6) factorizes and in a few lines one obtains

$$\langle 1 \rangle_{\text{F}} = \left| \det_{e_\alpha > 0} M \right|^2 = \det_{e_\alpha > 0} D, \quad (3.19)$$

which is the expected result for a vector-like theory. Up to contact terms and with an appropriate assignment of field components, there is in fact a complete matching between the chiral and the vector theory in the vacuum sector. Presumably this is also the case in the other sectors, but the issue will not be pursued here.

#### 4. Conditions on the fermion integration measure

According to the universality hypothesis, the details of the lattice theory should not influence the continuum limit, apart from finite renormalizations, as long as a few basic principles are respected. One of them is that the theory should be formulated locally with no long-range couplings in the action. Symmetries are also very important and the universality of the continuum limit is more likely to be guaranteed if they are preserved on the lattice.

The conditions on the fermion integration measure listed below have been devised with this in mind. They should be regarded as a maximal set of requirements which one may reasonably hope to fulfil and one can be quite confident that the correct continuum limit will be obtained if they are all satisfied.

(1) *Differentiability with respect to the gauge field.* The expectation value  $\langle \mathcal{O} \rangle_{\text{F}}$  of arbitrary (finite) products  $\mathcal{O}$  of the fermion fields and the link variables should be smooth functions of the gauge field. This is a somewhat technical requirement, but there are a few instances where the smoothness of the fermion integrals seems to be essential. In particular, the derivation of the field equation discussed below is invalid if this is not guaranteed.

As explained in sect. 8, this condition assumes a simple form in terms of the basis vectors  $v_j(x)$  and in the following we shall say that the fermion integration measure is smooth if the chosen basis has the properties stated there.

(2) *Locality of the field equations.* In euclidean field theory the field equations are linear relations between operator insertions in correlation functions. If the action is local, these operators are local composite fields and the locality properties of the theory are thus directly reflected by the field equations.

This leads us to require that the fermion integration measure should be such that the locality of the field equations is guaranteed. In particular, this should be so for the field equations associated with the gauge field, which one derives from the functional integral (2.30) by calculating the change of the integrand under field variations of the type considered in subsect. 3.3. Relatively little work is required for this if the field product  $\mathcal{O}$  does not involve the fermion fields, because only the sectors with vanishing index contribute in this case and one can then make use of eqs. (3.8) and (3.1) to show that

$$\langle \{\delta_\eta S_G + \sum_{x \in \Gamma} \bar{\psi}(x) \delta_\eta D\psi(x) + i\mathcal{L}_\eta\} \mathcal{O} \rangle = \langle \delta_\eta \mathcal{O} \rangle. \quad (4.1)$$

For local variations  $\eta_\mu(x)$  the first two terms in this equation are manifestly local. To ensure the locality of the field equations we thus require that the current  $j_\mu(x)$  is a local function of the gauge field  $\dagger$ .

If one considers more general field products  $\mathcal{O}$ , the field equations are not quite as easy to derive, but in all cases it turns out that the locality of the current implies the locality of the field equations.

(3) *Gauge invariance.* To preserve the gauge invariance of the theory we require that  $\langle \mathcal{O} \rangle_F$  is a gauge-invariant function of the gauge field if  $\mathcal{O}$  is a gauge-invariant product of the link variables and the fermion fields. In particular, the partition function  $\langle 1 \rangle_F$  should be invariant and from our discussion in subsect. 3.5 it is immediately clear that this condition will be fulfilled if

$$\partial_\mu^* j_\mu(x) = \mathcal{A}_L(x). \quad (4.2)$$

It is possible to prove that no further conditions arise when one considers arbitrary products  $\mathcal{O}$  of fields, i.e. the gauge invariance of the theory is guaranteed if eq. (4.2) holds. One of the consequences of this equation and the integrability condition (3.12) is, incidentally, that the current  $j_\mu(x)$  itself has to be gauge-invariant.

---

$\dagger$  The notion of locality used in this paper is the same as in refs. [21,24,26]. Details are given in appendix B for the case of the Dirac operator. Note that the term only makes sense if the lattice size  $L$  is much larger than the localization range of the fields that one is interested in



(4) *Lattice symmetries.* In the continuum limit the imaginary part of the effective action transforms in a particular way under the space-time symmetries. On the lattice one would like to preserve these symmetries as far as possible so as to reduce any remaining phase ambiguity in the fermion integration measure.

Lattice translations, hyper-cubic rotations, reflections at the lattice planes and charge conjugation will be referred to as the “lattice symmetries” in the following. We now demand that the measure term  $\mathfrak{L}_\eta$  transforms in the same way under these symmetries as the imaginary part of the first term on the right-hand side of eq. (3.8). This is equivalent to requiring the current  $j_\mu(x)$  to transform like the axial current

$$j_\mu^5(x) = \frac{1}{2} \{ \bar{\psi}(x) \gamma_5 \gamma_\mu U(x, \mu) \psi(x + \hat{\mu}) + \bar{\psi}(x + \hat{\mu}) \gamma_5 \gamma_\mu U(x, \mu)^{-1} \psi(x) \} \quad (4.3)$$

in ordinary lattice gauge theories with Wilson-Dirac fermions.

## 5. Statement of results

In the remainder of this paper we shall show that fermion integration measures satisfying conditions (1)–(4) exist in all topological sectors provided

$$N_e \text{ is even for all odd } e, \quad (5.1)$$

where  $N_e$  denotes the number of fermion flavours with  $|e_\alpha| = e$ . This includes the multiplet (2.14) and all cases with only even charges. In the vacuum sector there is actually no restriction on the charge assignment apart from the anomaly cancellation condition and it is currently not clear whether the constraint (5.1) reflects a fundamental limitation in finite volume or just a temporary technical difficulty.

For clarity the main steps of the construction are presented below in the form of three theorems together with some key formulae. All proofs are postponed to the later sections which should be consulted for full details.

### 5.1 Reconstruction theorem

While the fermion integration measure is a relatively complicated object, requiring the specification of a basis  $v_j(x)$  of left-handed fields modulo measure preserving basis transformations, the associated current  $j_\mu(x)$  is invariant under such transformations and is clearly much more tractable. The following theorem says that the measure can be reconstructed from the current under certain conditions.

**Theorem 5.1.** *Suppose  $j_\mu(x)$  is a given current with the following properties.*

(a)  $j_\mu(x)$  is defined for all admissible gauge fields and depends smoothly on the link variables.

(b)  $j_\mu(x)$  is gauge-invariant and transforms as an axial vector current under the lattice symmetries, as described in sect. 4.

(c) The linear functional  $\mathfrak{L}_\eta = \sum_{x \in \Gamma} \eta_\mu(x) j_\mu(x)$  satisfies the integrability condition (3.12).

(d) The anomalous conservation law  $\partial_\mu^* j_\mu(x) = \mathcal{A}_L(x)$  holds.

Then there exists a smooth fermion integration measure in the vacuum sector such that the associated current coincides with  $j_\mu(x)$ . The same is true in all other sectors if the charges satisfy the constraint (5.1). In each case the measure is uniquely determined up to a constant phase factor.

We are thus left with the problem to find a local current  $j_\mu(x)$  with the properties listed above. Since the notion of locality which is being adopted here allows for exponentially decaying tails (with a fixed localization range in lattice units), the current can have non-local contributions that are of this order in the lattice size  $L$ . In the following our strategy will be to provide an explicit expression for the current in infinite volume and to prove that a solution in finite volume can be obtained by adding an exponentially small correction.

## 5.2 Anomaly cancellation

Before proceeding with the construction of the current it is however useful to discuss the significance of the anomaly cancellation condition (1.1) in the present framework. For simplicity we consider the theory in infinite volume in this subsection. The properties of the Dirac operator listed in appendix B then imply that the anomaly

$$\mathcal{A}(x) = -\frac{1}{2} \text{tr} \{ \gamma_5 T D(x, x) \} \quad (5.2)$$

is a gauge-invariant local field. Moreover, using the Ginsparg-Wilson relation, the anomaly can be shown to be a topological field satisfying

$$\sum_{x \in \mathbb{Z}^4} \delta_\eta \mathcal{A}(x) = 0 \quad (5.3)$$

for any local deformation  $\eta_\mu(x)$  of the gauge field. It follows from this and a general

theorem established in ref. [26] that

$$\mathcal{A}(x) = \gamma \epsilon_{\mu\nu\rho\sigma} F_{\mu\nu}(x) F_{\rho\sigma}(x + \hat{\mu} + \hat{\nu}) + \partial_\mu^* k_\mu(x), \quad (5.4)$$

where  $\gamma$  is a constant and  $k_\mu(x)$  a gauge-invariant local current.

We now show that  $\gamma = 0$  by noting that the Dirac operator is equal to the same analytic expression for each fermion flavour  $\alpha$ , with the link variables  $U(x, \mu)$  replaced by  $U(x, \mu)^{e_\alpha}$ . The field tensor scales with the charge and there is another power of the charge coming from the generator  $T$  in eq. (5.2). The contribution to the constant  $\gamma$  of the fermion with flavour  $\alpha$  is hence proportional to  $e_\alpha^3$  and after summing over all flavours one gets zero because of eq. (1.1).

The anomaly thus cancels up to a divergence term. At first sight one might think that this is not enough to achieve the gauge invariance of the theory, but we only need to satisfy eq. (4.2) for this and it is then conceivable that the gauge field dependence of the measure exactly compensates for the divergence term. The important point to note here is that one would be unable to cancel the term proportional to  $\gamma$  in this way. The construction of a fermion integration measure complying with conditions (1)–(4) is hence only possible for anomaly-free fermion multiplets.

### 5.3 Solution of the integrability condition in infinite volume

One of the technical advantages which one has in infinite volume is that the gauge fields can be represented in a natural way through vector fields. The relevant lemma has been proved in ref. [26] and is quoted here for the reader's convenience.

**Lemma 5.2.** *Suppose  $U(x, \mu)$  is an admissible gauge field on the infinite lattice. Then there exists a vector field  $A_\mu(x)$  such that*

$$U(x, \mu) = e^{iA_\mu(x)}, \quad |A_\mu(x)| \leq \pi(1 + 8\|x\|), \quad (5.5)$$

$$F_{\mu\nu}(x) = \partial_\mu A_\nu(x) - \partial_\nu A_\mu(x). \quad (5.6)$$

Moreover, any other field with these properties is equal to  $A_\mu(x) + \partial_\mu \omega(x)$ , where the gauge function  $\omega(x)$  takes values that are integer multiples of  $2\pi$ .

The idea is now to construct a solution of the integrability condition first in terms of the vector field. So let us assume that  $A_\mu(x)$  is any given field representing an admissible gauge field  $U(x, \mu)$  as in lemma 5.2. The curve

$$U_t(x, \mu) = e^{itA_\mu(x)}, \quad 0 \leq t \leq 1, \quad (5.7)$$

contracts this field to the classical vacuum configuration in such a way that the field tensor remains bounded by  $\epsilon$  for all  $t$ . For any variation  $\eta_\mu(x)$  of the gauge potential with compact support, a linear functional  $\mathfrak{L}_\eta^*$  may thus be defined through

$$\begin{aligned} \mathfrak{L}_\eta^* = & i \int_0^1 dt \operatorname{Tr} \{ \hat{P}_- [\partial_t \hat{P}_-, \delta_\eta \hat{P}_-] \} + \\ & \int_0^1 dt \sum_{x \in \mathbb{Z}^4} \{ \eta_\mu(x) \bar{k}_\mu(x) + A_\mu(x) \delta_\eta \bar{k}_\mu(x) \}, \end{aligned} \quad (5.8)$$

where  $\bar{k}_\mu(x)$  is any gauge-invariant local current, which transforms as an axial vector field under the lattice symmetries and which satisfies  $\partial_\mu^* \bar{k}_\mu(x) = \mathcal{A}(x)$ . An example of such a field is obtained by averaging the current  $k_\mu(x)$  introduced in subsect. 5.2 over the lattice symmetries, with the appropriate weights so as to project to the axial vector component. Note that an explicit although very complicated expression for  $k_\mu(x)$  in terms of the first and second variations of the anomaly has been derived in ref. [26]. The existence of a current  $\bar{k}_\mu(x)$  with the required properties is thus guaranteed.

**Theorem 5.3.** *The linear functional  $\mathfrak{L}_\eta^* = \sum_{x \in \mathbb{Z}^4} \eta_\mu(x) j_\mu^*(x)$  defined above has the following properties.*

- (a)  $\mathfrak{L}_\eta^*$  is invariant under gauge transformations  $A_\mu(x) \rightarrow A_\mu(x) + \partial_\mu \omega(x)$ , for arbitrary gauge functions  $\omega(x)$  that are polynomially bounded at infinity.
- (b) The current  $j_\mu^*(x)$  is a local field, which depends smoothly on the gauge field and which transforms as an axial vector current under the lattice symmetries.
- (c)  $\mathfrak{L}_\eta^*$  is a solution of the integrability condition

$$\delta_\eta \mathfrak{L}_\zeta^* - \delta_\zeta \mathfrak{L}_\eta^* = i \operatorname{Tr} \{ \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \} \quad (5.9)$$

in infinite volume for all compactly supported variations  $\eta_\mu(x)$  and  $\zeta_\mu(x)$ .

- (d) The anomalous conservation law  $\partial_\mu^* j_\mu^*(x) = \mathcal{A}(x)$  holds.

An important consequence of the gauge invariance of  $\mathfrak{L}_\eta^*$  is that the current  $j_\mu^*(x)$  may be considered to be a function of the gauge field  $U(x, \mu)$  rather than the vector field  $A_\mu(x)$ , since the mapping between the two is one-to-one modulo gauge transformations. It can be shown that the locality, differentiability and symmetry properties of the current are the same independently of which point of view is adopted [26].

#### 5.4 Construction of the current $j_\mu(x)$ in finite volume

We now return to the theory in finite volume and first note that  $j_\mu^*(x)$  becomes a gauge-invariant local field on the finite lattice if attention is restricted to periodic gauge fields. As asserted by the following theorem, this current has all the required properties up to exponentially small finite-lattice corrections.

**Theorem 5.4.** *If the lattice is sufficiently large compared to the localization range  $\varrho$  of the Dirac operator, say  $L/\varrho \geq n$ , there exists a current  $j_\mu(x)$  which satisfies*

$$|j_\mu(x) - j_\mu^*(x)| \leq \kappa L^\nu e^{-L/\varrho} \quad (5.10)$$

and which fulfils conditions (a)–(d) of theorem 5.1. The bound (5.10) holds uniformly in the gauge field, i.e. the constants  $n$ ,  $\kappa$  and  $\nu$  are independent of the field.

Together with theorem 5.1 this result implies that fermion integration measures satisfying conditions (1)–(4) exist on large lattices. Note that the difference between  $j_\mu(x)$  and  $j_\mu^*(x)$  vanishes exponentially in the continuum limit, because  $\varrho$  is a fixed number in lattice units while  $L$  is a physical length scale. The detailed form of these corrections is hence of little interest.

The theorems quoted in this section are not easy to prove. Most of the difficulties can be traced back to the fact that the space of admissible gauge fields is topologically non-trivial in finite volume. Differential geometry and the theory of fibre bundles are the adequate tools to deal with this problem and the reader who wishes to go through the details in sects. 7–11 will be assumed to be familiar with the relevant mathematical terminology.

## 6. Proof of theorem 5.3

We first remark that the projector  $\hat{P}_-$  has the same locality properties as the Dirac operator. In particular, the kernel of  $\delta_\eta \hat{P}_-$  falls off exponentially away from the support of  $\eta_\mu(x)$  and the trace in eq. (5.8) is hence rapidly convergent in position space. One of the consequences of this technical observation is that  $\mathfrak{L}_\eta^*$  is a well-defined and smooth function of the gauge potential  $A_\mu(x)$ . We now establish the other properties of  $\mathfrak{L}_\eta^*$  in the order stated in the theorem.

(a) *Gauge invariance.* Taking the gauge covariance of the projector  $\hat{P}_-$  and the gauge invariance of the current  $\bar{k}_\mu(x)$  into account, it is easy to show that the change of

$\mathfrak{L}_\eta^*$  under gauge transformations  $A_\mu(x) \rightarrow A_\mu(x) + \partial_\mu\omega(x)$  is given by

$$\int_0^1 dt \operatorname{Tr}\{\hat{P}_- [[\omega T, \hat{P}_-], \delta_\eta \hat{P}_-]\} + \int_0^1 dt \sum_{x \in \mathbb{Z}^4} \partial_\mu \omega(x) \delta_\eta \bar{k}_\mu(x). \quad (6.1)$$

Expanding the commutators and using the identity

$$\hat{P}_- \delta_\eta \hat{P}_- \hat{P}_- = 0, \quad (6.2)$$

the first term can be rewritten in the form

$$\int_0^1 dt \sum_{x \in \mathbb{Z}^4} \omega(x) \delta_\eta \mathcal{A}(x), \quad (6.3)$$

where  $\mathcal{A}(x)$  denotes the anomaly in infinite volume. Recalling  $\partial_\mu^* \bar{k}_\mu(x) = \mathcal{A}(x)$  and performing a partial summation it is now clear that the two terms in eq. (6.1) cancel each other.

(b) *Locality and symmetry properties of  $j_\mu^*(x)$ .* From what has been said at the beginning of this section, and since  $\bar{k}_\mu(x)$  is a smooth local function of the gauge field, it is evident that the same is true for  $j_\mu^*(x)$ . Moreover under the lattice symmetries it transforms as an axial vector field. To prove this for space-time reflections one has to take into account that

$$\operatorname{Tr}\{\hat{P}_+ [\partial_t \hat{P}_+, \delta_\eta \hat{P}_+]\} = -\operatorname{Tr}\{\hat{P}_- [\partial_t \hat{P}_-, \delta_\eta \hat{P}_-]\}. \quad (6.4)$$

In all other cases the transformations commute with the projection to the left-handed fields and the covariance of the current is deduced straightforwardly.

(c) *Integrability condition.* Starting from the definition (5.8) of  $\mathfrak{L}_\eta^*$ , one quickly finds that the second term does not contribute to the curvature  $\delta_\eta \mathfrak{L}_\zeta^* - \delta_\zeta \mathfrak{L}_\eta^*$ . Another simple observation is that all terms of the form

$$\operatorname{Tr}\{\delta_1 \hat{P}_- \delta_2 \hat{P}_- \delta_3 \hat{P}_-\} \quad (6.5)$$

can be shown to vanish by inserting  $1 = (\hat{\gamma}_5)^2$  and using the fact that  $\hat{\gamma}_5$  anti-commutes with any variation of the projector  $\hat{P}_-$ . Taking this into account, there are only two terms which contribute to the curvature,

$$i \int_0^1 dt \operatorname{Tr}\{\hat{P}_- [\delta_\eta \partial_t \hat{P}_-, \delta_\zeta \hat{P}_-] - \hat{P}_- [\delta_\zeta \partial_t \hat{P}_-, \delta_\eta \hat{P}_-]\}, \quad (6.6)$$

and these may be rewritten in the form

$$i \int_0^1 dt \partial_t \text{Tr} \{ \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \}. \quad (6.7)$$

After integration one then ends up with eq. (5.9) since the contribution from the lower end of the integration range is equal to zero.

(d) *Anomalous conservation law.* Setting  $\eta_\mu(x) = -\partial_\mu \omega(x)$  (where  $\omega(x)$  is any lattice function with compact support) the left-hand side of eq. (5.8) becomes

$$\sum_{x \in \mathbb{Z}^4} \omega(x) \partial_\mu^* j_\mu^*(x). \quad (6.8)$$

On the other side we insert the identities

$$\delta_\eta \hat{P}_- = it [\omega T, \hat{P}_-], \quad \delta_\eta \bar{k}_\mu(x) = 0, \quad (6.9)$$

and in a few steps obtain a sum of two terms,

$$- \int_0^1 dt t \text{Tr} \{ \omega T \partial_t \hat{P}_- \} + \int_0^1 dt \sum_{x \in \mathbb{Z}^4} \omega(x) \mathcal{A}(x). \quad (6.10)$$

Expressing the trace in the first term through the anomaly (5.2), the terms nearly cancel after a partial integration and one is left with the contribution

$$\sum_{x \in \mathbb{Z}^4} \omega(x) \mathcal{A}(x)|_{t=1} \quad (6.11)$$

from the upper end of the integration range. Comparing with eq. (6.8) this shows that the divergence of  $j_\mu^*(x)$  is equal to the anomaly and thus completes the proof of the theorem.

## 7. Topology of the field space in finite volume

We now begin with the detailed discussion of the theory in finite volume and first determine the structure of the space of admissible gauge fields. As will be explained in the next section, the existence of smooth fermion integration measures depends on whether a certain  $U(1)$  bundle over this space is trivial. Evidently, to be able to address this problem, one needs to know the topology of the base manifold.

### 7.1 Preliminaries

In the following the lattice is assumed to be finite with periodic boundary conditions as specified in sect. 2. The space of admissible gauge fields is denoted by  $\mathfrak{U}$  and the gauge group  $\mathfrak{G}_0$  is taken to be the subset of gauge transformations  $\Lambda(x)$  satisfying  $\Lambda(x) = 1$  at  $x = 0 \bmod L$ .

For any given gauge field  $U(x, \mu)$ , the Wilson lines winding around the lattice along the coordinate axes are defined by

$$W_\mu(x) = \prod_{s=0}^{L-1} U(x + s\hat{\mu}, \mu). \quad (7.1)$$

They are gauge-invariant, but cannot be expressed through the field tensor  $F_{\mu\nu}(x)$  and thus carry independent information on the gauge field.

**Lemma 7.1.** *Any two admissible fields  $U(x, \mu)$  and  $\tilde{U}(x, \mu)$  satisfying*

$$F_{\mu\nu}(x) = \tilde{F}_{\mu\nu}(x) \quad \text{and} \quad W_\mu(x) = \tilde{W}_\mu(x) \quad (7.2)$$

*are gauge equivalent.*

*Proof:* If we introduce a new field through

$$V(x, \mu) = U(x, \mu)\tilde{U}(x, \mu)^{-1}, \quad (7.3)$$

it is obvious that the associated plaquette loops and Wilson lines are all equal to 1. The product  $\Lambda(x)$  of the link variables  $V(x, \mu)$  along any lattice path from  $x$  to the origin  $x = 0$  is hence independent of the chosen path and periodic in  $x$ . In other words,  $\Lambda(x)$  is an element of the gauge group  $\mathfrak{G}_0$  which transforms  $V(x, \mu)$  to 1 and thus  $U(x, \mu)$  to  $\tilde{U}(x, \mu)$ .  $\square$

The subspace  $\mathfrak{U}_0$  of all admissible gauge fields with vanishing field tensor contains the pure gauge configurations, but there are also non-trivial configurations with



Wilson lines different from 1. It is straightforward to show, however, that the Wilson lines  $W_\mu(x)$  do not depend on  $x$ . The gauge-invariant content of such fields is hence encoded in the constant phase factors  $w_\mu = W_\mu(x)$ .

**Lemma 7.2.** *The gauge fields with vanishing field tensor are of the form*

$$U(x, \mu) = \Lambda(x)U_{[w]}(x, \mu)\Lambda(x + \hat{\mu})^{-1}, \quad (7.4)$$

where  $\Lambda(x)$  is an element of  $\mathfrak{G}_0$  and the field  $U_{[w]}(x, \mu)$  is defined by

$$U_{[w]}(x, \mu) = \begin{cases} w_\mu & \text{if } x_\mu = 0 \pmod L, \\ 1 & \text{otherwise,} \end{cases} \quad (7.5)$$

for any given set of phase factors  $w_\mu \in U(1)$ . Moreover the representation (7.4) is unique and establishes the isomorphism  $\mathfrak{U}_0 \cong U(1)^4 \times \mathfrak{G}_0$ .

*Proof:* From the definition (7.5) it is obvious that  $U_{[w]}(x, \mu)$  is a gauge field with vanishing field tensor and Wilson lines  $W_\mu(x)$  equal to  $w_\mu$ . According to lemma 7.1, any other field  $U(x, \mu)$  with these properties is gauge equivalent to  $U_{[w]}(x, \mu)$ . This proves eq. (7.4) and it is now also evident that  $w_\mu$  and  $\Lambda(x)$  are uniquely determined by the gauge field.  $\square$

## 7.2 Flux sectors

As already mentioned in subsect. 2.2, the field space  $\mathfrak{U}$  is a union of disconnected subspaces  $\mathfrak{U}[m]$  labelled by the magnetic flux quantum numbers  $m_{\mu\nu}$ . We now prove this and provide some further information on the flux sectors.

**Lemma 7.3.** *Let  $U(x, \mu)$  be an admissible gauge field and define the associated magnetic flux  $\phi_{\mu\nu}(x)$  through eq. (2.10). Then there exists an anti-symmetric integer tensor  $m_{\mu\nu}$  such that  $\phi_{\mu\nu}(x) = 2\pi m_{\mu\nu}$  for all  $x$ .*

*Proof:* If we define a vector potential  $a_\mu(x)$  through

$$a_\mu(x) = \frac{1}{i} \ln U(x, \mu), \quad -\pi < a_\mu(x) \leq \pi, \quad (7.6)$$

it is straightforward to show that

$$F_{\mu\nu}(x) = \partial_\mu a_\nu(x) - \partial_\nu a_\mu(x) + 2\pi z_{\mu\nu}(x), \quad (7.7)$$

where  $z_{\mu\nu}(x)$  takes integer values. Only the second term in this equation contributes to the magnetic flux which is hence an integer multiple of  $2\pi$ .

The periodicity of the field tensor implies that  $\phi_{\mu\nu}(x)$  is independent of the coordinates  $x_\mu$  and  $x_\nu$ . To prove that the flux is also independent of the complementary components of  $x$ , we note that

$$\epsilon_{\mu\nu\rho\sigma}\partial_\nu F_{\rho\sigma}(x) = 0. \quad (7.8)$$

This is a straightforward consequence of lemma 5.2 and particularly of eq. (5.6). Using the periodicity of the field tensor again and partial summations, the change  $\partial_\rho\phi_{\mu\nu}(x)$  of the flux in any direction orthogonal to the  $(\mu, \nu)$ -plane is then easily shown to vanish.  $\square$

As long as only admissible fields are considered, the field tensor is a continuous function of the link variables and the magnetic flux quantum numbers consequently cannot change under continuous deformations of the field. The sectors  $\mathfrak{U}[m]$  of all fields with a given set of flux quantum numbers  $m_{\mu\nu}$  are thus disconnected from each other. There are at most a finite number of sectors since

$$|m_{\mu\nu}| < \frac{\epsilon}{2\pi}L^2 \quad (7.9)$$

as one may easily prove by combining eqs. (2.9) and (2.10). Conversely if  $m_{\mu\nu}$  is any prescribed, anti-symmetric integer tensor satisfying this bound, there exist admissible fields with these flux quantum numbers. An example of such a field is

$$V_{[m]}(x, \mu) = \exp \left\{ -\frac{2\pi i}{L^2} \left[ L\delta_{\tilde{x}_\mu, L-1} \sum_{\nu>\mu} m_{\mu\nu}\tilde{x}_\nu + \sum_{\nu<\mu} m_{\mu\nu}\tilde{x}_\nu \right] \right\}, \quad (7.10)$$

where the abbreviation  $\tilde{x}_\mu = x_\mu \bmod L$  has been used. This field is periodic and can be shown to have constant field tensor equal to  $2\pi m_{\mu\nu}/L^2$ .

### 7.3 Topology of $\mathfrak{U}[m]$

We now determine the structure of the flux sector  $\mathfrak{U}[m]$  for any given set of flux quantum numbers  $m_{\mu\nu}$ . As will be shown below, one of the factors of this manifold consists of the space  $\mathfrak{A}[m]$  of all periodic vector potentials  $A_\mu^T(x)$  satisfying

$$\partial_\mu^* A_\mu^T(x) = 0, \quad \sum_{x \in \Gamma} A_\mu^T(x) = 0, \quad (7.11)$$

$$|\partial_\mu A_\nu^T(x) - \partial_\nu A_\mu^T(x) + 2\pi m_{\mu\nu}/L^2| < \epsilon. \quad (7.12)$$

The index “T” reminds us that these fields are transverse and also serves to distinguish them from the vector potential  $A_\mu(x)$  which has been introduced in sect. 5. Note that  $\mathfrak{A}[m]$  is a convex space. In particular, it is contractible and thus topologically trivial.

**Lemma 7.4.** *The fields  $U(x, \mu)$  in the sector  $\mathfrak{A}[m]$  are of the form*

$$U(x, \mu) = V_{[m]}(x, \mu) \mathring{U}(x, \mu) e^{iA_\mu^T(x)}, \quad (7.13)$$

where  $\mathring{U}(x, \mu)$  has vanishing field tensor and  $A_\mu^T(x)$  is an element of  $\mathfrak{A}[m]$ . Moreover this representation is unique and establishes the isomorphism  $\mathfrak{A}[m] \cong \mathfrak{A}_0 \times \mathfrak{A}[m]$ .

*Proof:* We first prove the uniqueness of the representation (7.13) by noting that the field tensor of  $U(x, \mu)$  is given by

$$F_{\mu\nu}(x) = \partial_\mu A_\nu^T(x) - \partial_\nu A_\mu^T(x) + 2\pi m_{\mu\nu}/L^2. \quad (7.14)$$

Together with the constraints (7.11) this equation implies that

$$A_\mu^T(x) = \sum_{y \in \Gamma} G_L(x-y) \partial_\lambda^* F_{\lambda\mu}(y), \quad (7.15)$$

where  $G_L(z)$  denotes the Green function of the lattice laplacian,

$$\partial_\mu^* \partial_\mu G_L(z) = \delta_{z0} - L^{-4}, \quad G_L(z + L\hat{\mu}) = G_L(z), \quad \sum_{z \in \Gamma} G_L(z) = 0. \quad (7.16)$$

In particular, the transverse field is uniquely determined and so are the other factors in eq. (7.13).

To show that any given admissible field  $U(x, \mu)$  with field tensor  $F_{\mu\nu}(x)$  and flux quantum numbers  $m_{\mu\nu}$  can be represented in this way, we turn the argument around and define  $A_\mu^T(x)$  through eq. (7.15). From the properties of the Green function it is then clear that this field satisfies eq. (7.11). Moreover, using eq. (7.8) (which holds for any admissible field) and the fact that the zero-momentum component of  $F_{\mu\nu}(x)$  is proportional to  $m_{\mu\nu}$ , it is straightforward to establish eq. (7.14). In particular,  $A_\mu^T(x)$  is contained in  $\mathfrak{A}[m]$  and

$$\mathring{U}(x, \mu) = V_{[m]}(x, \mu)^{-1} U(x, \mu) e^{-iA_\mu^T(x)} \quad (7.17)$$

has vanishing field tensor.  $\square$

Taken together the results obtained in this section imply that

$$\mathfrak{U}[m] \cong \mathrm{U}(1)^4 \times \mathfrak{G}_0 \times \mathfrak{A}[m]. \quad (7.18)$$

Since  $\mathfrak{G}_0$  is a product of  $\mathrm{U}(1)$  factors, the sectors  $\mathfrak{U}[m]$  are thus either empty or equal to a multi-dimensional torus times a contractible space.

## 8. Fermion integration measures and $\mathrm{U}(1)$ bundles over $\mathfrak{U}$

One of the conditions on the fermion integration measure listed in sect. 4 is that the fermion expectation values  $\langle \mathcal{O} \rangle_{\mathrm{F}}$  should be smooth functions of the gauge field. In this section the implications of this requirement for the basis vectors  $v_j(x)$  are worked out and we shall then be able to reformulate the condition in geometrical terms which will later allow us to make use of some known results of the theory of fibre bundles.

### 8.1 Smooth measures

Evidently the differentiability of the fermion integrals  $\langle \mathcal{O} \rangle_{\mathrm{F}}$  will be guaranteed if the basis vectors  $v_j(x)$  are globally defined and smooth, but since the space of admissible gauge fields is topologically non-trivial one may be unable to find such a basis.

We can, however, cover the space of admissible fields with open contractible patches  $X_a$ , labelled by an index  $a$ , and choose a smooth basis  $v_j^a(x)$  on each of these patches. Since the projector to the left-handed fields depends smoothly on the gauge field and since the field manifold  $\mathfrak{U}$  is smooth, this is always possible. On the intersection  $X_a \cap X_b$  of any two patches, we then have two bases which are related to each other by a unitary transformation

$$v_j^b(x) = \sum_k v_k^a(x) \tau(a \rightarrow b)_{kj} \quad (8.1)$$

as in eq. (2.26). The transition matrices  $\tau(a \rightarrow b)$  satisfy the cocycle condition

$$\tau(a \rightarrow c) = \tau(a \rightarrow b)\tau(b \rightarrow c) \quad \text{on} \quad X_a \cap X_b \cap X_c \quad (8.2)$$

and thus define a unitary principal bundle over  $\mathfrak{U}$ .

The fermion integration measure changes by a phase factor equal to the determinant of the transition matrix if one passes from one basis to another. We now only need to make sure that

$$\det \tau(a \rightarrow b) = 1 \tag{8.3}$$

on the intersections  $X_a \cap X_b$ . The integration measure is then independent of the patch and the fermion integrals  $\langle \mathcal{O} \rangle_{\mathbb{F}}$  are thus globally defined and smooth.

If  $v_j^a(x)$  is any patched basis, not necessarily satisfying eq. (8.3), it is clear from the above that the phase factors

$$g_{ab} = \det \tau(a \rightarrow b) \tag{8.4}$$

define a  $U(1)$  bundle over  $\mathfrak{U}$ . Moreover, under a change of basis, these phase factors transform according to

$$g_{ab} \rightarrow h_a g_{ab} h_b^{-1} \quad \text{on } X_a \cap X_b, \tag{8.5}$$

where  $h_a$  is the determinant of the transformation matrix on patch  $X_a$ . Different choices of the basis vectors thus give rise to isomorphic bundles. The converse is also true since for any set of smooth phase factors  $h_a$  one can always find a corresponding basis transformation.

It should now be evident that smooth fermion integration measures exist if (and only if) this bundle is trivial. Whether this is the case depends on the properties of the projector to the left-handed fields and the base manifold  $\mathfrak{U}$ . If the bundle is non-trivial one has an anomaly and it is then not possible to construct fermion integration measures satisfying condition (1) †.

### 8.2 Geometrical interpretation of the measure term $\mathfrak{L}_\eta$

For any given basis  $v_j^a(x)$  the measure term is defined through

$$\mathfrak{L}_\eta^a = i \sum_j (v_j^a, \delta_\eta v_j^a) \tag{8.6}$$

on patch  $X_a$ . The transition rule for passing from one patch to another is

$$\mathfrak{L}_\eta^a = \mathfrak{L}_\eta^b - i g_{ab}^{-1} \delta_\eta g_{ab} \quad \text{on } X_a \cap X_b \tag{8.7}$$

---

† The discussion in this section is closely related to the recent work of Neuberger [27] on the gauge anomaly in the overlap formalism. An explicit example is given there demonstrating the presence of a non-integrable phase if the fermion multiplet is not anomaly-free

and  $\mathfrak{L}_\eta^a$  is thus a connection on the  $U(1)$  bundle constructed above. Note that  $\mathfrak{L}_\eta^a$  is independent of the patch if the smoothness condition (8.3) is fulfilled. The associated current  $j_\mu(x)$  is then a globally defined smooth function of the gauge field.

As we have previously remarked, the curvature (3.12) of the measure term is invariant under basis transformations. Evidently the curvature is just the field strength of the connection  $\mathfrak{L}_\eta^a$ . The Wilson lines constructed from  $\mathfrak{L}_\eta^a$  winding around the base manifold  $\mathfrak{U}$  in a particular direction are further invariants that carry important information on the measure term.

To make this completely clear let us consider a closed curve

$$U_t(x, \mu), \quad 0 \leq t \leq 2\pi, \quad (8.8)$$

of admissible fields. If we temporarily assume that the smoothness condition (8.3) is satisfied, the Wilson line associated with the curve is given by

$$\mathfrak{W} = \exp \left\{ i \int_0^{2\pi} dt \mathfrak{L}_\eta \right\}, \quad \eta_\mu(x) = -i U_t(x, \mu)^{-1} \partial_t U_t(x, \mu). \quad (8.9)$$

The patch label has been dropped here, because the measure term does not depend on it if (8.3) holds. Note that the variation  $\eta_\mu(x)$  may be regarded as the tangential vector along the curve. In particular,

$$\mathfrak{L}_\eta = i \sum_j (v_j^a, \partial_t v_j^a), \quad (8.10)$$

and it is then easy to check that the Wilson line does not depend on the choice of basis. As shown by the following lemma,  $\mathfrak{W}$  may in fact be expressed directly through the projector to the left-handed fields.

**Lemma 8.1.** *The Wilson line defined above is given by*

$$\mathfrak{W} = \lim_{n \rightarrow \infty} \det \{ 1 - P_{t_0} + P_{t_n} P_{t_{n-1}} \dots P_{t_0} \}, \quad t_k = 2\pi k/n, \quad (8.11)$$

where  $P_t$  is equal to the projector  $\hat{P}_-$  along the curve (8.8).

*Proof:* Since we are considering a closed curve, we have  $P_{t_n} = P_{t_0}$  and the determinant on the right-hand side of eq. (8.11) is hence equal to the determinant of the product  $P_{t_n} P_{t_{n-1}} \dots P_{t_0}$  in the subspace of left-handed fields at  $t = 0$ . To compute the determinant, we insert the representation

$$P_{t,L}(x, y) = \sum_j v_j^a(x) \otimes v_j^a(y)^* \quad (8.12)$$

for the kernels of the projectors  $P_t$  in position space. One then obtains a product of matrices with entries that are the scalar products of the basis vectors at subsequent values of  $t$ . At large  $n$  these matrices may be expanded according to

$$\delta_{lj} - (2\pi/n)(v_i^a, \partial_t v_j^a)_{t=t_k} + O(1/n^2) \quad (8.13)$$

and for the determinant the expression

$$\exp\left\{i(2\pi/n) \sum_{k=0}^{n-1} (\mathfrak{L}_\eta)_{t=t_k} + O(1/n)\right\} \quad (8.14)$$

is thus obtained, which converges to  $\mathfrak{W}$  in the limit  $n \rightarrow \infty$ .  $\square$

## 9. Abelian gauge fields on the $n$ -dimensional torus

In each topological sector the submanifold of admissible fields where the transverse field  $A_\mu^T(x)$  vanishes is an  $n$ -dimensional torus

$$T^n = U(1) \times U(1) \times \dots \times U(1) \quad (n \text{ factors}). \quad (9.1)$$

Since  $\mathfrak{A}[m]$  is contractible, a well-known theorem on fibre bundles may be invoked which says that the  $U(1)$  bundle constructed in the preceding section is trivial if its restriction to  $T^n$  is trivial.  $U(1)$  bundles over  $T^n$  can be completely classified and it is now helpful to discuss this and the gauge fields that live on them in some detail. The results quoted below are generally valid and do not refer to any particular properties of the bundle other than those specified in the following paragraphs. We shall then return to the case of interest in sects. 10 and 11.

### 9.1 $U(1)$ bundles over $T^n$

The points  $u$  of the torus (9.1) may be locally parametrized through

$$u = (e^{it_1}, \dots, e^{it_n}), \quad (9.2)$$

where the coordinates  $(t_1, \dots, t_n)$  range in some small contractible region in  $\mathbb{R}^n$ . A particular choice of this region defines a coordinate patch  $X_a$  on  $T^n$  and the set of all these patches provides an atlas for this manifold.

U(1) bundles over  $T^n$  may be defined by specifying a set of smooth transition functions  $g_{ab} \in \text{U}(1)$  on  $X_a \cap X_b$  such that the cocycle condition

$$g_{ac} = g_{ab}g_{bc} \quad \text{on} \quad X_a \cap X_b \cap X_c \quad (9.3)$$

is satisfied. Two bundles with transition functions  $g_{ab}$  and  $\tilde{g}_{ab}$  are isomorphic if

$$\tilde{g}_{ab} = h_a g_{ab} h_b^{-1} \quad (9.4)$$

for some smooth gauge transformation functions  $h_a \in \text{U}(1)$  on  $X_a$ . Isomorphic bundles can be continuously deformed into each other and are thus topologically indistinguishable. In particular, any bundle which is isomorphic to the bundle with transition functions  $g_{ab} = 1$  is referred to as trivial.

### 9.2 Gauge fields and topological classification of U(1) bundles

A gauge field on a given U(1) bundle consists of a set of locally defined smooth vector fields  $\mathfrak{B}_k^a$  such that

$$\mathfrak{B}_k^a = \mathfrak{B}_k^b - i g_{ab}^{-1} \partial_k g_{ab} \quad \text{on} \quad X_a \cap X_b \quad (9.5)$$

(the index  $k$  and the derivative  $\partial_k$  refer to the coordinates  $t_1, \dots, t_n$ ). The associated field tensor  $\mathfrak{C}_{kl} = \partial_k \mathfrak{B}_l^a - \partial_l \mathfrak{B}_k^a$  is invariant under these transformations and is thus independent of the patch label. It can be shown that gauge fields exist on any bundle and the following result is also well-known.

**Lemma 9.1.** *The magnetic flux*

$$\mathfrak{J}_{kl} = \int_0^{2\pi} dt_k dt_l \mathfrak{C}_{kl} \quad (\text{no sum over } k \text{ and } l) \quad (9.6)$$

*through the  $(k, l)$ -planes is quantized in units of  $2\pi$  and only depends on the underlying bundle. Moreover any two bundles with the same flux quantum numbers are isomorphic to each other.*

A given bundle is hence trivial if the integrals  $\mathfrak{J}_{kl}$  are equal to zero for some particular gauge field. It is in fact sufficient to show that  $|\mathfrak{J}_{kl}| < 2\pi$  since the magnetic flux is quantized.



### 9.3 Gauge fields on the trivial bundle

Gauge fields on the bundle with transition functions  $g_{ab} = 1$  may be represented by smooth periodic vector fields  $\mathfrak{B}_k(t)$  on  $\mathbb{R}^n$  with period  $2\pi$ . Such fields are not uniquely determined by the associated field tensor, but one can always find a periodic gauge potential for a given field tensor if a few obvious conditions are fulfilled. The following lemma provides a particular solution of this problem.

**Lemma 9.2.** *Suppose  $\mathfrak{C}_{kl}(t)$  is a smooth periodic tensor field satisfying*

$$\mathfrak{C}_{kl} = -\mathfrak{C}_{lk}, \quad \partial_k \mathfrak{C}_{lj} + \partial_l \mathfrak{C}_{jk} + \partial_j \mathfrak{C}_{kl} = 0. \quad (9.7)$$

*If the associated magnetic fluxes (9.6) vanish, there exists a smooth periodic vector field  $\mathfrak{B}_k(t)$  such that  $\mathfrak{C}_{kl} = \partial_k \mathfrak{B}_l - \partial_l \mathfrak{B}_k$  and*

$$|\mathfrak{B}_k(t)| \leq \pi(n-1) \sup_{r,k,l} |\mathfrak{C}_{kl}(r)|. \quad (9.8)$$

*Proof:* If  $n = 1$  there is nothing to prove since  $\mathfrak{B}_k(t) = 0$  is a possible choice for the gauge field. Now let us assume that the lemma has been established in dimension  $n - 1$  and that  $\mathfrak{C}_{kl}(t)$  is a given tensor field in  $n$  dimensions with the required properties. Evidently, when restricted to the hyper-plane  $t_n = 0$ , this field satisfies the premises of the lemma in  $n - 1$  dimensions and we may conclude that there exists a periodic vector field  $\mathfrak{B}_k^\circ(t)$  depending on  $t_1, \dots, t_{n-1}$  such that

$$\mathfrak{C}_{kl}(t)|_{t_n=0} = \partial_k \mathfrak{B}_l^\circ(t) - \partial_l \mathfrak{B}_k^\circ(t) \quad \text{for all } k, l < n. \quad (9.9)$$

In the following it will be convenient to consider  $\mathfrak{B}_k^\circ(t)$  to be a field on  $\mathbb{R}^n$  which is independent of  $t_n$ .

We now introduce the field

$$\mathfrak{b}_k(t) = - \int_0^{2\pi} \frac{dr_n}{2\pi} \mathfrak{C}_{nk}(r), \quad r = (t_1, \dots, t_{n-1}, r_n), \quad (9.10)$$

which is also independent of  $t_n$  and periodic. Using the properties of the field tensor, it is easy to show that  $\partial_k \mathfrak{b}_l - \partial_l \mathfrak{b}_k = 0$  and the line integral

$$\mathfrak{B}_n(t) = \int_0^t dr_k \mathfrak{b}_k(r) \quad (9.11)$$

is hence independent of the integration path. Note that  $\mathfrak{B}_n(t)$  is periodic in all coordinates  $t_k$  since the flux integrals  $\mathfrak{J}_{nk}$  vanish.

Next we define the components of the vector field with index  $k < n$  through

$$\mathfrak{B}_k(t) = \int_0^{t_n} dr_n \mathfrak{C}_{nk}(r) + t_n \mathfrak{b}_k(t) + \mathfrak{B}_k^\circ(t), \quad (9.12)$$

where  $r$  is as in eq. (9.10). This field is periodic and it is evident that

$$\partial_n \mathfrak{B}_k(t) = \mathfrak{C}_{nk}(t) + \mathfrak{b}_k(t). \quad (9.13)$$

Together with eq. (9.11) (which implies  $\partial_k \mathfrak{B}_n = \mathfrak{b}_k$ ), this proves that  $\mathfrak{C}_{nk} = \partial_n \mathfrak{B}_k - \partial_k \mathfrak{B}_n$  and it is easy to check that the other components of the field tensor are also correctly obtained.

To show that the so constructed gauge field satisfies the bound (9.8), one proceeds inductively, assuming the bound holds for the field  $\mathfrak{B}_k^\circ(t)$  with  $n$  replaced by  $n - 1$ . Straightforward estimates of the right-hand sides of eqs. (9.10)–(9.12), taking the periodicity of the fields into account, then lead to the desired bound.  $\square$

Another result on which we shall rely later is that the gauge-invariant content of a given gauge field  $\mathfrak{B}_k(t)$  is completely determined by the associated field tensor and the Wilson lines

$$\mathfrak{W}_k(t) = \exp \left\{ i \int_0^{2\pi} ds \mathfrak{B}_k(t) |_{t_k \rightarrow t_k + s} \right\} \quad (9.14)$$

winding around the torus. This is a well-known result and we simply quote

**Lemma 9.3.** *Any two smooth periodic gauge fields  $\mathfrak{B}_k(t)$  and  $\tilde{\mathfrak{B}}_k(t)$  with the same field tensors and the same Wilson lines are related to each other by a gauge transformation,*

$$\tilde{\mathfrak{B}}_k(t) = \mathfrak{B}_k(t) - ih(t)^{-1} \partial_k h(t), \quad (9.15)$$

where  $h(t) \in \text{U}(1)$  is a smooth periodic function of the coordinates  $t_1, \dots, t_n$ .

Note that the Wilson lines coincide at all  $t$  if do at  $t = 0$  and if the field tensors of the two fields are the same, since

$$\mathfrak{W}_k(t) = \mathfrak{W}_k(0) \exp \left\{ i \int_0^t dr_l \int_0^{2\pi} ds \mathfrak{C}_{lk}(r) |_{r_k \rightarrow r_k + s} \right\}, \quad (9.16)$$

where the line integral from 0 to  $t$  is taken along an arbitrary path.

## 10. Proof of theorem 5.1

We first show that smooth measures exist and shall then apply a basis transformation to match  $j_\mu(x)$  with the current  $\tilde{j}_\mu(x)$  derived from the measure.

### 10.1 Existence of a smooth measure

As discussed in sect. 8, we can always choose a patched basis  $v_j^a(x)$  of left-handed fields which is locally smooth. We now prove that the associated U(1) bundle is trivial. Since the space  $\mathfrak{A}[m]$  of transverse vector fields is contractible, it suffices to consider the bundle over the submanifold of admissible fields with  $A_\mu^T(x) = 0$ . Our task is then to show that the connection

$$\mathfrak{B}_k^a(t) = i \sum_j (v_j^a, \partial_{t_k} v_j^a) \quad (10.1)$$

has vanishing magnetic flux quantum numbers (see sect. 9 for unexplained notations and the relevant lemma). Once this is achieved the existence of a basis satisfying the smoothness condition (8.3) and thus of a smooth measure is guaranteed.

If we define the measure term

$$\tilde{\mathfrak{L}}_\eta^a = i \sum_j (v_j^a, \delta_\eta v_j^a) \quad (10.2)$$

as usual, with a tilde to distinguish it from the linear functional  $\mathfrak{L}_\eta$ , the connection may be represented through

$$\mathfrak{B}_k^a(t) = \tilde{\mathfrak{L}}_\eta^a, \quad \eta_\mu(x) = -iU(x, \mu)^{-1} \partial_{t_k} U(x, \mu). \quad (10.3)$$

The corresponding expression for the field tensor is

$$\mathfrak{E}_{kl}(t) = \delta_\eta \tilde{\mathfrak{L}}_\zeta^a - \delta_\zeta \tilde{\mathfrak{L}}_\eta^a, \quad (10.4)$$

where  $\zeta_\mu(x)$  is defined in the same way as  $\eta_\mu(x)$  with  $\partial_{t_k}$  replaced by  $\partial_{t_l}$ .

We now recall that the measure term satisfies the local integrability condition (3.12). The same is true for  $\mathfrak{L}_\eta$  and we thus conclude that

$$\mathfrak{E}_{kl}(t) = \delta_\eta \mathfrak{L}_\zeta - \delta_\zeta \mathfrak{L}_\eta. \quad (10.5)$$

In particular, since  $j_\mu(x)$  is globally defined and smooth, the magnetic flux integrals (9.6) are equal to zero and the U(1) bundle associated with the basis  $v_j^a(x)$  is hence trivial.

## 10.2 Basis transformation

We may now assume that the basis  $v_j^a(x)$  satisfies the smoothness condition (8.3). The associated current  $\tilde{j}_\mu(x)$  is then independent of the patch label  $a$  and smoothly dependent on the link variables. Our aim in the following is to prove that

$$\tilde{\mathfrak{L}}_\eta = \mathfrak{L}_\eta - i\hbar^{-1}\delta_\eta h, \quad (10.6)$$

where  $h$  is some globally defined smooth phase factor. It is then evident that a measure with the required properties is obtained by performing a basis transformation

$$v_j^a(x) \rightarrow \begin{cases} v_1^a(x)h & \text{if } j = 1, \\ v_j^a(x) & \text{otherwise.} \end{cases} \quad (10.7)$$

This measure is, incidentally, uniquely determined up to constant phase factor in each topological sector, because any basis transformation which preserves the measure term has to have constant determinant.

Recalling lemma 9.3 and our discussion above, it is clear that  $\mathfrak{L}_\eta$  and  $\tilde{\mathfrak{L}}_\eta$  are related by a basis transformation if (and only if) the associated Wilson lines (8.9) are the same. The lemma has been formulated for gauge fields on the  $n$ -dimensional torus, but it extends to the field manifold  $\mathfrak{U}[m]$  since the factor  $\mathfrak{A}[m]$  is contractible.

There are two different types of Wilson lines that have to be computed. The first of them are associated with the gauge loops

$$U_t(x, \mu) = \Lambda_t(x)V_{[m]}(x, \mu)\Lambda_t(x + \hat{\mu})^{-1}, \quad 0 \leq t \leq 2\pi, \quad (10.8)$$

in field space, where the transformation  $\Lambda_t(x)$  is defined by

$$\Lambda_t(x) = \exp\{it\delta_{\tilde{x}y}\}, \quad (10.9)$$

with  $y$  being some fixed lattice point and  $\tilde{x} = x \bmod L$  as before. The curve parameter  $t$  is just one of the coordinates  $t_k$  on the torus while all other coordinates are set zero. As discussed at the end of sect. 9, it is not necessary to work out the Wilson lines at other values of the coordinates since they are related to each other through eq. (9.16).

The other non-contractible loops that we need to consider are given by

$$U_t(x, \mu) = V_{[m]}(x, \mu) \exp\{it\delta_{\mu\nu}\delta_{\tilde{x},0}\}, \quad 0 \leq t \leq 2\pi, \quad (10.10)$$

where  $\nu$  is a fixed index. A somewhat surprising fact is that the Wilson lines around all these loops can be computed exactly in terms of the anomaly  $\mathcal{A}_L(x)$ . This is so for both currents,  $j_\mu(x)$  and  $\tilde{j}_\mu(x)$ , and in the following two subsections we shall show that the Wilson lines which one obtains are the same, thus completing the proof of theorem 5.1.

### 10.3 Computation of Wilson lines (gauge loops)

In the case of the gauge loop (10.8) we have

$$\eta_\mu(x) = -iU_t(x, \mu)^{-1} \partial_t U_t(x, \mu) = -\partial_\mu \delta_{\tilde{x}\tilde{y}} \quad (10.11)$$

and it follows from this and property (d) of the current  $j_\mu(x)$  that

$$\mathfrak{L}_\eta = \mathcal{A}_L(y). \quad (10.12)$$

The anomaly is gauge-invariant and hence independent of  $t$ . We thus obtain

$$\mathfrak{W} = \exp \{i2\pi \mathcal{A}_L(y)_{t=0}\} \quad (10.13)$$

for the Wilson line associated with  $\mathfrak{L}_\eta$ .

To compute the Wilson line associated with the measure term  $\tilde{\mathfrak{L}}_\eta$  we start from lemma 8.1 and note that the projector  $P_t$  is given by

$$P_t = R[\Lambda_t] P_0 R[\Lambda_t]^{-1}. \quad (10.14)$$

The lemma then implies

$$\tilde{\mathfrak{W}} = \lim_{n \rightarrow \infty} \det \{1 - P_0 + (P_0 R[\Lambda_{\Delta t}]^{-1} P_0)^n\}, \quad \Delta t = 2\pi/n, \quad (10.15)$$

and it is immediately clear from this expression that

$$\tilde{\mathfrak{W}} = \exp \{-i2\pi \text{Tr}_L [\omega T P_0]\}, \quad \omega(x) = \delta_{\tilde{x}\tilde{y}}. \quad (10.16)$$

Recalling the definitions of the projector  $P_0$  and the anomaly  $\mathcal{A}_L(x)$ , it is now obvious that the Wilson lines associated with  $j_\mu(x)$  and  $\tilde{j}_\mu(x)$  are the same.

#### 10.4 Computation of Wilson lines (non-gauge loops)

To calculate the Wilson lines along the loop (10.10) we make use of the symmetry transformation

$$U(x, \mu) \rightarrow U'(x, \mu) = U(-x - \hat{\mu}, \mu)^{-1}. \quad (10.17)$$

This is a proper rotation of the lattice which maps the field tensor  $F_{\mu\nu}(x)$  to

$$F'_{\mu\nu}(x) = F_{\mu\nu}(-x - \hat{\mu} - \hat{\nu}). \quad (10.18)$$

In particular, the flux sectors  $\mathfrak{U}[m]$  are invariant under this transformation.

From Lemma 7.1 and the definition (7.10) of  $V_{[m]}(x, \mu)$  we now infer that

$$V'_{[m]}(x, \mu) = \Omega_0(x) V_{[m]}(x, \mu) \Omega_0(x + \hat{\mu})^{-1} \quad (10.19)$$

for some gauge transformation function  $\Omega_0(x)$  satisfying  $\Omega_0(0) = 1$ . It follows from this that the fields  $U_t(x, \mu)$  along the curve transform according to

$$U'_t(x, \mu) = \Omega_t(x) U_{2\pi-t}(x, \mu) \Omega_t(x + \hat{\mu})^{-1}, \quad (10.20)$$

$$\Omega_t(x) = \Omega_0(x) \exp \{it\delta_{\hat{x}, 0}\}. \quad (10.21)$$

Up to a gauge transformation the curve is thus mapped onto itself with the reversed orientation.

Taking the gauge invariance and the other symmetry properties of the current  $j_\mu(x)$  into account, an immediate consequence of these observations is that

$$j_\mu(x)|_{t \rightarrow 2\pi-t} = -j_\mu(-x - \hat{\mu}). \quad (10.22)$$

Along the curve the field variation  $\eta_\mu(x)$  is given by

$$\eta_\mu(x) = -iU_t(x, \mu)^{-1} \partial_t U_t(x, \mu) = \delta_{\mu\nu} \delta_{\hat{x}, 0} \quad (10.23)$$

and it is now straightforward to show that

$$\int_0^{2\pi} dt \mathfrak{L}_\eta = \int_0^\pi dt \sum_{x \in \Gamma} \delta_{x, 0} \partial_\mu^* j_\mu(x). \quad (10.24)$$

Using property (d) of the current, the result

$$\mathfrak{W} = \exp \left\{ i \int_0^\pi dt \sum_{x \in \Gamma} \delta_{x, 0} \mathcal{A}_L(x) \right\} \quad (10.25)$$

is thus obtained.

Our starting point for the computation of the Wilson line  $\widetilde{\mathfrak{W}}$  associated with the measure term is again lemma 8.1. The symmetry discussed above implies that

$$P_t Q_t = Q_t P_{2\pi-t}, \quad Q_t Q_{2\pi-t} = 1, \quad (10.26)$$

where  $Q_t$  is a unitary operator which acts on fermion fields according to

$$Q_t \psi(x) = R[\Omega_t(-x)] \gamma_5 \psi(-x). \quad (10.27)$$

If we set  $n = 2r$  in eq. (8.11) it follows from this that

$$\begin{aligned} \widetilde{\mathfrak{W}} = \lim_{n \rightarrow \infty} \det \{ & 1 - P_0 + P_0(Q_{t_1})^{-1} P_{t_1} Q_{t_1} \dots (Q_{t_r})^{-1} P_{t_r} Q_{t_r} \\ & \times P_{t_{r-1}} P_{t_{r-2}} \dots P_{t_1} P_0 \}. \end{aligned} \quad (10.28)$$

We may now insert the representation (8.12) for the projectors and in a few steps one then ends up with the expression

$$\widetilde{\mathfrak{W}} = \mathfrak{W} \times \det \{ 1 - P_0 + P_0(Q_0)^{-1} P_0 \} \det \{ 1 - P_\pi + P_\pi Q_\pi P_\pi \}. \quad (10.29)$$

The operator  $Q_0$  maps the space of left-handed fields at  $t = 0$  onto itself and its square is equal to 1. Similarly  $Q_\pi$  operates in the space of left-handed fields at  $t = \pi$  and its square is also equal to 1. The determinants in eq. (10.29) thus contribute to the sign of the Wilson line and we are left with the problem to prove that the product of these sign factors is positive.

First note that the determinants are products of  $N$  sign factors, one for each fermion flavour. In all cases where the charge  $e_\alpha$  is even we have

$$U_0(x, \mu)^{e_\alpha} = U_\pi(x, \mu)^{e_\alpha}, \quad \Omega_0(x)^{e_\alpha} = \Omega_\pi(x)^{e_\alpha}, \quad (10.30)$$

and the contribution of these fermions is thus equal to 1. If one has a pair of charges  $e_\alpha = -e_{\alpha'}$  one can use charge conjugation to show that the corresponding factors cancel each other and the same is trivially true for any pair of equal charges. This proves that  $\widetilde{\mathfrak{W}} = \mathfrak{W}$  if the charges  $e_\alpha$  satisfy the constraint (5.1).

In the vacuum sector  $\mathfrak{U}[0]$  the situation is simplified by the fact that the sign factors are the same for any pair of odd charges. The anomaly cancellation condition (1.1) implies that the total number of odd charges is even so that the Wilson lines coincide in the vacuum sector independently of whether the constraint (5.1) is satisfied or not.

## 11. Proof of theorem 5.4

### 11.1 Properties of $j_\mu^*(x)$ in finite volume

For any admissible gauge field  $U(x, \mu)$  in finite volume, the current  $j_\mu^*(x)$  is well-defined and periodic in  $x$ . The current thus becomes a local composite field on the finite lattice and we now proceed to study the associated linear functional

$$\mathfrak{K}_\eta = \sum_{x \in \Gamma} \eta_\mu(x) j_\mu^*(x). \quad (11.1)$$

Note that  $\mathfrak{K}_\eta$  is not quite the same as  $\mathfrak{L}_\eta^*$ , since the latter is defined for source fields with compact support while we here assume that  $\eta_\mu(x)$  is a periodic field.

To work out the curvature of  $\mathfrak{K}_\eta$  it is helpful to define the truncated fields

$$\eta_\mu^n(x) = \begin{cases} \eta_\mu(x) & \text{if } x - Ln \in \Gamma, \\ 0 & \text{otherwise,} \end{cases} \quad (11.2)$$

for any integer vector  $n$ . Translation invariance and periodicity then imply

$$\delta_\eta \mathfrak{K}_\zeta - \delta_\zeta \mathfrak{K}_\eta = \sum_{n \in \mathbb{Z}^4} \{ \delta_{\eta^n} \mathfrak{L}_{\zeta^0}^* - \delta_{\zeta^0} \mathfrak{L}_{\eta^n}^* \} \quad (11.3)$$

and after inserting eq. (5.9) one obtains

$$\delta_\eta \mathfrak{K}_\zeta - \delta_\zeta \mathfrak{K}_\eta = i \text{Tr} \{ Q_\Gamma \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \}. \quad (11.4)$$

The projector which appears in this equation is defined by

$$Q_\Gamma \psi(x) = \begin{cases} \psi(x) & \text{if } x \in \Gamma, \\ 0 & \text{otherwise,} \end{cases} \quad (11.5)$$

and the trace is taken in infinite volume. Evidently the right-hand sides of eqs. (3.12) and (11.4) are different and  $j_\mu^*(x)$  itself is, therefore, not an acceptable choice for the current  $j_\mu(x)$  in finite volume.

We now introduce the kernel

$$P(x, y) = \frac{1}{2}(1 - \gamma_5) \delta_{xy} + \frac{1}{2} \gamma_5 D(x, y) \quad (11.6)$$



of the projector  $\hat{P}_-$  in infinite volume and note that

$$\begin{aligned} \mathrm{Tr}_L \{ \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \} &= \sum_{x \in \Gamma} \sum_{y, z \in \mathbb{Z}^4} \sum_{n \in \mathbb{Z}^4} \mathrm{tr} \{ P(x, y) \\ &\times [\delta_\eta P(y, z) \delta_\zeta P(z, x + Ln) - \delta_\zeta P(y, z) \delta_\eta P(z, x + Ln)] \}. \end{aligned} \quad (11.7)$$

The trace in eq. (11.4) coincides with the  $n = 0$  term in this sum and

$$\mathfrak{K}_{\eta\zeta} = i \mathrm{Tr}_L \{ \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \} - i \mathrm{Tr} \{ Q_\Gamma \hat{P}_- [\delta_\eta \hat{P}_-, \delta_\zeta \hat{P}_-] \} \quad (11.8)$$

is hence equal to the sum of all the other terms. Using the locality properties of the Dirac operator, it is possible to deduce the bounds †

$$\|P(x, y)\| \leq \kappa_1 (1 + \|x - y\|^{\nu_1}) e^{-\|x - y\|/\varrho}, \quad (11.9)$$

$$\|\delta_\eta P(x, y)\| \leq \kappa_2 (1 + \|x - y\|^{\nu_2}) e^{-\|x - y\|/\varrho} \|\eta\|_\infty, \quad (11.10)$$

where  $\varrho$  denotes the localization range of the Dirac operator and  $\kappa_i$  and  $\nu_i \geq 0$  are some constants that do not depend on the gauge field. It follows from this that

$$|\mathfrak{K}_{\eta\zeta}| \leq \kappa_3 L^{\nu_3} e^{-L/\varrho} \|\eta\|_\infty \|\zeta\|_\infty \quad (11.11)$$

and the functional  $\mathfrak{K}_\eta$  thus satisfies the integrability condition in finite volume up to exponentially small terms.

In the following we construct a gauge-invariant linear functional  $\mathfrak{S}_\eta$  which transforms in the same way as  $\mathfrak{K}_\eta$  under the lattice symmetries and which satisfies

$$\delta_\eta \mathfrak{S}_\zeta - \delta_\zeta \mathfrak{S}_\eta = \mathfrak{K}_{\eta\zeta}, \quad |\mathfrak{S}_\eta| \leq \kappa_4 L^{\nu_4} e^{-L/\varrho} \|\eta\|_\infty. \quad (11.12)$$

It is then evident that  $\mathfrak{L}_\eta = \mathfrak{K}_\eta + \mathfrak{S}_\eta$  is a solution of the integrability condition (3.12) and that the associated current  $j_\mu(x)$  has all the required properties apart from the fact that condition (d) of theorem 5.1 is not obviously fulfilled.

---

† The supremum norm is given by  $\|\eta\|_\infty = \sup_{x, \mu} |\eta_\mu(x)|$  and all other norms are the usual ones in the appropriate spaces of indices

### 11.2 Construction of $\mathfrak{S}_\eta$ at $A_\mu^T(x) = 0$

The submanifold of fields satisfying  $A_\mu^T(x) = 0$  is an  $n$ -dimensional torus which may be parametrized by coordinates  $t_1, \dots, t_n$  as in sect. 9. If we set

$$\mathfrak{C}_{kl}(t) = \mathfrak{R}_{\eta\zeta}, \quad (11.13)$$

where  $\eta_\mu(x)$  and  $\zeta_\mu(x)$  are the field variations associated with  $t_k$  and  $t_l$ , it is clear from the above that this tensor is exponentially small at large  $L$ . Moreover one knows that  $\mathfrak{C}_{kl}(t)$  is the field tensor of a connection on some  $U(1)$  bundle over the torus, because this is the case for both terms in eq. (11.8).

It follows from this that the flux integrals (9.6) are less than  $2\pi$  in magnitude and hence equal to zero if  $L$  exceeds a certain multiple of  $\varrho$ . On these lattices  $\mathfrak{C}_{kl}(t)$  thus satisfies all premises of lemma 9.2 and we conclude that there exists a linear functional  $\mathfrak{S}_\eta$  which satisfies eq. (11.12) along the submanifold of fields with  $A_\mu^T(x) = 0$ . Note that only the variations  $\eta_\mu(x)$  are admitted here which correspond to variations of the parameters  $t_1, \dots, t_n$ . These are precisely those for which  $\eta_\mu(x) = \eta_\mu^L(x)$ , where

$$\eta_\mu^L(x) = L^{-4} \sum_{y \in \Gamma} \eta_\mu(y) + \sum_{y \in \Gamma} \partial_\mu G_L(x-y) \partial_\nu^* \eta_\nu(y) \quad (11.14)$$

denotes the longitudinal part of any given variation (the Green function  $G_L(z)$  has been introduced in sect. 7).

The solution  $\mathfrak{S}_\eta$  which one obtains in this way does not have any special symmetry properties, but since  $\mathfrak{R}_{\eta\zeta}$  is gauge-invariant and transforms in the appropriate way under the lattice symmetries, we can enforce the proper transformation behaviour by averaging  $\mathfrak{S}_\eta$  over these symmetries and the gauge group.

### 11.3 Extension of $\mathfrak{S}_\eta$ to all admissible fields

According to lemma 7.4 any given field  $U(x, \mu)$  in the sector  $\mathfrak{U}[m]$  may be represented in a one-to-one manner by a field  $V_{[m]}(x, \mu) \mathring{U}(x, \mu)$  contained in the submanifold of fields considered above and the transverse field  $A_\mu^T(x)$ . The field variations  $\eta_\mu(x)$  accordingly split into a longitudinal variation  $\eta_\mu^L(x)$ , defined through eq. (11.14), and a transverse variation  $\eta_\mu^T(x) = \eta_\mu(x) - \eta_\mu^L(x)$ .

We now consider the curve

$$U_t(x, \mu) = V_{[m]}(x, \mu) \mathring{U}(x, \mu) e^{itA_\mu^T(x)}, \quad 0 \leq t \leq 1, \quad (11.15)$$

and define the functional

$$\mathfrak{S}_\eta = \mathfrak{S}_{\eta^L}|_{t=0} + i \int_0^1 dt \left[ \text{Tr}_L \{ \hat{P}_- [\partial_t \hat{P}_-, \delta_\eta \hat{P}_-] \} - \text{Tr} \{ Q_\Gamma \hat{P}_- [\partial_t \hat{P}_-, \delta_\eta \hat{P}_-] \} \right]. \quad (11.16)$$

The first term in this equation coincides with the linear functional constructed in the preceding subsection. As for the second term we remark that the square bracket is proportional to  $\mathfrak{R}_{\eta\zeta}$  with  $\zeta_\mu(x) = A_\mu^T(x)$ . From eq. (7.15) and the fact that the field tensor is bounded by  $\epsilon$  one infers that  $\|A^T\|_\infty \leq \kappa_5 L^4$  and  $\mathfrak{S}_\eta$  is thus exponentially small. Following the lines in part (c) of the proof of theorem 5.3 given in sect. 6, it is also easy to check that  $\mathfrak{S}_\eta$  has the right curvature.

#### 11.4 Final steps

At this point we have constructed a current  $j_\mu(x)$  which satisfies the bound (5.10) and which fulfils conditions (a)–(c) of theorem 5.1. To show that the last condition is also fulfilled, we substitute  $\eta_\mu(x) = -\partial_\mu \omega(x)$  in eq. (3.12) and make use of the gauge transformation properties of the current and the projector to the left-handed fields. This leads to the identity

$$\delta_\zeta \left\{ \sum_{x \in \Gamma} \omega(x) [\partial_\mu^* j_\mu(x) - \mathcal{A}_L(x)] \right\} = 0 \quad (11.17)$$

from which one infers that  $\partial_\mu^* j_\mu(x) - \mathcal{A}_L(x)$  only depends on the topological sector but not on the particular gauge field that has been chosen. Because of translation invariance a dependence on  $x$  is then also excluded.

It follows from this and the index theorem [16,18] that

$$L^4 [\partial_\mu^* j_\mu(x) - \mathcal{A}_L(x)] = - \sum_{y \in \Gamma} \mathcal{A}_L(y) \quad (11.18)$$

is an integer. On large lattices this integer has to vanish since

$$|\mathcal{A}_L(x) - \mathcal{A}(x)| \leq \kappa_6 L^{\nu_6} e^{-L/\varrho} \quad (11.19)$$

and since the anomaly in infinite volume cancels up to a divergence term. We thus conclude that the current  $j_\mu(x)$  satisfies condition (d) when  $L$  exceeds a certain multiple of  $\varrho$ .

## 12. Concluding remarks

Chiral gauge theories with anomaly-free multiplets of Weyl fermions are well-defined to all orders of perturbation theory, but it is not obvious that they can be consistently formulated at the non-perturbative level. The construction presented in this paper provides an affirmative answer to this question for the case of abelian gauge theories. Moreover it shows that one can introduce a momentum cutoff in these theories without breaking the gauge invariance or giving up the requirement of locality, which has long been thought to be impossible.

While the general structure of the lattice theories that we have described is simple, the definition of the fermion integration measure turned out to be non-trivial because of the gauge anomaly. We have shown that the measure can be characterized through a local current satisfying certain conditions and then gave a constructive proof that these conditions can be fulfilled. It is easy to convince oneself, using similar arguments as in the proof of theorem 5.1, that the current is uniquely determined up to terms which amount to adding counterterms to the gauge field action with the appropriate symmetry and locality properties. In other words, this is the usual regularization ambiguity which one has in any lattice theory.

The relative normalizations and phases of the different topological sectors [the weight factors  $w[m]$  in eq. (2.28)] however remain undetermined at this point. This problem is not specific to the lattice approach and it would be sufficient to know the normalizations in the semi-classical approximation in the continuum theory to be able to fix these factors. What seems to be lacking at present is a theoretical principle which restricts the relative weights of the different sectors.

Evidently one would be interested in extending the present work to non-abelian gauge theories. While the discussion in sects. 2–4 carries over with little change, it is not obvious how precisely the anomaly cancellation works out, because the general structure of the non-abelian anomaly is currently not known on the lattice. Moreover the topology of the field space is presumably not as simple as in the abelian case and the absence of global topological obstructions may consequently be difficult to prove. It is conceivable, however, that some of these problems can be bypassed if one succeeds in deriving a closed expression for the imaginary part of the effective action along the lines of refs. [13,14].

I would like to thank Peter Hasenfratz, Pilar Hernández, Karl Jansen, Ferenc Niedermayer and Peter Weisz for helpful discussions. I am also grateful for hospitality at the Max-Planck-Institute in Munich and the Institute for Theoretical Physics at the University of Bern, where part of this work has been completed.

## Appendix A

All fields considered in this paper live on a four-dimensional hyper-cubic euclidean lattice with lattice spacing  $a = 1$ . Flavour indices  $\alpha, \beta, \dots$  run from 1 to  $N$  and Lorentz indices  $\mu, \nu, \dots$  from 0 to 3. Unless stated otherwise the Einstein summation convention is applied to the latter. The symbol  $\epsilon_{\mu\nu\rho\sigma}$  stands for the totally anti-symmetric tensor with  $\epsilon_{0123} = 1$  and  $\delta_{xy}$  is equal to 1 if  $x = y$  and zero otherwise. The conventions for the Dirac matrices are

$$(\gamma_\mu)^\dagger = \gamma_\mu, \quad \{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}, \quad \gamma_5 = \gamma_0\gamma_1\gamma_2\gamma_3. \quad (\text{A.1})$$

In particular,  $\gamma_5$  is hermitean and  $(\gamma_5)^2 = 1$ .

The forward and backward nearest-neighbour difference operators  $\partial_\mu$  and  $\partial_\mu^*$  act lattice functions  $f(x)$  according to

$$\partial_\mu f(x) = f(x + \hat{\mu}) - f(x), \quad (\text{A.2})$$

$$\partial_\mu^* f(x) = f(x) - f(x - \hat{\mu}), \quad (\text{A.3})$$

where  $\hat{\mu}$  denotes the unit vector in direction  $\mu$ .

## Appendix B

In view of the discussion in subsect. 2.3, it suffices to consider the lattice Dirac operator in infinite volume. For all admissible gauge fields we then require that the following properties hold.

(a) *Locality and differentiability.* Ideally one would like the Dirac operator to be strictly local, which would imply that the non-zero contributions to the sum (2.15) come from the points  $y$  in a finite neighbourhood of  $x$ . Moreover the kernel  $D(x, y)$  should be a smooth function of the gauge field variables residing there.

This sort of locality is, however, incompatible with the Ginsparg-Wilson relation [22] and a more general notion of locality is hence adopted here, where the kernel is allowed to have exponentially decaying tails at large distances [21,24]. More precisely

we demand that  $D$  is a sum of strictly local operators,

$$D(x, y) = \sum_{k=1}^{\infty} D_k(x, y), \quad (\text{B.1})$$

with localization regions whose diameter  $d_k$  grows at most linearly with  $k$ . Moreover these kernels and their derivatives  $D_k(x, y; z_1, \mu_1; \dots; z_n, \mu_n)$  with respect to the gauge field variables  $U(z_1, \mu_1), \dots, U(z_n, \mu_n)$  are required to satisfy the bounds

$$\|D_k(x, y; z_1, \mu_1; \dots; z_n, \mu_n)\| \leq C_n k^{p_n} e^{-\theta k}, \quad (\text{B.2})$$

where the constants  $C_n, p_n \geq 0$  and  $\theta > 0$  are independent of the gauge field.

The important point to note here is that at large separations only the terms with large  $k$  contribute. As a consequence we have

$$\|D(x, y)\| \leq C (1 + \|x - y\|^p) e^{-\|x - y\|/\varrho} \quad (\text{B.3})$$

for some constants  $C$  and  $p \geq 0$ . The localization range

$$\varrho = \sup_{k \geq 1} \{d_k / (\theta k)\} \quad (\text{B.4})$$

is a fixed number in lattice units and is thus microscopically small compared to the physical distances in the theory. From the point of view of the continuum limit this kind of locality is hence as good as strict locality.

(b) *Gauge covariance and lattice symmetries.* Under gauge transformations and the lattice symmetries (translations, rotations, reflections, charge conjugation), the Dirac operator and the operators defined by the kernels  $D_k(x, y)$  should transform in the same way as the Wilson-Dirac operator  $D_w$  defined below.

(c) *Free fermion limit.* When the gauge field is set to the classical vacuum configuration,  $U(x, \mu) = 1$ , it follows from (a) and (b) that

$$D(x, y) = \int_{-\pi}^{\pi} \frac{d^4 p}{(2\pi)^4} e^{ip(x-y)} \tilde{D}(p), \quad (\text{B.5})$$

where  $\tilde{D}(p)$  is an analytic function in the momenta  $p_\mu$  with period  $2\pi$ . To obtain the correct spectrum of fermions we require that  $\tilde{D}(p)$  is invertible for all non-zero momenta (mod  $2\pi$ ), while for  $p \rightarrow 0$  it should be equal to  $i\gamma_\mu p_\mu + O(p^2)$ .

(d) *Chiral symmetry and hermiticity.* To preserve chiral symmetry on the lattice, the Dirac operator should be a solution of the Ginsparg-Wilson relation (1.2). This identity alone does not imply any hermiticity properties of  $D$ , but it is consistent to require that  $D^\dagger = \gamma_5 D \gamma_5$ .

(e) *Flavour coherence.* The last requirement is that the Dirac operator should be diagonal in flavour space with diagonal entries that are obtained from the same analytic expression by substituting  $U(x, \mu) \rightarrow U(x, \mu)^{e_\alpha}$  in the sector with flavour  $\alpha$ .

A relatively simple solution of the Ginsparg-Wilson relation has been derived by Neuberger by applying the overlap formalism to vector-like gauge theories [17]. Explicitly this operator is given by

$$D = 1 - A(A^\dagger A)^{-1/2}, \quad A = 1 - D_w, \quad (\text{B.6})$$

where  $D_w$  denotes the standard Wilson-Dirac operator

$$D_w = \frac{1}{2} \{ \gamma_\mu (\nabla_\mu^* + \nabla_\mu) - \nabla_\mu^* \nabla_\mu \}. \quad (\text{B.7})$$

Note that one has to insert the representation (2.13) in the definition

$$\nabla_\mu \psi(x) = R[U(x, \mu)] \psi(x + \hat{\mu}) - \psi(x), \quad (\text{B.8})$$

$$\nabla_\mu^* \psi(x) = \psi(x) - R[U(x - \hat{\mu}, \mu)]^{-1} \psi(x - \hat{\mu}), \quad (\text{B.9})$$

of the gauge covariant forward and backward difference operators. It is trivial to show that Neuberger's operator satisfies (b)–(e) and property (a) has recently been established for small  $\epsilon$  [21].

## References

- [1] Y. Shamir, Nucl. Phys. B (Proc. Suppl.) 47 (1996) 212
- [2] A. Borrelli, L. Maiani, G. C. Rossi, R. Sisto and M. Testa, Phys. Lett. B221 (1989) 360; Nucl. Phys. B333 (1990) 335
- [3] M. Testa, The Rome approach to chirality, Talk given at the APCTP - ICTP Joint International Conference (AIJIC 97) on Recent Developments in Nonperturbative Quantum Field Theory, Seoul 1997, hep-lat/9707007
- [4] W. Bock, M. F. L. Golterman and Y. Shamir, Gauge fixing approach to lattice chiral gauge theories, talk given at the 31st International Ahrenshoop Symposium on the Theory of Elementary Particles, Buckow 1997, hep-lat/9804015
- [5] M. Göckeler, A. Kronfeld, G. Schierholz and U. J. Wiese, Nucl. Phys. B404 (1993) 839
- [6] P. Hernández and R. Sundrum, Nucl. Phys. B455 (1995) 287; *ibid* B472 (1996) 334
- [7] G. 't Hooft, Phys. Lett. B349 (1995) 491
- [8] G. T. Bodwin, Phys. Rev. D54 (1996) 6497
- [9] D. B. Kaplan, Phys. Lett. B288 (1992) 342; Nucl. Phys. B (Proc. Suppl.) 30 (1993) 597
- [10] R. Narayanan and H. Neuberger, Phys. Rev. Lett. 71 (1993) 3251; Nucl. Phys. B412 (1994) 574; *ibid* B443 (1995) 305
- [11] S. Randjbar-Daemi and J. Strathdee, Phys. Lett. B348 (1995) 543; Nucl. Phys. B443 (1995) 386; *ibid* B466 (1996) 335; Phys. Lett. B402 (1997) 134
- [12] H. Leutwyler, Phys. Lett. B152 (1985) 78
- [13] L. Alvarez-Gaumé, S. Della Pietra and V. Della Pietra, Phys. Lett. B166 (1986) 177; Commun. Math. Phys. 109 (1987) 691
- [14] L. Alvarez-Gaumé, An introduction to anomalies, Lectures given at the International School on Mathematical Physics, Erice 1985, *in*: Fundamental problems of gauge field theory, eds. G. Velo and A. S. Wightman (Plenum Press, New York, 1986)
- [15] P. Hasenfratz, Nucl. Phys. B (Proc. Suppl.) 63A-C (1998) 53; Nucl. Phys. B525 (1998) 401
- [16] P. Hasenfratz, V. Laliena and F. Niedermayer, Phys. Lett. B427 (1998) 125
- [17] H. Neuberger, Phys. Lett. B417 (1998) 141; *ibid* B427 (1998) 353
- [18] M. Lüscher, Phys. Lett. B428 (1998) 342
- [19] P. H. Ginsparg and K. G. Wilson, Phys. Rev. D25 (1982) 2649
- [20] Y. Kikukawa and A. Yamada, Weak coupling expansion of massless QCD with a Ginsparg-Wilson fermion and axial U(1) anomaly, hep-lat/9806013



- [21] P. Hernández, K. Jansen and M. Lüscher, Locality properties of Neuberger's lattice Dirac operator, hep-lat/9808010
- [22] I. Horvath, Ginsparg-Wilson relation and ultralocality, hep-lat/9808002
- [23] P. Hasenfratz and F. Niedermayer, private communication
- [24] F. Niedermayer, Exact chiral symmetry, topological charge and related topics, Talk given at the International Symposium on Lattice Field Theory, Boulder 1998, hep-lat/9810026
- [25] R. Narayanan, Phys. Rev. D58 (1998) 97501
- [26] M. Lüscher, Topology and the axial anomaly in abelian lattice gauge theories, hep-lat/9808021
- [27] H. Neuberger, Geometrical aspects of chiral anomalies in the overlap, hep-lat/9802033