

OVERVIEW OF NEUTRINO OSCILLATION PHYSICS*

V. BARGER

Physics Department, University of Wisconsin, Madison, WI 53706

Recent evidence for neutrino oscillations has revolutionized the study of neutrino masses and mixing. This report gives an overview of what we are learning from the neutrino oscillation experiments, the prospects for the near term, and the bright future of neutrino mass studies.

1 Neutrino Masses

Tree-level mass generation occurs through the Higgs mechanism. The Dirac mass m_D arises in a lepton conserving ($\Delta L = 0$) interaction and requires a right-handed neutrino. A Majorana mass m_M occurs through a $\Delta L = 2$ process with only a left-handed light neutrino field and a heavy isosinglet intermediate field N^c . Then the see-saw mechanism with $m_D \sim 10^2$ GeV and $m_M > 10^{12}$ GeV generates light neutrinos

$$m_\nu = m_D^2/m_M \quad (1)$$

that are nearly Majorana. In the case that $m_D \sim m_M \sim$ eV, as can be realized in some models, active-sterile neutrino oscillations can take place.

Neutrino mass can alternatively be generated radiatively by new interactions, such as the R -parity violating $\nu b \tilde{b}$ interaction. Recent reviews of theoretical models for neutrino mass generation are given in Ref. 1.

2 Three-Neutrino Oscillations in Vacuum and Matter

The relation of the three-neutrino flavor eigenstates to the mass eigenstates is

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}, \quad (2)$$

where U is the 3×3 Maki-Nakagawa-Sakata (MNS) mixing matrix. It can be parametrized by

$$U = \begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{13}s_{23}c_{12}e^{i\delta} & c_{23}c_{12} - s_{13}s_{23}s_{12}e^{i\delta} & c_{13}s_{23} \\ s_{23}s_{12} - s_{13}c_{23}c_{12}e^{i\delta} & -s_{23}c_{12} - s_{13}c_{23}s_{12}e^{i\delta} & c_{13}c_{23} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\phi_2} & 0 \\ 0 & 0 & e^{i(\phi_3+\delta)} \end{pmatrix} \quad (3)$$

*Talk presented at the 7th International Symposium on Particles, Strings and Cosmology (PASCOS 99), Granlibakken, California, December 1999

where $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$. The extra diagonal phases are present for Majorana neutrinos but do not affect oscillation phenomena.

With three neutrinos there are two independent δm^2 and $\delta m_a^2 \gg \delta m_b^2$ is indicated by the oscillation evidence. The vacuum oscillation probabilities are

$$P(\nu_\alpha \rightarrow \nu_\beta) = A_{\alpha\beta} \sin^2 \Delta_a - B_{\alpha\beta} \sin^2 \Delta_b + \epsilon_{\alpha\beta} J \sin 2\Delta_b, \quad (4)$$

where $\Delta_a \equiv \delta m_a^2 L / 4E_\nu$. $A_{\alpha\beta}$ is the amplitude of the leading oscillation, $B_{\alpha\beta}$ the amplitude of the sub-leading oscillation and J the CP-violating amplitude; all are determined by the U matrix elements. The physical variable is L/E_ν , where L is the baseline from source to detector and E_ν is the neutrino energy.

In matter, ν_e scatter differently from ν_μ and ν_τ , and the effective neutrino mixing amplitude in matter can be very different from the vacuum amplitude.² For the leading oscillation the matter and vacuum oscillation mixings are related in the approximation of constant matter density by

$$\sin^2 2\theta_{13}^m = \frac{\sin^2 2\theta_{13}}{(\cos 2\theta_{13} - A/\delta m_a^2)^2 + \sin^2 2\theta_{13}}, \quad (5)$$

where

$$A = 2\sqrt{2} G_F Y_e \rho(x) E_\nu. \quad (6)$$

Here Y_e is the electron fraction and $\rho(x)$ is the density at path-length x . The $\nu_e \rightarrow \nu_\mu$ (or $\nu_e \rightarrow \nu_\tau$) oscillation argument in matter of constant density is

$$\Delta_a^m = \frac{1.27 \delta m_a^2 (\text{eV}^2) L (\text{km})}{E_\nu (\text{GeV})} \sqrt{\left(\frac{A}{\delta m_a^2} - \cos 2\theta_{13}\right)^2 + \sin^2 2\theta_{13}}. \quad (7)$$

Resonance enhancements in matter are possible for $\delta m_a^2 > 0$, while suppression occurs for $\delta m_a^2 < 0$. It is significant that the resonant energies correspond to neutrino energies relevant to the atmospheric and solar anomalies.

$$\text{Earth : } E_\nu \simeq 15 \text{ GeV} \left(\frac{\delta m_a^2}{3.5 \times 10^{-3} \text{eV}^2} \right) \left(\frac{1.5 \text{ gm/cm}^3}{\rho Y_e} \right) \quad (8)$$

$$\text{Sun : } E_\nu \simeq 10 \text{ MeV} \left(\frac{\delta m_b^2}{10^{-5} \text{eV}^2} \right) \left(\frac{10 \text{ g/cm}^3}{\rho Y_e} \right). \quad (9)$$

2.1 Atmospheric Neutrino Oscillations

The Kamiokande, SuperKamiokande (SuperK), Macro and Soudan atmospheric neutrino measurements³ show a μ/e ratio that is about 0.6 of expectations. The SuperK experiment has established the dependence of the e and μ event rates on zenith angle, or equivalently the baseline L . $N(e)$ is

independent of L and validates the ν_e flux calculation (within 20%). $N(\mu)$ depletion increases with L . The muon event distributions are will described by $\nu_\mu \rightarrow \nu_\tau$ vacuum oscillations with a ν_μ *survival* probability

$$P(\nu_\mu \rightarrow \nu_\mu) = 1 - A_{\mu\tau} \sin^2(1.27\delta m_a^2 L/E_\nu) \quad (10)$$

with $\delta m_a^2 = 3.5 \times 10^{-3} \text{ eV}^2$ and maximal or near maximal amplitude

$$A_{\mu\tau} = 1_{-0.2}^{+0.0} \quad (\text{i.e., } |\theta_{32} - 45^\circ| < 13^\circ). \quad (11)$$

With further data accumulation³ a slightly lower central value is now indicated ($\delta m_a^2 = 2.8 \times 10^{-3} \text{ eV}^2$). The $\nu_\mu \rightarrow \nu_\tau$ oscillations are not resolved due to smearing of L and inferred E_ν values, and equally good fits to the SuperK data are found with oscillation and neutrino decay ($\nu_2 \rightarrow \bar{\nu}_4 + J$) models.⁴ For now we assume the simplest interpretation of the SuperK data, namely oscillations. We note that in the CHOOZ reactor experiment $\bar{\nu}_e$ disappearance is not observed at the δm_a^2 scale and the corresponding constraint on 3-neutrino mixing for $\delta m_a^2 = 3.5 \times 10^{-3} \text{ eV}^2$ is

$$A_{\mu e} < 0.2, \quad |U_{e3}| < 0.23, \quad \theta_{13} < 13^\circ. \quad (12)$$

2.2 Solar Neutrino Oscillations

The solar neutrino experiments³ sample different ν_e energy ranges and find different flux deficits compared to the Standard Solar Model (SSM)⁵ as follows:

$\nu_e \text{ } ^{71}\text{Ga} \rightarrow \text{} ^{71}\text{Ge} e$	GALLEX	0.60 ± 0.06
	SAGE	0.52 ± 0.06
$\nu_e \text{ } ^{37}\text{Cl} \rightarrow \text{} ^{37}\text{Ar} e$	Homestake	0.33 ± 0.03
$\nu e \rightarrow \nu e$	SuperK	0.47 ± 0.02

(13)

Thus the ν_e survival probability is inferred to be energy dependent.

Global oscillation fits⁶ have been made using floating ^8B and hep flux normalizations which are somewhat uncertain in the SSM. The relative normalizations from the fits range from 0.5 to 1.2 for the ^8B flux and 1 to 25 for the hep flux. These global fits include the data on (i) total rates, assuming all the experiments are okay — the different Cl suppression ratio plays a vital role; (ii) the night-day asymmetry, which is observed at the 2σ level³ — the large angle matter solution gives night rates $>$ day rates; (iii) seasonal dependence beyond $1/r^2$, which can occur for vacuum solutions. The oscillation analyses⁶ generally agree on the allowed δm_{21}^2 and $\sin^2 2\theta_{12}$ regions for acceptable solutions. Typical candidate solar solutions are given in Table 1.

In the case of vacuum oscillations (VO) several discrete regions of δm_{21}^2 are possible.

Table 1. Representative solutions to the solar neutrino anomaly.

solution	$\sin^2 2\theta_{12}$	δm_{21}^2 (eV ²)
SAM	$\sim 5 \times 10^{-3}$	$\sim 5 \times 10^{-6}$
LAM	~ 1	$\sim 3 \times 10^{-5}$
LOW	~ 1	$\sim 10^{-7}$
VO	~ 1	$\sim 10^{-10}$

2.3 3-Neutrino Mixing Matrix

Once the solar oscillation solution is pinned down, and θ_{12} is thus determined, we will have approximate knowledge of the mixing angles of the 3-neutrino matrix, with $\theta_{23} \sim \pi/4$ and $\theta_{13} \sim 0$ from the atmospheric and CHOOZ data. Upcoming experiments are expected to shed light on the solar solution. In the SNO experiment, which is now taking data, and the forthcoming ICARUS experiment, the high energy ν_e CC events may distinguish LAM, SAM, and LOW solutions with large hep flux contributions from the VO or the SAM sterile neutrino solutions.⁷ Also, the neutral-current to charged-current ratio will distinguish active from sterile oscillations. The Borexino experiment can measure the VO seasonal variation of the ⁷Be line flux. The KamLand reactor experiment to measure the $\bar{\nu}_e$ survival probability will be sensitive to the LAM and LOW solar solutions.⁸

The CP phase δ may be measurable at a neutrino factory if the solar solution is LAM.^{9,10} The CP violation comes in only at the sub-leading oscillation scale.¹¹ An apparent CP-odd asymmetry is induced by matter.

2.4 Models

For maximal mixing in both atmospheric and solar sectors, there is an unique mixing matrix¹²

$$U = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 1/2 & 1/2 & -1/\sqrt{2} \\ 1/2 & 1/2 & 1/\sqrt{2} \end{pmatrix}. \quad (14)$$

In this bimaximal mixing model, there would be no CP-violating effects. However, because $U_{e3} = 0$, long-baseline experiments would have some sensitivity to the sub-leading LAM solar scale oscillations.

Many unification models predict that the neutrino masses are Majorana and hierarchical, there is no cosmologically significant dark matter, and the SAM solar solution (small θ_{12} mixing) obtains.^{1,13}

2.5 Beyond 3 Neutrinos

The LSND evidence for $\nu_\mu \rightarrow \nu_e$ oscillations with $\delta m^2 \sim 1 \text{ eV}^2$, $\sin^2 2\theta \sim 10^{-2}$ requires a δm^2 scale distinct from the atmospheric and solar oscillation scales, and thus a sterile neutrino state would be needed to explain all the oscillation phenomena. Then to also satisfy limits from CDHS accelerator and Bugey reactor experiments, the mass hierarchy must be two separated pairs.¹⁴ Such a scenario would allow even more interesting effects at a neutrino factory, such as large CP violation, since both the leading and sub-leading oscillation scales would be accessible. The MiniBooNE experiment will settle whether the LSND evidence is real. Other interest in sterile neutrinos comes from r -process nucleosynthesis if it occurs in supernovae.¹⁵

3 Long-Baseline Experiments

Long-baseline experiments are needed to (i) confirm the atmospheric evidence for $P(\nu_\mu \rightarrow \nu_\mu)$ at accelerators; (ii) resolve the leading $\nu_\mu \rightarrow \nu_\mu$ oscillation and exclude the neutrino decay possibility; (iii) precisely measure $|\delta m_a^2|$; (iv) exclude $\nu_\mu \rightarrow \nu_s$ disappearance, although SuperK has now shown that oscillations to sterile neutrinos are excluded at 99% CL;³ (v) measure $|U_{e3}|$ from $\nu_e \rightarrow \nu_\mu$ appearance, which requires a muon decay source for the neutrino beam; (vi) determine the sign of δm_a^2 from matter effects in the Earth's crust; and (vii) search for CP violation.

The first long-baseline experiments will measure the energy dependence of the produced muons and measure the neutral-current to charged-current ratio, to partially address the first four issues listed above. The K2K experiment from KEK to SuperK is in operation, with a baseline $L = 250 \text{ km}$ and mean neutrino energy $\langle E_\nu \rangle = 1.4 \text{ GeV}$. The MINOS experiment from Fermilab to Soudan, with $L = 732 \text{ km}$ and possible energies of $\langle E_\nu \rangle = 3, 6, 12 \text{ GeV}$ will begin in 2002. A 10% precision on $|\delta m_a^2|$ may ultimately be possible at MINOS. The ICANOE and OPERA long-baseline experiments from CERN to Gran Sasso with $L \simeq 743 \text{ km}$ have been approved.

Muon storage rings could provide intense neutrino beams ($\sim 10^{19}$ – 10^{21} per year) that would yield thousands of charged-current neutrino interactions in a reasonably sized detector (10–50 kt) anywhere on Earth.^{16,17} These neutrino factories would have pure neutrino beams ($\nu_e, \bar{\nu}_\mu$ from stored μ^+ and

$\bar{\nu}_e, \nu_\mu$ from stored μ^-) with 50% ν_e or $\bar{\nu}_e$ components. Detection of wrong-sign muons (the muons with opposite sign to the charge current from the beam muon neutrino) would signal $\nu_e \rightarrow \nu_\mu$ or $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$ appearance oscillations. We now discuss the capability of a neutrino factory with 2×10^{20} muons a year and a 10 kt detector to resolve the issues raised in the preceding section.

With an $E_\mu = 30$ GeV storage ring at a baseline of $L = 2800$ km, a statistical precision of a few % on $\sin^2 2\theta_{23}$ is possible in ν_μ survival measurements. This accuracy in measuring $\sin^2 2\theta_{23}$ would differentiate the bimaximal model prediction¹² of $\sin^2 2\theta_{23} = 1$ from the democratic model prediction¹⁸ of $\sin^2 2\theta_{23} = 8/9$.

With stored muon energies $E_\mu = 10$ to 50 GeV and baselines of $L = 732$ to 7332 km and 1 kt detector, there would be hundreds of events per year from $\nu_\mu \rightarrow \nu_\tau$ oscillations for an intensity of 2×10^{20} neutrinos.

The wrong-sign muon event rates are approximately proportional to $\sin^2 2\theta_{13}$. For non-zero $\sin^2 2\theta$ the observation of $\nu_e \rightarrow \nu_\mu$ and $\bar{\nu}_e \rightarrow \bar{\nu}_\mu$ appearance oscillations at baselines long enough to have significant matter effects will allow^{10,17,19} a determination of the sign of δm_{32}^2 , and thus determine the pattern of the masses (a 1 + 2 mass hierarchy versus a 2 + 1 hierarchy for three neutrinos). A proof of the principle that the sign of δm^2 can be so determined has been given^{10,17} for a baseline $L = 2800$ km. In μ^+ appearance, $\delta m_{32}^2 > 0$ gives a smaller rate and harder spectrum than $\delta m_{32}^2 < 0$, while the results are opposite in μ^- appearance.¹⁷

In optimizing E_μ and L for long-baseline experiments to find the sign of δm_{32}^2 , $L = 732$ km is too short (matter effects are small) and $L = 7332$ km is too far (event rates are low). The sensitivity to determine the sign of δm_{32}^2 improves linearly with E_μ . There is a tradeoff between energy, detector size and muon beam intensity.¹⁰

4 Absolute Neutrino Masses

Oscillation phenomena determine only mass-squared differences, leaving the absolute mass scale unknown. However, because the atmospheric and solar δm^2 values are $\ll (1 \text{ eV})^2$, all mass eigenvalues are approximately degenerate if at the ~ 1 eV scale. Thus all neutrino mass eigenvalues are bounded²⁰ by the tritium limit from the Troitsk and Mainz experiments,

$$m_j < 3 \text{ eV} \quad \text{for } j = 1, 2, 3. \quad (15)$$

Neutrinoless double-beta decay ($0\nu\beta\beta$) provides a probe of Majorana neutrino mass.²¹ The rate is proportional to the $\nu_e\nu_e$ element of the neutrino mass

matrix. The present limit from the Heidelberg experiment²² is

$$M_{\nu_e\nu_e} < 0.2 \text{ eV} \times f, \quad (16)$$

where the factor f represents uncertainty in the nuclear matrix elements, which might be as large as a factor of 3. The $0\nu\beta\beta$ limit translates to a bound on the summed neutrino Majorana masses of²³

$$\sum m_\nu < 0.75 \text{ eV} \times f \quad (17)$$

in the SAM solar solution. No similar constraints apply to the LAM, LOW or VO solutions where the bound can be satisfied by having opposite CP parity of ν_1 and ν_2 mass eigenstates. Future experiments²⁴ may probe to

$$|M_{\nu_e\nu_e}| = 0.01 \text{ eV}, \quad (18)$$

which would provide sensitivity down to

$$\sum m_\nu = 0.08 \text{ eV} \times f \quad (19)$$

in the SAM solution.

Measurements of the power spectrum by the MAP and PLANCK satellites may determine $\sum m_\nu$ down to $\sim 0.4 \text{ eV}$.²⁵ The heights of the acoustic peaks can also decide how the mass is distributed among the neutrino eigenstates.²⁶

5 Summary

We have entered an exciting new era in the study of neutrino masses and mixing. From the SuperK evidence on atmospheric neutrino oscillations, we already have a surprising amount of information about the neutrino mixing matrix (near maximal $\sin^2 2\theta_{23}$ and near minimal $\sin^2 2\theta_{13}$). The SuperK, SNO, Borexino, KamLand, and ICARUS experiments are expected to differentiate among the candidate solar oscillation possibilities and determine $\sin^2 2\theta_{12}$. MiniBooNE will tell us whether a sterile neutrino is mandated. Neutrino factories will study the leading oscillations, determine the sign of δm_a^2 , measure U_{e3} , and possibly detect CP violation. The GENIUS $0\nu\beta\beta$ experiment and the MAP and PLANCK satellite measurements of the power spectrum will probe the absolute scale of neutrino masses. There is a synergy of particle, physics, nuclear physics, and cosmology occurring in establishing the fundamental properties of neutrinos. A theoretical synthesis should emerge from these experimental pillars.

A more complete version of this review, including figures and more extensive references, can be found in Ref. 27.

Acknowledgments

I thank my collaborators K. Whisnant, T. Weiler, S. Pakvasa, S. Geer, R. Raja, J. Learned, P. Lipari, and M. Lusignoli. This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

References

1. R.N. Mohapatra, hep-ph/9910365, to appear in “Current Aspects of Neutrino Physics”, ed. by D. Caldwell (Springer-Verlag, 2000); G. Altarelli and F. Feruglio, Phys. Rep. **320**, 295 (1999); P. Ramond, hep-ph/0001009; and references therein.
2. L. Wolfenstein, Phys. Rev. **D17**, 2369 (1978). V. Barger, S. Pakvasa, R.J.N. Phillips, and K. Whisnant, Phys. Rev. **D22**, 2718 (1980). S.P. Mikheyev and A. Smirnov, Yad. Fiz. **42**, 1441 (1985) [Sov. J. Nucl. Phys. **42**, 913 (1986)].
3. For a recent review of the atmospheric and solar neutrino data, see T. Kajita, these proceedings.
4. V. Barger, J.G. Learned, P. Lipari, M. Lusignoli, S. Pakvasa and T.J. Weiler, Phys. Lett. **B462**, 109 (1999); J. Valle, Phys. Lett. **B131**, 87 (1983).
5. J.N. Bahcall, S. Basu, and M.H. Pinsonneault, Phys. Lett. **B 433**, 1 (1998).
6. See M.C. Gonzalez-Garcia and C. Peña-Garay, hep-ph/0002186; A. de Gouvea, A. Friedland, and A. Murayama, hep-ph/0002064; J.N. Bahcall, P. Krastev, and A.Yu. Smirnov, Phys. Rev. **D58**, 096016 (1998); Phys. Rev. **D60**, 093001; and references therein.
7. J.N. Bahcall, P. Krastev, and A.Yu. Smirnov, hep-ph/0002293
8. A. de Gouvea, A. Friedland, and H. Murayama, hep-ph/9910286.
9. A. De Rujula, M.B. Gavela, and P. Hernandez, Nucl. Phys. **B547**, 21 (1999); A. Cervera, A. Donini, M.B. Gavela, J.J. Gomez Cadenas, P. Hernandez, O. Mena, and S. Rigolin, hep-ph/0002108; M. Freund, M. Lindner, S.T. Petcov, and A. Romanino, hep-ph/9912457.
10. V. Barger, S. Geer, R. Raja, and K. Whisnant, hep-ph/0003184.
11. V. Barger, K. Whisnant, R.J.N. Phillips, Phys. Rev. Lett. **45**, 2084 (1980); S. Pakvasa, in *Proc. of the XXth International Conference on High Energy Physics*, ed. by L. Durand and L.G. Pondrom, AIP Conf. Proc. No. 68 (AIP, New York, 1981), Vol. 2, p. 1164.

12. V. Barger, S. Pakvasa, T.J. Weiler, and K. Whisnant, Phys. Lett. **B 437**, 107 (1998); A.J. Baltz, A. Goldhaber, and M. Goldhaber, Phys. Rev. Lett. **81**, 5730 (1998); S. Vissani, hep-ph/9708483.
13. See e.g. S.F. King, Phys. Lett. **B439**, 350 (1998); Nucl. Phys. **562**, 57 (1999); C.H. Albright and S.M. Barr, Phys. Rev. **D58**, 013002 (1998); F. Wilczek, *Proc. of Neutrino 98*, p. 511; and references therein.
14. S.M. Bilenky, C. Giunti, and W. Grimus, Eur. Phys. J. **C 1**, 247 (1998); V. Barger, S. Pakvasa, T.J. Weiler, and K. Whisnant, Phys. Rev. **D58**, 093016 (1998).
15. G.C. McLaughlin, J.M. Fetter, A.B. Balantekin, and G.M. Fuller, Phys. Rev. **C59**, 2873 (1999); D.O. Caldwell, G.M. Fuller and Y.-Z. Qian, astro-ph/9910175.
16. S. Geer, Phys. Rev. **D57**, 6989 (1998).
17. V. Barger, S. Geer, and K. Whisnant, Phys. Rev. **D61**, 053004 (2000); V. Barger, S. Geer, R. Raja, and K. Whisnant, hep-ph/9911524, Phys. Rev. D (in press).
18. H. Fritzsch and Z. Xing, Phys. Lett. **B372**, 265 (1996); **440**, 313 (1998); E. Torrente-Lujan, Phys. Lett. **B389**, 557 (1996); M. Fukugita, M. Tanimoto, and T. Yanagida, Phys. Rev. **D57**, 4429 (1998).
19. I. Mocioiu and R. Shrock, hep-ph/0002149.
20. V. Barger, T.J. Weiler, and K. Whisnant, Phys. Lett. **B442**, 255 (1998).
21. M. Doi *et al.*, Phys. Lett. **B 102**, 323 (1981); H. Minakata and O. Yasuda, Phys. Rev. **D 56**, 1692 (1997); H. Georgi and S.L. Glashow, hep-ph/9808293; F. Vissani, JHEP **9906**, 022 (1999); J. Ellis and S. Lola, Phys. Lett. **B458**, 310 (1999); S.M. Bilenki *et al.*, Phys. Lett. **B465**, 193 (1999).
22. L. Baudis *et al.*, Phys. Rev. Lett. **83**, 41 (1999).
23. V. Barger and K. Whisnant, Phys. Lett. **B 456**, 194 (1999).
24. H.V. Klapdor-Kleingrothaus, J. Hellmig, and M. Hirsch, J. Phys. **G 24**, 483 (1998); M. Danilov *et al.*, hep-ex/0002003.
25. D.J. Eisenstein, W. Hu, and M. Tegmark, Phys. Rev. Lett. **80**, 5255 (1998); astro-ph/9807130; R.A.C. Croft, W. Hu, and R. Davé, Phys. Rev. Lett. **83**, 1092 (1999).
26. V. Barger, T. Falk, and R.-J. Zhang (paper in preparation).
27. V. Barger, hep-ph/0003212, to appear in *Proc. of the 5th International Conference on Physics Potential and Development of $\mu^+\mu^-$ Colliders*, San Francisco, December 1999.