

Polarization of vacuum in the decays of neutral kaons

D.L. Khokhlov

Sumy State University, R.-Korsakov St. 2,

Sumy 40007, Ukraine

E-mail: khokhlov@cafe.sumy.ua

Abstract

It is considered the state of K^0 before the decay as that embedded into the vacuum of $+\bar{K}^0 - \bar{K}^0$. Since the state of K_S^0 has the lifetime τ_S , and the state of K_L^0 has the lifetime τ_L , the vacuum of $+\bar{K}^0 - \bar{K}^0$ is polarized. Due to the polarization of the vacuum, $2\tau_S/\tau_L$ of K_S^0 decay with the delay $\approx \tau_L$. Thus CP-invariance holds true in the decays of neutral kaons.

As known [1] strange K^0 and \bar{K}^0 mesons do not possess certain lifetimes relative to the weak decays, since weak interactions do not conserve the strangeness number. There exist two independent linear combinations of K^0 and \bar{K}^0

$$|K_S^0\rangle = \frac{1}{\sqrt{2}}(K^0 - \bar{K}^0) \quad (1)$$

$$|K_L^0\rangle = \frac{1}{\sqrt{2}}(K^0 + \bar{K}^0) \quad (2)$$

They correspond to the particles with the lifetimes $\tau_S = 8.9 \times 10^{-11}$ s and $\tau_L = 5.2 \times 10^{-8}$ s respectively. The states K_S^0 and K_L^0 are of CP-invariance with the eigenvalues +1 and -1 respectively. K_S^0 decays into the system of two pions

$$K_S^0 \rightarrow \pi\pi \quad (3)$$

with the CP-eigenvalue +1, and K_L^0 decays into the system of three pions

$$K_L^0 \rightarrow \pi\pi\pi \quad (4)$$

with the CP-eigenvalue -1.

K^0 is considered as a superposition of K_S^0 and K_L^0

$$|K^0\rangle = \frac{1}{\sqrt{2}}(K_S^0 + K_L^0). \quad (5)$$

Evolution of K^0 is given by

$$|K^0(t)\rangle = \frac{1}{\sqrt{2}}(K_S^0 e^{-t/2\tau_S} + K_L^0 e^{-t/2\tau_L}). \quad (6)$$

One can expect that, within the time $t < \tau_S$, K^0 decays into two pions, and within the time $\tau_S < t < \tau_L$, K^0 decays into three pions. But, within the time $\tau_S < t < \tau_L$, there exists the probability of the decays of K^0 into two pions

$$\frac{\Gamma(K^0(\tau_S < t < \tau_L) \rightarrow \pi^+\pi^-)}{\Gamma(K^0(\tau_S < t < \tau_L) \rightarrow all)} \approx 2 \times 10^{-3} \quad (7)$$

$$\frac{\Gamma(K^0(\tau_S < t < \tau_L) \rightarrow \pi^0\pi^0)}{\Gamma(K^0(\tau_S < t < \tau_L) \rightarrow all)} \approx 10^{-3}. \quad (8)$$

Decays $K^0 \rightarrow \pi\pi$ within the time $\tau_S < t < \tau_L$ are treated as CP-violation.

Consider the decay of K^0 . In view of (1), (2), (5), K^0 decays in combination with $-\bar{K}^0$ as K_S^0 and in combination with $+\bar{K}^0$ as K_L^0 . From this one can consider K^0 before the decay as that embedded into the vacuum of $+\bar{K}^0 - \bar{K}^0$. Since the probabilities of the decay of K^0 in combinations with $+\bar{K}^0$ given by $1/\tau_L$ and in combinations with $-\bar{K}^0$ given by $1/\tau_S$ are different, the vacuum of $+\bar{K}^0 - \bar{K}^0$ is polarized. Due to the polarization of the vacuum K^0 is surrounded by τ_S/τ_L of $+\bar{K}^0$ and by $1/2 - \tau_S/\tau_L$ of $-\bar{K}^0$. The decay of K^0 occurs via the birth of the pair of $+\bar{K}^0 - \bar{K}^0$ from vacuum. From this the decay of K^0 in combination with $+\bar{K}^0$ is accompanied by the birth of $-\bar{K}^0$. The decay of K^0 in combination with $-\bar{K}^0$ is accompanied by the birth of $+\bar{K}^0$. Taking into account that combination of K^0 with $-\bar{K}^0$ corresponds to the state of K_S^0 and combination of K^0 with $+\bar{K}^0$ corresponds to the state of K_L^0 , write the state of K^0 in the form

$$|K^0\rangle = \frac{1}{\sqrt{2}} \left[\left(1 - 2\frac{\tau_S}{\tau_L}\right) K_S^0 + \left(2\frac{\tau_S}{\tau_L}\right) K_L^0 \right] + \frac{1}{\sqrt{2}} \left[\left(1 - 2\frac{\tau_S}{\tau_L}\right) K_L^0 + \left(2\frac{\tau_S}{\tau_L}\right) K_S^0 \right]. \quad (9)$$

$1 - 2\tau_S/\tau_L$ of K_S^0 and $2\tau_S/\tau_L$ of K_L^0 start to decay at $t = 0$. $1 - 2\tau_S/\tau_L$ of K_L^0 start to decay after the decay of $1 - 2\tau_S/\tau_L$ of K_S^0 , i.e. with the delay $\approx \tau_S$. $2\tau_S/\tau_L$ of K_S^0 start to decay after the decay of $2\tau_S/\tau_L$ of K_L^0 , i.e. with the delay $\approx \tau_L$.

From the above consideration it follows that, within the time $t < \tau_S$, $1 - 2\tau_S/\tau_L$ of K_S^0 decay. That is, within the time $t < \tau_S$, the number of K^0 decayed into two pions is estimated as

$$\frac{\Gamma(K^0(t < \tau_S) \rightarrow \pi\pi)}{\Gamma(K^0(t < \tau_L) \rightarrow all)} = \frac{1}{2} - \frac{\tau_S}{\tau_L}. \quad (10)$$

Due to the delay $\approx \tau_L$ caused by the polarization of the vacuum, within the time $\tau_S < t < \tau_L$, $2\tau_S/\tau_L$ of K_S^0 decay. That is, within the time $\tau_S < t < \tau_L$, the number of K^0 decayed into two pions is estimated as

$$\frac{\Gamma(K^0(\tau_S < t < \tau_L) \rightarrow \pi\pi)}{\Gamma(K^0(\tau_S < t < \tau_L) \rightarrow all)} = 2\frac{\tau_S}{\tau_L} = 3.4 \times 10^{-3}. \quad (11)$$

Thus the decays of K^0 into two pions within the time $\tau_S < t < \tau_L$ can be explained by the delay due to the polarization of the vacuum of $+\bar{K}^0 - \bar{K}^0$. Hence CP-invariance holds true in the decays of neutral kaons.

References

- [1] E.D. Commins, P.H. Bucksbaum, *Weak interactions of leptons and quarks* (Cambridge University Press, 1983)