Should E_8 SUSY Yang-Mills be Reconsidered

as a Family Unification Model?

Stephen L. Adler

Institute for Advanced Study

Princeton, NJ 08540

Send correspondence to:

Stephen L. Adler Institute for Advanced Study Einstein Drive, Princeton, NJ 08540 Phone 609-734-8051; FAX 609-924-8399; email adler@ias. edu

ABSTRACT

We review earlier proposals for E_8 family unification, and discuss why recent work of Kovner and Shifman on condensates in supersymmetric Yang-Mills theories suggests the reconsideration of E_8 supersymmetric Yang-Mills as a family unification theory.

One of the outstanding mysteries of the current standard model is the triple repetition of fundamental fermions. Many different ideas [1] have been proposed to explain why there are three (or in some models more) families; here we focus on the possibility, already addressed in the earlier literature, that the family structure has a group theoretic origin, with all three families embedded in a large representation of a family unification group. Since there are 15 (or if right handed neutrinos are included) 16 Weyl spinor fields in each family, a group representation of dimension at least 45 or 48 is required. So we are necessarily considering a large group representation, and if we invoke naturalness to require that it be a low-lying representation of its Lie group or algebra, then we are necessarily considering a large group. A particularly interesting candidate is the group E_8 , which has a 248 dimensional Lie algebra and, as the largest exceptional group, a unique position in the standard Cartan classification of Lie groups. Our aim in this paper, which is unapologetically speculative and programmatic, is to review earlier work on E_8 unification, to explain difficulties encountered, and to argue that recent developments suggest that there may be mechanisms that can overcome these difficulties. Thus the time may be ripe to reconsider E_8 , and specifically E_8 supersymmetric Yang-Mills theory, as a family unification model.

Unification theories based on simple Lie groups follow a basic paradigm established by the SU(5) and SO(10) models. The gauge bosons are as usual in the adjoint representation of the group, and left-handed Weyl fermions are placed in one or more additional representations, chosen to give cancellation of anomalies together with the standard model fermion structure under breaking of the unification group to $SU(3) \times SU(2) \times U(1)$. Turning to E_8 , this is the unique simple Lie group in which the adjoint representation, of dimension 248, is also the fundamental representation. Hence the natural implementation of the basic paradigm is to place left-handed Weyl fermions in the 248 representation, giving a model in which the gauge bosons or gluons, and the fermionic matter fields, are both in the adjoint 248 representation. Since in four dimensions supersymmetric Yang-Mills theory can be constructed with adjoint fermions that are either Majorana or Weyl [2], in this E_8 model the fermions and gluons are in the same supermultiplet, achieving a complete unification of matter fields and force-carrying fields. The point that an E_8 unification model is automatically supersymmetric was made independently more than twenty years ago by Baaklini [3], by Bars and Günaydin [4], and by Konshtein and Fradkin [5], was followed up on in a paper of Koca [6], and was briefly noted in Slansky's comprehensive review [7] of group theory for model building.

Another interesting feature of E_8 is that it naturally contains three families. Most of the recent discussions of single family grand unification are based on either the group SO(10)[8] or the group E_6 [9]. In SO(10) unification the 16 Weyl fermions of a family (including a right handed neutrino) are placed in a 16 representation, while in unification in the larger group E_6 , of which SO(10) is a subgroup, these fermions are placed in a 27 representation. Under the decomposition $E_8 \supset SU(3) \times E_6$, the 248 of E_8 branches [7] as

$$248 = (8,1) + (1,78) + (3,27) + (\overline{3},\overline{27}) \quad , \tag{1a}$$

while under $E_8 \supset SU(3) \times SO(10) \times U(1)$, the 248 branches [7] as

$$248 = (1, 16)(3) + (1, \overline{16})(-3) + (3, 16)(-1) + (\overline{3}, \overline{16})(1) + (3, 10)(2) + (\overline{3}, 10)(-2) + (3, 1)(-4) + (\overline{3}, 1)(4) + (8, 1)(0) + (1, 45)(0) + (1, 1)(0) ,$$

$$(1b)$$

with the U(1) generator in parentheses. Thus, the 248 of E_8 naturally contains three 27's of E_6 and three 16's of SO(10), and so can unify the three families into a single representation. The point that E_8 Yang-Mills theory can contain SU(3) as a family group was made by Bars and Günaydin [4] and was emphasized in Slansky's review [7] and also by Barr [10]. In the different dynamical context of supersymmetric nonlinear σ models, the point that E_8 can naturally lead to three families was made in papers of Ong [11], Buchmüller and Napoly [12], Itoh, Kugo, and Kunitomo [13], and Ellwanger [14].

Despite the attractive features of automatic supersymmetry and natural inclusion of three families, the reason that E_8 has not been further pursued as a unification group is that in addition to three families, it contains three mirror families. Thus, under $E_8 \supset SU(3) \times E_6$, in addition to three 27's there are three $\overline{27}$'s, while under $E_8 \supset SU(3) \times SO(10) \times U(1)$, in addition to three 16's there are three $\overline{16}$'s. The presence of mirror families leads to potential phenomenological and theoretical difficulties.

The phenomenological difficulty is that since the masses of mirror families break $SU(2) \times U(1)$ electroweak symmetry, they must be of order the electroweak symmetry breaking scale, at most a few hundred GeV. Hence, although they need not have been produced in current accelerator experiments, they will manifest themselves indirectly through electroweak radiative corrections, and should be copiously produced once the large hadron collider (LHC) is operative. A detailed review of experimental signatures for mirror fermions has been given by Maalampi and Roos [15] (see also Montvay [16] and Triantaphyllou [17]; the latter also has discussed a possible role for mirror fermions in dynamical electroweak symmetry breaking.) One potential phenomenological objection to mirror fermions is that under the assumptions that their masses are much larger than the Z boson mass and are degenerate within righthanded doublets, each family of mirror fermions would make a contribution of $2/(3\pi)$ to the electroweak S parameter (see Peskin and Takeuchi [18], and the review by Erler and Langacker [19]), in strong disagreement with experiment. However, this is not as definitive as it seems; when the degeneracy assumption is dropped the contribution of a mirror family to S can have either sign (or be zero), and recent analyses of the electroweak precision data by Novikov, Okun, Rozanov, and Vysotsky [20] and by He, Polonsky, and Su [20] conclude that additional chiral generations are not currently excluded, with Novikov et al. finding a chi-squared minimum between one and two extra generations. An second analysis by Choudhury, Tait and Wagner [21], focusing on additional mirror bottom quarks, also finds an improved fit to the electroweak data. In an E_8 unification model, each fermion family is accompanied by a family of vector gluons which will also, at least [22] in the case of nonmass degenerate vector doublets, make contributions to the S parameter, and therefore will further weaken the constraints coming from the electroweak data. Thus the mirror structure predicted by an E_8 model may well be consistent with current data.

The theoretical difficulty is that under the most attractive channel rule, a theory with equal numbers of ordinary and mirror families would in general be expected to form a chiral symmetry breaking family-mirror family condensate, and so one would naively expect no low energy families to survive in the low energy effective action. This expectation has become virtual dogma in model building, where it is usually stated that in a model with n_f families and $n_{\bar{f}}$ mirror families, the difference $n_f - n_{\bar{f}}$ gives the number of surviving low energy families if positive, and the number of surviving low energy mirror families if negative. However, this dogma must be treated with some skepticism, since there are known instances (see Seiberg [23] and Holdom and Roux [24]) where the most attractive channel rule breaks down. In the specific context of supersymmetric E_8 Yang-Mills theory, the issue is whether an E_8 singlet gluino condensate $\langle \lambda \lambda \rangle$ forms, as suggested by an effective action argument of Veneziano and Yankielowicz [25]. The presence of such a condensate would prevent the appearance of fermions (which are the E_8 gluinos) in the low-energy effective action.

Recently, in a very interesting paper, Kovner and Shifman [26] have argued that the Veneziano-Yankielowicz effective action must be modified so as to explicitly exhibit the $Z_{2T(G)}$ discrete chiral symmetry, which is the nonanomalous remnant of the anomalous U(1)axial symmetry generated by phase rotations of the gluino fields. (Here $\ell = 2T(g)$ is the Dynkin index of the adjoint representation, which equals 60 for the adjoint 248 representation of E_8 .) They show that there is a simple modification of the Veneziano-Yankielowicz action which has the required discrete symmetry, and that this action predicts that there is a phase in which the discrete chiral symmetry is unbroken, and thus in which the usual singlet gluino condensate does not develop. While the two independent arguments advanced by Kovner and Shifman to support their suggestion for a new phase are now discounted (one of these was based on problems with the Witten index for certain groups, which are now resolved [27]; the other on a mismatch between the strong and weak coupling instanton calculations of the gluino condensate, which has been given another explanation [28]), their effective action argument for the existence of a phase without a gluino condensate is still viable, and their conjecture of a new phase for supersymmetric gluodynamics is open, although still debated [29, 30].

In particular, although Csáki and Murayama [30] have used discrete anomaly matching to argue against the Kovner-Shifman vacuum, their argument assumes that the ground state spectrum consists of hypercolor (here E_8) singlets. Thus it does not rule out the possibility that the Kovner-Shifman vacuum is in a trivial, deconfined phase with the same particle spectrum as the starting E_8 gauge theory, before symmetry breaking arising from perturbations to the SUSY gluodynamics structure is taken into account. A deconfined phase would obey all anomaly matching constraints, and even if not generic for SUSY Yang-Mills gluodynamics, its presence just in special cases including E_8 would suffice for the arguments we are making.

If supersymmetric Yang-Mills for the E_8 group is in the Kovner-Shifman vacuum, then the principal theoretical objection to E_8 as a unification group disappears, since the theory in isolation would remain a supersymmetric theory (as assured by Witten index arguments [27,31]) with massless gluinos in the Kovner-Shifman phase. Of course, to get a realistic theory breaking of both E_8 symmetry and supersymmetry is needed. As noted by Shifman and Vainshtain [29] (in the course of a discussion of the Witten index, but their remark is more generally relevant) the Kovner-Shifman vacuum is "potentially unstable under various deformations." One obvious deformation that could be relevant is the embedding of supersymmetric E_8 in supergravity. When the gravitino and graviton fields are integrated out at tree level, one obtains [32] a supersymmetric four-gluino effective action that could be the trigger for dynamical symmetry breaking of either or both the E_8 internal symmetry and supersymmetry. Supersymmetry breaking could also arise from supersymmetry breaking in another sector of the theory (such as the second E_8 expected in string theory) communicated by the supergravity interaction between the two; a general review of this approach is given in Weinberg [33], and an application of gravity-mediated supersymmetry breaking to sequential breaking of E_8 to E_6 and then to SO(10) is discussed by Mahapatra and Deo [34].

As noted by Bars and Günaydin [4], an E_8 unification theory cannot have elementary Higgs scalars without losing the property of asymptotic freedom, because the Dynkin index of the smallest candidate Higgs representation (the 3875) is already too large. Hence in an asymptotically free E_8 theory, all symmetry breaking (other than that communicated

by gravity mediation from another sector) must be dynamical, through the formation of suitable condensates of the gluinos (and of gluinos and gluons as well, if condensate formation preserves supersymmetry). Chiral symmetry breaking by condensate formation was reviewed some time ago by Peskin [35], and recently there has been much interest in the role of nonsinglet condensates that break gauge symmetry, in the context of "color superconductivity" in high density QCD [36]. In order to give the mirror fermions larger masses than the top quark, there must be a condensate that introduces an asymmetry between the fermions and their mirror partners. One candidate arises from the fact that in $SU(3) \times E_6$ one has $(\overline{3},\overline{27})\times(\overline{3},\overline{27})\supset(6_s,27_s)$. Since under the decomposition $E_6\supset SO(10)\times U(1)$ one has $27\supset$ 1(4), a gluino-gluino condensate with nonvanishing vacuum expectation of the 1(4) would preserve SO(10) symmetry, while breaking the U(1) factor and introducing an asymmetry between the three fermion families and their mirror families. Moreover, since under the family group decomposition $SU(3) \supset SU(2) \times U(1)$ the 6_s contains a singlet of SU(2), this expectation would split two degenerate families apart from a third, approximating what is observed. (From the viewpoint of E_8 , the condensate we are proposing is contained in $3875_s \subset 248 \times 248$, which is the second most attractive symmetric channel according to the most attractive channel rule.) To break SO(10) down to the standard model further condensates would be needed; we note that all of the Higgs representations used in models for the breaking of SO(10) unification are contained in the 248×248 of E_8 , and so could be generated by the formation of non-singlet gluino-gluino condensates. As a final remark on symmetry breaking, we mention that a much studied alternative to dynamical generation of Higgs condensates is their generation by dimensional reduction from a higher dimensional gauge theory; for a recent discussion of this mechanism as applied to E_8 and three family

unification, in the context dimensional reduction over coset spaces, see Manousselis and Zoupanos [37].

The phenomenology of a supersymmetric E_8 -based grand unification and family unification model will differ significantly from that expected in the minimal supersymmetric standard model (MSSM) and its extensions. As in the MSSM, the superpartners for the gauge bosons in the E_8 model are spin-1/2 fermions, and *R*-parity conservation [38] implies that the lightest superpartner will be stable. However, in the E_8 theory, in addition to there being mirror fermions, the superpartners for the quarks and leptons are vectors rather than scalars. Thus there are potentially observable signatures for E_8 unification at the LHC and other future facilities.

Acknowledgments

This work was supported in part by the Department of Energy under Grant #DE– FG02–90ER40542. I wish to thank Csaba Csáki, Maurice Goldhaber, Paul Langacker, Michael Peskin, Nathan Seiberg, Jacob Sonnenschein, Edward Witten, and Ren-Jie Zhang for helpful discussions or email correspondence.

References

- [1] Three prominent types of ideas that have been advanced to explain family structure are (1) composite models, in which the quarks and leptons are not elementary, and the different families are different internal symmetry states of their constituents, (2) string-inspired models, in which the different families correspond to string excitations with different topological quantum numbers of the background manifold on which the string spacetime is compactified, and (3) as discussed here, group theoretic family unification models, in which the quarks and leptons are elementary (at least below the Planck scale) and the family structure arises because the fermions lie in a large group representation. For early discussions of family groups, see P. Ramond, 'The Family Group in Grand Unified Theories', invited talk at the Sanibel Symposia, 1979 (reissued as ArXiv: hep-ph/9809459); and A. Zee, Unity of Forces in the Universe Vol. I (World Scientific, 1982), Sec. VIII. A pedagogical account of composite and string-inspired models is given in Chapt. 8 and Chapt. 17, respectively, of R. N. Mohapatra, Unification and Supersymmetry, Second Ed. (Springer, 1992). For a recent discussion of empirical systematics of the family problem, and further references, see M. Goldhaber, Proc. Nat. Acad. Sci. USA 99 (2002) 33 and arXiv: hep-ph/0201208.
- [2] L. Brink, J. H. Schwarz and J. Scherk, Nucl. Phys. B121 (1977), 77, show that the Fierz identity needed to verify supersymmetry in four dimensional Yang-Mills theory holds for Weyl as well as Majorana fermions. One can also directly transform supersymmetric Yang-Mills theory from Majorana to Weyl fermions, using the Pauli-

Gürsey transformation (which is a special instance of a Bogoliubov transformation);
see W. Pauli, Nuovo Cimento 6 (1957) 204 and F. Gürsey, Nuovo Cimento 7 (1958)
411; R. E. Marshak, Riazuddin and C. P. Ryan, *Theory of Weak Interactions in Particle Physics* (Wiley-Interscience, 1969), pp. 71-72.

- [3] N. S. Baaklini, Phys. Lett. 91B (1980) 376.
- [4] I. Bars and M. Günaydin, Phys. Rev. Lett. 45 (1980) 859.
- [5] S. E. Konshtein and E. S. Fradkin, Pis'ma Zh. Eksp. Teor. Fiz. 32 (1980) 575
 [English translation: JETP Lett. 32 (1981) 557].
- [6] M. Koca, Phys. Lett. 107B (1981) 73.
- [7] R. Slansky, Phys. Reports **79** (1981) 1.
- [8] For an introduction to SO(10) unification and reprints of seminal papers, see A. Zee,
 Ref. [1], Sec. IV. For a more detailed pedagogical account, see R. N. Mohapatra,
 Ref. [1], Chapt. 7.
- [9] For an introduction to E₆ unification and reprints of seminal papers, see A. Zee, Ref.
 [1], Sec. V. For a review of E₆ phenomenology, see J. L. Hewett ad T. G. Rizzo, Phys. Rep. 183 (1989) 193.
- [10] S. M. Barr, Phys. Rev. D37 (1988) 204.
- [11] C. L. Ong, Phys. Rev. D31 (1985) 3271.
- [12] W. Buchmüller and O. Napoly, Phys. Lett. 163B (1985) 161.

- [13] K. Itoh, T. Kugo and H. Kunitomo, Prog. Theor. Phys. 75 (1986) 386.
- [14] U. Ellwanger, Nucl. Phys. B356 (1991) 46.
- [15] J. Maalampi and M. Roos, Phys. Reports 186 (1990) 53.
- [16] I. Montvay, Phys. Lett. 205B, 315 (1988).
- [17] G. Triantaphyllou, Int. Journ. Mod. Phys. A (2000) 265; J. Phys. G: Nucl. Part. Phys. 26 (2000) 99. These papers focus on the phenomenology of mirrors. For a discussion of dynamical symmetry breaking by mirrors with new strong interactions, see G. Triantaphyllou, 'Dynamical Symmetry Breaking with Mirror Fermions', talk at the Corfu Summer Institute on Elementary Particle Physics, 1998, arXiv: hepph/9908251; Mod. Phys. Lett. A16 (2001) 53; 'The Weak Scale: Dynamical Determination Versus Accidental Stabilization', talk at the Corfu Summer Institute on Elementary Particle Physics, 2001, arXiv: hep-ph/0109023.
- [18] M. E. Peskin and T. Takeuchi, Phys. Rev. D 46 (1992) 381.
- [19] J. Erler and P. Langacker, 'Electroweak Model and Constraints on New Physics', in Particle Data Group, *Review of Particle Physics*, 2000, 2001 (available online at http://pdg.lbl.gov/).
- [20] V. A. Novikov, L. B. Okun, A. N. Rozanov and M. I. Vysotsky, 'Extra generations and discrepancies of electroweak precision data', arXiv: hep-ph/0111028; H.-J. He, N. Polonsky, and S. Su, 'Extra Families, Higgs Spectrum and Oblique Corrections', arXiv: hep-ph/0102144.

- [21] D. Choudhury, T. M. P. Tait and C. E. M. Wagner, 'Beautiful Mirrors and Precision Electroweak Data', arXiv: hep-ph/0109097.
- [22] R.-J. Zhang and S. L. Adler, unpublished. Degenerate vector doublets make a vanishing contribution to the S parameter, but nondegenerate doublets should contribute a logarithm similar in structure, but with a different numerical coefficient that has not been computed, to that found in the fermion case.
- [23] N. Seiberg, 'The Power of Holomorphy Exact Results in 4D SUSY Field Theories', arXiv: hep-th/9408013, Secs. 3 and 5. See also K. Intriligator and N. Seiberg, 'Lectures on Supersymmetric Gauge Theories and Electric-Magnetic Duality', arXiv: hep-th/9509066, Secs. 4.3 and 5.4; to appear in the Proceedings of the Trieste '95 spring school, TASI '95, Trieste '95 summer school, and Cargese '95 summer school.
- [24] B. Holdom and F. S. Roux, Phys. Rev. D59 (1999) 015006.
- [25] G. Veneziano and S. Yankielowicz, Phys. Lett. 113B (1982) 231; see also T. R.
 Taylor, G. Veneziano and S. Yankielowicz, Nucl. Phys. B218 (1983) 493; H. P.
 Nilles, Phys. Lett. 112B (1982) 455 and Phys. Lett. 115B (1982) 193.
- [26] A. Kovner and M. Shifman, Phys. Rev. D56 (1997) 2396.
- [27] E. Witten, 'Supersymmetric Index in Four-Dimensional Gauge Theories', arXiv: hepth/0006010.
- [28] T. J. Hollowood, V. V. Khoze, W. Lee and M. P. Mattis, Nucl. Phys. B 570 (2000) 241.

- [29] I. I. Kogan, A. Kovner and M. Shifman, Phys. Rev. D57 (1998) 5195; M. Shifman and A. Vainshtein, 'Instantons Versus Supersymmetry: Fifteen Years Later', arXiv: hep-th/9902018; A. Ritz and A. Vainshtein, Nucl. Phys. B 566 (2000) 311; Y. Shadmi and Y. Shirman, Rev. Mod. Phys. 72 (2000) 25, note 11 on p. 33; V. S. Kaplunovsky, J. Sonnenschein, and S. Yankielowicz, Nucl. Phys. B552 (1999) 209.
- [30] C. Csáki and H. Murayama, Nucl. Phys. B515 (1998) 114.
- [31] For a survey of the Witten index analysis of supersymmetric gauge theories, see Secs.
 29.1 and 29.4 of S. Weinberg *The Quantum Theory of Fields*, Vol. III: Supersymmetry (Cambridge University Press, 2000).
- [32] S. L. Adler, Ann. Phys. 290 (11) 2001.
- [33] For a survey of gravity mediated supersymmetry breaking, see Sec. 31.7 of Weinberg, Ref. [31].
- [34] S. Mahapatra and B. B. Deo, Phys. Rev. D38 (1988) 3554.
- [35] M. E. Peskin, 'Chiral Symmetry and Chiral Symmetry Breaking', in *Recent Advances in Field Theory and Statistical Mechanics* (Les Houches, 1982), J.-B. Zuber and R. Stora, eds. (North-Holland, 1984).
- [36] For a recent review, see M. Alford, 'Color Superconducting Quark Matter', in Annual Review of Nuclear and Particle Science, V. 51 (Annual Reviews, 2001).
- [37] P. Manousselis and G. Zoupanos, 'Dimensional Reduction over Coset Spaces and Supersymmetry Breaking', arXiv: hep-ph/0111125.

[38] The *R*-parity is defined by $\Pi_R = (-1)^F (-1)^{3(B-L)}$, with $(-1)^F$ the fermion parity which is +1 for all bosons (irrespective of whether vector or scalar) and -1 for all fermions. Hence in the E_8 theory, as in the MSSM, all standard model particles have R = 1 and all superpartners have R = -1. For further details and references, see Weinberg, Ref. [31], p. 184 and Mohapatra, Ref. [1], p. 277.