

Scalar mesons and the muon anomaly

Stephan Narison^a

^a Laboratoire de Physique Mathématique et Théorique, Université de Montpellier II, Case 070, Place Eugène Bataillon, 34095 - Montpellier Cedex 05, France.
E-mail: qcd@lpm.univ-montp2.fr

We evaluate systematically some contributions of the QCD scalar mesons, including radiative decay-productions, not considered with a better attention until now in the evaluation of the hadronic contributions to the muon anomaly. The sum of the scalar contributions to be added to the existing Standard Model predictions a_μ^{SM} are estimated to be $a_\mu^S = 11.75(8.25) \times 10^{-10}$, where the errors are dominated by the ones from the experimental widths of these scalar mesons. This result suggests that the value of a_μ^{SM} and its errors might have been underestimated in previous works. The inclusion of these new effects leads to a perfect agreement (0.1σ) of the measured value a_μ^{exp} and a_μ^{SM} from τ -decay and implies a 1.7σ discrepancy between a_μ^{exp} and a_μ^{SM} from $e^+e^- \rightarrow$ hadrons data. More refined unbiased theoretical estimates require improved measurements of the scalar meson masses and widths. The impact of our results to a_μ^{SM} is summarized in the conclusions.

arXiv:hep-ph/0303004v1 2 Mar 2003

1. INTRODUCTION

Improved accurate experimental measurement [1] and recent theoretical estimates of the muon anomaly [2,3,4,5] are now available. The theoretical accuracy in [2,3] ¹ is mainly attributed to the use of the new CMD-2 [6] around the ρ mass and on BES data [7] around the J/Ψ region. The impact of the former data on the determination of the muon anomaly is intuitively more important due to the low-energy dominance of the anomaly kernel function [8,9]. However, what is more intriguing within this increasing precision is the discrepancy between the results from e^+e^- and τ -decay data [2,3], which was not the case in the previous determinations using preliminary data [4,5]. In this short note, we study some other sources of contributions, not considered with a better attention until now, from the scalar mesons. These scalar ($\bar{q}q$, gluonia,...) are conceptually fundamental consequences of QCD. Though their existence is not precisely confirmed, there are increasing evidences of their findings in different e^+e^- and hadronic experiments [10].

2. ISOSCALAR SCALAR MESONS

The interest in these $I = 0$ scalar mesons are that they cannot be obtained from usual ChPT approaches. They are related to the QCD scale anomaly:

$$\theta_\mu^\mu = \frac{1}{4}\beta(\alpha_s)G^2 + \sum_i [1 + \gamma_m(\alpha_s)]m_i\bar{\psi}_i\psi_i, \quad (1)$$

where $G_{\mu\nu}^a$ is the gluon field strengths, ψ_i is the quark field; $\beta(\alpha_s)$ and $\gamma_m(\alpha_s)$ are respectively the QCD β -function and quark mass-anomalous dimension. In this case, arguments based on $SU(2)$ symmetry or its violation used to estimate some processes like e.g. radiative processes cannot be applied [11]. Using QCD spectral sum rules (QSSR) [12,13] and low-energy theorems (LET) for estimating the mass and its width, it has been shown in [14,15] that the wide $\sigma(0.60)$ meson is the best candidate meson (gluonium) associated to the previous interpolating current (see also [16]). Its mass is due to the gluon component and its large width into $\pi\pi$ is due to a large violation of the OZI rule ² (analogue of the η' -meson of the $U(1)_A$ sector [17]). The observed $\sigma(0.60)$ and $f_0(0.98)$ can be explained by a “max-

¹Though agreeing in the total sum, the results from these two estimates differ in each energy region.

²This feature also implies that lattice calculations of the gluonium mass in pure Yang-Mills or quenched approximation are bad approximations and might miss this “unusual” glueball.

imal quarkonium-gluonium mixing scheme" [14,15]. In the following, we study the σ and other scalar mesons contributions to the muon anomaly.

3. THE $e^+e^- \rightarrow S\gamma$ PROCESSES

Using analytic properties of the photon propagator, the general contribution to the muon anomaly from the process:

$$e^+e^- \rightarrow \gamma^* \rightarrow \text{hadrons} \quad (2)$$

can be written in a closed form as [8]:

$$a_l^{\text{had}}(l.o) = \frac{1}{4\pi^3} \int_{4m_\pi^2}^{\infty} dt K_l(t) \sigma_H(t), \quad (3)$$

where:

• $K_l(t \geq 0)$ is the QED kernel function [9]:

$$K_l(t) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (t/m_l^2)(1-x)}, \quad (4)$$

with the analytic form:

$$K_l(t \geq 4m_l^2) = z_l^2 \left(1 - \frac{z_l^2}{2}\right) + (1+z_l)^2 \times \left(1 + \frac{1}{z_l^2}\right) \left[\log(1+z_l) - z_l + \frac{z_l^2}{2}\right] + \left(\frac{1+z_l}{1-z_l}\right) z_l^2 \log z_l, \quad (5)$$

with:

$$y_l = \frac{t}{4m_l^2}, \quad z_l = \frac{(1-v_l)}{(1+v_l)} \quad \text{and} \quad v_l = \sqrt{1 - \frac{4m_l^2}{t}}. \quad (6)$$

$K_l(t)$ is a monotonically decreasing function of t . For large t , it behaves as:

$$K_l(t > m_l^2) \simeq \frac{m_l^2}{3t}, \quad (7)$$

which will be an useful approximation for the analysis in the large t regime. Such properties then emphasize the importance of the low-energy contribution to $a_l^{\text{had}}(l.o)$ ($l \equiv e, \mu$), where the QCD analytic calculations cannot be applied.

• $\sigma_H(t) \equiv \sigma(e^+e^- \rightarrow \text{hadrons})$ is the $e^+e^- \rightarrow \text{hadrons}$ total cross-section which can be related to the hadronic two-point spectral function $\text{Im}\Pi(t)_{em}$ through the optical theorem:

$$R_{e^+e^-} \equiv \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = 12\pi \text{Im}\Pi(t)_{em}, \quad (8)$$

where:

$$\sigma(e^+e^- \rightarrow \mu^+\mu^-) = \frac{4\pi\alpha^2}{3t}. \quad (9)$$

Here,

$$\begin{aligned} \Pi_{em}^{\mu\nu} &\equiv i \int d^4x e^{iqx} \langle 0 | \mathcal{T} J_{em}^\mu(x) (J_{em}^\nu(x))^\dagger | 0 \rangle \\ &= -(g^{\mu\nu} q^2 - q^\mu q^\nu) \Pi_{em}(q^2) \end{aligned} \quad (10)$$

is the correlator built from the local electromagnetic current:

$$J_{em}^\mu(x) = \frac{2}{3} \bar{u} \gamma^\mu u - \frac{1}{3} \bar{d} \gamma^\mu d - \frac{1}{3} \bar{s} \gamma^\mu s + \dots \quad (11)$$

Using Vector Meson Dominance Model (VDM) in a Narrow Width Approximation (NWA), one obtains [18]:

$$a_\mu^{\text{VDM}}(l.o) \simeq \frac{3}{\pi} K \left(\frac{M_V^2}{m_l^2} \right) \frac{\Gamma_{V \rightarrow e^+e^-}}{M_V} \frac{\Gamma_{V \rightarrow \gamma X}}{\Gamma_{V \rightarrow \text{all}}}. \quad (12)$$

The $V \rightarrow \gamma X$ ($X \equiv \sigma, f_0$) coupling has been estimated from the $X\gamma\gamma$ one given in [14,15] and in the PDG data [19] and using the VDM relation:

$$g_{X\gamma V} \simeq \frac{\sqrt{2}\gamma_V}{e} g_{X\gamma\gamma}, \quad (13)$$

where e is the electric charge and we use the normalization $\gamma_\rho \simeq 2.51 \pm 0.02$. For the scalar mesons, we use:

$$\begin{aligned} M_\sigma &\simeq (0.6 \sim 0.8) \text{ GeV}, \\ \Gamma_{\sigma \rightarrow \gamma\gamma} &\simeq (0.2 \sim 5.) \text{ keV}, \\ \Gamma_{f_0(1.4) \rightarrow \gamma\gamma} &\simeq (0.7 \sim 5.) \text{ keV}, \end{aligned} \quad (14)$$

where the low values of the widths come from [14,15] and the high ones from [19,20]. The ones of the other ω and ϕ radiative widths come from PDG [19]. We also use in MeV units [19]:

$$\begin{aligned} \Gamma_{\omega(1.42) \rightarrow e^+e^-} &\simeq 0.08 \times 10^{-3}, \quad \Gamma_{\text{tot}} \simeq 174(59) \\ \Gamma_{\rho(1.45) \rightarrow e^+e^-} &\simeq 0.44 \times 10^{-3}, \quad \Gamma_{\text{tot}} \simeq 310(60) \\ \Gamma_{\phi(1.68) \rightarrow e^+e^-} &\simeq 0.48 \times 10^{-3}, \quad \Gamma_{\text{tot}} \simeq 150(50), \end{aligned} \quad (15)$$

We deduce the results in Table 1, where the lower value corresponds to the set $(M_\sigma, \Gamma_{\sigma \rightarrow \gamma\gamma}) = (0.8, 0.2)$ and the upper value to the set $(0.6, 5.)$. It is clear from Table 1 that the results are very sensitive to the value of the σ mass and $\gamma\gamma$ width. Though, in the "maximal quark-gluonium mixing scheme" of the σ meson, one favours a small $\gamma\gamma$ width [14,15], the data [19,20] and some other QCD models still allow higher values. For a conservative and unbiased estimate, we translate the total sum in Table 1 into:

$$a_\mu^{S\gamma} = 6.45(6.13) \times 10^{-10}. \quad (16)$$

In order to include this result into the one in [2], and for avoiding an eventual double counting, we "de-rescale" their result on the top of the ω by 0.888 and on the ϕ by the factor 0.984 used there for taking into account the missing modes. In this way, we obtain the corresponding "pure" hadronic decay:

$$a_\mu(\omega + \phi \rightarrow 3\pi + K\bar{K}) = 67.04(1.50) \times 10^{-10}, \quad (17)$$

Using the ω and ϕ into $\eta\gamma$ and $\pi^0\gamma$ radiative decay widths from [19], we deduce:

$$a_\mu(\omega + \phi \rightarrow \eta\gamma + \pi^0\gamma) = 4.36(0.16) \times 10^{-10}. \quad (18)$$

Table 1

 $e^+e^- \rightarrow \text{scalar}+\gamma$ contributions to a_μ^{had} *).

Processes	$a_\mu^{\text{had}} \times 10^{10}$
$\rho \rightarrow \sigma\gamma$	0. \sim 0.05
$\omega \rightarrow \sigma\gamma$	0. \sim 0.07
$\phi \rightarrow \sigma\gamma$	0.03 \sim 10.1
$\phi \rightarrow f_0(.98)\gamma$	0.01 \pm 0.00
$\phi \rightarrow a_0(.98)\gamma$	0.03 \pm 0.01
$\rho(1.45) \rightarrow \sigma\gamma$	0.01 \sim 0.42
$\omega(1.42) \rightarrow \sigma\gamma$	0.10 \sim 0.71
$\phi(1.68) \rightarrow \sigma\gamma$	0.14 \sim 1.17
$\phi(1.68) \rightarrow f_0(1.4)\gamma$	–
Total	0.32 \sim 12.57

*) For completeness, we also include the contribution of the isovector $a_0(980)$.

Taking into account these different effects, we conclude from this analysis that the inclusion of these new radiative decays increases the results in [2] by:

$$\Delta a_\mu^{X\gamma} = 6.1(6.1) \times 10^{-10}, \quad (19)$$

and may indicate that the scaling factors used in [2] are not sufficient for taking into account the different missing modes of the ω and ϕ mesons. Therefore, adding our result in Eq. (19), to the recent estimate of $a_\mu^{\text{had}}(l.o)$ in [2,3,4,5], the lowest order hadronic contributions to a_μ becomes ³ in units of 10^{-10} :


$$\begin{aligned} a_\mu^{\text{had}}(l.o) &= 712.4(6.8)(6.1) [\tau], \\ &= 689.7(6.4)(6.1) [e^+e^-], \end{aligned} \quad (20)$$

where the first error is the ones from [2,3,4], and the second one comes from the present analysis. Our results suggest that the estimate of $a_\mu^{\text{had}}(l.o)$ and its errors might have been underestimated in the previous determinations.

4. NEW CONTRIBUTIONS OF THE SCALAR MESON TO a_μ

Here, we study the effect of scalar mesons via a Higgs-like triangle diagram shown in Fig. 1. This new hadronic contribution to a_μ^{had} cannot be estimated using the usual electromagnetic spectral function associated to one virtual photon. The evaluation

³For the τ -decay data, we use the average of the results (in units of 10^{-10}) 709.0(5.9) [2] and 703.6(7.6) [4]. For the e^+e^- data, we use the average of the most recent estimates 684.0(6.5) [2] and 683.1(6.2) [3], which are however lower by about 1.4σ than the previous estimates in [4,5] based on preliminary data.

Figure 1. Scalar meson contribution to a_l .

of this diagram gives the contribution (see e.g. [21]):

$$a_\mu^{Su} = \frac{|g_F|^2}{16\pi^2} 2 \int_0^1 dx \frac{x^2(2-x)}{x^2 + (M_S^2/m_\mu^2)(1-x)}, \quad (21)$$

where M_S is the scalar mass and g_F is the “effective” Sl^+l^- coupling. However, the coupling of these scalar mesons to e^+e^- is not known, and in the case of the isoscalar, the naïve argument based on chiral symmetry (Higgs-type coupling) may not be applied due to the presence of gluon fields in the $U(1)_V$ dilaton current given in Eq. (1). Note that these scalar mesons might also be produced in the s -channel in e^+e^- experiments, most probably via two-photon exchange, which should differ from the usual spectral function of a one photon exchange used to estimate $a_\mu^{\text{had}}(l.o)$, and therefore prevent from some eventual double counting of the $e^+e^- \rightarrow \text{hadrons}$ data. However, we are not also aware of existing analyses of the angular distribution which can differentiate a scalar from a vector meson in the low-energy e^+e^- region concerned here [22], while present LEP limits on electroweak scalar particles (sleptons, squarks) given in [23] may not be applied here.. If one esti-

Table 2

Contribution of Fig. 1 to a_μ using the upper bound on the leptonic widths compiled by PDG [19].

	$a_\mu \times 10^{10}$	$\Gamma_{S \rightarrow e^+e^-}$ [eV]
$\sigma(.60)$	0. \sim 8.9	≤ 20 *)
$f_0(.98)$	0. \sim 1.1	≤ 8.4
$a_0(.98)$	0. \sim 0.2	≤ 1.5
$f_0(1.37)$	0 \sim 1.1	≤ 20
Total	0. \sim 11.4	

*) We assume that the bound for the σ is the same as the one from $f_0(1.37)$.

mates this coupling from the experimental bound on the $S \rightarrow e^+e^-$ width [19]⁴, and use the positivity of the contribution, one gets the result in Table 2, where an universal coupling to l^+l^- ($l \equiv e, \mu$) has been assumed⁵. In order to see the strength of this experimental bound, one may assume that the coupling of the scalar to l^+l^- is dominated by the one through two photons. Therefore, an *alternative rough estimate* of this effect can be obtained by relating it to the light-by-light (LL) scattering diagram contribution [25], where we obtain⁶:

$$a_\mu^{Sll}(LL) \approx e^2 a_\mu^{LL} \approx 0.1 \times 10^{-10}. \quad (22)$$

This result indicates that the present experimental upper bound might be very weak, and needs to be improved. Within the present accuracy, we adopt the most conservative final average estimate which takes into account all range given in Eqs. (22) and Table 2:

$$a_\mu^{Sll} = 5.65(5.55) \times 10^{-10}. \quad (23)$$

Adding the results in Eqs. (19) and (23), one finally deduces the sum of the additional contributions due to scalar mesons:

$$a_\mu^S = 11.75(8.25) \times 10^{-10}, \quad (24)$$

to be added to the SM predictions a_μ^{SM} .

5. CONCLUSIONS

The inclusion of the scalar meson contribution in Eq. (19) modifies the recent estimate of $a_\mu^{had}(l.o)$ from [2,3,4] into the one in Eq. (20), while the direct exchange of the scalar meson shown in Fig. 1 gives the new contribution in Eq. (23). The total sum of the scalar contributions is given in Eq. (24). Adding our result in Eqs. (20) and (23), we deduce in units of 10^{-10} :

$$\begin{aligned} a_\mu^{had} &\equiv a_\mu^{had}(l.o) + a_\mu^{Sll} \\ &= 718.1(6.8)(8.3) [\tau], \\ &= 695.3(6.4)(8.3) [e^+e^-], \end{aligned} \quad (25)$$

where the first error is the ones from [2,3,4], and the second one comes from the present analysis. Our results suggest that the value and errors of the hadronic contributions to a_μ^{SM} might have been underestimated in the previous determinations. From the above results, we deduce our final estimate of the muon anomaly in units of 10^{-10} :

$$\begin{aligned} a_\mu^{SM} &= 11\,659\,202.3(11.3) [\tau], \\ &= 11\,659\,179.9(11.1) [e^+e^-], \end{aligned} \quad (26)$$

⁴Note that the difference $a_e^{exp} - a_e^{SM} = (34.9 \pm 28.4) \times 10^{-12}$ from the electron anomaly [24] leads to a weaker upper bound.

⁵The bound is weaker for a Higgs-type coupling.

⁶We use the total LL contribution in order to take into account other mesons contributions

compared with the recent data [1]:

$$a_\mu^{exp} = 11\,659\,204.0(8.6) \times 10^{-10}. \quad (27)$$

Then, we deduce in units of 10^{-10} :

$$\begin{aligned} a_\mu^{exp} - a_\mu^{SM} &= 1.7(14.2) [\tau], \\ &= 24.1(14.0) [e^+e^-]. \end{aligned} \quad (28)$$

One can see that the inclusion of these new effects implies a perfect agreement (0.1σ) between a_μ^{exp} and a_μ^{SM} from τ -decay while the disagreement between a_μ^{SM} from e^+e^- and a_μ^{exp} becomes 1.7σ instead of 3σ obtained in previous works [2,3]. More refined theoretical estimates require improved measurements of the masses and widths of the scalar mesons, which are also necessary for making a progress in our QCD understanding of their nature. In addition to some problems mentioned in [2] (absolute normalization of the cross-section, radiative corrections, $SU(2)$ breakings,...), the discrepancy between the τ -decay and e^+e^- data might also be due to some interferences of isoscalar $I = 0$ mesons (scalar or/and meson with exotic quantum numbers, e.g. $J^{PC} = 1^{-+}$ not produced in τ -decay) with vector mesons. In this respect, we expect that the result from e^+e^- is less cleaner than the one from τ -decay. We plan to come back to this point in a future work.

ACKNOWLEDGEMENTS

Stimulating exchanges with William Marciano and Simon Eidelman are gratefully acknowledged.

REFERENCES

1. C.S. Ozben et al., hep-ex/0211044; G.W. Bennett et al., *Phys. Rev. Lett.* **89** (2002) 1001804.
2. M. Davier et al., hep-ph/0208177.
3. K. Hagiwara et al., hep-ph/0209187.
4. S. Narison, *Phys. Lett.* **B 513** (2001) 53; Erratum **526** (2002) 414; hep-ph/0108065; hep-ph/0203053.
5. J.F. de Trocóniz and J.F. Ynduráin, *Phys. Rev.* **D 65** (2001) 093001.
6. The CMD-2 collaboration, R.R. Akhmetshin et al., *Phys. Lett.* **B 527** (2002) 161.
7. The BES collaboration, J.Z. Bai et al., *Phys. Rev. Lett.* **84** (200) 594; **88** (2002) 101802.
8. C. Bouchiat and L. Michel, *J. Phys. Radium* **22** (1961) 121; M. Gourdin and E. de Rafael, *Nucl. Phys.* **B 10** (1969) 667.
9. B.E. Lautrup and E. de Rafael, *Phys. Rev.* **174** (1968) 1835.
10. L. Montanet, *Nucl. Phys. (Proc. Suppl.)* **B 86** (2000) 381; U. Gastaldi, *Nucl. Phys. (Proc. Suppl.)* **B 96** (2001) 234; I. Bediaga, *Nucl. Phys. (Proc. Suppl.)* **B 96** (2001) 225; *Possible existence of σ meson and its implications*, *Kyoto, KEK Proceedings 2000-4*, ed. Ishida et al.

11. V.A. Novikov et al., *Nucl. Phys.* **B 165** (1980) 67; J.Ellis and M.S. Chanowitz, *Phys. Lett.* **B 40** (1972) 397; *Phys. Rev.* **D 7** (1973) 2490; R.J. Crewther, *Phys. Rev. Lett.* **28** (1972) 1421.
12. M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Nucl. Phys.* **B 147** (1979) 385, 448.
13. S. Narison, *QCD as a Theory of Hadrons Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol.* **17** (2002) 1 [hep-ph/0205006]; *QCD Spectral Sum Rules. World Sci. Lect. Notes Phys.* **26** (1989) 1.
14. S. Narison and G. Veneziano, *Int. Mod. Phys.* **A 4** (1989) 2751; A. Bramon and S. Narison, *Mod. Phys. Lett.* **A4** (1989) 1113.
15. S. Narison, *Nucl. Phys.* **B 509** (1998) 312; *Nucl. Phys.* **A 675** (2000) 54c; *Nucl. Phys. Proc. Suppl.* **96** (2001) 244; hep-ph/0009108; hep-ph/08081; H.G. Dosch and S. Narison, hep-ph/0208271.
16. W. Ochs and P. Minkowski, hep-ph/0209225; hep-ph/0209223; J. Ellis and J. Lanik, *Phys. Lett.* **B 150** (1985) 289.
17. E. Witten, *Nucl. Phys.* **B 156** (1979) 269; G. Veneziano, *Nucl. Phys.* **B 159** (1979) 213; G.M. Shore, S. Narison and G. Veneziano, *Nucl. Phys.* **B 546** (1999) 235; *Nucl. Phys.* **B 433** (1995) 209.
18. S. Narison, *Thèse de Doctorat 3ème cycle*, Marseille (1976).
19. PDG02, K. Hagiwara et al., *Phys. Rev.* **D 66** (2002) 010001 and references quoted therein.
20. G. Mennessier, *Z. Phys.* **C 16** (1983) 241 and private communications.
21. J. Calmet et al., *Rev. Mod. Phys.* **49** (1977) 21.
22. Private communications from Jacques Layssac and Fernand Michel Renard.
23. L3 collaboration, M. Acciarri et al., *Phys. Lett.* **B 414** (1997) 373. I thank Jean Loïc Kneur for bringing this paper to my attention.
24. W.J. Marciano and T. Kinoshita, *World Sci. Advanced Series on Directions in High Energy Physics*, **Vol 7** (1990); A. Czarnecki and W.J. Marciano, *Nucl. Phys. (Proc. Suppl.)* **B 76** (1999) 245.
25. M. Knecht and A. Niffeler, *Phys. Rev.* **D 65** (2002) 073034; M. Knecht et al., *Phys. Rev. Lett.* **88** (2002) 071802; M. Hayakawa and T. Kinoshita, hep-ph/0112102; J. Bijnens et al., *Nucl. Phys.* **B 626** (2002) 410.