

In-medium Yang-Mills equations: a derivation and canonical quantization

M. K. Djongolov¹⁾, S.Pisov²⁾, V.Rizov²⁾

1) Department of Physics and Astronomy, University of Tennessee,

e-mail: martin@spock.phys.utk.edu

2) University of Sofia, Faculty of Physics,

e-mail: pisov@phys.uni-sofia.bg; rizov@phys.uni-sofia.bg

Abstract. The equations for Yang-Mills field in a medium are derived in a linear approximation with respect to the gauge coupling parameter and the external field. The obtained equations closely resemble the macroscopic Maxwell equations. A canonical quantization is performed for a family of Fermi-like gauges in the case of constant and diagonal (in the group indices) tensors of electric permittivity and magnetic permeability. The physical subspace is defined and the gauge field propagator is evaluated for a particular choice of the gauge.

1. Introduction

In some developments of the Yang-Mills theories one arrives at situations when the encountered states can be effectively described as media, possessing nonabelian polarization properties. We list a few of these examples for sake of brevity.

Among the first prototypes for a nonabelian polarizable medium one may take the QCD vacuum state in the aspect considered by Savvidy [1], Fukuda and Kazama [2], Adler [3], Nielsen and Olesen [4]. In [5] and [6] Baker and Zachariasen used the Yang-Mills vacuum as a polarizable medium in order to derive a confining potential. They also solved the Dyson equations for the gluon propagator $\Delta_{\mu\nu}(q)$ for $q^2 \rightarrow 0$ evaluating the color susceptibility of the vacuum in the weak field approximation. The polarization properties of the nonabelian $SU(2)$ vacuum were exploited in [7, 8] for an approximate classical description of some nonperturbative effects such as solutions corresponding to nonvanishing gauge field "condensates".

Another and more recent set of examples comes from the theoretical study of some not yet confirmed states of the hadron matter - hadron matter at very high densities and quark-gluon plasma (QGP) at very high temperatures where one expects the quarks and gluons to be in a deconfined phase. In a systematic study performed in the framework of kinetic theory the color conductivity and other transport coefficients of nonabelian plasma were derived by Heinz in [9, 10] in the linear response approximation. The response of QGP beyond the linear approximation was considered in [11] yielding a color dielectric susceptibility modified by the nonlinear corrections. Typical nonabelian medium effects on the color dielectric constant ϵ were further investigated in [12] also on the base of kinetic theory.

From the analysis of the phenomenon of color superconductivity of hadron matter at high density [13, 14], it has been found that the medium surrounding the liberated quarks exhibits a polarization behavior characterized by an effective color dielectric constant $\epsilon \gg 1$, the magnetic permeability remaining close to one. Another estimate $\epsilon \gg 1$ has been done in [15] evaluating the photon self-energy for a color superconductor.

Finally, we mention models such as the chromodielectric model proposed by Friedberg and Lee [16, 17] and developed in [18, 19, 20, 21]. Aimed to describe an effective mechanism for the confinement phenomena, this model exploits a dielectric function depending on a classical field in order to account for the interaction of the QCD matter.

Having in mind these examples, our view-point is that, if in the interactions of a medium with Yang-Mills fields one encounters manifestations of its polarization-type response to the gauge fields, then it is worth developing an in-medium Yang-Mills theory and apply it to situations where such a response is found to be essential (like, e.g. the quark-gluon plasma).

In this paper we derive the classical Yang-Mills equations in a medium in the linear approximation (i.e. we derive the analogs of the macroscopic Maxwell equations for a gauge theory) and perform a canonical quantization of the arising theory. In the derivation of the classical in-medium gauge field equations we follow the method of Akhiezer and Peletminsky [22] in their derivation of the macroscopic Maxwell equations. In Section 2 we introduce some of the notations and write down the equations for a system of interacting spinor and gauge fields in the presence of a weak external gauge field. In Section 3 these equations are averaged by means of a nonequilibrium statistical operator in the linear approximation with respect to both the gauge coupling constant and the external field. The generalized conductivity tensors are defined in terms of the relevant retarded Green's functions. Section 4 contains the introduction of the nonabelian ("color") electric permittivity and magnetic permeability tensors, ϵ^{ab} and μ^{ab} , as well as the nanabelian polarization tensors. Defining the constitutive equations for the medium, we obtain relations between ϵ^{ab} , μ^{ab} , and the conductivity tensors, again in the linear approximation. These relations appear as natural generalizations of their analogs in the classical macroscopic electrodynamics. At the end of this section the in-medium (macroscopic) Yang-Mills equations are written down in a form closely analogous to its electro-dynamical counterpart. We follow Brevik and Lautrup [23] and Nakanishi [24, 25] in their treatment of the electromagnetic field which avoids work with constrained systems. Although in [23] one deals with in-medium electromagnetic field, and [24, 25, 26] consider the electromagnetic field in vacuum, these papers use a common approach, namely a Lagrangian formulation involving an auxiliary scalar field whose action on the Fock space is designed to separate the physical subspace. Section 5 provides a lagrangian formulation of the previously obtained equations introducing a set of $\dim(G)$ auxiliary fields B^a (G being the gauge group). For sake of simplicity we consider constant and diagonal tensors ϵ and μ . From the equal time commutation relations, in Section 6 we evaluate the commutators at arbitrary space-time points. In Section 7 the fields' Fourier transforms are introduced. In the resulting Fock space a physical subspace is defined by means of the operators B^a . Finally, for a particular choice of the gauge we evaluate the gauge field propagator.

2. Field equations

Let G be a compact r -dimensional simple matrix Lie group. Let T^a ($a = 1, \dots, r$) be a basis of hermitian matrices in its Lie algebra \mathcal{G} with commutation relations

$$[T^a, T^b] = if^{abc}T^c,$$

and normalized by

$$\text{tr}(T^a T^b) = \frac{1}{2} \delta^{ab}.$$

Lower case Latin letters a, b, c, \dots are used for the group /algebra indices and here and below summation over repeated indices is understood.

Let $\hat{\Psi}(x)$ be a quantized spinor field with values in a vector space \mathcal{W} , i.e. $\hat{\Psi}(x) \in \hat{\mathcal{S}} \otimes \mathcal{W}$, where $\hat{\mathcal{S}}$ is the space of operator-valued spinors over Minkowski space M , $x \in M$. We suppose that \mathcal{W} carries a linear representation $\omega \rightarrow R(\omega)$ of G , $\omega \in G$, and that \mathcal{W} is equipped with a scalar product such that $R(\omega)$ is unitary.

Given a quantized Yang-Mills gauge potential $\hat{\mathcal{A}}_\mu(x) = \hat{A}_\mu^a(x) T^a$ ($\mu = 0, \dots, 3$), the gauge covariant derivative of the spinor field $\hat{\Psi}(x)$ reads (in coordinates $(x^\mu) = (x^0, \vec{x})$)

$$\mathcal{D}_\mu(\hat{\mathcal{A}})\hat{\Psi}(x) = \partial_\mu \hat{\Psi}(x) + ig R(\hat{\mathcal{A}}_\mu(x)) \hat{\Psi}(x) = \partial_\mu \hat{\Psi}(x) + ig \hat{A}_\mu^a(x) R(T^a) \hat{\Psi}(x),$$

where $\mathcal{D}_\mu(\hat{\mathcal{A}})$ denotes covariant differentiation with respect to $\hat{\mathcal{A}}_\mu$, and g is a gauge coupling constant ($R(T^a)$ stands for the generators of G in the representation R). We suppose, in addition, that the spinor field $\hat{\Psi}$ is put in an external (classical) gauge field $f_{\mu\nu}$ with gauge potential $a_\mu = a_\mu^c T^c$, i.e.

$$f_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu + ig[a_\mu, a_\nu],$$

and $f_{\mu\nu}$ is determined by an external current

$$j^\mu = j^{a,\mu} T^a,$$

so that

$$\mathcal{D}_\mu(a) f^{\mu\nu} = \partial_\mu f^{\mu\nu} + ig[a_\mu, f^{\mu\nu}] = g j^\nu. \quad (1)$$

Fixing the units such that $\hbar = c = 1$, the Minkowski metric $(g_{\mu\nu}) = \text{diag}(1, -1, -1, -1)$, and denoting the operators $(\partial_n) = (-\partial^n) = \vec{\nabla}$, we also introduce the components of the analogs of the electric and magnetic fields

$$\hat{E}^{a,k} = -\hat{\mathcal{F}}^{a,0k}, \quad (2)$$

$$\hat{B}^{a,k} = -\frac{1}{2} \epsilon^{kmj} \hat{\mathcal{F}}^{a,mj},$$

where $a = 1, \dots, r$; $k = 1, 2, 3$, $\epsilon^{123} = 1$ and

$$\hat{\mathcal{F}}^{\mu\nu} = \partial^\mu \hat{\mathcal{A}}^\nu - \partial^\nu \hat{\mathcal{A}}^\mu + ig[\hat{\mathcal{A}}^\mu, \hat{\mathcal{A}}^\nu] = \hat{F}^{a,\mu\nu} T^a.$$

The gauge potentials $\hat{\mathcal{A}}_\mu$ are taken in the temporal gauge, $\hat{\mathcal{A}}_0 = 0$, satisfying $\partial_k \hat{\mathcal{A}}_k = 0$. We fix also $\partial_\mu a^\mu = 0$. The equations for $\hat{\mathcal{A}}(x)$ are

$$\partial_0 \hat{E}^{a,k}(x) = -\epsilon^{kmj} (\mathcal{D}^m(\hat{\mathcal{A}}) \hat{B}^j(x))^a - g(\hat{I}^{a,k}(x) + j^{a,k}(x^0, \vec{x})), \quad (3)$$

$$\partial_0 \hat{B}^{a,k}(x) = \epsilon^{kmj} (\mathcal{D}^m(\hat{\mathcal{A}}) \hat{E}^j(x))^a, \quad (4)$$

$$(\mathcal{D}^k(\hat{\mathcal{A}}) \hat{B}^k(x))^a = 0, \quad (5)$$

$$(\mathcal{D}^k(\hat{\mathcal{A}}) \hat{E}^k(x))^a = g(\hat{I}^{a,0}(x) + j^{a,0}(x^0, \vec{x})). \quad (6)$$

Here $\hat{I}_\nu(x) = \hat{I}_\nu^a(x)T^a$ is the spinor current with components

$$\hat{I}_\nu^a(x) = \hat{\Psi}^\dagger(x)\gamma_\nu R(T^a)\hat{\Psi}(x). \quad (7)$$

The equation for $\hat{\Psi}(x)$ reads

$$i\gamma^0\partial_0\hat{\Psi}(x) + i\gamma^n\mathcal{D}_n(\hat{\mathcal{A}})\hat{\Psi}(x) + i\gamma^0\mathcal{D}_0(a)\hat{\Psi}(x) + i\gamma^k\mathcal{D}_k(a)\hat{\Psi}(x) - m\hat{\Psi}(x) = 0, \quad (8)$$

where $\gamma^\lambda\gamma^\mu + \gamma^\mu\gamma^\lambda = 2g^{\lambda\mu}\mathbf{1}$. It follows from equation (8) that

$$\mathcal{D}^\mu(\hat{\mathcal{A}})\hat{I}_\mu = -ig[a^\mu, \hat{I}_\mu]. \quad (9)$$

In absence of the exterior gauge field the energy functional has the form

$$\hat{H}_0 = \int dx^3 \left\{ \frac{1}{2} \sum_{a=1}^r (\vec{E}^a \cdot \vec{E}^a + \vec{B}^a \cdot \vec{B}^a) + \hat{\Psi}^\dagger \gamma^0 (-i\gamma^k \mathcal{D}_k(\hat{\mathcal{A}}) \hat{\Psi} + m \hat{\Psi}) \right\}.$$

3. Response of the system to the exterior gauge field

3.1. The density operator and the averaged field equations

We suppose that at infinite past ($x^0 = -\infty$) in the absence of the exterior field the system of interacting fermion and gauge fields is in equilibrium at temperature $1/\beta$, described by the density operator $\hat{\rho}_0 = \exp(-\beta\hat{H})$. At time $x^0 = -\infty$ the exterior field a^μ is switched in adiabatically and the system is described by a nonequilibrium density operator $\hat{\rho}(x^0)$ satisfying

$$i\partial_0\hat{\rho}(x^0) = [\hat{H}(x^0), \hat{\rho}(x^0)], \quad (10)$$

with the initial condition $\hat{\rho}(-\infty) = \hat{\rho}_0$. The total Hamiltonian is

$$\hat{H}(x^0) = \hat{H}_0 + g\hat{V}(x^0), \quad (11)$$

where $\hat{V}(x^0)$ describes the interaction of the fermions with the gauge potential a^μ ,

$$\hat{V}(x^0) = \int d^3x \hat{I}^{c,\mu}(\vec{x}) a_\mu^c(x^0, \vec{x}). \quad (12)$$

It is convenient to go to the interaction (Dirac) representation, defined by the splitting of the Hamiltonian

$$\hat{H} = \hat{H}(x^0) = \hat{H}_0 + g\hat{V}(x^0).$$

In this representation the density operator $\hat{\rho}_{\mathcal{D}}(x^0)$ satisfies

$$i\partial\hat{\rho}_{\mathcal{D}}(x^0) = g[\hat{V}_{\mathcal{D}}(x^0), \hat{\rho}_{\mathcal{D}}(x^0)], \quad (13)$$

and the initial condition $\hat{\rho}_{\mathcal{D}}(-\infty) = \hat{\rho}_0$, where

$$\hat{V}_{\mathcal{D}}(x^0) = \int d^3x \hat{J}^\mu(x) a_\mu(x), \quad (14)$$

and \hat{J}^μ is the fermion current in the Dirac representation

$$\hat{J}^\mu(x) \equiv \hat{I}_{\mathcal{D}}^\mu(x) = \exp(ix^0\hat{H}_0)\hat{I}^\mu(x)\exp(-ix^0\hat{H}_0). \quad (15)$$

Combining equation (13) and the initial condition in the integral equation

$$\hat{\rho}_{\mathcal{D}}(x^0) = \hat{\rho}_0 - ig \int_{-\infty}^{x^0} dt [\hat{V}_{\mathcal{D}}(t), \rho_{\mathcal{D}}(t)],$$

after the first iteration one finds the well known representation

$$\hat{\rho}_{\mathcal{D}}(x^0) = \hat{\rho}_0 - ig \int_{-\infty}^{+\infty} dt \theta(x^0 - t) [\hat{V}_{\mathcal{D}}(t), \hat{\rho}_0]. \quad (16)$$

In other terms, the linear approximation of $\hat{\rho}_{\mathcal{D}}$ with respect to the exterior fields (and in first order in g) is

$$\hat{\rho}_{\mathcal{D}}(x^0) = \hat{\rho}_0 - ig \int d^4 y \theta(x^0 - y^0) [\hat{J}_{\mu}(y), \hat{\rho}_0] a^{\mu}(y). \quad (17)$$

By $\langle \dots \rangle$ and $\langle \dots \rangle_0$ we shall denote mean values with respect to $\hat{\rho}_{\mathcal{D}}$ and $\hat{\rho}_0$, respectively, i.e.

$$\langle A(x) \rangle = \text{tr}(\hat{\rho}_{\mathcal{D}}(x^0) \hat{A}(x)), \quad \langle A(x) \rangle_0 = \text{tr}(\hat{\rho}_0 \hat{A}(x)).$$

Applying the density operator $\hat{\rho}_{\mathcal{D}}$ on equations (3-6) one finds that the mean values

$$A_k^a(x) = \langle \hat{A}_k^a(x) \rangle, \quad E_n^a(x) = \langle \hat{E}_n^a(x) \rangle \quad \text{and} \quad B_n^a(x) = \langle \hat{B}_n^a(x) \rangle,$$

obey the following equations (valid up to first order in g)

$$\partial_0 \vec{E}^a = (\vec{\mathcal{D}}(A) \times \vec{B})^a - g(\vec{J}^a + \vec{j}^a), \quad (18)$$

$$\partial_0 \vec{B}^a = (\vec{\mathcal{D}}(A) \times \vec{E})^a, \quad (19)$$

$$(\vec{\mathcal{D}}(A) \cdot \vec{B})^a = 0, \quad (20)$$

$$(\vec{\mathcal{D}}(A) \cdot \vec{E})^a = g(J_0^a + j_0^a), \quad (21)$$

and

$$-\partial_0 \vec{A}^a = \vec{E}^a, \quad (22)$$

with the latter coming from $\hat{E}^{a,k} = -\hat{F}^{a,0k}$. Here we use the notation

$$J^{a,k}(x) \equiv \langle \hat{J}^{a,k}(x) \rangle$$

for the mean values of the fermion currents, and also

$$(\vec{\mathcal{D}}(A) \times \vec{B})^{c,n} = \epsilon^{nmj} \partial_m B^{c,j} - g \epsilon^{nmj} [A^m, B^j]^c,$$

$$(\vec{\mathcal{D}}(A) \times \vec{E})^{c,n} = \epsilon^{nmj} \partial_m E^{c,j} - g \epsilon^{nmj} [A^m, E^j]^c.$$

3.2. Retarded Green's functions and tensors of conductivity

For the mean value of an operator $A_{\mathcal{D}}$ one obtains in the first approximation

$$\begin{aligned} \langle \hat{A}(x^0, \vec{x}) \rangle &\equiv \text{tr}(\hat{A}(x^0, \vec{x}) \rho_{\mathcal{D}}(x^0)) \\ &= \langle \hat{A}(x^0, \vec{x}) \rangle_0 - ig \int_{-\infty}^{\infty} dt \langle [\hat{A}(x^0, \vec{x}), \hat{V}_{\mathcal{D}}(x^0)] \rangle_0 \theta(x^0 - t). \end{aligned} \quad (23)$$

Using the form of $V_{\mathcal{D}}$ from equation (14) and the notation

$$G_{A,j\mu}(x, y) \equiv -i \theta(x^0 - y^0) \langle [\hat{A}(x), \hat{j}_{\mu}(y)] \rangle_0, \quad (24)$$

for the retarded Green's function, one rewrites equation (23) as

$$\langle \hat{A}(x) \rangle = \langle \hat{A}(x) \rangle_0 + g \int d^4 y G_{A,j\mu}(x, y) a^{\mu}(y). \quad (25)$$

For two operators $\hat{A}(x)$ and $\hat{B}(x)$ which are covariant under space translations, the corresponding Green's function $G_{AB}(x, y) = -i\theta(x^0 - y^0) \langle [\hat{A}(x), \hat{B}(y)] \rangle_0$ depends only on $x - y$ and in this case we shall simply write $G_{AB}(x, y) \equiv G_{AB}(x - y)$. Let \hat{A} and \hat{B} transform under the time reflection antiunitary operator \mathbf{U} , according to

$$\mathbf{U}\hat{A}(x)\mathbf{U}^{-1} = \epsilon_A \hat{A}^*(i_t x), \quad \mathbf{U}\hat{B}(x)\mathbf{U}^{-1} = \epsilon_B \hat{B}^*(i_t x)$$

where $*$ denotes hermitian conjugation, $i_t x = (-x^0, \vec{x})$, and $|\epsilon_A| = |\epsilon_B| = 1$. If, in addition, \hat{A} and \hat{B} are hermitian, then the invariance under time reflection and space translations implies the property

$$G_{AB}(k_0, \vec{k}) = \epsilon_A \epsilon_B G_{BA}(k_0, -\vec{k}) \quad (26)$$

of the Fourier transform of G_{AB}

$$G_{AB}(k_0, \vec{k}) = \int e^{ik_0 x_0 - i\vec{k} \cdot \vec{x}} G_{AB}(x_0, \vec{x}) dx_0 d\vec{x}$$

(denoted by the same symbol). Substituting in equation (25) the components $\hat{J}_\mu(x)$ of the fermionic current equations (15), (7), we obtain

$$J_\mu^a(x) = \langle \hat{J}_\mu^a(x) \rangle = g \int d^4 y G_{\mu\nu}^{ab}(x - y) a^{b,\nu}(y), \quad (27)$$

$a = 1, \dots, r$, where

$$G_{\mu\nu}^{ab}(x - y) = -i\theta(x^0 - y^0) \langle [\hat{J}_\mu^a(x), \hat{J}_\nu^b(y)] \rangle_0 = -i\theta(x^0 - y^0) \langle [\hat{I}_\mu^a(x), \hat{I}_\nu^b(y)] \rangle_0 \quad (28)$$

and we have used the fact that for a neutral medium in equilibrium, the mean values $\langle \hat{J}_\mu^a(x) \rangle_0 = 0$. We shall use the Fourier transform of equation (27)

$$J_\mu^a(k_0, \vec{k}) = g G_{\mu\nu}^{ab}(k_0, \vec{k}) a^{b,\nu}(k_0, \vec{k}). \quad (29)$$

Thus the averaged (or induced) current J_μ^a depends linearly on the external field up to first order in g . In the same approximation and using local commutativity, one derives from equation (9) that

$$\mathcal{D}^\mu(A) G_{\mu\nu}^{ab}(x) = 0 \quad \text{and} \quad \mathcal{D}^\nu(A) G_{\mu\nu}^{ab}(x) = 0. \quad (30)$$

In the sequel we shall look for relations which hold up to first order in g and/or are linear in the external gauge field. Since the r.h.s of equation (27) (or equation (29)) is proportional to g , we shall use the consequences of equation (30) written in its approximate forms $\partial^\mu G_{\mu\nu}^{ab}(x) = 0$ and $\partial^\nu G_{\mu\nu}^{ab}(x) = 0$. For the Fourier transforms this approximation gives

$$\begin{aligned} k^0 G_{00}^{ab}(k_0, \vec{k}) + k^n G_{n0}^{ab}(k_0, \vec{k}) &= 0, \\ k^0 G_{00}^{ab}(k_0, \vec{k}) + k^n G_{0n}^{ab}(k_0, \vec{k}) &= 0, \\ k^m G_{mn}^{ab}(k_0, \vec{k}) + k^0 G_{0n}^{ab}(k_0, \vec{k}) &= 0, \\ k^n G_{mn}^{ab}(k_0, \vec{k}) + k^0 G_{m0}^{ab}(k_0, \vec{k}) &= 0, \\ (\vec{k} = (k^n) = -(k_n)). & \end{aligned} \quad (31)$$

It is consistent (up to first order in g) to ignore the commutator term in

$$e_k^b(x) = -f_{0k}^b(x) = -(\partial_0 a_k^b(x) - \partial_k a_0^b(x) + ig[a_0(x), a_k(x)]^b),$$

and to write the Fourier transform of $e_k^b(x)$ as

$$\begin{aligned} e^{b,n}(k_0, \vec{k}) &= \int \exp(ik_0 x^0 + ik_n x^n) e^{b,n}(x^0, \vec{x}) dx^0 d^3x \\ &= -i(k^n a^{b,0}(k_0, \vec{k}) - k^0 a^{b,n}(k_0, \vec{k})). \end{aligned} \quad (32)$$

Then it follows from equation (29) and equation (31) that

$$J_0^a(k_0, \vec{k}) = -ig \frac{k_m}{k_0^2} G_{ml}^{ab}(k_0, \vec{k}) e^{b,l}(k_0, \vec{k}), \quad (33)$$

$$J_j^a(k_0, \vec{k}) = -ig \frac{1}{k_0} G_{jl}^{ab}(k_0, \vec{k}) e^{b,l}(k_0, \vec{k}). \quad (34)$$

For a homogeneous and isotropic medium $G_{mn}^{ab}(k_0, \vec{k})$ may be written as a sum of tensors

$$G_{mn}^{ab}(k_0, \vec{k}) = -ik_0 [\tilde{\sigma}_l^{ab}(k) \frac{k_m k_n}{k^2} + \tilde{\sigma}_t^{ab}(k) (\delta_{mn} - \frac{k_m k_n}{k^2})], \quad (35)$$

where $\tilde{\sigma}_l^a$ and $\tilde{\sigma}_t^a$ are "color" tensors depending on $k = |\vec{k}|$ and arise as analogs of the so called exterior longitudinal and transverse electric conductivity coefficients [22]. This form of G_{mn}^{ab} generates a decomposition of J_0^a and J_j^a into longitudinal and transverse parts with respect to \vec{k} ,

$$J_0^a(k_0, \vec{k}) = \frac{g}{k_0} \tilde{\sigma}_l^{ab}(k) (\vec{k} \cdot \vec{e}^b), \quad (36)$$

$$J_j^a(k_0, \vec{k}) = g \tilde{\sigma}_l^{ab}(k) \frac{(\vec{k} \cdot \vec{e}^b)}{k^2} k_j + g \tilde{\sigma}_t^{ab}(k) \frac{((\vec{k} \times \vec{e}^b) \times \vec{k})_j}{k^2}. \quad (37)$$

4. Tensors of generalized electric and magnetic permeability

Following the derivation of [22], we introduce the analog of the polarization tensor

$$\mathcal{P}_{\mu\nu}(x) = -\mathcal{P}_{\nu\mu}(x) = \mathcal{P}_{\mu\nu}^a T^a, \quad (38)$$

through a gauge covariant relation between $\mathcal{P}_{\mu\nu}$ and the averaged current J_ν ,

$$J_\nu(x) = \mathcal{D}^\mu \mathcal{P}_{\mu\nu}(x) = \partial^\mu \mathcal{P}_{\mu\nu}(x) - ig[A^\mu(x), \mathcal{P}_{\mu\nu}(x)], \quad (39)$$

which is the substitute of $J_\nu = \partial^\mu P_{\mu\nu}$ in the case of classical electrodynamics. We also define the corresponding generalization of the 3-vectors of the electric and magnetic polarizations, respectively,

$$P^{a,n} = \mathcal{P}^{a,0n}, \quad (40)$$

and

$$M^{a,n} = -\frac{1}{2} \epsilon^{nlj} \mathcal{P}^{a,lj}. \quad (41)$$

Written in components, the defining equation (39) reads

$$\begin{aligned} J^{a,n} &= \partial_0 P^{a,n} + \epsilon^{nmj} [\partial_m M^{a,j} + g f^{abc} (A^{b,m} M^{c,j})] \\ &\equiv \partial_0 P^{a,n} + (\vec{\mathcal{D}}(A) \times \vec{M})^{a,n}, \end{aligned} \quad (42)$$

$$J_0^a = -\vec{\nabla} \cdot \vec{P}^a - g f^{abc} (A^{b,m} P^{c,j}) \epsilon^{nmj} \equiv -(\vec{\mathcal{D}}(A) \cdot \vec{P})^a. \quad (43)$$

($\vec{P}^a = P^{a,n}$, $\vec{M}^a = M^{a,n}$). Applying again the analogy with the case of classical electrodynamics, we fix the arbitrariness in $\mathcal{P}_{\mu\nu}$ assuming a linear relation between (the Fourier transforms of) \vec{P}^a and \vec{M}^a , on the one hand, and the averaged "electric" and "magnetic" fields, \vec{E}^a and \vec{B}^a , on the other:

$$\vec{P}^a(k_0, \vec{k}) = \kappa^{ab}(k_0, \vec{k})\vec{E}^b(k_0, \vec{k}), \quad (44)$$

$$\vec{M}^a(k_0, \vec{k}) = (\mu^{-1})^{ab}(k_0, \vec{k})\chi^{bc}(k_0, \vec{k})\vec{B}^c(k_0, \vec{k}). \quad (45)$$

Equations (44) and (45) are the analogs of the constitutive relations of a medium, where the "color" tensors $\kappa = (\kappa^{ab})$ and $\chi = (\chi^{ab})$ appear as generalizations of the electric and magnetic susceptibilities, respectively, while $\mu = (\mu^{ab})$ is defined as the *tensor of generalized magnetic permeability* related with χ by the defining equality

$$\mu^{ab} = \delta^{ab} + g\chi^{ab}.$$

We also define (the Fourier transforms of) the generalized electric and magnetic induction by

$$D^{a,n}(k_0, \vec{k}) = E^{a,n}(k_0, \vec{k}) + gP^{a,n}(k_0, \vec{k}), \quad (46)$$

and

$$H^{a,n}(k_0, \vec{k}) = B^{a,n}(k_0, \vec{k}) - gM^{a,n}(k_0, \vec{k}), \quad (47)$$

respectively. From equations (44) and (45) one finds

$$B^{a,n} = \mu^{ab}H^{b,n}, \quad D^{a,n} = \epsilon^{ab}E^{b,n}, \quad (48)$$

where $\epsilon^{ab} = \delta^{ab} + g\kappa^{ab}$ will be called *tensor of generalized electric permittivity*. If we keep the analogy with the electromagnetic properties of the vacuum, we have to prescribe the property of generalized permeability μ_0 and permittivity ϵ_0 to the empty space with $\mu_0 = \text{const}$ and $\epsilon_0 = \text{const}$. Then the definitions of μ and ϵ will read

$$\mu^{ab}/\mu_0 = \delta^{ab} + g\chi^{ab},$$

and

$$\epsilon^{ab}/\epsilon_0 = \delta^{ab} + g\kappa^{ab}.$$

The system of units will be fixed such that $\epsilon_0 = 1$ and $\mu_0 = 1$.

In terms of the induction 3-vectors \vec{D}^a and \vec{H}^a , the Fourier transforms of the averaged field equations (18) - (22) become

$$\begin{aligned} ik_0\vec{A}^a(k_0, \vec{k}) &= \vec{E}^a(k_0, \vec{k}), \\ ik_0\vec{D}^a(k_0, \vec{k}) + i\vec{k} \times \vec{H}^a(k_0, \vec{k}) \\ &+ \frac{g}{(2\pi)^4} \int d^4 p f^{bca} \vec{A}^b(k_0 - p_0, \vec{k} - \vec{p}) \times \vec{H}^c(p_0, \vec{p}) = g\vec{j}^a(k_0, \vec{k}), \\ i\vec{k} \cdot \vec{D}(k_0, \vec{k}) + \frac{g}{(2\pi)^4} \int d^4 p f^{abc} \vec{A}^b(k_0 - p_0, \vec{k} - \vec{p}) \cdot \vec{D}^c(p_0, \vec{p}) &= gj_0^c(k_0, \vec{k}), \\ ik_0\vec{B}^c(k_0, \vec{k}) - i\vec{k} \times \vec{E}^a(k_0, \vec{k}) - \frac{g}{(2\pi)^4} \int d^4 p f^{abc} \vec{A}^b(k_0 - p_0, \vec{k} - \vec{p}) \times \vec{E}^c(p_0, \vec{p}) &= 0, \\ i\vec{k} \cdot \vec{B}(k_0, \vec{k}) + \frac{g}{(2\pi)^4} \int d^4 p f^{abc} \vec{A}^b(k_0 - p_0, \vec{k} - \vec{p}) \cdot \vec{B}^c(p_0, \vec{p}) &= 0. \end{aligned} \quad (49)$$

Our next goal is to express the electric permittivity ϵ and the magnetic permeability μ in terms of the exterior longitudinal and transverse electric conductivity tensors $\tilde{\sigma}_l^a$ and $\tilde{\sigma}_t^a$, i.e. in terms of the Green's functions.

The first step in this direction is to find a relation between $\vec{E}^a(k)$, $\vec{B}^a(k)$ and the external field. To this end we first consider the systems (49) and (1) writing the latter in terms of the components $e^{c,n} = -f^{c,0n}$ and $b^{c,n} = -\frac{1}{2}\epsilon^{n,jm}f^{c,jm}$ ($c = 1, \dots, r; n = 1, 2, 3$) of the external gauge field, namely

$$\begin{aligned}
 & e^{c,n}(k_0, \vec{k}) \\
 &= ik_0 a^{c,n}(k_0, \vec{k}) - ik^n a^{c,0}(k_0, \vec{k}) + \frac{g}{(2\pi)^4} \int d^4 p f^{cbd} a^{b,0}(k_0 - p_0, \vec{k} - \vec{p}) e^{d,n}(p_0, \vec{p}), \\
 & ik_0 \vec{e}^c(k_0, \vec{k}) + i\vec{k} \times \vec{h}^c(k_0, \vec{k}) \\
 & \quad + \frac{g}{(2\pi)^4} \int d^4 p f^{cbd} \vec{a}^b(k_0 - p_0, \vec{k} - \vec{p}) \times \vec{h}^d(p_0, \vec{p}) = g\vec{j}^c(k_0, \vec{k}), \\
 & i\vec{k} \cdot \vec{e}^c(k_0, \vec{k}) + \frac{g}{(2\pi)^4} \int d^4 p f^{cbd} \vec{a}^b(k_0 - p_0, \vec{k} - \vec{p}) \times \vec{e}^d(p_0, \vec{p}) = gj_0^c(k_0, \vec{k}), \\
 & ik_0 \vec{h}^c(k_0, \vec{k}) - i\vec{k} \times \vec{e}^c(k_0, \vec{k}) - \frac{g}{(2\pi)^4} \int d^4 p f^{cbd} \vec{a}^b(k_0 - p_0, \vec{k} - \vec{p}) \times \vec{e}^d(p_0, \vec{p}) = 0, \\
 & \quad i\vec{k} \cdot \vec{h}^c(k_0, \vec{k}) + \frac{g}{(2\pi)^4} \int d^4 p f^{cbd} \vec{a}^b(k_0 - p_0, \vec{k} - \vec{p}) \cdot \vec{h}^d(p_0, \vec{p}) = 0.
 \end{aligned} \tag{50}$$

Restricting ourselves to relations holding up to linear terms in the external field, we neglect the integral terms in the system (50) reducing it to the form

$$\begin{aligned}
 e^{c,n}(k) &= ik_0 a^{c,n}(k) - ik^n a^{c,0}(k), \\
 -ik_0 \vec{e}^c(k) - ik \times \vec{h}^c(k) &= -g\vec{j}^c(k), \\
 i\vec{k} \cdot \vec{e}^c(k) &= gj_0^c(k), \\
 -ik_0 \vec{h}^c(k) + i\vec{k} \times \vec{e}^c(k) &= 0, \\
 i\vec{k} \cdot \vec{h}^c(k) &= 0.
 \end{aligned} \tag{51}$$

($k = (k_0, \vec{k})$). Combining the systems (49) and (51) provides an expression for $\vec{D}^a = \epsilon^{ab}\vec{E}^b$ and $\vec{B}^a = \mu^{ab}\vec{H}^b$ in terms of the external field. In order to keep our approximation, the bilinear integral terms in equation (49) are neglected and thus one finds

$$\begin{aligned}
 ik_0 \vec{A}^c(k) &= \vec{E}^c(k), \\
 -ik_0 \vec{D}^c(k) - i\vec{k} \times \vec{H}^c(k) &= -g\vec{j}^c(k), \\
 i\vec{k} \cdot \vec{D}^c(k) &= gj_0^c(k), \\
 -ik_0 \vec{B}^c(k) + i\vec{k} \times \vec{E}^c(k) &= 0, \\
 i\vec{k} \cdot \vec{B}^c(k) &= 0,
 \end{aligned} \tag{52}$$

$$\tag{53}$$

This gives for the longitudinal component (with respect to \vec{k})

$$\vec{D}_{\parallel}^c(k_0, \vec{k}) \equiv \frac{\vec{k} \cdot \vec{D}^c(k_0, \vec{k})}{|\vec{k}|} \vec{k} = \epsilon^{cd}(k_0, \vec{k}) \vec{E}_{\parallel}^d(k_0, \vec{k}) \equiv \vec{e}_{\parallel}^c(k_0, \vec{k}). \tag{54}$$

Equations (48), (51) and (52) yield that the transverse components of $\vec{E}^a(k)$ and $\vec{e}^a(k)$ are related by

$$[k_0^2 \epsilon^{ab}(k) - \vec{k}^2 (\mu^{-1}(k))^{ab}] \vec{E}_\perp^b(k) = (k_0^2 - \vec{k}^2) \vec{e}_\perp^a(k), \quad (55)$$

where for a vector \vec{v}^a ($a = 1, \dots, r$) we define

$$\vec{v}_\perp^a = \frac{1}{\vec{k}^2} (\vec{k} \times \vec{v}^a) \times \vec{k}.$$

In a similar way, from equations (48), (51) and (52), we infer $\vec{B}_\parallel = 0$ and

$$[k_0^2 \epsilon^{ab}(k) - \vec{k}^2 (\mu^{-1}(k))^{ab}] \vec{B}_\perp^b(k) = (k_0^2 - \vec{k}^2) \vec{h}^a(k). \quad (56)$$

Equations (54), (55) and (56) provide the desired relation between $\vec{E}^a(k)$, $\vec{B}^a(k)$ and the external fields $\vec{e}^a(k)$, $\vec{h}^a(k)$.

The aim of the second step is, using the relations (54)-(56), to express $\tilde{\sigma}_t^{ab}$ and $\tilde{\sigma}_l^{ab}$ in equations (36), (37) in terms of ϵ^{ab} and μ^{ab} . To this end, following our approximation, one neglects the bilinear terms in equations (42, 43) and arrives at

$$J_0^a(x) = -\vec{\nabla} \cdot \vec{P}^a(x), \quad (57)$$

$$\vec{J}^a(x) = \partial_0 \vec{P}^a(x) + \vec{\nabla} \times \vec{M}^a(x),$$

or

$$J_0^a(k) = -i\vec{k} \cdot \vec{P}^a(k), \quad (58)$$

$$\vec{J}^a(k) = -ik_0 \vec{P}^a(k) + i\vec{k} \times \vec{M}^a(k),$$

in terms of the Fourier transforms. From equations (36), (58), (44) and (54), we derive

$$k_0 \kappa^{ab}(k_0, \vec{k}) = ig(\tilde{\sigma}_t)^{ac}(k_0, \vec{k}) \epsilon^{cb}(k_0, \vec{k}),$$

or, in a compact notation,

$$\mathbf{1} - \epsilon^{-1} = ig^2 \tilde{\sigma}_t. \quad (59)$$

Separating the longitudinal and transverse parts of (37), (58) and making use of equations (44), (45), (55) and (56) one finds the tensor equality

$$\mathbf{1} - \mu^{-1} = g\mu^{-1} \cdot \chi = i \frac{k_0}{\vec{k}^2} g^2 \left\{ \tilde{\sigma}_t \left(\mathbf{1} - ig^2 k_0 \frac{\tilde{\sigma}_t}{(k_0^2 - \vec{k}^2)} \right)^{-1} - \left(\mathbf{1} - i \frac{g^2}{k_0} \tilde{\sigma}_l \right)^{-1} \tilde{\sigma}_l \right\}. \quad (60)$$

Equations (59) and (60) resemble their counterparts in electrodynamics and, by analogy, we call

$$\sigma_l = \left(\mathbf{1} - i \frac{g^2}{k_0} \tilde{\sigma}_l \right)^{-1} \tilde{\sigma}_l \quad (61)$$

and

$$\sigma_t = \tilde{\sigma}_t \left(\mathbf{1} - ig^2 k_0 \frac{\tilde{\sigma}_t}{(k_0^2 - \vec{k}^2)} \right)^{-1} \quad (62)$$

tensors of intrinsic longitudinal and transverse conductivity, respectively.

At the end of this section we write down the system (49) of averaged equations in terms of the space-time coordinates

$$\begin{aligned}
 (\vec{\mathcal{D}}^a(A) \times \vec{H})^a - \partial_0 \vec{D}^a &= g \vec{j}^a, \\
 (\vec{\mathcal{D}}^a(A) \cdot \vec{D})^a &= g j_0^a, \\
 (\vec{\mathcal{D}}^a(A) \times \vec{E})^a - \partial_0 \vec{B}^a &= 0, \\
 (\vec{\mathcal{D}}^a(A) \cdot \vec{B})^a &= 0.
 \end{aligned} \tag{63}$$

These equations (as well as (49)) hold up to first order in g . The system (63) together with equations (48), where the tensors ϵ^{ab} and μ^{ab} are given functions of (x^0, \vec{x}) , will be called *Yang-Mills equations in medium*.

5. Lagrangian and canonical quantization

For a medium in rest with respect to a fixed Lorenz frame we define the antisymmetric tensors $H^{a,\mu\nu}$ by

$$\begin{aligned}
 H^{a,0\kappa} &= -D^{a,\kappa}, \\
 H^{a,mn} &= -\epsilon^{mnk} H^{a,k}, \quad n, m = 1, 2, 3.
 \end{aligned} \tag{64}$$

The components of the tensors (64) read

$$H^{a,\mu\nu} = (\mu^{-1})^{ab} F^{b,\mu\nu} + (\epsilon - \mu^{-1})^{ab} V^\tau (g^{\mu\sigma} V^\nu - g^{\nu\sigma} V^\mu) F_{\sigma\tau}^b, \tag{65}$$

if the medium moves uniformly with velocity V^μ ($V^2 = 1$). For vanishing exterior current j_μ^a the lagrangian

$$\mathcal{L}_0 = -\frac{1}{4} F_{\mu\nu}^a H^{a,\mu\nu} \tag{66}$$

reproduces the first two equations of the system equation (63) written in a covariant form

$$D(A)_\mu H^{a,\mu\nu} = 0, \tag{67}$$

valid for any reference frame. Since in equations (65) and (66) the tensor $F_{\mu\nu}^a$ reads

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g f^{abc} A_\mu^b A_\nu^c, \tag{68}$$

it satisfies the covariant form of the last two equations of the system equation (63), i.e.

$$D(A)_\lambda F_{\mu\nu}^a + D(A)_\mu F_{\nu\lambda}^a + D(A)_\nu F_{\lambda\mu}^a = 0. \tag{69}$$

In order to perform the canonical quantization, we add a gauge fixing term \mathcal{L}_{gf} to \mathcal{L}_0 ,

$$\mathcal{L}_{gf} = \frac{\xi}{2} (B^a B^a) - B^a (\partial^\mu A_\mu^a + (n^2 - 1)^{ab} V^\mu \partial_\mu V^\lambda A_\lambda^b), \tag{70}$$

where ξ is a real parameter and B^a ($a = 1, \dots, r$) are real auxiliary scalar fields. The tensor $(n^2)^{ab} = \epsilon^{ac} \mu^{cb}$ is the generalization of the classical refraction coefficient. The form (70) of \mathcal{L}_{gf} is taken in analogy with the case of electromagnetic field in a medium [23]. In the sequel we will consider the case $\epsilon^{ab} = \epsilon \delta^{ab}$, $\mu^{ab} = \mu \delta^{ab}$, where ϵ and μ are constants. Then $(n^2)^{ab} = \epsilon \mu \delta^{ab}$ and the notation $\beta = n^2 - 1$ will be used (here and

below we shall write n^2 for $\epsilon\mu$). We split the total lagrangian $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{gf}$ into two parts, $\mathcal{L} = \mathcal{L}_{free} + \mathcal{L}_{int}$, where \mathcal{L}_{int} contains all terms containing the gauge coupling constant g . The part \mathcal{L}_{free} equals

$$\mathcal{L}_{free} = -\frac{1}{4}\tilde{F}_{\mu\nu}^a\tilde{H}^{a,\mu\nu} + \mathcal{L}_{gf}, \quad (71)$$

where

$$\tilde{F}_{\lambda\mu}^a = \partial_\lambda A_\mu^a - \partial_\mu A_\lambda^a \quad (72)$$

and

$$\tilde{H}^{a,\mu\nu} = (\mu^{-1})^{ab}\tilde{F}^{b,\mu\nu} + (\epsilon - \mu^{-1})^{ab}V^\tau(g^{\mu\sigma}V^\nu - g^{\nu\sigma}V^\mu)\tilde{F}_{\sigma\tau}^b. \quad (73)$$

The equations of motion following from the free field part \mathcal{L}_{free} are

$$\square A_\lambda^a + \beta(V \cdot \partial)^2 A_\lambda^a + (\mu - \xi)\partial_\lambda B^a \equiv Q A_\lambda^a + (\mu - \xi)\partial_\lambda B^a = 0, \quad (74)$$

$$\square B^a + \beta(V \cdot \partial)^2 B^a \equiv Q B^a = 0, \quad (75)$$

$$\xi B^a = \partial \cdot A^a + \beta(V \cdot \partial)(V \cdot A^a), \quad (76)$$

where $Q = \square + \beta(V \cdot \partial)^2$. We choose $A_\lambda^a(x)$ as generalized coordinates, the conjugate momenta being

$$\pi^{a,\mu}(x) = \frac{\partial \mathcal{L}_0}{\partial(\partial_0 A_\mu^a)}(x),$$

or

$$\pi^{a,0} = -[1 + \beta V_0^2]B^a, \quad (77)$$

$$\pi^{a,k} = \beta V_0 V_k B^a + \frac{\beta}{\mu} V_0 V_n \tilde{F}_{kn}^a + [(1 + 2\beta V_0^2)\delta_{kn} - \beta V_k V_n] \tilde{F}_{0n}^a, \quad (78)$$

($k = 1, 2, 3$). From the equal time canonical commutation relations

$$[A_\lambda(x), A_\mu(y)]_{x_0=y_0} = 0, \quad (79)$$

$$[\pi^{a,\lambda}(x), \pi^{b,\mu}(y)]_{x_0=y_0} = 0, \quad (79)$$

$$[\pi^{a,\lambda}(x), A_\mu^b(y)]_{x_0=y_0} = -i\delta^{ab}\delta_\mu^\lambda\delta(\vec{x} - \vec{y}), \quad (80)$$

one derives

$$[B^a(x), A_\lambda^b(y)]_{x_0=y_0} = \frac{i\delta^{ab}\delta_{0\lambda}}{(1 + \beta V_0^2)}\delta(\vec{x} - \vec{y}), \quad (81)$$

$$[B^a(x), \partial_0 A_\lambda^b(y)]_{x_0=y_0} = (1 - \delta_{0\lambda})\frac{i\delta^{ab}}{(1 + \beta V_0^2)}\frac{\partial}{\partial y^n}\delta(\vec{x} - \vec{y}), \quad (82)$$

as well as

$$[\partial_0 A_0^a(x), A_0^b(y)]_{x_0=y_0} = \frac{i\delta^{ab}\delta(\vec{x} - \vec{y})}{(1 + \beta V_0^2)^2}[\xi - \mu\beta^2\frac{V_0^2\vec{V}^2}{(1 + \alpha)}], \quad (83)$$

$$[\partial_0 A_0^a(x), A_n^b(y)]_{x_0=y_0} = -\frac{i\mu\beta V_0 V_n \delta^{ab}}{(1 + \beta)(1 + \beta V_0^2)}\delta(\vec{x} - \vec{y}), \quad (84)$$

$$[\partial_0 B^a(x), A_0^b(y)]_{x_0=y_0} = 2i\frac{\beta V_0}{(1 + \beta V_0^2)^2}\delta^{ab}(V_n\frac{\partial}{\partial x^n})\delta(\vec{x} - \vec{y}), \quad (85)$$

$$[\partial_0 B^a(x), A_n^b(y)]_{x_0=y_0} = \frac{i\delta^{ab}}{(1 + \beta V_0^2)}\frac{\partial}{\partial x^n}\delta(\vec{x} - \vec{y}). \quad (86)$$

Following [23] we introduce the tensor

$$b_\tau^\sigma = \delta_\tau^\sigma + (n-1)V^\sigma V_\tau,$$

satisfying

$$(b^l)^\sigma_\tau = \delta_\tau^\sigma + (n^l-1)V^\sigma V_\tau$$

for every integer number l . In terms of b_τ^σ , the gauge fixing term \mathcal{L}_{gf} reads

$$\mathcal{L}_{gf} = \frac{\xi}{2} B^a B^a - B^a \partial^\mu (b^2)_\mu^\tau A_\tau$$

and the operator Q takes the form

$$Q = \partial^\mu (b^2)_\mu^\tau \partial_\tau.$$

6. Commutation relations at non-equal times

We define the functions

$$F(x) \equiv i \int \frac{d^4 k}{(2\pi)^3} \exp(-ik \cdot x) \delta(Q(k)) \epsilon(V \cdot k), \quad (87)$$

and

$$H(x) \equiv i \int \frac{d^4 k}{(2\pi)^3} \exp(-ik \cdot x) \delta'(Q(k)) \epsilon(V \cdot k), \quad (88)$$

with the notations

$$Q(k) \equiv k^2 + \beta(V \cdot k)^2 = (1 + \beta V_0^2)(k_0 - k_+(k))(k_0 - k_-(k)), \quad (89)$$

and

$$k_\pm(\vec{k}) \equiv \frac{1}{(1 + \beta V_0^2)} \left[\beta V_0(\vec{k} \cdot \vec{V}) \pm \sqrt{(1 + \beta V_0^2)\vec{k}^2 - \beta(\vec{k} \cdot \vec{V})^2} \right]. \quad (90)$$

Note that $k_+(\vec{k})$ is negative for reference frames in which $\beta V^2 > 1$. However, it always holds that $k_\pm(\vec{k}) = -k_\mp(-\vec{k})$.

The functions F and H have the properties $QF = 0$, $QH = F$, as well as

$$\begin{aligned} \partial_0 F(0, \vec{x}) &= \frac{1}{(1 + \beta V_0^2)} \delta(\vec{x}), \\ F(0, \vec{x}) &= 0, \end{aligned} \quad (91)$$

$$\begin{aligned} H(0, \vec{x}) = \partial_0 H(0, \vec{x}) = \partial_0^2 H(0, \vec{x}) &= 0, \\ \partial_0^3 H(0, \vec{x}) &= \frac{1}{(1 + \beta V_0^2)^2} \delta(\vec{x}). \end{aligned} \quad (92)$$

Equations (87) and (90) yield

$$F(x) = \frac{i}{(2\pi)^3} \int \frac{d^3 k}{(1 + \beta V_0^2)} \frac{1}{|k_+(\vec{k}) - k_-(\vec{k})|} (e^{-ik'x} - e^{-ik''x}), \quad (93)$$

where $k' = (k_+(\vec{k}), \vec{k})$, $k'' = (k_-(\vec{k}), \vec{k})$. The current

$$J_\mu(y, x) = F(y-x) \mathcal{D}^{\leftrightarrow}_{x,\mu} g(x) + H(y-x) \mathcal{D}^{\leftrightarrow}_{x,\mu} Qg(x) \quad (94)$$

is conserved, $\frac{\partial}{\partial x^\mu} J^\mu(y, x) = 0$, for every solution $g(x)$ of

$$Q^2 g(x) = 0.$$

The operator $\mathcal{D}_{x,\mu}$ is defined as

$$\mathcal{D}_{x,\mu} \equiv \frac{\partial}{\partial x^\mu} + \beta V_\mu (V^\lambda \frac{\partial}{\partial x^\lambda}). \quad (95)$$

Therefore the integral

$$\int_\Sigma d\sigma^\mu(z) J_\mu(y, z)$$

($d\sigma^\mu(z)$ is the element on the surface Σ) depends only on y . For a space-like Σ with $d\sigma^\mu(z) = (1, 0, 0, 0)d^3z$ we define

$$G(y) \equiv \int_{z^0=y^0} d^3z J_0(y, z). \quad (96)$$

It follows from equation (94) and the properties (91) - (92) that

$$G(y) = g(y) = \int_{z^0=y^0} d^3z \left[F(y-z) \mathcal{D}^{\leftrightarrow}_{z,0} g(z) + H(y-z) \mathcal{D}^{\leftrightarrow}_{z,0} Qg(z) \right]. \quad (97)$$

From (74) and (75) one derives

$$Q^2 A_\lambda^a(x) = 0. \quad (98)$$

Therefore one obtains

$$A_\lambda^a(x) = \int_{x^0=z^0} d^3z \left[F(x-z) \mathcal{D}^{\leftrightarrow}_{z,0} A_\lambda^a(z) + H(x-z) \mathcal{D}^{\leftrightarrow}_{z,0} Q A_\lambda^a(z) \right]. \quad (99)$$

This expression allows one to calculate the commutators among the fields $A_\lambda^a(x)$ for arbitrary times using the equal time commutators appearing in the r.h.s of the identify

$$\begin{aligned} [A_\lambda^a(x), A_\mu^b(y)] = & \int_{z_0=y_0} d^3z \left\{ F(x-z) \mathcal{D}^{\leftrightarrow}_{z,0} [A_\lambda^a(z), A_\mu^b(y)] \right. \\ & \left. + H(x-z) \mathcal{D}^{\leftrightarrow}_{z,0} (\xi - \mu) \frac{\partial}{\partial z^\mu} [B^a(z), A_\nu^b(y)] \right\}. \end{aligned} \quad (100)$$

The result reads

$$[A_\lambda^a(x), A_\mu^b(y)] = i\mu \delta^{ab} (g_{\lambda\mu} - \frac{\beta}{1+\beta} V_\lambda V_\mu) F(x-y) - i(\mu - \xi) \delta^{ab} \frac{\partial^2 H(x-y)}{\partial x^\lambda \partial x^\mu}. \quad (101)$$

Using equation (76), i.e.

$$\xi B^a(x) = \mathcal{D}_{x,\mu} A^{a,\mu}(x)$$

with $\mathcal{D}_{x,\mu}$ given by (95), one obtains from equation (100)

$$[A_\lambda^a(x), B^b(y)] = -i\delta^{ab} \frac{\partial}{\partial x^\lambda} F(x-y), \quad (102)$$

and

$$[B^a(x), B^b(y)] = 0. \quad (103)$$

The commutation relations (101-103) generalize those derived for the electromagnetic field in a medium [23] in a natural way.

One may use the technique of [23] to derive the commutation relations equation (101) in a much simpler way. Indeed, the change of variables

$$x^\lambda \rightarrow y^\lambda = (b^{-1})^\lambda_\mu x^\mu, \quad (104)$$

$$A_\lambda^a(x) \rightarrow \Gamma_\lambda^a(y) = \frac{1}{\sqrt{\mu}} A_\sigma^a(x) b_\lambda^\sigma, \quad (105)$$

$$B^a(x) \rightarrow \Delta^a(y) = \sqrt{\mu} B^a(x),$$

and

$$g \rightarrow g' = \sqrt{\mu} g, \quad \xi \rightarrow \xi' = \xi/\mu, \quad (106)$$

brings the lagrangian

$$\mathcal{L} = -\frac{1}{4} F^{\alpha,\mu\nu}(x) \mathcal{H}_{\mu\nu}^\alpha(x) - \xi B^a(x) [\partial \cdot A^a(x) + (n^2 - 1)(V \cdot \partial) V \cdot A^a(x)] + \frac{1}{2} B^a(x) B^a(x) \quad (107)$$

to the form

$$\mathcal{L} = -\frac{1}{4} G^{\alpha,\mu\nu}(y) G_{\mu\nu}^\alpha(y) - \xi' \Delta^a(y) \frac{\partial}{\partial y^\mu} \Gamma^{\alpha,\mu}(y) + \frac{1}{2} \Delta^a(y) \Delta^a(y), \quad (108)$$

where

$$G_{\mu\nu}^\alpha(y) = \frac{\partial \Gamma_\nu^\alpha(y)}{\partial y^\mu} - \frac{\partial \Gamma_\mu^\alpha(y)}{\partial y^\nu} - g' f^{abc} \Gamma_\mu^b(y) \Gamma_\nu^c(y). \quad (109)$$

In the approximation where one takes $g' \rightarrow 0$ this lagrangian describes a system of $r = \dim(G)$ noninteracting U(1)-gauge fields, i.e. a system of r free "electromagnetic" fields with the corresponding gauge fixing terms containing the fields Δ^a . After quantization of each type $\Gamma_\mu^a(x)$, ($a = 1, \dots, r$), of "electromagnetic" fields one obtains the commutation relations [23]

$$[\Gamma_\lambda^a(y), \Gamma_\nu^b(y')] = i g_{\lambda\nu} \delta^{ab} \mathcal{D}_0(y - y') - i(1 - \xi') \frac{\partial^2}{\partial y^\lambda \partial y^\nu} E(y - y'),$$

where

$$D_0(y) = i \int \frac{d^4 k}{(2\pi)^3} e^{-iky} \epsilon(k_0) \delta(k^2), \quad (110)$$

$$E(y) = i \int \frac{d^4 k}{(2\pi)^3} e^{-iky} \epsilon(k_0) \delta'(k^2).$$

Going back to the fields A_λ^a and the variables (x^μ) , one recovers equation (101) having in mind that

$$D_0(b^{-1} \cdot x) = F(x), \quad \text{and} \quad E(b^{-1} \cdot x) = H(x). \quad (111)$$

7. Physical subspace of the Fock space

We introduce the Fourier integral for the fields $A_\lambda^d(x)$ and $B^a(x)$,

$$A_\lambda^d(x) = \frac{1}{(2\pi)^{3/2}} \int d^4 p \theta(p_0) \left[e^{ipx} a_\lambda^{+d}(p) + e^{-ipx} a_\lambda^{-d}(p) \right], \quad (112)$$

$$B^a(x) = \frac{1}{(2\pi)^{3/2}} \int d^4 p \theta(p_0) \left[e^{ipx} b^{+a}(p) + e^{-ipx} b^{-a}(p) \right]. \quad (113)$$

As a consequence of the equations of motion one obtains

$$\theta(k^0)[Q(k)a_\lambda^{\pm d}(k) \pm i(\mu - \xi)b^{\pm d}(k)] = 0, \quad (114)$$

$$\theta(k^0)Q(k)b^{\pm d}(k) = 0, \quad (115)$$

$$\pm i\theta(k^0)[k^\lambda a_\lambda^{\pm d}(k) + (V \cdot k)V^\lambda a_\lambda^{\pm d}(k)] = \theta(k^0)\xi b^{\pm d}(k). \quad (116)$$

Inserting equation (112) into (101) gives the commutation relations

$$[a_\lambda^{+d}(k), a_\mu^{+c}(q)] = 0, [a_\lambda^{-d}(k), a_\mu^{-c}(q)] = 0 \quad (117)$$

and

$$\begin{aligned} [a_\sigma^{-d}(p), a_\tau^{+c}(q)] = & -\mu(g_{\sigma\tau} - \frac{\beta}{1+\beta}V_\sigma V_\tau)\delta^{dc}\delta(Q(p))\delta(p-q) \\ & - (\mu - \xi)\delta^{dc}p_\sigma p_\tau \delta'(Q(p))\delta(p-q). \end{aligned} \quad (118)$$

For the operators $b^{\pm d}(p)$ one finds

$$[b^{+d}(p), b^{+c}(q)] = 0, [b^{-d}(p), b^{-c}(q)] = 0 \quad (119)$$

and

$$[a_\lambda^{-d}(p), b^{+c}(q)] = -\delta^{dc}p_\lambda \delta(Q(p))\delta(p-q). \quad (120)$$

It is convenient for each four vector p to introduce a local frame consisting of the orthonormal vectors V , ($V^2 = 1$), $N(p)$ and $e^{(r)}(p)$, ($r=1,2$), satisfying

$$N(p)^2 = (e^{(r)}(p))^2 = -1, \quad (121)$$

where

$$N(p)_\mu \equiv \frac{\tilde{p}_\mu - (V \cdot \tilde{p})V_\mu}{\sqrt{(V \cdot \tilde{p})^2 - \tilde{p}^2}} \quad (122)$$

and $\tilde{p}_\mu \equiv p_\mu + \beta V_\mu(V \cdot p)$. We decompose the operators $a_\lambda^{\pm d}(p)$ with respect to the base vectors V , $N(p)$, $e^{(r)}(p)$,

$$a_\lambda^{\pm d}(p) = \sum_{r=1}^2 \alpha_r^{\pm d}(p)e_\lambda^{(r)}(p) + \alpha_3^{\pm d}N_\lambda + \alpha_0^{\pm d}V_\lambda. \quad (123)$$

The commutation relations for the new operators $\alpha_\lambda^{\pm d}(p)$ read

$$[\alpha_0^{-d}(p), \alpha_0^{+c}(q)] = -\frac{\mu\delta^{dc}}{1+\beta}\delta(Q(p))\delta(p-q) - (\mu - \xi)\delta^{dc}(V \cdot p)^2\delta'(Q(p))\delta(p-q), \quad (124)$$

$$[\alpha_3^{-d}(p), \alpha_3^{+c}(q)] = \mu\delta^{dc}\delta(Q(p))\delta(p-q) - (\mu - \xi)\delta^{dc}[(V \cdot p)^2 - p^2]\delta'(Q(p))\delta(p-q), \quad (125)$$

$$[\alpha_k^{-d}(p), \alpha_l^{+c}(q)] = \mu\delta^{dc}\delta(Q(p))\delta(p-q), \quad (126)$$

$$[\alpha_0^{-d}(p), \alpha_3^{+c}(q)] = -(\mu - \xi)\delta^{dc}\sqrt{(V \cdot p)^2 - p^2}(V \cdot p)\delta'(Q(p))\delta(p-q), \quad (127)$$

$$[\alpha_0^{-d}(p), \alpha_k^{+c}(q)] = 0, \quad (128)$$

$$[\alpha_3^{-d}(p), \alpha_k^{+c}(q)] = 0, \quad (129)$$

$$[\alpha_k^{-d}(p), b^{+c}(q)] = 0, \quad (130)$$

$$[\alpha_3^{-d}(p), b^{+c}(q)] = -i\delta^{dc}\sqrt{(V \cdot p)^2 - p^2}\delta(Q(p))\delta(p-q), \quad (131)$$

$$[\alpha_0^{-d}(p), b^{+c}(q)] = -i\delta^{dc}(V \cdot p)\delta(Q(p))\delta(p-q). \quad (132)$$

The vacuum vector $|0\rangle$ of the Fock space \mathcal{H}_F is defined by

$$\alpha_\lambda^{-d}(p)|0\rangle = 0.$$

It follows from the commutation relations equations (124)-(132) that the operators $\alpha_0^{+d}(p)$ and $\alpha_3^{+d}(p)$ do not generate states with positive definite norm squared due to the terms with $\delta'(Q(p))$. The physical subspace of \mathcal{H}_F is defined by

$$\mathcal{H}_{phys} = \left\{ |\Psi\rangle \in \mathcal{H}_F; b^{-d}(p)|\Psi\rangle = 0 \right\}. \quad (133)$$

According to equation (116) the condition $b^{-d}(p)|\Psi\rangle = 0$ is equivalent to

$$\tilde{p}a_\lambda^{-d}(p)|\Psi\rangle = 0,$$

implying

$$L^{-d}(p)|\Psi\rangle = 0, \quad (134)$$

where

$$L^{\pm d}(p) = \sqrt{(V \cdot p)^2 - p^2} \alpha_0^{\pm d}(p) - (V \cdot p) \alpha_3^{\pm d}(p) \quad (135)$$

for any $|\Psi\rangle \in \mathcal{H}_{phys}$. The states with norm squared ≥ 0 are generated by $\alpha_{1,2}^{+d}(p)$ and $L^{+d}(p)$ which are the operators commuting with $b^{-c}(q)$. However, the subspace \mathcal{H}_0 of \mathcal{H}_{phys} generated by $L^{+d}(p)$ consists of zero norm vectors. Therefore, the space of vectors with positive norm squared is defined by the quotient

$$\mathcal{H}' = \mathcal{H}_{phys}/\mathcal{H}_0. \quad (136)$$

The energy-momentum tensor $T_{\lambda\mu}$ derived from the lagrangian $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{gf}$ is

$$T_{\lambda\mu} = -\mathcal{H}_{\lambda\tau}^a \partial_\mu A^{a,\tau} + (b^{-2} \cdot A^a)_\lambda \partial_\mu B^a - g_{\lambda\mu} \mathcal{L}, \quad (137)$$

$(b^{-2})_\lambda^\sigma = (\delta_\lambda^\sigma + \beta V_\lambda V^\sigma)$, and may be written as

$$T_{\lambda\mu} = -\mathcal{H}_{\lambda\tau}^a F_\mu^{a,\tau} + \frac{1}{4} g_{\lambda\mu} (F_{\alpha\beta}^a H^{a,\beta\alpha})$$

up to a total divergence $\sim \partial^\tau (\mathcal{H}_{\lambda\tau}^a A_\mu^a)$. For the matrix elements of the corresponding four-momentum operator one finds

$$\langle \Phi | P_\lambda | \Phi' \rangle = \int d^3 k k_\lambda \langle \Phi | \sum_{r=1}^2 \alpha_r^{+a}(k) \alpha_r^{-a}(k) | \Phi' \rangle, \quad (138)$$

$|\Phi\rangle, |\Phi'\rangle \in \mathcal{H}'$.

In the gauge $\xi = \mu$ the free propagator $\langle 0 | T(A_\mu^b(x) A_\nu^d(y)) | 0 \rangle$ reads

$$(1/i) \mathcal{D}_{\mu\nu}^{bd}(x-y) \equiv \langle 0 | T(A_\mu^b(x) A_\nu^d(y)) | 0 \rangle = -i\mu (g_{\mu\nu} - \frac{\beta}{1+\beta} V_\mu V_\nu) \delta^{bd} F_c(x-y), \quad (139)$$

where

$$F_c(x-y) = \int \frac{d^4 k}{(2\pi)^4} \frac{e^{-ik(x-y)}}{Q(k) + i\epsilon}. \quad (140)$$

If $\beta = n^2 - 1 \gg 1$, i.e. $n^2 = \epsilon\mu \gg 1$, the tensor components in the r.h.s of equation (139) reduce to the matrix $diag[0, -1, -1, -1]$ in the co-moving frame where $(V^\mu) = (1, \vec{0})$. Estimates for $\mu \approx 1$ and $\epsilon \gg 1$ are given in [13, 14]. For such values of the permittivity only transverse and longitudinal virtual gluons contribute to the amplitude of processes of the type $f + f' \rightarrow f + f'$ where f, f' are fermions.

8. Summary

In this paper the in-medium Yang-Mills equations are derived for a gauge group G in the linear approximation. In this approximation they resemble the macroscopic Maxwell equations for $\dim(G)$ copies of the electromagnetic field. The medium characteristics - the conductivity tensor σ , the electric permittivity tensor ϵ and the magnetic permeability tensor μ are expressed in terms of retarded Green's functions in a way analogous to the case of the in-medium electrodynamics. The canonical quantization is performed for constant and diagonal ϵ and μ in order to simplify the Euler-Lagrange equations. A set of $\dim(G)$ scalar auxiliary fields B^a is introduced to avoid working with constrained systems. The commutation relations among the gauge potentials A_μ^a and the field B^a are derived in two ways for a family of Fermi-like gauges. The first one is straightforward but lengthy and employs an integral representation of the solutions of equation (98), while the second one is direct and based on a simple transformation of the coordinates and fields. For the gauge choice $z = \mu$ we write down the "gluon" propagator $\mathcal{D}_{\mu\nu}^{bd}$. If $\epsilon \gg 1$ as suggested by some studies of color superconductivity, $\mathcal{D}_{\mu\nu}^{bd}$ reduces essentially to its 3-space components. Thus it modifies e.g. the spinor-spinor scattering amplitude compared to their form evaluated via the standard gluon propagator. Computations in this direction will be published in a forthcoming paper.

References

- [1] G. K. Savvidy, Phys. Lett. B **71** (1977) 133.
- [2] R. Fukuda and Y. Kazama, Phys. Rev. Lett. **45** (1980) 1142.
- [3] S. L. Adler, Phys. Rev. D **23** (1981) 2905.
- [4] H. Nielsen and P. Olsen, Nucl. Phys. B **144** (1978) 376.
- [5] M. Baker and F. Zachariasen, Phys. Lett. B **108** (1982) 206.
- [6] M. Baker and F. Zachariasen, Nucl. Phys. B **226** (1983) 455.
- [7] D. Sivers, Phys. Rev. D **27** (1983) 947.
- [8] J. Ralston and D. Sivers, Phys-Rev. D **28** (1983) 953.
- [9] U. Heinz, Am. Phys. **161** (1985) 48.
- [10] U. Heinz, Am. Phys. **168** (1986) 148.
- [11] Chen Jisheng, Zheng Xiaoping and Li Jiarong, *The nonlinear permittivity including non-abelian self-interaction of plasmons in QGP*, hep-ph/9907369.
- [12] Ji-sheng Chen, Jia-rong Li and Peng-fei Zhuang, *Non-abelian medium effects in QGP*, hep-ph/0112128.
- [13] D. H. Rischke, D. T. - Song and M. A. Stephanov, Phys. Rev. Lett. **87** (2001) 062001.
- [14] D. Litim and C. Manuel, Phys. Rev. D **64** (2001) 094013.
- [15] J. Ruppert, G. Nayak, D. Dietrich, H. Stöcker and W. Greiner, Phys. Lett. B **520** (2001) 233.
- [16] R. Friedberg and T. D. Lee, Phys. Rev., D **15** (1977) 1694.
- [17] R. Friedberg and T. D. Lee, Phys. Rev., D **16** (1977) 1696.
- [18] U. Kalmbach, T. Vetter, T. S. Biró and U. Mosel, Nucl. Phys. A **563** (1993) 584.
- [19] T. Vetter, T. S. Biró and U. Mosel, Phys. Lett. A **581** (1995) 598.
- [20] S. Loh, T. S. Biró, U. Mosel and M. H. Thoma, Phys. Lett. B **387** (1996) 685.
- [21] C. T. Traxler, U. Mosel and T. S. Biró, Phys. Rev. C **59** (1999) 1620.
- [22] A. I. Akhiezer and S. V. Peletminsky, *Methods of Statistical Physics*, Pergamon, London, 1981.
- [23] I. Brevik and B. Lautrup, *Quantum Electrodynamics in Material Media*. Mat. Fys. Medd. Dan.

Vid. Selsk. 38, (1970) NORDITA.

[24] N. Nakanishi, *Progr. Theor. Physics*, **35** (1966) 1111.

[25] N. Nakanishi, *Suppl. Progr. Theor. Physics*, **51** (1972) 1.

[26] D. Zwanziger, *Phys. Rev. D* **17** (1978) 457.