

Bose-Einstein correlations in the Quantum Clan Model

O.V.Utyuzh^{1,a}, G.Wilk^{1,b} and Z.Włodarczyk^{2,c}

¹ The Andrzej Soltan Institute for Nuclear Studies;
Hoża 69; 00-681 Warsaw, Poland

² Institute of Physics, Świętokrzyska Academy,
Świętokrzyska 15; 25-406 Kielce, Poland

Received 1 January 2004

Abstract. We propose novel numerical method of *modelling* Bose-Einstein correlations (BEC) observed among identical (bosonic) particles produced in multiparticle production reactions. We argue that the most natural approach is to work directly in the momentum space in which the Bose statistics of secondaries reveals itself in their tendency to bunch in a specific way in the available phase space. Because such procedure is essentially identical to the clan model of multiparticle distributions proposed some time ago, therefore we call it the *Quantum Clan Model*.

Keywords: Bose-Einstein correlations; Statistical models; Fluctuations

PACS: 25.75.Gz 12.40.Ee 03.65.-w 05.30.Jp

The phenomenon of Bose-Einstein correlations (BEC) is so widely know [1] that we shall start directly with problem of its *proper numerical modelling*, such which would account from the very beginning for the quantum statistical bosonic character of identical secondaries produced in hadronization process. To our knowledge this problem was so far considered only in [2](cf., however, [3]). All other approaches claiming to model BEC numerically [4] use as their starting point the outcomes of existing Monte-Carlo event generators (MCG) describing multiparticle production process [5] and *modify* them in a suitable way to fit the BEC data. These modifications are called *afterburners*. They inevitably lead to such unwanted features as violation of energy-momentum conservation or to changes in the original multiparticle spectra. Actually, it is worth to mention at this point that in [6] we have proposed afterburner free from such unwanted effects. It was based on different concept of introducing quantum mechanical (QM) effects in the otherwise purely probabilistic distributions from those proposed in [7]. Namely, each MCG provides us usually with a given number of particles, each one endowed with one of (+/–/0) charge and with well defined spatio-temporal position and energy-momentum. On

the other hand experiment provides us information on only the first and last characteristics. The spatio-temporal information is not available directly (in fact, the universal hope expressed in [1, 4] is it can be deduced from the previous two via the measured BEC). Our reasoning was as follows: (i) BEC phenomenon is of the QM origin, therefore one has to introduce in the otherwise purely classical distributions provided by MCG a new element mimicking QM uncertainties; (ii) this cannot be done with energy-momenta because they are measured and therefore fixed; (iii) the next candidate, i.e., spatio-temporal characteristics, can be changed but this was already done in [7, 4]; (iv) one is thus left with charges and in [6] we have simply assigned (on event-by-event basis) new charges to the particles selected by MCG conserving, however, the original multiplicity of (+/- /0). This has been done in such way as to make particles of the same charge to be located maximally near to each other in the phase space by exploring natural fluctuations of spatio-temporal and energy-momentum characteristic resulting from MCG. In this way automatically conserve all energy-momenta and do not change multiparticle distributions and do it already on the *level of single event* provided by MCG . However, the new assignment of charges introduces a profound change in the structure of the original MCG. Generally speaking (cf. [6] for details) it is equivalent to introduction in the MCG used the bunching of particles of the same charge.

This observation will be the cornerstone of our new proposition. Let us remind that idea of bunching of particles as quantum statistical (QS) effect is not new [8]. It was used in connection with BEC for the first time in [9] and later it was a cornerstone of the *clan model* of multiparticle distributions $P(n)$ leading in natural way to their negative binomial (NB) form observed in experiment [10]. It was introduced in the realm of BEC in [11] and [2, 3]. Because our motivation comes basically from [2] let us outline shortly its basic points. It deals with the problem of how to distribute in a least biased way a given number of bosonic secondaries, $\langle n \rangle = \langle n^{(+)} \rangle + \langle n^{(-)} \rangle + \langle n^{(0)} \rangle$, $\langle n^{(+)} \rangle = \langle n^{(-)} \rangle = \langle n^{(0)} \rangle$. Using information theory approach (cf., [12]) their rapidity distribution was obtained in form of grand partition function with temperature T and chemical potential μ . In addition, the rapidity space was divided into *cells* of size δy (fitted parameter) each. It turned out that whereas the very fact of existence of such cells was enough to obtain reasonably good multiparticle distributions, $P(n)$, (actually, in the NB-like form), their size, δy , was crucial for obtaining the characteristic form of the 2-body BEC function $C_2(Q = |p_i - p_j|)$ (peaked and greater than unity at $Q = 0$ and then decreasing in a characteristic way towards $C_2 = 1$ for large values of Q) out of which one usually deduces the spatio-temporal characteristics of the hadronization source [1] (see [2] for more details). The outcome was obvious: to get C_2 peaked and greater than unity at $Q = 0$ and then decreasing in a characteristic way towards $C_2 = 1$ for large values of Q one must have particles located in cells in phase space which are of nonzero size¹.

¹It means then that from C_2 one gets not the size of the hadronizing source but only size of the emitting cell, in [2] $R \sim 1/\delta y$, cf. [13]. In the quantum field theoretical formulation of BEC this directly corresponds to the necessity of replacing delta functions in commutator relations by a well




Fig. 1. The proposed algorithm is similar to the classical clan model proposed in [10] but its clans contain particles of the same charge and (almost) the same energies and are distributed according to geometrical distribution what results in overall Pòlya-Aepli distribution, $P_{PA}(n)$ [16], instead of NB one, $P_{NB}(n)$.

To illustrate our proposition let us assume that mass M hadronizes into $N = \langle n \rangle$ bosonic particles (we take them as pions of mass m) with equal numbers of (+/-/0) charges and with limited transverse momenta p_T . Suppose that their multiplicity distribution $P(n)$ follows a NB-like form (i.e., it is broader than Poissonian) and that their two-particle correlation function of identical particles, $C_2(Q)$, has the specific BEC form mentioned above. To model such process accounting from the very beginning, for the bosonic character of produced particles we propose the following steps (illustrated by comparison to some selected LEP e^+e^- data [17], cf., Fig. 1):

(1) Using some (assumed) function $f(E)$ select a particle of energy $E_1^{(1)}$ and charge $Q^{(1)}$. The actual form of $f(E)$ should reflect somehow our *a priori* knowledge of the particular collision process under consideration. In what follows we shall assume that $f(E) = \exp(-E/T)$, with T being parameter (playing in our example the role of "temperature").

(2) Treat this particle as seed of the first *elementary emitting cell* (EEC) (introduced in [11]) and add to it, until the first failure, other particles of the same charge $Q^{(1)}$ selected according to distribution $P(E) = P_0 \cdot f(E)$, where P_0 is another parameter (playing the role of "chemical potential" $\mu = T \cdot \ln P_0$). This assures that the number of particles in this EEC, k_1 , will follow geometrical (or Bose-Einstein)

defined peaked functions introducing in this way same dimensional scale to be obtained from fits to data [14]. This fact was known even before but without any phenomenological consequences [15].

distribution, i.e., that

$$\langle k_1 \rangle = P(E)/([1 + P(E)]),$$

and accounts therefore for their bosonic character. As result $C_2(Q) > 1$ but only *at one point*, namely for $Q = 0$.




Fig. 2. Examples of confrontation with the experimental data [17]. Left panel: fit to charge multiplicity distribution. Right panel: results for $C_2(Q = |p_1 - p_2|)$ correlation function (one dimensional phase space was used here only). Two different sets of parameters have been used. Notice that whereas they lead to essentially similar $P(n)$ the resulting $C_2(Q)$ are drastically different.

(3) To get the experimentally observed width of $C_2(Q)$ one has to allow that particles in each EEC can have (slightly) different energies from energy of the particle being its seed. To do it one must allow that each additional particle selected in point (2) above have energy $E_i^{(1)}$ selected from some distribution function peaked at $E_1^{(1)}$, $G(E_1^{(1)} - E_i^{(1)})$, where the width of this distribution, σ , is another free parameter².

(4) Repeat points (1) - (3) as long as there is enough energy left. Correct in every event for every energy-momentum nonconservation caused by the selection procedure adopted and assure that $N^{(+)} = N^{(-)}$.

As result in each event we get a number of EEC with particles of the same charge and (almost) the same energy, i.e., picture closely resembling classical *clans* of [10] (with no effects of statistics imposed, see Fig. 1). Our clans (containing

²It reflects situation encountered in [14] where, as we have already mentioned before, the observed shape of $C_2(Q)$ was coming from the assumed shape of function replacing Dirac delta function in energy, i.e., introducing smearing in energy.

now identical bosonic particles subjected to quantum statistics and therefore named *quantum clans*) are distributed in the same way as the particles forming seeds for EEC, i.e., according to Poisson distribution. With particles in each clan distributed according to geometrical distribution they lead therefore to the overall distribution in our Quantum Clan Model case being of the so called Pölya-Aeppli type [16]. This distribution strongly resembles the Negative Binomial distributions obtained in the classical clan model [10] where particles in each clans were assumed to follow *logarithmic* distribution instead (with differences occurring for small multiplicities [16]). The first preliminary results presented in Fig. 2 are quite encouraging (especially when one remembers that so far effects of resonances and all kind of final state interactions to which C_2 is sensitive were neglected here). It remains now to be checked what two-body BEC functions for other components of the momentum differences and how they depend on the EEC parameters: T , P_0 and σ . So far the main outcome is that BEC are due to EEC's only and therefore provide us mainly with their characteristics. This should clear at least some of many apparently "strange" results obtained from BEC recently [18].

Acknowledgment(s)

Two of us (OU and GW) are grateful for support from the Hungarian Academy of Science and for warm hospitality at the BudaRhic2004 meeting. Partial support of the Polish State Committee for Scientific Research (KBN) (grant 621/E-78/SPUB/CERN/P-03/DZ4/99 (GW)) is acknowledged.

Notes

- a. utyuzh@fuw.edu.pl
- b. wilk@fuw.edu.pl
- c. wlod@pu.kielce.pl

References

1. Cf., for example, T. Csörgő, in *Particle Production Spanning MeV and TeV Energies*, eds. W.Kittel et al., NATO Science Series C, Vol. 554, Kluwer Acad. Pub. (2000), p. 203 (see also: hep-ph/0001233); W. Kittel, Acta Phys. Polon. **B32** (2001) 3927; G. Alexander, Rept. Prog. Phys. **66** (2003) 481. (and references therein).
2. T. Osada, M. Maruyama and F. Takagi, Phys. Rev. **D59** (1999) 014024.
3. W. Zajc, Phys. Rev. **D53** (1987) 3396; R.L. Ray, Phys. Rev. **C57** (1998) 2352; J.G.Cramer, *Event Simulation of High-Order Bose-Einstein and Coulomb Correlations*, Univ. of Washington preprint (1996, unpublished).
4. L. Lönnblad and T. Sjöstrand, Eur. Phys. J. **C2** (1998) 165; K. Fiałkowski, R. Wit and J. Wosiek, Phys. Rev. **D58**(1998) 094013; T.Wibig, Phys. Rev.

- D53** (1996) 3586. B. Andersson, Acta Phys. Polon. **B29** (1998) 1885; K. Geiger, J. Ellis, U. Heinz and U.A. Wiedemann, Phys. Rev. **D61**(2000) 054002.
5. K.J. Eskola, Nucl. Phys. **A698** (2002) 78; S.A.Bass et al. Prog. Part. Nucl. **41** (1998) 225.
 6. O.V. Utyuzh, G. Wilk and Z. Włodarczyk, Phys. Lett. **B522** (2001) 273 and Acta Phys. Polon. **B33**(2002) 2681.
 7. A. Białas and A. Krzywicki, Phys. Lett. **B354**, (1995) 134; H. Merlitz and D. Pelte, Z. Phys. **A351**(1995) 187 and **A357** (1997) 175; U.A. Wiedemann et al., Phys. Rev. **C56** (1997) R614; T. Csörgő and J. Zimányi, Phys. Rev. Lett. **80** (1998) 916 and Heavy Ion Phys. **9** (1999) 241.
 8. W.J. Knox, Phys. Rev. **D10** (1974) 65; E.H. De Groot and H. Satz, Nucl. Phys. **B130** (1977) 257; J. Kripfganz, Acta Phys. Polon. **B8**(1977) 945; A.M. Cooper, O. Miyamura, A. Suzuki and K. Takahashi, Phys. Lett. **B87** (1979) 393; F. Takagi, Prog. Theor. Phys. Suppl. **120** (1995) 201.
 9. E.E. Purcell, *Nature* **178** (1956) 1447; A. Giovannini and H.B. Nielsen, *Stimulated emission model for multiplicity fluctuations*, The Niels Bohr Institute preprint, NBI-HE-73-17 (unpublished) and *Stimulated emission effect on multiplicity distribution* in: Proc. of the IV Int. Symp. on Multip. Hadrodynamics, Pavia 1973, Eds. F. Duimio, A. Giovannini and S. Rattii, p. 538.
 10. A. Giovannini and L. Van Hove, Z. Phys. **C30** (1986) 381; see also: P. Carruthers and C.S. Shih, Int. J. Mod. Phys. **A2** (1986) 1447.
 11. M. Biyajima, N. Suzuki, G. Wilk and Z. Włodarczyk, Phys. Lett. **B386** (1996) 297.
 12. F.S. Navarra, O.V. Utyuzh, G. Wilk and Z. Włodarczyk, Phys. Rev. **D67** (2003) 114002.
 13. W.A. Zajc, *A pedestrian's guide to interferometry*, in "Particle Production in Highly Excited Matter", eds. H.H.Gutbrod and J.Rafelski, Plenum Press, New York 1993, p. 435.
 14. G.A. Kozlov, O.V. Utyuzh and G. Wilk, Phys. Rev. **C68** (2003) 024901 and Ukr. J. Phys. **48** (2003) 1313.
 15. K. Zalewski, Lecture Notes in Physics **539**(2000) 291.
 16. J. Finkelstein, Phys. Rev. **D37** (1988) 2446; Ding-wei Huang, Phys. Rev. **D58** (1998) 017501.
 17. P.Abreu et al. (DELPHI Collab.), Phys. Lett. **B286** (1992) 201 and Phys. Lett. **B247** (1990) 137.
 18. J.T. Mitchell, J. Phys. **G30** (2004) S819; H. Appelhäuser, *ibidem*, S935.