Second order brane cosm ology with radion stabilization

GL. Alberghi, 1, 2, 3, D. Bom bardelli, 1, 3, y R. Casadio, 1, 3, z and A. Tronconi1, 3, x

¹Department of Physics, University of Bologna, V ia Innerio 46, 40126 Bologna, Italy.

²Department of Astronomy, University of Bologna, Italy.

³I.N.F.N., Sezione di Bologna, Italy.

We study cosmology in the ve-dimensional Randall-Sundrum brane-world with a stabilizing e ective potential for the radion and matter localized on the branes. The analysis is performed by employing a perturbative expansion in the ratio =V between the matter energy density on the branes and the brane tensions around the static Randall-Sundrum solution (which has = 0 and brane tensions V). This approach ensures that the matter evolves adiabatically and allows us to not approximate solutions to second order in =V. Some particular cases are then analyzed in details.

PACS num bers: 12.60.-i, 04.50.+ h, 98.80.C q, 98.80.-k

I. IN TRODUCTION

H igher and higher precision data which are about to be collected in new experiments of particle physics and astrophysics in the next few years convey considerable attention to theories with extra dimensions. The main role of such theories, originally introduced in the 20's by K aluza and K lein [1, 2], is to provide a connection between particle physics and gravity at some level. At a deeper level, string theory uni es all the interactions by means of some n-dimensional manifold (with n > 4) where the fundamental objects are supposedly living; at a more phenomenological level, models which assume the existence of extra dimensions, no matter their origin, are considered in order to solve some puzzles of particle physics, cosmology and astrophysics, giving rise to many possible observable consequences.

Originally proposed in order to solve the problem of the large hierarchy between G ravity and Standard M odel scales, the Randall-Sundrum model of Ref. [3] (RS I) has acquired considerable relevance due to its stringy inspiration. It represents the prototype of the so-called braneworld and diers from previous models in that it constrains standard matter on a four-dimensional manifold (the brane) just letting gravity (and exotic m atter) propagate everywhere. The RS I solution to the hierarchy problem needs one additional com pacti ed (orbifolded) spatial dimension with two branes located at its xed points, plus a negative cosm ological constant lling the space between such branes (the bulk). The bulk cosm ological constant warps the extra dimension and generates the e ective four-dimensional physical constants we measure. It was soon realized that the modi cations to four-dimensional gravity induced by the fth dimension may be reduced to such a short distance e ect to

be unobservable even in the presence of just one brane and in nite compacti cation radius (the RS IIm odel of Ref. [4]).

The cosm ological features of the RS m odels are now adays being investigated even m ore than its particle physics consequences, due to the re ned results lately obtained and to the m a jor problem s recent astrophysical data have revealed: the possible late time acceleration from supernovae, CMBR spectrum, dark m atter and dark energy quests suggest either a full revision of the m odern theoretical physics approach or the possibility of the existence of further, up to now ignored, ingredients such as the extra dimensions.

In particular the single brane RS II cosm ological dynam ics [5, 6] is known to generate $(=V)^2$ corrections to standard Friedm ann and acceleration equations, where is the energy density of the uid lling the brane and V is the constant brane tension. These corrections are neg-V, the regime in which the RS II m odel ligible when is reliable and leads to standard cosm ic evolution. The two brane RS I setup is much more involved: a stabilization mechanism for the distance between the branes, such as that of R ef. [7], is necessary to get the correct hierarchy in the absence of matter. Moreover, a bulk potential for the radion (the metric degree of freedom associated with the ffh dimension) is necessary to achieve solvable junction conditions when m atter is present on the boundaries [8]. In this case cosm ological solutions are known up to order =V [9]. Such solutions are in fact insu cient to grasp the particular features of the background m etric evolution originated by the extra dimension.

The aim of this article is to go beyond the rst order approximation in brane cosmology for RS I models with two branes. The calculations are carried out assuming an elective bulk potential for the radion and will show how ² contributions to the four-dimensional Hubble parameterm ay a lect the model (orm ay be unobservable). Such terms are expected as ngerprints of the flh dimension in analogy with the single brane RS II fram ework. Furtherm ore some hints about the possibility of an accelerated expansion driven by exotic uids with p = w and w > 0 will be presented, thus suggesting the necessity to

E lectronic address: alberghi@ bo.infn.it

^yE lectronic address: bom bardelli@ bo.infn.it

^zE lectronic address: casadio@ bo.infn.it

^xE lectronic address: tronconi@ bo.infn.it

go beyond the second order approxim ation.

The paper is organized as follows: in Section II, we present the complete setup of the model under consideration; in Section III the second order ansatz is described and E instein equations are perturbatively solved; in Section IV cosm ological consequences of the solutions are analyzed and compared to the known brane-world solutions; in Section V the analysis of the approximations is performed and, nally, in Section V I, some conclusions are drawn. For the ve-dimensionalmetric g_{AB} we shall use the signature (+; ; ; ;), so that g detage) > 0.

II. EINSTEIN EQUATIONS

Let us consider a RS Im odelperturbed by the presence of matter on the two branes. The bulk metric is given by

$$ds^{2} \quad g_{AB} dx^{A} dx^{B}$$

= n² (y;t) dt² a² (y;t) dxⁱ dxⁱ b² (y;t) dy² :(1)

The Einstein tensor for this metric is

ab

$$G_{44}=3 \quad \frac{a^0}{a} \quad \frac{a^0}{a} + \frac{n^0}{n} \quad \frac{b^2}{n^2} \quad \frac{a}{a} \quad \frac{a}{a} \quad \frac{n}{n} + \frac{a}{a} \quad (2d)$$

where a prime denotes a derivative with respect to y and a dot a derivative with respect to the universal time t. The energy-momentum tensor in the bulk is that of an anti-de Sitter space with a correction coming from the e ective potential for the radion [10]

$$T_{B}^{A} = g_{B}^{A} + T_{B}^{A}$$
 (b); (3)

where, as usual,

$$T_{AB}(b) = \frac{2}{p_{\overline{g}}} \frac{L_{eff}}{g^{AB}} : \qquad (4)$$

W e shall consider the particular case of a harm onic potential for the radion, with an e ective Lagrangian of the form

$$L_{eff} = {}^{p}\overline{g}!^{2}(b \quad b)^{2} \qquad {}^{p}\overline{g}U: \qquad (5)$$

The two 3-branes have opposite tensions and their contribution to the total energy-momentum tensor is given by

$$\Gamma_{iB}^{A} = \frac{(V \ \underline{N})}{b}$$
diag (V_i + _i;V_i + p_i;V_i + p_i;V_i + p_i;0) ;(6)

where i = p, n, and $y_p = 0$ ($y_n = 1=2$) is the position of the positive (negative) tension brane. The Einstein equations in the bulk,

$$G_{AB} = k^2 T_{AB} ; \qquad (7)$$

form a system of four di erential equations for the three independent functions f = (n;a;b), which, at rst sight, appears over-determ ined. How ever, on using the B ianchi identity $r_A G^{A 0} = 0$, it is straightforward to show that only three equations are independent and that the system is indeed well de ned. O ne can therefore choose to solve

or equivalently (see Refs. [5, 6])

with

F (y;t) =
$$\frac{(a a^0)^2}{b^2} - \frac{(a \underline{a})^2}{n^2}$$
: (10)

If T_{0}^{0} does not depend on y, the rst equation in (9) can be replaced by

$$F(y;t) + \frac{k^2}{6} a^4 T_0^0 = e(t);$$
 (11)

whereas, in general, one should use the form

$$\frac{a^{2}}{a^{2}} = \frac{n^{2} a^{02}}{b^{2} a^{2}} = \frac{k^{2}}{6}n^{2} T_{0}^{0} + \frac{k^{2} n^{2}}{6a^{4}} = \frac{k^{4}}{a^{4}} T_{0}^{0} dy$$

$$= \frac{n^{2}}{a^{4}} e(t) ; \qquad (12)$$

where c(t) is related to the boundary conditions.

In the presence of the e ective potential (5) the equation of motion for the (unspecified) stabilizing eld gives rise to the conservation equation $r_A T^{A4} = 0$, that is

$$b^{0} = (b_{0} \quad b) \quad \frac{n^{0}}{n} + 3\frac{a^{0}}{a} + 2\frac{b^{0}}{b} ;$$
 (13)

which is identically satis ed by the solutions of the system (9). One can therefore replace one of the equations

in (9) with Eq. (13), as we shall nd convenient in the following.

M oreover, bulk solutions must satisfy the boundary equations given by the junction conditions on the two branes,

$$\overset{8}{\underset{y! \ y_{i}^{+}}{\underset{y! \ y_{i}^{+}}{\overset{a^{0}}{a}}} = \frac{k^{2}}{6} (V_{i} + i) b_{y=y_{i}}$$
(14)
$$\overset{8}{\underset{y! \ y_{i}^{+}}{\underset{y! \ y_{i}^{+}}{\overset{n^{0}}{n}}} = \frac{k^{2}}{6} [V_{i} (2 + 3w_{i})_{i}] b_{y=y_{i}};$$

where we assumed an equation of state for the vacuum perturbations of the form $p_i = w_i$. When i! 0 the RS I solution is fully recovered and one nds the usual warped static metric with c(t) = 0 and

$$n_{RS} (y;t) = a_{RS} (y;t) = \exp(m b_0 jy)$$

 $b_{RS} = b_0$; (15)

which also require the well known ne-tuning

$$V_{\rm p} = V_{\rm n} = \frac{6m}{k^2}$$
; $= \frac{6m^2}{k^2}$: (16)

III. THE CARDASSIAN EXPANSION

The Cardassian expansion can be adopted in brane cosm ology to investigate solutions to the E instein equations by taking as a starting point the static RS Im etric with $_{\rm i} = 0$ reviewed in the previous section. In fact, in the low density regime

$$\frac{i}{\dot{y}_{ij}}$$
 1; (17)

one can express the corrections to the solution (15) to all orders in $_i=V_i$ by assuming that the metric functions f = (n;a;b) can be written as [11]

$$f = f_{RS} + f$$
; (18)

with f P $P_{n_i;n_j \ 1} C_{n_in_j \ i \ j}$.

In order to keep track of the various orders in the above expansion, it is useful to introduce an expansion param – eter by replacing i ! i (and setting = 1 at the end of the computation). We shall also nd convenient to write the perturbed metric as

$$n(y;t) = \exp(m \log y) [1 + f_n(y;t)]$$
 (19a)

$$a(y;t) = a_h(t) \exp(m b_j y) [1 + f_a(y;t)] (19b)$$

$$b(y;t) = b_0 + f_b(y;t);$$
 (19c)

so that the hom ogeneous scale factor a_h (t) is factored out, and b_0 is the equilibrium point corresponding to the RS I model [see Eq. (15) above]. The Einstein equations (9) to second order in can then be completely expressed in terms of the functions

$$f = \begin{pmatrix} h & i \\ f_{p}^{(1)}(y) & p + f_{p}^{(1)}(y) & n \\ h & i \\ + {}^{2} f_{p}^{(2)}(y) & p + f_{p}^{(2)}(y) & 1 \\ + {}^{2} f_{p}^{(2)}(y) & p + f_{p}^{(2)}(y) & 1 \\ + {}^{2} f_{p}^{(2)}(y) & p + f_{p}^{(2)}(y) & 1 \\ + {}^{2} f_{p}^{(2)}(y) & 1$$

$$= h_{h,p}^{(1)} + h_{h,n}^{(1)} + \frac{2}{h_{h,p}^{(2)}} + h_{h,n}^{(2)} + h_{h,n}^{(2)} + h_{h,m}^{(2)} + h_{h$$

c(t)
$$\frac{e}{a_{h}^{4}}$$

= $c_{p}^{(1)}{}_{p} + c_{h}^{(1)}{}_{n}$
+ ${}^{2} c_{p}^{(2)}{}_{p}^{2} + c_{h}^{(2)}{}_{n}^{2} + c_{m}^{(2)}{}_{p}{}_{n}$; (20c)

and their derivatives. Note that the time dependence is just carried by the functions $_{i} = _{i}(t)$ and that, of all the coe cients appearing above, only the f $_{i}^{(n)}$'s in Eq. (20a) depend on y, whereas the others are constant.

In order to proceed with the Cardassian expansion, one also needs to expand _i. This can be obtained by taking the limit y ! y_i in the equation G₀₄ = 0 to recover the usual conservation equation

$$\underline{i} = 3H (\underline{y};t) (1 + w_{i}) \underline{i}$$
: (21)

From the above, it immediately follows that the time evolution of the matter densities is adiabatic, since H $\underline{a}=\underline{a}$ $\mathbf{1}=2$ and, therefore,

$$\frac{j_{i}j_{j}}{j_{i}} = \frac{j_{i}}{j_{i}} \frac{j_{i}}{j_{i}} = 1:$$
 (22)

If we now assume that, to second order in ,

$$\begin{array}{ccccccccc} & h & i \\ H^{2}(y;t)' & h_{p}^{(1)}(y)_{p} + h_{n}^{(1)}(y)_{n} & (23) \\ & h & i \\ & + {}^{2} & h_{p}^{(2)}(y)_{p} + h_{n}^{(2)}(y)_{n} + h_{m}^{(2)}(y)_{p} & n \end{array}$$

the coe cients of the above expansion can be related to the corresponding ones in Eqs. (20b) and (20a) for = a by equating the two expressions for H 2 at y = y_i up to second order,

$$H^{2}(y_{i};t) = H_{h}(t) + \frac{f_{a}(y_{i};t)}{1 + f_{a}(y_{i};t)} : (24)$$

W ith the help of Eq. (21), one nally obtains

$$h_{i}^{(1)}(y) = h_{h,i}^{(1)}$$
 (25a)

$$h_{i}^{(2)}(y) = h_{h,i}^{(2)} \quad 6(1 + w_{i}) h_{h,i}^{(1)} f_{a,i}^{(1)}(y)$$
 (25b)

$$h_{m}^{(2)}(y) = h_{h,m}^{(2)} \quad 6(1 + w_{p}) h_{h,n}^{(1)} f_{a,p}^{(1)}(y)$$

$$6(1 + w_{n}) h_{h,p}^{(1)} f_{a,n}^{(1)}(y) : \qquad (25c)$$

A . First order results

In order to solve the bulk equations order by order, one has to substitute the previous expansion in the dynam ical equations. This will allow us to determ ine explicitly the ∞ explicitly on ∞ the boundary conditions are in –

posed. Let us begin with rst order equations. At order, the constraint (13) reads

$$f_{b,p}^{(1)^{0}} + f_{b,n}^{(1)^{0}} + 4m b_{b} f_{b,p}^{(1)} + f_{b,n}^{(1)} = 0 : (26)$$

If we allow $_{\rm p}$ and $_{\rm n}$ to be arbitrary functions of the time, the above equation splits into two independent equations for the coe cients of $_{\rm i}$, and one nds

$$f_{b;i}^{(1)}(y) = b_i^{(1)} \exp (4 m b_0 y)$$
; (27)

where $\boldsymbol{b}_i^{(1)}$ are constant coe cients to be determ ined.

The functions $f_{a;i}^{(1)}(y)$'s can now be calculated by solving Eq. (12). Since the contribution of the stabilizing potential vanishes at order , the integral-di erential equation (12) becomes the rst order linear di erential equation

$$e^{2m b_0 y} = 2m^2 b_i^{(1)} e^{4m b_0 y} = b_i^{(1)} e^{4m b_0 y} = b_i^{(1)} e^{4m b_0 y} + b_0 h_{h,i}^{(1)} e^{2m b_0 y} + 2m f_{a,i}^{(1)^0} = 0 :$$
(28)

On solving for each coe cient of $_{i}$ independently, one obtains

$$f_{a,i}^{(1)} = \frac{1}{8} e^{2m \ b_0 \ y} e^{2m \ b_0 \ y} \frac{c_i^{(1)}}{m^2} 2\frac{b_i^{(1)}}{b_0} 2\frac{h_{h,i}^{(1)}}{m^2} + c_{a,i}^{(1)}$$
(29)

F inally, one can expand the equation $G_{44} = k^2 T_{44}$ to storder,

and solve the two equations for $f_{n\,;i}^{(1)}\left(y\right)$. The result is

$$f_{n,i}^{(1)} = \frac{1}{2} e^{2m \ b_0 \ y} \ e^{2m \ b_0 \ y} \ \frac{k^2 \ !^2}{3m^2} \ b_0 \ b_i^{(1)} \ \frac{3 \ c_i^{(1)}}{4m^2} \ \frac{b_i^{(1)}}{2b_0} + \frac{(2 + 3w_i)}{2m^2} \ h_{h,i}^{(1)} \ + \ c_{n,i}^{(1)} \ (31)$$

W e are now left with six num erical coe cients

$$h_{h;i}^{(1)}$$
; $b_{i}^{(1)}$; $c_{i}^{(1)}$; (32a)

and four integration constants

$$c_{a;i}^{(1)}$$
; $c_{n;i}^{(1)}$: (32b)

In order to x the above, one has to use the junction conditions. These four conditions, written in terms of the coe cients of $_{i}$, form a system of eight equations: the discontinuity constraints for $a^0(y;t)$ at $y = y_i$ im ply

$$\begin{cases} 8 \\ p & 3c_{p}^{(1)} & 3h_{h,p}^{(1)} + k^{2}m + 3 \\ n & c_{n}^{(1)} & h_{h,n}^{(1)} = 0 \end{cases}$$

$$\begin{cases} 3 \\ p & c_{p}^{(1)}e^{m b_{0}} & h_{h,p}^{(1)} \\ + \\ n & 3c_{n}^{(1)}e^{2m b_{0}} & 3h_{h,n}^{(1)}e^{m b_{0}} & k^{2}m = 0; \end{cases}$$

$$(33)$$

whose solution is given by

$$h_{h,p}^{(1)} = \frac{k^2 m e^{m b_0}}{3 (e^{m b_0} 1)}; \qquad h_{h,n}^{(1)} = \frac{k^2 m e^{m b_0}}{3 (e^{m b_0} 1)} (34a)$$

$$c_p^{(1)} = \frac{k^2 m}{3 (e^{m b_0} 1)}; \quad c_n^{(1)} = \frac{k^2 m e^{m b_0}}{3 (e^{m b_0} 1)}; \quad (34b)$$

The two analogous constraints for $n^0(y;t)$ are both equivalent to the equation

$$\begin{array}{c} h & i \\ e^{m \ b_0} \ m \ (3 \ w_p \ 1) + 4 \ !^2 \ b_0 \ b_p^{(1)} \ e^{m \ b_0} \ 1 \ p \ (35) \\ h & i \\ + \ m \ (3 \ w_n \ 1) + 4 \ !^2 \ b_0 \ b_n^{(1)} e^{m \ b_0} \ e^{m \ b_0} \ 1 \ t = 0 \end{array}$$

which yields

$$b_{p}^{(1)} = \frac{m (3w_{p} \ 1)}{4!^{2}b_{0} (e^{m \ b_{0}} \ 1)}$$

$$b_{n}^{(1)} = \frac{m (3w_{n} \ 1)e^{m \ b_{0}}}{4!^{2}b_{0} (e^{m \ b_{0}} \ 1)} ;$$
(36)

W e see that the junction conditions are not su cient to determ ine the integration constants (32b). Such freedom is in fact related to the gauge freedom in the choice of the initial value for the scale factor and time variable. W ithout loss of generality, and to simplify the second order calculations, we then set $c_{a,i}^{(1)} = 0$. The values of the $c_{n,i}^{(1)}$'s are related to the choice of the time variable. Since one usually considers the negative tension brane in RS I as the four-dimensional \visible Universe", it is natural to use the propertime on this brane and choose the $c_{n,i}^{(1)}$'s so as to have n (y_n ;) = 1. This can be achieved by setting

$$c_{n,p}^{(1)} = \frac{e^{2m \ b_0} \ 3m^2 \ (3w_p \ 1) + 2t_0^2 \ k^2 \ !^2 \ (3w_p + 2)}{48m \ b_0^2 \ !^2 \ (e^{m \ b_0} \ 1)}$$

$$c_{n,n}^{(1)} = \frac{e^{m \ b_0} \ 3m^2 \ (3w_n \ 1) + 2t_0^2 \ k^2 \ !^2 \ 1 \ 2e^{m \ b_0} \ (3w_n + 2)}{48m \ b_0^2 \ !^2 \ (e^{m \ b_0} \ 1)}$$
(37)

and de ning the time coordinate as

$$d = \exp - \frac{m b_0}{2} dt:$$
 (38)

W ith this choice, the cosm ological Friedmann equations can be easily compared to standard ones.

Let us now comment on the rst order results. As far as the radion perturbation is concerned, we found

$$f_{b} = \frac{m e^{4 m b_{0} y}}{4 b_{0} !^{2}} (1 \quad 3w_{p})_{p} + e^{m b_{0}} (1 \quad 3w_{n})_{n} + O(2);$$
(39)

which was expected, as it is due to the known coupling of the radion with the trace of the energy-momentum tensor of brane matter. Traceless uids, such as radiation, have no rst order e ect on the excitation of the radion. If one

lls the branes with som e pressureless uid, the distance between the two branes grows. Thise ect, being counterintuitive for the attractive nature of New tonian gravity, is in fact a consequence of the form of the stabilizing potential. Its non trivial contribution to the bulk energymom entum tensor at order is

$$T_{44} = 2\dot{\beta}!^2 f_b$$
: (40)

The rst order expressions are identical to those for a static solution, as they can be obtained by neglecting $_i$. Every kind of matter on the branes thus acts so as to detune the brane tensions from the bulk cosm ological constant and can be balanced by som e constant pressure along the flh dimension. Such pressure is given by the rst order contribution of (40) which increases when f_b decreases.

Note that (39) is proportional to the inverse of $!^2$, which represents the elective spring constant coming from some stabilization mechanism. When such a constant diverges, the correction f_b vanishes and the length of the flh dimension is xed as expected, even if there is a nite, !-independent T_{44} pressure term. On the other hand, the correction to the scale factor,

$$f_{a} = \frac{e^{2 m b_{0} y}}{48 m b_{0}^{2} !^{2} (e^{m b_{0}} 1)} 2b_{0}^{2} k^{2} !^{2} e^{2 m b_{0} y} 2e^{m b_{0}} + 3m^{2} (3w_{p} 1)e^{2 m b_{0} y} p + e^{m b_{0}} 2b_{0}^{2} k^{2} !^{2} e^{2 m b_{0} y} 2 + 3m^{2} (3w_{n} 1)e^{2 m b_{0} y} n + O(2);$$
(41)

never vanishes when matter is present on the branes, even if that is trace-less. Furthermore a nite, non vanishing f_a can be obtained in the limit of in nite spring constant, regardless of the matter equation of state.

$$f_{n} = \frac{e^{2 m b_{0} y}}{48 m b_{0}^{2} ! ^{2} (e^{m b_{0}} 1)} 2b_{0}^{2} k^{2} ! ^{2} (3w_{p} + 2) 2e^{m b_{0}} e^{2 m b_{0} y} + 3m^{2} (3w_{p} 1)e^{2 m b_{0} y} p$$

$$+ e^{m b_{0}} 2b_{0}^{2} k^{2} ! ^{2} (3w_{n} + 2) 2 e^{2 m b_{0} y} + 3m^{2} (3w_{n} 1)e^{2 m b_{0} y} n + c_{hp}^{(1)} p + c_{hp}^{(1)} n + O(2);$$

$$(42)$$

does not vanish when the branes are led with trace-less matter. Note, however, that a vanishing correction can be obtained for some exotic uid with $w_i = 2=3$ and negative pressure in the limit !! 1 . Apart from these exceptions, one has non negligible corrections everywhere in the bulk.

In order to compare the rst order results with the RS II case of a single brane, we must instead use the proper time on the positive tension brane. This is achieved by setting

$$c_{h,p}^{(1)} = \frac{3 \text{ m}^{2} (3 \text{w}_{p} \quad 1) + 2 \text{ l}_{0}^{2} \text{ k}^{2} !^{2} 1 \quad 2 \text{ e}^{\text{m} b_{0}} \quad (3 \text{w}_{p} + 2)}{48 \text{ m} b_{0}^{2} !^{2} (\text{e}^{\text{m} b_{0}} \quad 1)}$$
(43)

$$c_{n,n}^{(1)} = \frac{e^{m b_0} 3m^2 (3w_n 1) + 2b_0^2 k^2 !^2 (3w_n + 2)}{48m b_0^2 !^2 (1 e^{b_0})};$$

and letting $b_0 \ ! \ 1$. The Friedmann equation is simply obtained by keeping O () terms in H 2 (y_i;t). Since the rst order four-dimensional H ubble parameter is hom ogeneous, it reacts to all the sources along the y direction. On the positive tension brane one has, to rst order in ,

$$H_{p}^{2} = \frac{m k^{2} e^{m b_{0}} p + e^{m b_{0}} n}{3 (e^{m b_{0}} 1)} e^{(b_{0}!)!} \frac{m k^{2}}{3} p ; (44)$$

regardless of the value of !. W hen the negative tension brane is moved to in nity, its contribution goes to zero and one recovers the usual rst order e ect in brane cosm ology. On the visible brane, the Friedmann equation is slightly modi ed by the rescaled time parameter ,

$$H_{n}^{2} = \frac{m k^{2} e^{2m b_{0}} p + n}{3 (e^{m b_{0}} 1)} e^{p!}!^{0} \frac{m k^{2} n}{3 (e^{m b_{0}} 1)} : (45)$$

The acceleration equation is hom ogeneous as the Friedm ann equation and becomes

$$\frac{a(y_{i};t)}{a(y_{i};t)} = \frac{m \ k^{2} \ e^{m \ b_{0}} \ (1 + w_{p}) \ p + e^{m \ b_{0}} \ (1 + w_{n}) \ n}{6 \ (1 \ e^{n \ b_{0}})} ;$$
(46)

which has the weighted brane uid energy densities as sources.

B. Second order results

W e are now ready to evaluate O (2) corrections to the vacuum solution RS I.The procedure will be analogous to the one used for rst order results in the previous section.

As a rst step, one can impose the constraint (13) in order to nd the dependence on y of the second order coe cients in f_b . We are then left with three inhom ogeneous equations obtained by setting to zero the coe cients of the independent m atter densities in

$$\begin{aligned} & f_{bp}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpn}^{(2)} \stackrel{0}{_{n}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpn}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpn}^{(2)} \stackrel{0}{_{n}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{n}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{p}}{}^{2} + f_{bpm}^{(2)} \stackrel{0}{_{p}}{}^{2} + \frac{m(1 - 3w_{n})^{2}e^{2mb_{0}}(3y-1)}{24b_{0}^{2}!^{4}(e^{mb_{0}}-1)} e^{2mb_{0}} e^{2mb_{0}}$$

which contains the storder parameters previously determ ined. The solutions are

$$f_{b,p}^{(2)} = e^{4m \ b_0 \ y} \left(b_p^{(2)} - \frac{(1 \ 3w_p)^2 \ e^{4m \ b_0 \ y}}{96b_0^3 \ ! \ ^4 \ (e^{m \ b_0} \ 1)^2} \right)^6 6m^2 + b_0^2 k^2 \ ! \ ^2 \ 2b_0^2 k^2 \ ! \ ^2 e^{m \ b_0 \ (2y \ 3)} \left(48a \right)$$
(48a)

$$\mathbf{f}_{b;n}^{(2)} = e^{4 \, \mathrm{m} \, b_0 \, \mathrm{y}} \, b_n^{(2)} - \frac{(1 - 3w_n)^2 \, e^{6 \, \mathrm{m} \, b_0 \, (\mathrm{y} - 1)}}{96 \, b_0^3 \, ! \, ^4 \, (e^{\mathrm{m} \, b_0} - 1)^2} \, e^{2 \, \mathrm{m} \, b_0 \, \mathrm{y}} \, 6 \, \mathrm{m}^2 + b_0^2 \, \mathrm{k}^2 \, ! \, ^2 - 2 \, b_0^2 \, \mathrm{k}^2 \, ! \, ^2 \tag{48b}$$

$$f_{bm}^{(2)} = e^{4m \ b_0 \ y} \ b_m^{(2)} - \frac{(1 \ 3w_p) \ (1 \ 3w_n) \ e^{m \ b_0 \ (6y \ 5)}}{48 \ b_0^3 \ ! \ ^4 \ (e^{m \ b_0} \ 1)^2} \ e^{2m \ b_0 \ y} \ 6m^2 + \ b_0^2 \ k^2 \ ! \ ^2 \ b_0^2 \ k^2 \ ! \ ^2 \ 1 + \ e^{m \ b_0} \ ; (48c)$$

where

$$b_p^{(2)}$$
; $b_n^{(2)}$; $b_m^{(2)}$; (49)

are integration constants to be determ ined from the junction conditions. Once we plug the $f_{b;i}^{(2)}$'s into the second order terms in Eq. (12), we get the equations for the $f_{a;i}^{(2)}$'s, which are not displayed for the sake of brevity. Finally, by solving $G_{44} = k^2 T_{44}$ one obtains the corrections $f_{n;i}^{(2)}$'s. The results will contain six integration constants from the solutions of the rst order di erential equations for $f_{a;i}^{(2)}$ and $f_{n;i}^{(2)}$,

$$c_{a;i}^{(2)}$$
; $c_{n;i}^{(2)}$; (50)

and nine parameters related to the radion, the fourdimensional Hubble parameter and c(t) respectively,

$$h_{h;i}^{(2)}$$
; $b_{i}^{(2)}$; $c_{i}^{(2)}$; (51)

where i runs overp, n and m for second order quantities. A nalogously to the rst order case, one can x the coefcients (51) by imposing the junction conditions, which form a system of nine independent equations. N onetheless, one is again not able to x the constants (50). Here are the solutions obtained at the end of the calculations described above:

$$f_{a,p}^{(2)} = \frac{1}{4608 \, \mathrm{m}^2 \, \mathrm{b}_0^4 \, \mathrm{!}^4 \, (\mathrm{e}^{\mathrm{m} \, \mathrm{b}_0} \, 1)^2} \, 2 \, \mathrm{e}^{4 \, \mathrm{m} \, \mathrm{b}_0 \, \mathrm{y}} \, 3 \, \mathrm{m}^2 \, (1 \quad 3 \, \mathrm{w}_p)^2 \, 8 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, 6 \, \mathrm{m}^2 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \\ + 8 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, \mathrm{e}^{\mathrm{m} \, \mathrm{b}_0 \, (6y+1)} \, 11 \, \mathrm{m}^2 \, (3 \, \mathrm{w}_p \, 1) + 2 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, \mathrm{!}^2 \\ + 4 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, \mathrm{e}^{\mathrm{m} \, \mathrm{b}_0 \, (2y+1)} \, 3 \, \mathrm{m}^2 \, (9 \, \mathrm{w}_p^2 + 18 \, \mathrm{w}_p \, 7) + 8 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, \mathrm{!}^2 \\ + 4 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, \mathrm{e}^{2 \, \mathrm{m} \, \mathrm{b}_0 \, (y+1)} \, 27 \, \mathrm{m}^2 \, (3 \, \mathrm{w}_p^2 + 2 \, \mathrm{w}_p \, 1) + 16 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, \mathrm{!}^2 + \\ + \, \mathrm{e}^{8 \, \mathrm{m} \, \mathrm{b}_0 \, \mathrm{y}} \, 45 \, \mathrm{m}^4 \, (1 \, 3 \, \mathrm{w}_p)^2 + 3 \, \mathrm{m}^2 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, (9 \, \mathrm{w}_p^2 \, 42 \, \mathrm{w}_p + 13) \, 4 \, \mathrm{d}_0^2 \, \mathrm{k}^4 \, \mathrm{!}^4 \\ 2 \, \mathrm{e}^{\mathrm{m} \, \mathrm{b}_0 \, (2y+1)} \, 81 \, \mathrm{m}^4 \, (9 \, \mathrm{w}_p^3 \, + 9 \, \mathrm{w}_p^2 \, \mathrm{w}_p \, 1) + 3 \, \mathrm{m}^2 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, (27 \, \mathrm{w}_p^2 \, + 42 \, \mathrm{w}_p \, 1) + 8 \, \mathrm{b}_0^4 \, \mathrm{k}^4 \, \mathrm{!}^4 \\ 2 \, \mathrm{e}^{\mathrm{m} \, \mathrm{b}_0 \, (4y+1)} \, 81 \, \mathrm{m}^4 \, (9 \, \mathrm{w}_p^3 \, + 9 \, \mathrm{w}_p^2 \, \mathrm{w}_p \, 1) \, 3 \, \mathrm{m}^2 \, \mathrm{b}_0^2 \, \mathrm{k}^2 \, \mathrm{!}^2 \, (9 \, \mathrm{w}_p^2 \, + 6 \, \mathrm{w}_p \, + 29) + 16 \, \mathrm{b}_0^4 \, \mathrm{k}^4 \, \mathrm{!}^4 \, \mathrm{e}^{\mathrm{c}_{2,p}} \, (52a)$$

$$\mathbf{f}_{a,m}^{(2)} = \frac{e^{2m b_0 (y - 1)}}{4608m ^2 b_0^4 ! ^4 (e^{m b_0} - 1)^2} n^2 2e^{2m b_0 (y + 1)} 3m ^2 (1 - 3w_n)^2 - 8l_0^2 k^2 ! ^2 - 6m ^2 - l_0^2 k^2 ! ^2 + 8b_0^2 k^2 ! ^2 e^{4m b_0 y} 11m ^2 (3w_n - 1) + 2l_0^2 k^2 ! ^2 - 4l_0^2 k^2 ! ^2 e^{2m b_0} - 3m ^2 (9w_n^2 + 18w_n - 7) + 8l_0^2 k^2 ! ^2 + 4b_0^2 k^2 ! ^2 e^{m b_0} - 27m ^2 (3w_n^2 + 2w_n - 1) + 16l_0^2 k^2 ! ^2 + e^{6m b_0 y} - 45m ^4 (1 - 3w_n)^2 + 3m ^2 b_0^2 k^2 ! ^2 (9w_n^2 - 42w_n + 13) - 4l_0^2 k^4 ! ^4 - 2e^{m b_0 y} - 81m ^4 (9w_n^3 + 9w_n^2 - w_n - 1) + 3m^2 b_0^2 k^2 ! ^2 (27w_n^2 + 42w_n - 1) + 8l_0^4 k^4 ! ^4 - 2e^{m b_0 (2y + 1)} - 81m ^4 (9w_n^3 + 9w_n^2 - w_n - 1) - 3m^2 b_0^2 k^2 ! ^2 (9w_n^2 + 6w_n + 29) + 16b_0^4 k^4 ! ^4 - (52b)$$

$$f_{a,m}^{(2)} = \frac{e^{m b_0 (2y - 1)}}{2304m 2b_0^4 ! ^4 (e^{m b_0} - 1)^2} 4b_0^2 k^2 ! ^2 e^{4m b_0 y} m^2 (3w_p - 1) (6w_p - 6w_n + 11) + 2b_0^2 k^2 ! ^2 + 4b_0^2 k^2 ! ^2 e^{m b_0 (4y + 1)} m^2 (3w_n - 1) (6w_n - 6w_p + 11) + 2b_0^2 k^2 ! ^2 + 4b_0^2 k^2 ! ^2 e^{m b_0} 3m^2 (18w_p^2 - 9w_p w_n + 9w_p - 9w_n - 1) + 4b_0^2 k^2 ! ^2 + 4b_0^2 k^2 ! ^2 e^{2m b_0} 3m^2 (18w_n^2 - 9w_p w_n + 9w_n - 9w_p - 1) + 4b_0^2 k^2 ! ^2 + e^{6m b_0 y} 45m^4 (3w_p - 1) (3w_n - 1) + 3m^2 b_0^2 k^2 ! ^2 (9w_p w_n - 21w_p - 21w_n + 13) - 4b_0^4 k^4 ! ^4 - e^{2m b_0 y} 9m^4 (3w_p - 1) (18w_p^2 + 9w_p w_n + 27w_p + 3w_n + 11) + 36m^2 b_0^2 k^2 ! ^2 (3w_p^2 - 3w_p w_n + w_p - 2w_n - 1) + 8b_0^4 k^4 ! ^4 - e^{2m b_0 (y+1)} 9m^4 (3w_n - 1) (18w_n^2 + 9w_p w_n + 27w_n + 3w_n + 11) + 36m^2 b_0^2 k^2 ! ^2 (3w_n^2 - 3w_p w_n + w_n - 2w_p - 1) + 8b_0^4 k^4 ! ^4 - e^{2m b_0 (2y+1)} 81m^4 (6w_p^3 + 6w_n^3 + 3w_p^2 w_n + 3w_p w_n^2 + 7w_p^2 + 7w_n^2 + 4w_p w_n - w_p - w_n - 2) + 6m^2 b_0^2 k^2 ! ^2 (18w_p^2 + 18w_n^2 - 9w_p w_n + 21w_p + 21w_n - 1) + 16b_0^4 k^4 ! ^4 + c_{a,m}^{(2)}$$

$$(52c)$$

$$f_{n,p}^{(2)} = \frac{e^{2^{m b_0 y}}}{4608m^2 b_0^4 !^4 (e^{n b_0} 1)^2} \qquad 8l_0^2 k^2 !^2 (3w_p + 2)e^{3m b_0} 3m^2 (3w_p 1) + 2l_0^2 k^2 !^2 (3w_p + 2) \\ 8l_0^2 k^2 !^2 e^{n b_0 (4y+1)} m^2 (81w_p^3 + 180w_p^2 + 24w_p 31) + 2l_0^2 k^2 !^2 (6w_p + 5) \\ + 2e^{2^{m b_0 y}} 6m^2 l_0^2 k^2 !^2 3m^2 (3w_p 1)^2 4l_0^2 k^2 !^2 (9w_p^2 + 6w_p 1) \\ 4l_0^2 k^2 !^2 e^{m b_0} 3m^2 (63w_p^2 + 30w_p 17) + 4l_0^2 k^2 !^2 (9w_p^2 + 6w_p 1) \\ + 4b_0^2 k^2 !^2 27m^2 (18w_p^3 + 27w_p^2 + 4w_p 5) + 8l_0^2 k^2 !^2 (9w_p^2 + 6w_p 1) \\ + e^{6^{m b_0 y}} 45m^4 (3w_p 1)^2 + 3m^2 b_0^2 k^2 !^2 (117w_p^2 + 30w_p 23) + 4l_0^2 k^4 !^4 (6w_p + 5) \\ 2e^{m b_0 (2y+1)} 81m^4 (9w_p^3 + 9w_p^2 w_p 1) 3m^2 b_0^2 k^2 !^2 (162w_p^3 + 279w_p^2 + 60w_p 25) \\ + 8b_0^4 k^4 !^4 (9w_p^2 + 6w_p 1) + 2e^{2^{m b_0} (y+1)} 9m^4 (81w_p^3 + 90w_p^2 15w_p 8) \\ + 3m^2 b_0^2 k^2 !^2 (162w_p^3 + 243w_p^2 + 12w_p 53) + 4l_0^4 k^4 !^4 (9w_p^2 + 24w_p + 14) + c_{n,p}^{(2)}$$
(52d)

$$f_{n,n}^{(2)} = \frac{e^{2m b_0 (y-1)}}{4608m 2b_0^4 ! ^4 (e^{m b_0} 1)^2} \qquad 4l_0^2 k^2 ! ^2 (6w_n + 5)e^{m b_0} 27m ^2 (3w_n^2 + 2w_n 1) + 8l_0^2 k^2 ! ^2 8l_0^2 k^2 ! ^2 e^{4m b_0 y} m ^2 (8lw_n^3 + 180w_n^2 + 24w_n 31) + 2l_0^2 k^2 ! ^2 (6w_n + 5) + 4b_0^2 k^2 ! ^2 e^{2m b_0} 3m ^2 (45w_n^2 + 24w_n 13) + 4l_0^2 k^2 ! ^2 (6w_n + 5) 2e^{m b_0 (2y+1)} 8lm ^4 (9w_n^3 + 9w_n^2 w_n 1) 3m^2 b_0^2 k^2 ! ^2 (162w_n^3 + 243w_n^2 + 48w_n 17) 8l_0^4 k^4 ! ^4 (6w_n + 5) + 2e^{2m b_0 (y+1)} 9m ^4 (3w_n 1)^2 3m^2 b_0^2 k^2 ! ^2 (45w_n^2 + 30w_n + 1) 4l_0^4 k^4 ! ^4 (6w_n + 5) + e^{6m b_0 y} 45m ^4 (3w_n 1)^2 + 3m ^2 b_0^2 k^2 ! ^2 (117w_n^2 + 30w_n 23) + 4l_0^4 k^4 ! ^4 (6w_n + 5) + 2e^{2m b_0 y} 81m^4 (9w_n^3 + 9w_n^2 w_n 1) + 3m ^2 b_0^2 k^2 ! ^2 (162w_n^3 + 243w_n^2 + 12w_n 53) + 8l_0^4 k^4 ! ^4 (6w_n + 5) + c_{n,n}^{(2)}$$
(52e)

IV. SECOND ORDER COSMOLOGY

Let us now look at the physical consequences of the above results as applied to two di erent cosm ological scenarios. The Friedm ann and acceleration equations will be showed and their phenom enology investigated in different regimes of ! and for di erent m atter equations of state on the branes.

W e begin from the case with two branes [12] for which we shall study how cosmology would be described by observers on the negative brane. In order to achieve that, one can use the corresponding proper time by xing the $c_{n,i}^{(1)}$'s as in Eq. (37), choosing

$$c_{n,p}^{(2)} = \frac{e^{2m b_0}}{4608m^2 b_0^4 ! ^4 (e^{m b_0} - 1)^2} 9m^4 (3w_p - 1) 3e^{2m b_0} (3w_p^2 + 4w_p + 1) + 18e^{m b_0} (3w_p^2 + 4w_p + 1) - 12w_p + 4$$

$$m^2 b_0^2 k^2 ! ^2 e^{2m b_0} (324w_p^3 + 153w_p^2 - 102w_p - 91) - 6E^{b_0} (162w_p^3 + 207w_p^2 + 12w_p - 65)$$

$$+ 54 (5w_p^2 + 2w_p - 3) + 4b_0^4 k^4 ! ^4 - 3e^{2m b_0} (6w_p^2 + 6w_p + 1) + 2 - 1 - 2E^{m b_0} (9w_p^2 + 6w_p - 1)$$

$$c_{n,m}^{(2)} = \frac{1}{4608m^2 b_0^4 ! ^4 (e^{m b_0} - 1)^2} 9m^4 (3w_n - 1) - 7e^{2m b_0} (3w_n - 1) + 18e^{m b_0} + 1 - (3w_n^2 + 4w_n + 1)$$

$$m^2 b_0^2 k^2 ! ^2 e^{2m b_0} (81w_p^2 - 90w_p - 75) + 2e^{m b_0} (162w_p^3 + 279w_p^2 + 192w_p - 5)$$

$$m \ b_0 k \ ! \ e^{-3} (81W_n - 90W_n - 75) + 2e^{-3} (162W_n + 279W_n + 192W_n - 5)$$

$$6 (162W_n^3 + 243W_n^2 + 60W_n - 37) + 4b_0^4 k^4 \ ! \ ^4 e^{m \ b_0} - 2^2 (6W_n + 5)$$
(53b)

$$\begin{aligned} c_{n,m}^{(2)} &= \frac{e^{m \ b_0}}{2304m \ ^2b_0^4 ! \ ^4 \ (e^{m \ b_0} \ 1)^2} \ 9m^4 \ e^{2m \ b_0} \ (3w_n \ 1) \ (18w_n^2 + 9w_pw_n \ 6w_p + 27w_n + 14) \\ &9e^{m \ b_0} \ (6w_p^3 + 6w_n^3 + 3w_p^2w_n + 3w_pw_n^2 + 7w_p^2 + 7w_n^2 + 4w_pw_n \ w_p \ w_n \ 2) + \ (3w_p \ 1) \ (18w_p^2 + 9w_pw_n + 27w_p + 3w_n + 11)] \ m^2b_0^2k^2 ! \ e^{2m \ b_0} \ (108w_n^3 + 54w_pw_n^2 + 234w_n^2 \ 9w_pw_n + 87w_p + 39w_n + 1) + 2e^{m \ b_0} \ (54w_p^3 \ 162w_n^3 + 27w_p^2w_n \ 81w_pw_n^2 + 117w_p^2 \ 351w_n^2 \ 45w_pw_n + 30w_p + 96w_n + 11) \ 6 \ (54w_p^3 + 27w_p^2w_n + 117w_p^2 \ 36w_pw_n + 30w_p \ 33w_n \ 19) \\ &4w_0^4k^4 ! \ e^{2m \ b_0} \ 2e^{m \ b_0} \ (3w_p + 2) \ (3w_n + 2) + 9w_pw_n + 3w_p + 3w_n \ 1 \end{aligned}$$

and nally rescaling n (y;t) to satisfy the condition n (1=2;) = 1. The Friedmann equation we are interested in is given by the second order expression of the Hubble parameter at y = 1=2 as a function of ,

$$H^{2}(1=2;) = \frac{k^{2}m}{3(e^{m b_{0}} 1)} + e^{2m b_{0}} p \quad (w_{n} + 1) (3w_{n} 1) \frac{3m^{2}k^{2} e^{m b_{0}} + 1}{32b_{0}^{2}!^{2} (e^{m b_{0}} 1)^{2}}$$

$$(\frac{k^{4}}{36} + \frac{m^{2}k^{2} e^{m b_{0}} + 1}{48b_{0}^{2}!^{2} (e^{m b_{0}} 1)^{2}} 18w_{p}^{2} 9w_{p}(w_{n} 1) 9w_{n} 1 + 2e^{m b_{0}} (9w_{n}^{2} + 9w_{n} 4) e^{m b_{0}} p^{n}$$

$$(\frac{k^{4}}{36} + \frac{m^{2}k^{2} (3w_{p} 1) e^{m b_{0}} + 1}{96b_{0}^{2}!^{2} (e^{m b_{0}} 1)^{2}} 3w_{p} + 7 4e^{m b_{0}} (3w_{p} + 4) e^{2m b_{0}} \frac{2}{p};$$

$$(54)$$

Note that the results (53a)-(54) do not depend on the integration constants $c_{a,i}^{(1)}$'s and $c_{a,i}^{(2)}$'s, which rejects the fact that the three-dimensional spatial curvature has been set to zero ab initio.

The Friedmann equation contains coe cients up to second order in the vacuum perturbations of both branes. First and second order contributions in Eq. (54) are consequences of the adiabatic regime of the verdimensional dynamics which determines the value of the integration constant c(t) in Eq. (12). The value of the latter is a ected by the presence of matter on both branes through the junction conditions and, for instance, up to 0 (²) is given by

$$\lim_{\substack{l \leq 1 \\ l \leq 1}} c^{(1)} = \frac{m k^2 p + e^{m b_0} n}{3 (e^{m b_0} 1)}$$
(55a)
$$\binom{m}{2} k^4 e^{2m b_0} + 2e^{m b_0} 1 q e^{m b_0} + 1^2 q e^{2m b_0} + 2e^{m b_0} 1 q e^{2m b_0} + 2e^{m b_0} 1 q$$

$$\lim_{1 \le 1} c^{(2)} = \frac{k^4}{36} \frac{e^{2m \ b_0} + 2e^{m \ b_0}}{(e^{m \ b_0} \ 1)^2} \frac{1}{p} + \frac{e^{m \ b_0} + 1}{e^{m \ b_0} \ 1} e^{m \ b_0} \frac{1}{p \ n} + \frac{e^{2m \ b_0} + 2e^{m \ b_0}}{(e^{m \ b_0} \ 1)^2} \frac{1}{p} \frac{2}{n} ;$$
(55b)

in the lim it of in nite spring constant. Furtherm ore, as previously noted, this e ect is also a consequence of the radion eld potential acting as a source in Eq. (12). One can expect to cancel some term s in Eq. (54) by arbitrarily increasing the spring constant of the e ective radion potential in order to decrease the radion shift from equilibrium. This mechanism partially works as the rst order contributions to the bulk potential vanish along the tim e-tim e direction, whereas only second order term s, which depend on the matter equation of state, cancel when !! 1 . One is then left with

$$\lim_{\substack{1 \leq 1 \\ 1 \leq 1}} H^{2} (1=2;) = \frac{m k^{2} n + e^{2m b_{0}} p}{3 (e^{m b_{0}} 1)}$$
$$\frac{k^{4}}{36} e^{m b_{0}} p n + \frac{k^{4}}{36} e^{2m b_{0}} \frac{2}{p} :(56)$$

The matter on the positive tension brane appears at second order with the role of som e \dark" uid and acts as a sort of B rans-D icke eld which adiabatically modi es the New ton constant perceived in the visible U niverse. This behavior is som ehow inherited by the dynam ics of the radion which is known to modulate the strength of gravity on the visible brane.

N ote that while rst order ∞e cients are positive definite, irrespective of the brane tension, the sign of second order ones depends on the matter equation of state. D i erently enough from unstabilized brane ∞m ology, this fact im plies that leading order ∞m ological equations have the correct behavior on both branes. Letting the energy density $_{\rm p}$! 0 in Eq. (54), one obtains

$$H^{2} (1=2;) = (w_{n} + 1) (1 \quad 3w_{n}) \frac{3m^{2}k^{2} e^{m b_{0}} + 1}{32b_{0}^{2}!^{2} (e^{m b_{0}} \quad 1)^{2}} \int_{n}^{2} + \frac{m k^{2} n}{3 (e^{m b_{0}} \quad 1)};$$
(57)

which has the usual rst order solution for both radiation and a cosm ological constant on the visible brane. A m atter dom inated Universe would otherw ise generate second order corrections.

We now come to the equation for the acceleration, which has the general form

$$\frac{a(1=2;)}{a(1=2;)} = \frac{m k^2}{6 (1 e^{0 b_0})} (3w_n + 1)_n + e^{2m b_0} (3w_p + 1)_p + (w_n + 1) (3w_n - 1) (3w_n + 2) \frac{3m^2 k^2 e^{m b_0} + 1}{32b_0^2 ! 2 (e^{m b_0} - 1)^2}_n^2 \frac{k^4}{144} (3w_n + 1) (3w_p + 1) + \frac{m^2 k^2}{96b_0^2 ! 2 (e^{m b_0} - 1)^2} e^{2m b_0} (3w_n - 1) (18w_n^2 + 9w_p w_n + 15w_p + 39w_n + 23) + 9e^{m b_0} (6w_p^3 + w_n^3 + 3w_p^2 w_n + 3w_p w_n^2 + 11w_p^2 + 11w_n^2 + 2w_p w_n - 4) + 54w_p^3 - 27w_p^2 w_n + 99w_p^2 - 18w_p w_n + 24w_p - 13 e^{2m b_0} p - n + (\frac{k^4}{144} (3w_p + 1)^2 + \frac{m^2 k^2 (3w_p - 1) e^{m b_0} + 1}{96b_0^2 ! ^2 (e^{m b_0} - 1)^2} e^{m b_0} (27w_p^2 + 54w_p + 23) - 9w_p - 5 e^{3m b_0} \frac{2}{p};$$
(58)

and, for ! ! 1, reduces to

$$\frac{a(1=2;)}{a(1=2;)} = \frac{m k^2}{6(1 e^{n b_0})} (3w_n + 1)_n + e^{2m b_0} (3w_p + 1)_p$$

$$\frac{k^4}{144} (3w_n + 1) (3w_p + 1)e^{2m b_0}_{p n} + \frac{k^4}{144} (3w_p + 1)^2 e^{3m b_0}_p^2 ; \qquad (59)$$

The coe cient of p^2 in Eq. (59) is positive or zero in this limit and provides an accelerating contribution to the equation. The coe cient of the mixed term has a positive value when just one uid has w $p_i < 1=3$. For p! 0, one is further left with

$$\frac{a(1=2;)}{a(1=2;)} = \frac{m \ k^2 \ (3w_n + 1)}{6 \ (1 \ e^{m \ b_0})} \ _n + \ (w_n + 1) \ (3w_n - 1) \ (3w_n + 2) \frac{3m^2 k^2 \ e^{m \ b_0} + 1}{32b_0^2 !^2 \ (e^{m \ b_0} - 1)^2} \ _n^2; \tag{60}$$

where the second order contribution is inversely proportional to ! and vanishes for radiation and a cosm obgical constant. This peculiarity leads to the standard cosm obgical evolution up to 0 ($\frac{2}{n}$) until the matter dom inated era.

A singularity in the lapse function n(y;t) for $b_0 ! 1$ prevents us from analyzing the correct limit when the distance between the branes becomes in nite, we shall thus comment on this problem in the next subsection. On setting $_p ! 0, Eq. (56)$ admits the nite but trivial limit

$$\lim_{b_0 \mid 1} H^2 (1=2;) = 0;$$
 (61a)

in which one also has

$$\lim_{b_0 \mid 1} \frac{a(1=2;)}{a(1=2;)} = 0 :$$
 (61b)

This result is due to the reduced strength of the gravitational interaction at in nity.

B. RS II

On can think of the RS II model as the limit of RS I in which the distance between the two branes becomes in nite, thus one expects that only $_{\rm p}$ contributes in this limit. The cosm ological proper time is now the one on the P lanck brane and is recovered upon choosing

$$c_{n,p}^{(2)} = \frac{1}{4608m^{2}b_{0}^{4}!^{4} (e^{m b_{0}} 1)^{2}} 9m^{4} (3w_{p} 1) 18e^{2m b_{0}} (3w_{p}^{2} + 4w_{p} + 1) + 18e^{m b_{0}} (3w_{p}^{2} + 4w_{p} + 1)$$

$$21w_{p} + 7] + m^{2}b_{0}^{2}k^{2}!^{2} 6e^{2m b_{0}} (162w_{p}^{3} + 243w_{p}^{2} + 60w_{p} 37)$$

$$2e^{m b_{0}} (162w_{p}^{3} + 279w_{p}^{2} + 192w_{p} 5) 81w_{p}^{2} + 90w_{p} + 75$$

$$+ 4b_{0}^{4}k^{4}!^{4} 1 2e^{m b_{0}}{}^{2} (6w_{p} + 5)$$
(62a)

$$c_{n,n}^{(2)} = \frac{e^{2m b_0}}{4608m^2 b_0^4!^4 (e^{m b_0} 1)^2} 9m^4 (3w_n 1) 4e^{2m b_0} (1 3w_n) + 18e^{m b_0} (3w_n^2 + 4w_n + 1) + 54w_n^2 + 63w_n + 21 m^2 b_0^2 k^2!^2 54e^{2m b_0} (5w_n^2 + 2w_n 3) 6e^{m b_0} (162w_n^3 + 207w_n^2 + 12w_n 65) + 324w_n^3 + 153w_n^2 102w_n 91 + 4b_0^4 k^4!^4 2 e^{2m b_0} 2e^{m b_0} (9w_n^2 + 6w_n 1) + 3(6w_n^2 + 6w_n + 1)$$
(62b)

$$c_{n,m}^{(2)} = \frac{e^{m b_0}}{2304m \ ^2b_0^4 ! \ ^4 \ (e^{m b_0} \ 1)^2} \ 9m^4 \ e^{2m b_0} \ (3w_n \ 1) \ (18w_n^2 + 9w_pw_n + 3w_p + 27w_n + 11) \\ + 9e^{m b_0} \ (6w_p^3 + 6w_n^3 + 3w_p^2w_n + 3w_pw_n^2 + 7w_p^2 + 7w_n^2 + 4w_pw_n \ w_p \ w_n \ 2) \\ (3w_p \ 1) \ (18w_p^2 + 9w_pw_n + 27w_p \ 6w_n + 14) \ + m^2b_0^2k^2 !^2 \ 6e^{2m b_0} \ (54w_n^2 + + 27w_pw_n^2 \\ + 117w_n^2 \ 36w_pw_n \ 33w_p + 30w_n \ 19) \ + 2e^{m b_0} \ (162w_p^3 \ 54w_n^3 + 81w_p^2w_n \ 27w_pw_n^2 \\ + 351w_p^2 \ 117w_n^2 \ 96w_p \ 30w_n \ 11) \ 108w_p^3 \ 54w_p^2w_n \ 234w_p^2 \ + 9w_pw_n \ + 39w_p \ 87w_n \ 1 \\ 4b_0^4k^4 ! \ 4 \ e^{2m b_0} \ (9w_pw_n \ + 3w_p \ 3w_n \ 1) \ + \ 1 \ 2e^{m b_0} \ (3w_p \ + 2) \ (3w_n \ + 2)$$

together with Eq. (43). The Friedmann equation, for a nite b_0 , reduces to

$$H^{2}(0;t) = \frac{m k^{2}}{3 (e^{m b_{0}} 1)} e^{m b_{0}} p + e^{m b_{0}} n + (w_{p} + 1) (1 3w_{p}) \frac{3m^{2}k^{2}e^{m b_{0}} e^{m b_{0}} + 1}{32b_{0}^{2}!^{2} (e^{m b_{0}} 1)^{2}} p^{2}$$

$$\left(\frac{k^{4}}{36} + \frac{m^{2}k^{2} e^{m b_{0}} + 1 e^{m b_{0}} (18w_{n}^{2} 9w_{p}w_{n} 9w_{p} + 9w_{n} 1) (9w_{p}^{2} + 9w_{p} 4)}{24b_{0}^{2}!^{2} (e^{m b_{0}} 1)^{2}} e^{m b_{0}} p^{n}$$

$$\left(\frac{k^{4}}{36} + \frac{m^{2}k^{2} e^{m b_{0}} + 1 (3w_{n} 1) e^{m b_{0}} (3w_{n} + 7) 4 (3w_{n} + 4)}{96b_{0}^{2}!^{2} (e^{m b_{0}} 1)^{2}} e^{2m b_{0}} p^{n} \right)$$

$$(63)$$

Г

which is similar to Eq. (54) but has the nite, non trivial limit

$$\lim_{b_0! \ 1} \ H^2(0;t) = \frac{m \ k^2}{3} \ p :$$
 (64)

This is precisely the standard Friedmann equation one has in four-dimensional cosmology. Furthermore, the equation for the acceleration to second order in terms of the time on the positive tension brane is given by

$$\frac{a\ (0;t)}{a\ (0;t)} = \frac{m\ k^{2}\ e^{m\ b_{0}}}{6\ (1\ e^{m\ b_{0}}\)} (3w_{p}+1)\ _{p} + (3w_{n}\ 1)e^{2m\ b_{0}}\ _{n} + (w_{p}+1)\ (3w_{p}\ 1)\ (3w_{p}+2)\frac{3m\ ^{2}k^{2}\ e^{m\ b_{0}}\ e^{m\ b_{0}}+1}{32b_{0}^{2}l^{2}\ (e^{m\ b_{0}}\ 1)^{2}}\ _{p}^{2}$$

$$= \frac{k^{4}}{144}(3w_{p}+1)\ (3w_{n}+1)\ \frac{m\ ^{2}k^{2}}{96b_{0}^{2}l^{2}\ (e^{m\ b_{0}}\ 1)^{2}}\ e^{2m\ b_{0}}\ (54w_{n}^{3}+27w_{p}w_{n}^{2}+99w_{n}^{2}\ 18w_{p}w_{n}$$

$$= 21w_{p}+24w_{n}\ 13)+9e^{m\ b_{0}}\ (6w_{p}^{3}+6w_{n}^{3}+3w_{p}^{2}w_{n}+3w_{p}w_{n}^{2}+11w_{p}^{2}+11w_{n}^{2}+2w_{p}w_{n}$$

$$+w_{p}+w_{n}\ 4)+\ (3w_{p}\ 1)\ (18w_{p}^{2}+9w_{p}w_{n}+39w_{p}+15w_{n}+23)\ e\ ^{m\ b_{0}}\ p\ n$$

$$+\frac{k^{4}}{144}(3w_{n}+1)^{2}\ \frac{m\ ^{2}k^{2}\ e^{m\ b_{0}}\ (3w_{n}\ 1)}{96b_{0}^{2}l^{2}(e^{m\ b_{0}}\ 1)^{2}}\ e^{m\ b_{0}}\ (9w_{n}+5)\ 27w_{n}^{2}+54w_{n}+23\ e\ ^{2m\ b_{0}\ 2}\ e\ ^{2m\ b_{0}\ 2}\ (65)$$

(66)

and, compatibly with Eq. (64), yields

 $\lim_{b_0! \ 1} \frac{a(0;t)}{a(0;t)} = -\frac{m \ k^2}{6} (3w_p + 1)_p;$

grows, which consequently leads to unobservable deviations from standard four-dimensionalGeneralRelativity.

V. APPROXIMATION ANALYSIS

The results obtained so far hold with the assumption that $_{i}$ M⁴ where M is, in general, the natural mass scale of the m odel. By taking all the m ass param eters to such a natural scale, one expects the solutions to second order provide a good approximation for Eq. (9). In this regime, however, second order e ects are certainly subleading and thus insu cient to signi cantly alter rst order behavior. W e shall hence present below a num erical analysis of the validity of our approximate solutions in the attempt to extend the range of the parameters in which ours results hold valid and widen the conclusions one can draw from second order expressions. In particular, we are interested in possible deviations from standard $\cos m$ obgical equations due to term s of 0 (²) in the rst. and third of Eqs. (8). Note that all num erical results will be obtained by setting the expansion parameter = 1 as previously prescribed.

In order to test our approximations, since Eqs. (9) are not analytically solvable, we substitute the second order solutions into the exact E instein equations to obtain an estimate of their non vanishing remainders which we then compare with the leading contributions (satisfying the corresponding approximate equations). Since the time dependence is contained in $_{i}$ in our expansions (20a)– (20c), it is also convenient to trade the time for . We thus divide Eq. (12) into the following veterms

$$\frac{1}{n^2}$$
 (y;) $\frac{1}{n^2}$ $\frac{a^2}{y;t}$ (67a)

$$_{2}(y;) \qquad \frac{1}{b^{2}(y;t)} = \frac{a^{0}(y;t)}{a(y;t)}^{2}$$
 (67b)

$$_{3}$$
 (y;) $\frac{k^{2}}{6}$ (+ U) (67c)

$$_{4}$$
 (y;) $\frac{k^{2}}{6a^{4}} (y;t) = a^{4} T_{0}^{0} dx$ (67d)

$$_{5}(y;) \frac{e(t)}{a^{4}(y;t)};$$
 (67e)

in which t = t() in the right hand sides is understood as the time when = n (p w ill be chosen either equal to n or zero). The sum,

evaluated on the second order solutions yields R (y;) = 0 (³) as a measure of the corresponding error. Sim ilarly, the third of Eqs. (9) may be written as the sum of the following six terms

$$\frac{1}{n^2(y;t)} = \frac{1}{n^2(y;t)} \frac{a(y;t)}{a(y;t)}$$
 (69a)

$$\frac{1}{b^2(y;t)} = \frac{1}{a(y;t)} \left(\frac{a^0(y;t)}{a(y;t)}\right)^2$$
 (69b)

$$_{3}(y;) \qquad \frac{1}{b^{2}(y;t)} \frac{a^{0}(y;t)}{a(y;t)} \frac{n^{0}(y;t)}{n(y;t)}$$
(69c)

$$_{4}$$
 (y;) $\frac{1}{n^{2}$ (y;t)} $\frac{\underline{a}^{2}$ (y;t)}{a^{2}} (y;t) \frac{\underline{a}(y;t) \underline{n}(y;t)}{a(y;t) n(y;t)} (69d)

$$_{5}(y;)$$
 $\frac{k^{2}}{3}(+U)$ (69e)

$$_{6}(y;)$$
 $\frac{2}{3}k^{2}!^{2}b(y;t)[b(y;t), b_{0}];$ (69f)

and

with R (y;) = O (3) for the same approximate solutions. One may now assume that approximate metric functions computed to O (2) are accurate approximations of exact solutions to Eqs. (9) if

 $j_1j_1j_1j_1j_2j_1$; (71)

for every term in Eqs. (67a)-(69f) evaluated to $0(^2)$. (It should actually be su cient to satisfy the above conditions for the leading terms of each equation.) Throughout this section, where unspecied, a natural choice [13] of dimensionful parameters is considered.

Fig.1 shows the functions 1 and 1 evaluated to second order in and the corresponding R and R for $_{i}$ = 10^1 M^4 and w_i = 0.5. The box in the rst row on 2 the left shows the absolute values of the leading term s $_{2}$ and 3, which are roughly one order of m agnitude larger than the remaining term s 1's displayed in the plot on the right along with R (the dotted line). In the second row of Fig.1, them odula of the leading term s $_{2, 3}$ and $_{5}$ are plotted on the left while the remaining coe cients 1 and R (y) are presented on the right. Thanks to the relatively large am plitudes of the leading term s, one can rely on the second order approximation even when the i's are not too small, implying that a great improvement over rst order results can be achieved in a regime where i 10 2 . In this case the errors produced by truncating the expansion to 0 $(^{2})$ are much smaller than one percent.

Let us further consider the particular case $_{\rm p}$ = 0. This choice is made in order to study the cosm obgical consequences experienced on the negative tension brane (at y = 1=2) generated by second order terms proportional to $_{\rm n}^2$. We therefore use the proper time on the negative tension brane [14]. In each graph in Fig. 2 we plot the squared Hubble parameter on the negative tension brane of Eq. (57) and the acceleration of Eq. (60) for a given energy density $_{\rm n}$ as a function of its equation of



FIG.1: The graphs on the left contain the plot of the absolute values of the leading term s am ong $_1$ (above) and $_1$ (below) to 0 (²). The graphs on the right show the subleading term s am ong $_1$ (above) and $_1$ (below) and the corresponding rem ainders R and R to 0 (²) (dotted lines). All plots are for $_i = 2 10^1$ M⁴ and $w_i = 0.5$ and cover all the bulk between the two branes.

state w_n . The plot on the left in the rst row is for the sm all density $_{n} = 10^{-2} M^{-4} M^{-4}$ and shows a behavior which is typical of standard four-dimensional cosm ology. In fact, only $w_n < 1=3$ leads to an accelerating phase. This trend is modi ed by higher densities, as it em erges in the remaining graphs of Fig. 2. In particular when 10^1 M⁴ (plot on the left in the second row) the n = 2 second order corrections seem to provide an accelerated regime for $0 < w_n < 1$. The acceleration appears am plied when $_{n} = 6 \quad 10^{1} \text{ M}^{4} < \text{M}^{4}$ (plot on the right in the second row) or higher. Note how ever that the Hubble param eter, a positive de nite quantity, constrains the region swept by w_n which is not allowed to reach unity. Finally the interm ediate case is showed in the plot on the right the rst row: due to the second order e ect, H 2 exhibits a dependence on w_n otherwise not present.

The four regimes described above have to be tested with particular accuracy because the acceleration and Hubble parameter plotted in Fig. 2 are not the leading terms in Eqs. (9) and could thus be comparable with the remainders. In this case, it is somehow possible that the remainders signi cantly modify the behavior. We rst note that H² 1 and a=a 1, as de ned above in Eqs. (67a) and (69a), hence we can use 1 and 1 in place of H² and a=a respectively. We then plot in Fig. 3

the ratio between the remainder R (evaluated to second order) and the squared Hubble parameter $_1$ evaluated to rst and second orders. The four plots show this ratio for dierent choices of n and w_n (the same as in Fig. 2 and in the same order) as a function of y. In particular, we choose $w_n = 0.95$ in order to explore regions where the acceleration has an unconventional behavior for $n = 10^{2} M^{4}$, $10^{1} M^{4}$, 2 $10^1 \,\mathrm{M}^4$ and $w_n = 0.65$ for n = 6 10¹ M⁴. A part from the last case, the corrections given by the neglected term s cannot signi cantly m odify H² on the negative tension brane. In the rst line ofFig.4, the ratios between R (to second order) and the acceleration 1 to 1st and second order are analogously plotted. For low enough energy densities, the second order expressions appear to be good approximations. On the other hand, in the second row it is shown that, in the unconventional regime of the acceleration (that is $_{n}$ = 2 10¹ M ⁴ and w_{n} = 0.95, and $_{n}$ = 6 10¹ M ⁴ and $w_n = 0.65$) the two terms are of the same order of m agnitude. This shows that one should go beyond $O(^2)$ in order to determ ine the true behavior of a=a for such equations of state.



FIG.2: Plots of H² (1=2;) (solid line) and a (1=2;)=a (1=2;) (dotted line) to 0 (2) at a given time, as functions of w_n, for $_n = 10^2$ M⁴, 10¹ M⁴, 2 10¹ M⁴ and 6 10¹ M⁴ (from top left to bottom right).

VI. CONCLUSIONS

We have computed approximate cosmological solutions of ve-dimensional Einstein equations for R and all-Sundrum models in the presence of a radion elective potential. The calculations were performed up to the second order in the energy densities of the matter on the branes and assuming an adiabatic evolution of the system. For the RS I model with matter localized only on the negative tension brane, we found negligible corrections for the Hubble parameter in the case of radiation or cosmological constant. For RS II, we found negligible corrections for the equations of state just described and in the limit when the distance between the branes is taken to in n-ity. There is some evidence of an accelerating phase for a wider range of values of w_n if the distance between the branes is nite, but one needs to carry the computation to higher orders, since such an e ect appears near the limit of validity of our perturbative expansion.

- [1] T. Kaluza, Sitzungsber. Preuss. Akad. W iss. Berlin (M ath.Phys.) 1921, 966 (1921).
- [2] O.Klein, Z.Phys. 37, 895 (1926).
- [3] L.Randall and R. Sundrum, Phys. Rev. Lett. 83, 3370 (1999).
- [4] L.Randall and R.Sundrum, Phys. Rev. Lett. 83, 4690 (1999).
- [5] P. Binetruy, C. Defayet, and D. langlois, Nucl. Phys. B 565, 269 (2000).
- [6] P.B inetruy, C.D efayet, U.E llwanger, and D. langlois, Phys.Lett.B 477, 285 (2000).
- [7] W D.Goldberger and M B.W ise, Phys. Rev. Lett. 83, 4922 (1999).
- [8] C.C. saki, M. Graesser, L. Randall, and J. Teming,

Phys.Rev.D 62,045015 (2000).

- [9] JM. Cline and H. Firouzjahi, Phys. Lett. B 495, 271 (2000).
- [10] W e assume such a potential is generated by the presence of an unspecied eld in the bulk which couples with the radion and serves the purpose of stabilizing the distance between the two branes of the RS Im odel.
- [11] Note that the approximation which makes use of an effective Lagrangian is only compatible with second order calculations.
- [12] The unperturbed brane distance is taken to be nite and equal to $b_0=2$.
- [13] In terms of the fundam ental scale M $\,$ one has m = M , k^2 = M 3 , b_0 = M $^{-1}$, and ! 2 = M 7 .



FIG.3: The two graphs in the rst row show the ratio between R to O (²) and $_1$ H² to O () (dotted line) and to O (²) (solid line) at a given time, for $_n = 10^2$ M⁴ and $w_n = 0.95$ (left) and for $_n = 10^1$ M⁴ and $w_n = 0.95$ (right). In the two graphs in the second row, the same ratios are are given for $_n = 2 \ 10^1$ M⁴ and $w_n = 0.95$ (left) and for $_n = 6 \ 10^1$ M⁴ and $w_n = 0.95$ (left). The plots cover all the bulk between the two branes.

[14] Let us recall that this is achieved by adopting the particular gauge choice for $c_{n\,;i}^{(1)}$ and $c_{n\,;i}^{(2)}$ which sets $f_{n\,;i}^{(1)}$ (t) =

 $f_{n,i}^{(2)}(t) = 0 \text{ and } c_{a,i}^{(1)} = c_{a,i}^{(2)} = 0.$



FIG.4: The two graphs in the nst row show the ratio between R to 0 (2) and $_1$ a=a to 0 () (dotted line) and to 0 (2) (solid line) at a given time, for $_n = 10$ 2 M 4 and $w_n = 0.95$ (left) and for $_n = 10$ 1 M 4 and $w_n = 0.95$ (right). In the two graphs in the second row, R to 0 (2) (solid line) is compared to the modula of $_1$ to 0 () (dashed line) and to 0 (2) (dotted line) for $_n = 2$ 10 1 M 4 and $w_n = 0.95$ (left), and for $_n = 6$ 10 1 M 4 and $w_n = 0.65$ (right). The plots cover all the bulk between the two branes.