

Curvaton Scenario in the Presence of Two Dilatons Coupled to the Scalar Curvature

Kazuharu BAMB¹ and Motohiko YOSHIMURA²

¹*Department of Earth and Space Science, Graduate School of Science,
Osaka University, Toyonaka 560-0043, Japan*

²*Department of Physics, Okayama University,
Okayama 700-8530, Japan*

Abstract

The possibility of the realization of a curvaton scenario is studied in a theory in which two dilatons are introduced along with coupling to the scalar curvature. It is shown that when the two dilatons have an approximate $O(2)$ symmetric coupling, a scalar field playing the role of the curvaton may exist in the framework of this theory without introducing any other scalar field for the curvaton. Thus the curvaton scenario can be realized, and in the simple version of the curvaton scenario, in which the curvaton potential is quadratic, the curvature perturbation with a sufficient amplitude and a nearly scale-invariant spectrum suggested by observations obtained from WMAP can be generated.

§1. Introduction

The hierarchy problem exists in both particle physics and cosmology. In the former case, this is the hierarchy problem between gravity and standard particle physics mass scales, while in the latter case, it is the presence of a very small but finite cosmological constant or the dark energy indicated by the analysis of type-Ia supernovae observations,¹⁾ whose magnitude is as small as 120 orders smaller than its theoretically natural value, i.e., the Planck scale (or a similar scale).²⁾ The origin of this small cosmological constant is not well understood yet. There has yet been no definite conclusion established in theoretical studies.

Recently, one of the present authors attempted to construct a model³⁾ which simultaneously solves the above two hierarchy problems by radically changing cosmology in the same spirit as the ideas of Dirac⁴⁾ and Brans and Dicke.^{5),6)} In this model, a theory with two dilaton fields coupled to the scalar curvature is considered. The existence of many dilatons is conceivable in light of higher-dimensional theories. Moreover, it has been argued that an effective gravity mass scale (the square root of the inverse of Newton's constant) increases in the inflationary stage⁷⁾ (for a review of inflation, see Refs. 8) and 9)), and that the small cosmological constant or the dark energy density in the present universe could be dynamically realized in the case that two dilatons have approximately an $O(2)$ symmetry, taking the fundamental mass scale to be of TeV. In such a TeV scale model, where the potential of the dilatons is of order $\sim (\text{TeV})^4$, however, it is conjectured that a naive estimate of the curvature perturbation gives a magnitude which is much smaller than the recent observational results of the anisotropy of the cosmic microwave background (CMB) radiation obtained from Wilkinson microwave background probe (WMAP).¹⁰⁾ Moreover, as is shown in §3, power-law inflation could be realized in this model. In the standard inflation models, however, inflation is driven by the potential energy of a scalar field, called an “inflaton”, as it slowly rolls down the potential hill. This slow roll over in the quasi-de Sitter stage is necessary to account for the nearly scale-invariant spectrum of the primordial curvature perturbation out of the quantum fluctuation of the inflaton, which is also suggested by the observational results obtained from WMAP.¹¹⁾

However, in recent years a new mechanism for generating the primordial curvature perturbation has been proposed, in which a late-decaying massive scalar field provides the dominant source of the curvature perturbation.^{12),13),14),15)} In this scenario, the dominant part of the curvature perturbation originates from the quantum fluctuations of a new scalar field, called the “curvaton”, which is different from the inflaton. In contrast to the situation with the usual mechanism, with this new mechanism, inflation need not be of the slow-roll variety. Hence, even in the case of power-law inflation, the nearly scale-invariant spectrum

of the curvature perturbation demanded by the observations can be realized. Instead, it is required that the curvaton potential be sufficiently flat during inflation.

The purpose of the present paper is to argue that the curvaton scenario can be realized in the above theory, in which two dilatons are introduced along with the coupling to the scalar curvature.³⁾ In particular, we show that there exists a scalar field corresponding to the curvaton in the framework of this theory without introducing any other scalar field that plays the role of the curvaton, and discuss the case in which the curvaton scenario is realized. Also, we are able to construct the curvature perturbation with sufficiently large amplitude and nearly scale-invariant spectrum suggested by the observational results obtained from WMAP.

This paper is organized as follows. In §2 we describe our model and derive field equations from its action. In §3 we show that power-law inflation can be realized in this model. Next, in the framework of this theory the existence of a scalar field corresponding to the curvaton is demonstrated in §4. Then, in §5, we study the case in which the curvaton scenario can be realized in this model. Finally, §6 is devoted to a conclusion. Throughout the present paper we use units in which $k_B = c = \hbar = 1$ and denote Newton's constant by $G = m_{\text{Pl}}^{-2}$, where $m_{\text{Pl}} = 1.2 \times 10^{19} \text{GeV}$ is the Planck mass.

§2. Model

2.1. Action

We introduce two scalar fields, φ_i ($i = 1, 2$). Furthermore, we introduce dilatonic coupling of these scalars to the scalar curvature. Our model action is the following:³⁾

$$S = \int d^4x \sqrt{-g} \left[-f(\varphi_i)R + \frac{1}{2}(\partial\varphi_i)^2 - V[\varphi_i] + \mathcal{L}_m \right], \quad (1)$$

$$f(\varphi_i) = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2, \quad (2)$$

$$V[\varphi_i] = V_0 \cos \frac{\varphi_r}{M} + \Lambda, \quad (3)$$

where R is the scalar curvature arising from the spacetime metric tensor $g_{\mu\nu}$, g is the determinant of $g_{\mu\nu}$, $f(\varphi_i)$ is the coupling between the dilaton and the scalar curvature [with ϵ_i (> 0) being dimensionless constants], $V[\varphi_i]$ is the potential of the dilaton (with $\varphi_r = \sqrt{\varphi_1^2 + \varphi_2^2}$ and $V_0 > \Lambda > 0$), and M denotes a mass scale. Here Λ is the collection of all the constants in the Standard Model Lagrangian \mathcal{L}_m such that the potential of the Standard Model Lagrangian vanishes at its minimum. We assume $O(2)$ symmetry for the potential $V[\varphi_i]$,

while we allow its violation by choosing $\epsilon_1 \neq \epsilon_2$. Moreover, we use the simplified notation $(\partial\varphi_i)^2 \equiv g^{\mu\nu}\partial_\mu\varphi_i\partial_\nu\varphi_i$.

The dilatonic coupling of the scalar field to the scalar curvature given by $f(\varphi_i)R$ is taken from the Brans-Dicke theory.⁵⁾ However, with regard to the choice of the potential $V[\varphi_i]$, we part with the Brans-Dicke theory, in which a single dilaton, φ , is introduced along with the null potential and $f(\varphi) = (1/2)\xi\varphi^2$, where ξ is a dimensionless constant.⁶⁾ For both simplicity and naturalness, we assume that all mass parameters are of the same order, and thus $V_0 \approx \Lambda = O[M^4]$ for the choice (3). In the present model, we use a common mass scale M of order m_{Pl} , although in the scenario proposed in Ref. 3), a common mass scale M of order TeV, favored by the present small value of the dark energy, is used.

2.2. Field equations

The field equations can be derived by taking variations of the above action in Eq. (1) with respect to the metric $g_{\mu\nu}$ and the dilatons φ_i as follows:¹⁶⁾

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{1}{2f} [T_{\mu\nu}^{(\text{m})} + T_{\mu\nu}^{(\varphi_i)}] + \frac{1}{f} (\nabla_\mu\nabla_\nu f - g_{\mu\nu}\square f), \quad (4)$$

with

$$T_{\mu\nu}^{(\varphi_i)} = \partial_\mu\varphi_i\partial_\nu\varphi_i - g_{\mu\nu} \left[\frac{1}{2}(\partial\varphi_i)^2 - V[\varphi_i] \right], \quad (5)$$

and

$$\square\varphi_i = -\frac{\partial V}{\partial\varphi_i} - \frac{\partial f}{\partial\varphi_i}R, \quad (6)$$

where ∇_μ is the covariant derivative operator associated with $g_{\mu\nu}$, and $\square \equiv g^{\mu\nu}\nabla_\mu\nabla_\nu$ is the covariant d'Alembertian for a scalar field. In addition, $R_{\mu\nu}$ is the Ricci curvature tensor, while $T_{\mu\nu}^{(\varphi_i)}$ is the contribution to the energy-momentum tensor from the scalars φ_i , and $T_{\mu\nu}^{(\text{m})}$ is the usual contribution of radiation, matter and other fields. Taking the trace of Eq. (4), we obtain

$$-R = \frac{1}{2f} [T^{(\text{m})} - (\partial\varphi_i)^2 + 4V(\varphi_i) - 6\square f], \quad (7)$$

where $T^{(\text{m})}$ is the trace of $T_{\mu\nu}^{(\text{m})}$.

We now assume the spatially flat Friedmann-Robertson-Walker (FRW) spacetime with the metric

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = dt^2 - a^2(t)d\mathbf{x}^2, \quad (8)$$

where $a(t)$ is the scale factor. In the FRW metric (8), the equations of motion for the background homogeneous scalar fields read

$$\ddot{\varphi}_i + 3H\dot{\varphi}_i = -\frac{\varphi_i}{\varphi_r}V' - 2\epsilon_i\varphi_i R, \quad (9)$$

where the dot denotes differentiation with respect to time, and the prime denotes differentiation with respect to φ_r . Here H is the Hubble parameter. Using the equations of motion for the dilaton fields (9) with Eq. (7), we find the dynamical equations in terms of the two field variables $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\tilde{f} = \epsilon_1\varphi_1^2 - \epsilon_2\varphi_2^2$ as

$$\begin{aligned} \ddot{f} + 3H\dot{f} = 2F^{-1} \left\{ (\epsilon_1\dot{\varphi}_1^2 + \epsilon_2\dot{\varphi}_2^2) - \frac{f}{\varphi_r}V' \right. \\ \left. + \frac{\epsilon_1^2\varphi_1^2 + \epsilon_2^2\varphi_2^2}{f} [T^{(m)} + 4V - (\dot{\varphi}_1^2 + \dot{\varphi}_2^2)] \right\}, \end{aligned} \quad (10)$$

$$\begin{aligned} \ddot{\tilde{f}} + 3H\dot{\tilde{f}} = 2F^{-1} \left\{ F(\epsilon_1\dot{\varphi}_1^2 - \epsilon_2\dot{\varphi}_2^2) - \left[\frac{\tilde{f}}{f} - 24\epsilon_1\epsilon_2(\epsilon_1 - \epsilon_2) \left(\frac{\varphi_1\varphi_2}{f} \right)^2 \right] \frac{f}{\varphi_r}V' \right. \\ \left. + \left(\frac{\epsilon_1^2\varphi_1^2 - \epsilon_2^2\varphi_2^2}{f} \right) [T^{(m)} + 4V - (\dot{\varphi}_1^2 + \dot{\varphi}_2^2) - 12(\epsilon_1\dot{\varphi}_1^2 + \epsilon_2\dot{\varphi}_2^2)] \right\}, \end{aligned} \quad (11)$$

$$F = 1 + 12 \frac{\epsilon_1^2\varphi_1^2 + \epsilon_2^2\varphi_2^2}{f}. \quad (12)$$

(The reason we have chosen the field variable \tilde{f} as the partner of the coupling f is stated in §A.2.) Furthermore, the gravitational field equations read

$$H^2 = \left(\frac{\dot{a}}{a} \right)^2 = \frac{1}{6f} \left[T_{00}^{(m)} + \frac{1}{2}(\dot{\varphi}_1^2 + \dot{\varphi}_2^2) + V \right] - H \frac{\dot{f}}{f}, \quad (13)$$

$$\begin{aligned} \dot{H} - 2H \frac{\dot{f}}{f} = -\frac{F^{-1}}{f} \left[\frac{1}{12} \left(4FT_{00}^{(m)} - T^{(m)} \right) + (\epsilon_1\dot{\varphi}_1^2 + \epsilon_2\dot{\varphi}_2^2) \right. \\ \left. + \frac{1}{4} \left(1 + 8 \frac{\epsilon_1^2\varphi_1^2 + \epsilon_2^2\varphi_2^2}{f} \right) (\dot{\varphi}_1^2 + \dot{\varphi}_2^2) \right. \\ \left. - \frac{f}{\varphi_r}V' + 4 \frac{\epsilon_1^2\varphi_1^2 + \epsilon_2^2\varphi_2^2}{f} V \right], \end{aligned} \quad (14)$$

where in deriving the expression (14) we have used Eq. (10).

§3. Power-law inflation

In this section, we show that power-law inflation can be realized in the presently considered model. Here we seek solutions with the ansatz

$$f = At^2, \quad L = Bt, \quad \varphi_r = Ct, \quad a \propto t^\omega, \quad (15)$$

which is valid for large t . Here, $L = \varphi_1 \dot{\varphi}_2 - \varphi_2 \dot{\varphi}_1$ is the angular momentum. The existence of the angular momentum is one of the important features of this model. We now assume that in the inflationary stage, the cosmic energy density is dominated by the dilatons, and hence $T_{\mu\nu}^{(m)}$ is negligible. Moreover, for the moment we ignore variation of the potential and hence we replace V by its average value, A .

Substituting the ansatz (15) into Eqs. (10), (13) and (14), we obtain

$$1 + 3\omega = F^{-1} \left[4 \left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} \right) \frac{A}{A} - \left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} - \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} \right) \frac{1}{A} \left(C^2 + \frac{B^2}{C^2} \right) \right], \quad (16)$$

$$\omega^2 + 2\omega = \frac{1}{12} \left[2 \frac{A}{A} + \frac{1}{A} \left(C^2 + \frac{B^2}{C^2} \right) \right], \quad (17)$$

$$\omega = F^{-1} \left[\frac{4}{5} \left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} \right) \frac{A}{A} + \frac{1}{20} \left(1 + 8 \frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} + 4 \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} \right) \frac{1}{A} \left(C^2 + \frac{B^2}{C^2} \right) \right], \quad (18)$$

where in deriving the expressions (16)–(18) we have used the following relation:

$$\dot{\varphi}_1^2 + \dot{\varphi}_2^2 = \dot{\varphi}_r^2 + \frac{L^2}{\varphi_r^2} = C^2 + \frac{B^2}{C^2}. \quad (19)$$

From Eqs. (17) and (18), we find

$$\frac{A}{A} = 2 \left(1 + 4 \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} \right)^{-1} \times \left[3(\omega^2 + 2\omega) \left(1 + 8 \frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} + 4 \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} \right) - 5\omega F \right], \quad (20)$$

$$\frac{1}{A} \left(C^2 + \frac{B^2}{C^2} \right) = 4 \left(1 + 4 \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} \right)^{-1}$$

$$\times \left[-24(\omega^2 + 2\omega) \left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} \right) + 5\omega F \right]. \quad (21)$$

Eliminating Λ/A and $(C^2 + B^2/C^2)/A$ from Eqs. (16), (20) and (21), we obtain

$$24 \left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} \right) \omega^2 - \left[12 \left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} \right) - 8 \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} + 3 \right] \omega - \left(1 + 4 \frac{\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2}{\dot{\varphi}_1^2 + \dot{\varphi}_2^2} \right) = 0. \quad (22)$$

We now assume $\varphi_1^2 \approx \varphi_2^2$ and $\dot{\varphi}_1^2 \approx \dot{\varphi}_2^2$ and approximately determine ω in terms of ϵ_i . It follows from these approximate relations that Eq. (22) can be rewritten in the form

$$24\beta\omega^2 - (12\beta - 4\alpha^2 + 3\alpha)\omega - \alpha(1 + 2\alpha) = 0, \quad (23)$$

with

$$\alpha \equiv \epsilon_1 + \epsilon_2, \quad (24)$$

$$\beta \equiv \epsilon_1^2 + \epsilon_2^2. \quad (25)$$

The solution of Eq. (23) is given by

$$\omega = \frac{1}{48\beta} \left[(12\beta - 4\alpha^2 + 3\alpha) + \sqrt{D} \right], \quad (26)$$

$$D = 9\alpha^2 - 24\alpha^3 + 168\alpha\beta + 16\alpha^4 + 96\alpha^2\beta + 144\beta^2, \quad (27)$$

where we have taken the positive solution of ω . In the case $\epsilon_i \ll 1$, considering only the terms of leading order in ϵ_i in expressions (26) and (27), we find

$$\omega \approx \frac{\alpha}{8\beta} = \frac{\epsilon_1 + \epsilon_2}{8(\epsilon_1^2 + \epsilon_2^2)}. \quad (28)$$

Hence, for $\epsilon_i \ll 1$, the index of the power ω can be much larger than unity, and thus the power-law inflation can certainly be realized. From this point, we consider only the case $\epsilon_i \ll 1$.

A large value of ω is favored to approximately mimic the exponential expansion of the cosmological scale factor. The gravity mass scale increases as $f \propto a^{2/\omega}$ during inflation. Thus, there is no difficulty in obtaining a large enough number of e -folds during inflation, and simultaneously resolving the mass hierarchy problem. This is a feature that also appears in the model of extended inflation¹⁷⁾ (see also Ref. 18)). The number of e -folds for a particular case in this model is estimated in §4.3. The requirement for power-law inflation in the

curvaton scenario is that the power-law exponent ω be much larger than unity, as is discussed in §5. Furthermore, we note the following point. The discussion in this section corresponds to analysis in the Jordan frame, in which there exist scalar fields nonminimally coupled to the scalar curvature. It is not obvious, however, that power-law inflation can be realized not only in the Jordan frame but also in the Einstein frame, in which there ordinarily exist scalar fields minimally coupled to the scalar curvature. For this reason, in Appendix B, we show that power-law inflation can also be realized in the Einstein frame.

Here we present approximate expressions of A , B , and C . Their detailed derivations are given in §A.1. These approximate expressions are as follows:

$$A \approx \frac{32}{3} \left(\frac{\beta}{\alpha} \right)^2 \Lambda, \quad (29)$$

$$B \approx \pm \frac{128\beta^2}{3\alpha^3} \sqrt{\frac{\alpha^2}{2\beta - \alpha^2}} \Lambda, \quad (30)$$

$$C \approx \sqrt{\frac{64\beta^2}{3\alpha^3}} \Lambda. \quad (31)$$

Finally, we discuss the following points. We first consider the allowed case in which we can ignore the variation of the potential $V[\varphi_i] = V_0 \cos(\varphi_r/M) + \Lambda$ in Eq. (3), and therefore replace V by its average value, Λ , as done in deriving the solutions of power-law inflation at the beginning of this section. As explained in §4.2, in this model, we consider the following case. In the inflationary stage, the field amplitude of the dilatons is given by $\varphi_1 = \gamma_1 M$, $\varphi_2 = \gamma_2 M$, where γ_1 and γ_2 are dimensionless parameters with time dependence. In addition, we assume $\gamma_1 \approx \gamma_2 \approx \gamma = \gamma(t)$; these relations are consistent with the assumption $\varphi_1^2 \approx \varphi_2^2$, used in deriving the approximate expressions of A , B , C and ω above. In this case, we have $\varphi_r = \sqrt{\varphi_1^2 + \varphi_2^2} \approx \gamma M$. Hence, if $\gamma(t) \ll 1$ during inflation, we can ignore variation of the potential. In this case, we find $V \approx V_0 + \Lambda$. Thus it follows from the assumption $V_0 \approx \Lambda = O[M^4]$ made in §2.1 that it is valid to recognize V as being approximately equal to Λ . As shown in §4.2, the case in which we are interested corresponds to just this case. In fact, for example, in the case $\epsilon_1 = 3.1 \times 10^{-3}$, $\epsilon_2 = 3.0 \times 10^{-3}$ and $\gamma = 3.5 \times 10^{-4}$, which is the case (v) in Table I of §4.2, the solution derived by partially taking the variation of the potential V into consideration is $\omega = 4.15 \times 10^1 \gg 1$. Hence power-law inflation can be practically realized. Furthermore, the value $\omega = 4.10 \times 10^1$, which is estimated by ignoring the potential variation and replacing V by its average value Λ , as shown in Table I, is very close to that in the above solution. Thus, if $\gamma(t) \ll 1$ during inflation, the above approximation in which we ignore the variation of the potential and regard V as being approximately Λ is valid. Here

we note that in deriving the above solution by partially taking the variation of the potential V into consideration, we have used $\varphi_r \approx \gamma M$ and the relation $t \approx \omega \sqrt{6\alpha\gamma^2}/M$, which is derived by using Eqs. (41) and (42), which are given in §4.2, and the relation $H = \omega/t$.

From the above consideration, we see that if $\gamma(t) \ll 1$ during inflation, then this case is just the case considered in the later discussion appearing in §4 and 5, as stated above, and power-law inflation can be practically realized in this model. In contrast to our scenario, in the scenario proposed in Ref. 3), which simultaneously solves the hierarchy problem between gravity and particle physics mass scales and the small cosmological constant or the dark energy problem, the following case is considered. In the inflationary stage, φ_r goes over the many local maxima of the potential and then changes for many periods, i.e., $\gamma(t) \gtrsim 1$ and $\gamma(t_f) - \gamma(t_i) \gg 1$, where t_f and t_i are the time at the end and at the beginning of inflation, respectively. Thus, the situation for the evolution of φ_r during inflation considered in the present model is different from that in the scenario proposed in Ref. 3).

Second, we consider the reason why power-law inflation occurs. In the inflationary stage, from Eq. (13) we find $H^2 \approx 1/(6f) [(\dot{\varphi}_1^2 + \dot{\varphi}_2^2)/2 + V] \approx \Lambda/(6f)$. Here, in deriving the first approximate equality, we have used $H^2 \gg H\dot{f}/f$. This relation follows from $H/(\dot{f}/f) = \omega/2 \gg 1$, where we have used $H = \omega/t$ and $f = At^2$. Moreover, in deriving the second approximate equality, we have used $(\dot{\varphi}_1^2 + \dot{\varphi}_2^2)/2 + V \approx [16\beta/(3\alpha)]\Lambda + \Lambda \approx [2/(3\omega)]\Lambda + \Lambda \approx \Lambda$, where in deriving the first approximate equality we have used Eqs. (19) and (A.12), in deriving the second approximate equality we have used Eq. (28), and in deriving the last approximate equality we have used $\omega \gg 1$. Furthermore, we have ignored variation of the potential and hence replaced V by its average value, Λ . From the above consideration, we understand that the contribution of the cosmological constant Λ to the power-law inflation is much larger than that of the kinetic term of the dilatons. Thus, the contribution to the power-law inflation comes mainly from the effect of cosmological constant, Λ .

§4. Extremely large mass hierarchy

In this section, we show that there exists a scalar field corresponding to the curvaton in the framework of this theory without introducing any other scalar field that plays the role of the curvaton. First, we discuss the fluctuation equation in terms of $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\tilde{f} = \epsilon_1\varphi_1^2 - \epsilon_2\varphi_2^2$ in order to derive these mass values in the inflationary stage. Next, we consider the case in which an extremely large hierarchy of these mass values in the inflationary stage can be realized. The realization of this extremely large hierarchy implies that the potential of the scalar field with the lighter mass is much flatter than that of the scalar field with the heavier mass. Furthermore, if the lighter mass is much smaller than the

Hubble parameter in the inflationary stage, the scalar field with the lighter mass can be the curvaton, as discussed in §5.2.

4.1. Linearized equations

To begin with, we discuss the fluctuation equation in terms of $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\tilde{f} = \epsilon_1\varphi_1^2 - \epsilon_2\varphi_2^2$ in order to derive these mass values in the inflationary stage. With the ansatz

$$\begin{aligned} f &= f_0 + \delta f, & |\delta f| &\ll |f_0|, \\ \tilde{f} &= \tilde{f}_0 + \delta\tilde{f}, & |\delta\tilde{f}| &\ll |\tilde{f}_0|, \end{aligned} \quad (32)$$

where f_0 and \tilde{f}_0 are the zeroth order quantities and satisfy Eqs. (10) and (11), respectively, we obtain the linearized equations

$$\begin{pmatrix} \ddot{\delta f} + 3H\dot{\delta f} \\ \ddot{\delta\tilde{f}} + 3H\dot{\delta\tilde{f}} \end{pmatrix} = -\mathcal{M}^2 \begin{pmatrix} \delta f \\ \delta\tilde{f} \end{pmatrix}, \quad (33)$$

where

$$\mathcal{M}^2 \equiv \begin{pmatrix} \mathcal{M}_{11}^2 & \mathcal{M}_{12}^2 \\ \mathcal{M}_{21}^2 & \mathcal{M}_{22}^2 \end{pmatrix}. \quad (34)$$

In deriving the expression (33), we have kept terms up to those of first order in δf and $\delta\tilde{f}$. A detailed derivation of Eq. (33) and the expressions of the components of \mathcal{M}^2 are given in §A.2.

Here we emphasize the reason why we need to estimate the mass values of the scalar quantities f and \tilde{f} . As stated at the beginning of this section, the final purpose of this section is to show that there exists a scalar field corresponding to the curvaton in the framework of this theory, without introducing any other scalar field that plays the role of the curvaton. The necessary condition for a scalar field to correspond to the curvaton is that its mass be much smaller than the Hubble parameter in the inflationary stage. Hence, in order to examine the existence of a scalar field corresponding to the curvaton, we consider not the mass scales of the dilatons φ_i themselves, but those of the scalar quantities $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\tilde{f} = \epsilon_1\varphi_1^2 - \epsilon_2\varphi_2^2$, which are expressed as linear combinations of φ_1^2 and φ_2^2 . We cannot easily estimate these mass values from Eqs. (10) and (11) themselves, however, because the parts corresponding to the mass terms of these equations are too complicated. Therefore we consider the linearized equations (33) by making use of the ansatz (32), and then we estimate the eigenmasses of f and \tilde{f} by carrying out the diagonalization of the matrix \mathcal{M}^2 in (34), as discussed in the next subsection.

4.2. Mass diagonalization

Next, we show that an extremely large hierarchy of the mass values of f and \tilde{f} in the inflationary stage can be realized. For this purpose, we investigate these eigenmasses by solving the following characteristic equation of \mathcal{M}^2 :

$$\det(\mathcal{M}^2 - \lambda E) = 0 \iff \lambda^2 - \text{Tr} \mathcal{M}^2 \lambda + \det \mathcal{M}^2 = 0, \quad (35)$$

where λ is an eigenvalue of \mathcal{M}^2 , and E is the unit matrix. The solutions of Eq. (35) are equivalent to the diagonal components of the diagonalized matrix of \mathcal{M}^2 , namely, the square of the eigenmasses of f and \tilde{f} . The solution of Eq. (35) is given by

$$\lambda_{\pm} = \frac{\text{Tr} \mathcal{M}^2}{2} \left[1 \pm \sqrt{1 - \frac{4 \det \mathcal{M}^2}{(\text{Tr} \mathcal{M}^2)^2}} \right], \quad (36)$$

where the signs \pm in λ_{\pm} correspond to those on the right-hand side of this equation.

The necessary conditions for an extremely large hierarchy of the eigenmasses of f and \tilde{f} to be realized are the following three: (I) $\text{Tr} \mathcal{M}^2 > 0$, (II) $\det \mathcal{M}^2 > 0$, and (III) $\Theta \equiv 4 \det \mathcal{M}^2 / (\text{Tr} \mathcal{M}^2)^2 \ll 1$. The first and second conditions are necessary in order that both the square of the eigenmasses can be positive, that is, $\lambda_+ > 0$ and $\lambda_- > 0$. The third condition is necessary in order that an extremely large hierarchy of the eigenmasses can be realized, that is, $\lambda_- / \lambda_+ \ll 1$. In the case that the above three conditions are satisfied, it follows from Eq. (36) that the square of the eigenmasses of f and \tilde{f} can be approximately expressed as

$$\lambda_+ \approx \text{Tr} \mathcal{M}^2 \left[1 - \frac{\det \mathcal{M}^2}{(\text{Tr} \mathcal{M}^2)^2} \right] > 0, \quad (37)$$

$$\lambda_- \approx \frac{\det \mathcal{M}^2}{\text{Tr} \mathcal{M}^2} > 0, \quad (38)$$

$$\frac{\lambda_-}{\lambda_+} \approx \frac{\det \mathcal{M}^2}{(\text{Tr} \mathcal{M}^2)^2} \left[1 + \frac{\det \mathcal{M}^2}{(\text{Tr} \mathcal{M}^2)^2} \right] \ll 1, \quad (39)$$

where each approximate equality in these relations follows from the above third condition, $\Theta = 4 \det \mathcal{M}^2 / (\text{Tr} \mathcal{M}^2)^2 \ll 1$. We therefore investigate the case in which the above three conditions are satisfied by estimating $\text{Tr} \mathcal{M}^2$ and $\det \mathcal{M}^2$.

Here we consider the following case. As stated in §2, in regard to the dilaton potential appearing in Eq. (3), we assume $V_0 \approx \Lambda \approx M^4$. Moreover, in the inflationary stage, the field amplitude of the dilatons is given by $\varphi_1 = \gamma_1 M$ and $\varphi_2 = \gamma_2 M$, where γ_1 and γ_2 are dimensionless parameters with time dependence. In addition, we assume $\gamma_1 \approx \gamma_2 \approx \gamma =$

$\gamma(t)$. (These relations are consistent with the assumption $\varphi_1^2 \approx \varphi_2^2$, made in deriving the approximate expressions of A , B , C and ω in §3.) Furthermore, we consider the case in which the value of ϵ_1 is close to that of ϵ_2 and introduce a dimensionless constant $\bar{\epsilon} \equiv (\epsilon_1 + \epsilon_2)/2 = \alpha/2 = O[\epsilon_i]$. In this case, from Eq. (2) we find

$$f \approx 2\bar{\epsilon}\gamma^2 M^2 = \alpha\gamma^2 M^2. \quad (40)$$

Hence, it follows from Eqs. (13) and (40) that the Hubble parameter in the inflationary stage is expressed as

$$H \approx \sqrt{\frac{M^4}{6f}} \approx \Upsilon M, \quad (41)$$

$$\Upsilon \equiv \frac{1}{\sqrt{12\bar{\epsilon}\gamma^2}} = \frac{1}{\sqrt{6\alpha\gamma^2}}. \quad (42)$$

In deriving the first approximate equality in Eq. (41) we have used two relations. The first is $(\dot{\varphi}_1^2 + \dot{\varphi}_2^2)/2 + V \approx [16\beta/(3\alpha)]\Lambda + \Lambda \approx [2/(3\omega)]\Lambda + \Lambda \approx \Lambda$, where, in deriving the first approximate equality, we have used Eqs. (19) and (A.12), in deriving the second approximate equality, we have used Eq. (28), and in deriving the last approximate equality, we have used $\omega \gg 1$. Furthermore, we have ignored variation of the potential, and hence replaced V by its average value, Λ . The second relation used here is $H^2 \gg H\dot{f}/f$. This relation follows from $H/(\dot{f}/f) = \omega/2 \gg 1$, where we have used $H = \omega/t$ and $f = At^2$ in the ansatz (15).

From Eq. (3) and the expressions (A.22), (A.24), (A.26) and (A.28), we can obtain approximate expressions for $\text{Tr } \mathcal{M}^2$ and $\det \mathcal{M}^2$. For convenience in describing these expressions, we represent $\text{Tr } \mathcal{M}^2$ and $\det \mathcal{M}^2$ as

$$\begin{aligned} \text{Tr } \mathcal{M}^2 &= \mathcal{M}_{11}^2 + \mathcal{M}_{22}^2 \\ &\equiv (\text{Tr } \mathcal{M}^2)^{(0)} + (\text{Tr } \mathcal{M}^2)^{(1)}, \end{aligned} \quad (43)$$

and

$$\begin{aligned} \det \mathcal{M}^2 &= \mathcal{M}_{11}^2 \mathcal{M}_{22}^2 - \mathcal{M}_{12}^2 \mathcal{M}_{21}^2 \\ &\equiv (\det \mathcal{M}^2)^{(0)} + (\det \mathcal{M}^2)^{(1)} + (\det \mathcal{M}^2)^{(2)}. \end{aligned} \quad (44)$$

Here, $(\text{Tr } \mathcal{M}^2)^{(0)}$ and $(\det \mathcal{M}^2)^{(0)}$ are the parts independent of the sine functions. Moreover, $(\text{Tr } \mathcal{M}^2)^{(1)}$ and $(\det \mathcal{M}^2)^{(1)}$ are the parts proportional to $\sin[\varphi_r/M + \arctan(X_1/X_2)]$ and $\sin[\varphi_r/M + \arctan(Y_1/Y_2)]$, respectively, and $(\det \mathcal{M}^2)^{(2)}$ is the part proportional to

$\sin [2\varphi_r/M + \arctan (Z_1/Z_2)]$. These approximate expressions for $\text{Tr } \mathcal{M}^2$ and $\det \mathcal{M}^2$ and the expressions for X_1, X_2, Y_1, Y_2, Z_1 and Z_2 are presented in §A.3. Here, as stated above, we have considered the case $V_0 \approx \Lambda \approx M^4$. Moreover, from Eq. (41) we have used the relation

$$M \approx \frac{H}{\Upsilon} = \frac{\omega}{\Upsilon} \frac{1}{t} \approx \frac{\alpha}{8\beta} \frac{1}{\Upsilon} \frac{1}{t}, \quad (45)$$

where the last approximate equality follows from Eq. (28).

We now approximately evaluate $\text{Tr } \mathcal{M}^2$ and $\det \mathcal{M}^2$ by using Eqs. (A.42)–(A.48) and then investigate the case in which the above three conditions can be satisfied. In order to confirm that the above three conditions can always be satisfied independently of the values of the sine functions, we investigate the case that the following three relations can be satisfied:

$$\text{Tr } \tilde{\mathcal{M}}^2 \equiv (\text{Tr } \mathcal{M}^2)^{(0)} - \frac{\sqrt{X_1^2 + X_2^2}}{t^2} > 0, \quad (46)$$

$$\det \tilde{\mathcal{M}}^2 \equiv (\det \mathcal{M}^2)^{(0)} - \frac{\sqrt{Y_1^2 + Y_2^2} + \sqrt{Z_1^2 + Z_2^2}}{t^4} > 0, \quad (47)$$

$$\tilde{\Theta} \equiv 4 \frac{(\det \mathcal{M}^2)^{(0)} + \left(\sqrt{Y_1^2 + Y_2^2} + \sqrt{Z_1^2 + Z_2^2} \right) t^{-4}}{\left(\text{Tr } \tilde{\mathcal{M}}^2 \right)^2} \ll 1, \quad (48)$$

where $\tilde{\Theta}$ is positive. The first relation means that the minimum of $\text{Tr } \mathcal{M}^2$ is positive. Moreover, the second relation means that a value smaller than the minimum of $\det \mathcal{M}^2$ is positive. Furthermore, the third relation means that a value larger than the maximum of Θ is much smaller than unity. The reason we have adopted $\det \tilde{\mathcal{M}}^2$ as a value smaller than the minimum of $\det \mathcal{M}^2$ and $\tilde{\Theta}$ as a value larger than the maximum of Θ is that we cannot obtain analytical expressions of the minimum and maximum of $\det \mathcal{M}^2$, because, as seen from Eqs. (A.45)–(A.48), $\det \mathcal{M}^2$ has two terms proportional to sine functions whose phases differ.

Moreover, as the effective values of λ_{\pm} and the ratio λ_-/λ_+ , we adopt the average values

$$\langle \lambda_+ \rangle \equiv (\text{Tr } \mathcal{M}^2)^{(0)} \left[1 - \frac{(\det \mathcal{M}^2)^{(0)}}{((\text{Tr } \mathcal{M}^2)^{(0)})^2} \right], \quad (49)$$

$$\langle \lambda_- \rangle \equiv \frac{(\det \mathcal{M}^2)^{(0)}}{(\text{Tr } \mathcal{M}^2)^{(0)}}, \quad (50)$$

$$\left\langle \frac{\lambda_-}{\lambda_+} \right\rangle \equiv \frac{(\det \mathcal{M}^2)^{(0)}}{((\text{Tr } \mathcal{M}^2)^{(0)})^2} \left[1 + \frac{(\det \mathcal{M}^2)^{(0)}}{((\text{Tr } \mathcal{M}^2)^{(0)})^2} \right], \quad (51)$$

where ‘ $\langle \rangle$ ’ denotes the average over many periods of the sine functions.

Table I. Examples of the values of the model parameters in the case that an extremely large hierarchy of the eigenmasses of f and \tilde{f} can be realized, that is, the three relations (46)–(48) can be satisfied. Here we have approximately evaluated ω , γ , and t^{-1} by using $\omega \approx \alpha/(8\beta)$, $\gamma \approx 1/\sqrt{6\alpha\Upsilon^2}$, and $t^{-1} \approx (8\beta/\alpha)\Upsilon M$, respectively.

| | ϵ_1 | ϵ_2 | Υ | ω | γ | t^{-1}/M |
|-------|----------------------|----------------------|-------------------|-------------------|----------------------|-------------------|
| (i) | 4.1×10^{-2} | 4.0×10^{-2} | 3.5×10^1 | 3.1 | 4.1×10^{-2} | 1.1×10^1 |
| (ii) | 2.6×10^{-2} | 2.5×10^{-2} | 3.0×10^1 | 4.9 | 6.0×10^{-2} | 6.1 |
| (iii) | 1.1×10^{-2} | 1.0×10^{-2} | 3.5×10^2 | 1.2×10^1 | 8.0×10^{-3} | 2.9×10^1 |
| (iv) | 6.6×10^{-3} | 6.5×10^{-3} | 4.5×10^2 | 1.9×10^1 | 7.9×10^{-3} | 2.4×10^1 |
| (v) | 3.1×10^{-3} | 3.0×10^{-3} | 1.5×10^4 | 4.1×10^1 | 3.5×10^{-4} | 3.7×10^2 |

Table II. Estimates of the quantities in the three relations (46)–(48) and of the effective values of the square of the eigenmasses of f and \tilde{f} and their ratio in the cases (i)–(v) displayed in Table I. Here we have defined $\Xi_1 \equiv \text{Tr} \tilde{\mathcal{M}}^2/M^2$ and $\Xi_2 \equiv \det \tilde{\mathcal{M}}^2/M^2$. Moreover, we have estimated these quantities by using Eqs. (A·43), (A·44) and (A·46)–(A·48).

| | Ξ_1 | Ξ_2 | $\tilde{\Theta}$ | $\langle \lambda_+ \rangle / M^2$ | $\langle \lambda_- \rangle / M^2$ | $\langle \lambda_- / \lambda_+ \rangle$ |
|-------|-------------------|----------------------|----------------------|-----------------------------------|-----------------------------------|---|
| (i) | 1.0×10^7 | 2.5×10^7 | 4.2×10^{-5} | 1.0×10^7 | 5.4×10^1 | 5.4×10^{-6} |
| (ii) | 1.0×10^6 | 5.0×10^6 | 3.0×10^{-4} | 1.0×10^6 | 4.0×10^1 | 4.0×10^{-5} |
| (iii) | 3.3×10^6 | 3.1×10^7 | 9.8×10^{-4} | 3.4×10^6 | 4.1×10^2 | 1.2×10^{-4} |
| (iv) | 8.2×10^7 | 2.1×10^9 | 2.7×10^{-5} | 8.2×10^7 | 2.9×10^2 | 3.6×10^{-6} |
| (v) | 4.1×10^9 | 5.8×10^{11} | 3.8×10^{-5} | 4.1×10^9 | 1.9×10^4 | 4.7×10^{-6} |

Table I displays some examples of the values of the model parameters in the case in which an extremely large hierarchy of the eigenmasses of f and \tilde{f} can be realized, that is, the three relations (46)–(48) can be satisfied. Here, from Eqs. (28), (42) and (45), we have approximately evaluated ω , γ , and t^{-1} by using $\omega \approx \alpha/(8\beta)$, $\gamma \approx 1/\sqrt{6\alpha\Upsilon^2}$, and $t^{-1} \approx (8\beta/\alpha)\Upsilon M$, respectively. Furthermore, Table II lists estimates of the quantities $\Xi_1 \equiv \text{Tr} \tilde{\mathcal{M}}^2/M^2$, $\Xi_2 \equiv \det \tilde{\mathcal{M}}^2/M^2$, and $\tilde{\Theta}$ in Eqs. (46)–(48) and estimates of the effective values of the square of the eigenmasses $\langle \lambda_+ \rangle$ and $\langle \lambda_- \rangle$, and the ratio $\langle \lambda_- / \lambda_+ \rangle$ in the cases displayed in Table I. Here we have estimated these quantities by using Eqs. (A·43), (A·44) and (A·46)–(A·48).

Figures 1, 2, and 3 depict estimates of $\Xi_1 = \text{Tr} \tilde{\mathcal{M}}^2/M^2$, $\Xi_2 = \det \tilde{\mathcal{M}}^2/M^2$, and $\tilde{\Theta}$, respectively. Moreover, Figs. 4 and 5 depict estimates of $\langle \lambda_+ \rangle / M^2$ and $\langle \lambda_- \rangle / M^2$, and the ratio $\langle \lambda_- / \lambda_+ \rangle$. In all the figures, the solid lines represent the case $\Upsilon = 1.5 \times 10^4$, and the dotted lines represent the case $\Upsilon = 3.5 \times 10^2$. Moreover, the left panels correspond to the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($0 \leq j \leq 89$), where j

is an integer, while the right panels correspond to the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 89$). In Fig. 1, for all the cases, we have $\Xi_1 > 0$, and hence the relation (46) is satisfied. In Fig. 2, in the case $\Upsilon = 1.5 \times 10^4$, for $\epsilon_2 \leq 2.9 \times 10^{-3}$ and $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\Xi_2 < 0$. Moreover, in the case $\Upsilon = 3.5 \times 10^2$, for $\epsilon_2 \leq 6.9 \times 10^{-3}$ and $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\Xi_2 < 0$. Hence, in these cases, the relation (47) is not satisfied. In Fig. 3, in the case $\Upsilon = 1.5 \times 10^4$, for $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\tilde{\Theta} < 0$. Moreover, in the case $\Upsilon = 3.5 \times 10^2$, for $\epsilon_2 \geq 4.6 \times 10^{-2}$, we have $\tilde{\Theta} < 0$. In the remaining cases, the relation $0 < \tilde{\Theta} \ll 1$ holds, and hence the relation (48) is satisfied. Consequently, from the above considerations, we see that all the relations (46)–(48) can be satisfied; that is, an extremely large hierarchy of the eigenmasses of f and \tilde{f} can be realized in the case $\Upsilon = 1.5 \times 10^4$ for $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($20 \leq j \leq 89$), and $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 34$), and in the case $\Upsilon = 3.5 \times 10^2$ for $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($60 \leq j \leq 89$), and $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 34$). Furthermore, from Figs. 4 and 5 we see that in the above cases, both $\langle \lambda_+ \rangle / M^2$ and $\langle \lambda_- \rangle / M^2$ are positive, and the ratio $\langle \lambda_- / \lambda_+ \rangle$ is much smaller than unity. From the above results, we see that in the case that the difference between the values of ϵ_1 and ϵ_2 is much smaller than ϵ_1 and ϵ_2 themselves, i.e., $|\epsilon_1 - \epsilon_2| \ll O[\epsilon_i]$ [in other words, the two dilatons have an approximate $O(2)$ symmetric coupling], an extremely large hierarchy of the eigenmasses of f and \tilde{f} can be realized.

Finally, we note that in all the above cases in which an extremely large hierarchy of the eigenmasses of f and \tilde{f} can be realized, the heavier eigenmass, $m_+ \equiv \sqrt{\langle \lambda_+ \rangle}$, is comparable to or larger than the Hubble parameter in the inflationary stage, while the lighter eigenmass, $m_- \equiv \sqrt{\langle \lambda_- \rangle}$, is much smaller than the Hubble parameter in the inflationary stage. For example, in the case (v) in Table I, we have $m_+/H \approx \sqrt{\langle \lambda_+ \rangle} / (\Upsilon M) = 4.3$ and $m_-/H \approx \sqrt{\langle \lambda_- \rangle} / (\Upsilon M) = 9.2 \times 10^{-3}$, where we have used Eq. (41). Thus, it is expected that the scalar field with the lighter eigenmass, m_- , which is much smaller than the Hubble parameter in the inflationary stage, can be regarded as corresponding to the curvaton.

4.3. Evolution of dilatons after inflation

In this subsection, we discuss the evolution of dilatons after inflation, in particular, when and how the inflation ends and how reheating occurs in this model.

As shown in the cases considered in Table I of the previous subsection, from this point, we mainly consider the following case $\varphi_r = \sqrt{\varphi_1^2 + \varphi_2^2} \approx \gamma M$, where $\gamma(t) \ll 1$ during inflation. In this case, during the inflationary stage the dilatons φ_i evolve as $\varphi_r = Ct$ in the ansatz (15) around the origin of the potential $V[\varphi_i] = V_0 \cos(\varphi_r/M) + \Lambda$ in Eq. (3). Although φ_r exists around the origin of the potential in the inflationary stage, after sufficient inflationary

expansion, the amplitude of φ_r becomes large, and then φ_r approaches the minimum of the potential, i.e., $\varphi_r = M\pi$, in which the value of the potential is $V = -V_0 + \Lambda \approx 0$, where the last approximate equality follows from $V_0 \approx \Lambda \approx M^4$. As φ_r nears the potential minimum, the potential steepens, and the evolution of φ_r becomes more rapid; as φ_r approaches and then, inevitably, overshoots the minimum of its potential, it begins to oscillate about $\varphi_r = M\pi$ on a time scale short compared to the Hubble time. Hence, after the oscillation epoch, φ_r does not evolve as $\varphi_r = Ct$ but remains near the potential minimum, $\varphi_r = M\pi$. An enormous vacuum energy of the dilatons then exists in the form of spatially coherent oscillations of the dilatons, corresponding to a condensate of zero-momentum φ_i particles. Particle creations, or, equivalently, the decay of φ_i particles into other, lighter fields to which it couples, will damp these oscillations. Furthermore, as the decay products thermalize, the universe is reheated.

Finally, we estimate the number of e -folds during inflation. This quantity is defined as⁹⁾

$$N \equiv \int_{t_i}^{t_f} H dt \quad (52)$$

$$= \omega \ln \left(\frac{t_f}{t_i} \right), \quad (53)$$

where in deriving Eq. (53) we have used the relation $H = \omega/t$. Moreover, it follows from the considerations at the beginning of this subsection that $\varphi_r(t_i) \approx \gamma(t_i)M$ and $\varphi_r(t_f) \approx M\pi$. On the other hand, from these relations and $\varphi_r = Ct$, we obtain

$$t_i = \frac{\varphi_r(t_i)}{C} \approx \frac{\gamma(t_i)}{C}M, \quad (54)$$

$$t_f = \frac{\varphi_r(t_f)}{C} \approx \frac{\pi}{C}M. \quad (55)$$

Substituting Eqs. (54) and (55) into Eq. (53), we obtain

$$N \approx \omega \ln \left[\frac{\pi}{\gamma(t_i)} \right]. \quad (56)$$

For example, in the case (v) of Table I, it follows from the values $\omega = 4.1 \times 10^1$ and $\gamma(t_i) \simeq 3.5 \times 10^{-4}$ and Eq. (56) that $N = 3.7 \times 10^2 > 70$. Thus, the expansion of the universe during the epoch of power-law inflation realized in this model is large enough to solve both the horizon and flatness problems.

§5. Curvaton scenario

In the previous section, we showed that when two dilatons have an approximate $O(2)$ symmetric coupling, a scalar field corresponding to the curvaton exists in this theory. Considering this point, in this section, we investigate the case in which the curvaton scenario can be realized in this model. In particular, we consider a simple version of the curvaton scenario following the outline given in Refs. 14) and 15).

5.1. Curvature perturbation

To begin with, we consider linear metric perturbations. (For a review of the theory of cosmological perturbations, see Ref. 19).) The curvature perturbation on uniform-density hypersurfaces ζ is related to the gauge-dependent curvature perturbation ψ on a generic slicing and the energy density perturbation $\delta\rho$ in that gauge through the following gauge-invariant formula:²⁰⁾

$$\zeta = -\psi - H \frac{\delta\rho}{\dot{\rho}}. \quad (57)$$

Furthermore, the time dependence of ζ on super-horizon scales is given by²⁰⁾

$$\dot{\zeta} = -\frac{H}{\rho + P} \delta P_{\text{nad}}, \quad (58)$$

where P is the pressure and $\delta P_{\text{nad}} = \delta P - c_s^2 \delta\rho$ is a non-adiabatic pressure perturbation. Here, $c_s^2 \equiv \dot{P}/\dot{\rho}$ is the adiabatic speed of sound. In the usual inflation scenario, in which ζ is generated during inflation through the perturbation of a single-component inflaton field, it becomes practically time independent soon after horizon exit and remains so until the approach of horizon entry. By contrast, in the curvaton scenario, the curvature perturbation generated in the inflationary stage is negligible, and it is generated later through a non-adiabatic pressure perturbation associated with the curvaton perturbation. Finally, we note that from this point, we mainly consider the curvature perturbation on super-horizon scales.

5.2. Curvaton field

Next, we investigate the case in which the curvaton scenario can be realized in this model. As discussed in §3, power-law inflation in which the power-law exponent ω is much larger than unity can be realized in this model, and hence the Hubble parameter H can be slowly varying:

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2} = \frac{1}{\omega} \ll 1. \quad (59)$$

The requirement for power-law inflation in the curvaton scenario is that the relation (59) be satisfied; that is, the power-law exponent ω should be much larger than unity. Furthermore,

as noted in the previous section, in the framework of this theory, the scalar field with the lighter eigenmass, m_- , which is much smaller than the Hubble parameter in the inflationary stage, can be regarded as corresponding to the curvaton. From this point, we therefore regard this field as the curvaton σ with mass $m_\sigma \equiv m_-$.

Here we assume a spatially flat FRW spacetime with the metric (8), and we take the Lagrangian of the curvaton σ to be given by

$$\mathcal{L}_\sigma = \frac{1}{2}(\partial\sigma)^2 - U[\sigma], \quad (60)$$

where $U[\sigma]$ is the potential of the curvaton. Furthermore, we assume that the curvature perturbations is negligible during inflation. After the smallest cosmological scale leaves the horizon, the curvature perturbation remains negligible until after the curvaton starts to oscillate. Consequently, the curvaton evolves in unperturbed spacetime. In order to follow this evolution, we assume that the curvaton has no significant coupling to other fields, or, to be more precise, that the effect of any coupling can be integrated out to give a possibly time-dependent effective potential U .

We can now split the curvaton field as

$$\sigma(\mathbf{x}, t) = \sigma_0(t) + \delta\sigma(\mathbf{x}, t), \quad (61)$$

where σ_0 is the classical field, while $\delta\sigma(\mathbf{x}, t)$ represents the quantum fluctuations around σ_0 .

The unperturbed curvaton field satisfies

$$\ddot{\sigma} + 3H\dot{\sigma} + U_\sigma = 0, \quad (62)$$

where the subscript σ denotes partial differentiation with respect to σ .

Expanding the perturbation of the curvaton $\delta\sigma$ in Fourier modes,

$$\delta\sigma(\mathbf{x}, t) = \int \frac{d^3k}{(2\pi)^{3/2}} e^{i\mathbf{k}\cdot\mathbf{x}} \delta\sigma_{\mathbf{k}}(t), \quad (63)$$

where \mathbf{k} is the comoving wave number, and k denotes its amplitude, $|\mathbf{k}|$, we can write the equation for the perturbation as

$$\delta\ddot{\sigma}_{\mathbf{k}} + 3H\delta\dot{\sigma}_{\mathbf{k}} + \left(\frac{k^2}{a^2} + U_{\sigma\sigma}\right) \delta\sigma_{\mathbf{k}} = 0. \quad (64)$$

Here we assume that the curvaton potential is sufficiently flat during inflation, i.e.,

$$|U_{\sigma\sigma}| \ll H^2, \quad (65)$$

and that on cosmological scales each Fourier component is in the vacuum state well before horizon exit. Then the vacuum fluctuation causes a classical perturbation $\delta\sigma_{\mathbf{k}}$ well after horizon exit, which satisfies Eq. (64), with a negligible gradient term,

$$\delta\ddot{\sigma}_{\mathbf{k}} + 3H\delta\dot{\sigma}_{\mathbf{k}} + U_{\sigma\sigma}\delta\sigma_{\mathbf{k}} = 0. \quad (66)$$

In the limit that the relation (65) is satisfied, the power spectrum of the perturbation of the curvaton on super-horizon scales is given by

$$\mathcal{P}_{\delta\sigma}^{1/2} = \left(\frac{H_*}{2\pi}\right) \left(\frac{k}{a_*H_*}\right)^{(n_{\delta\sigma}-1)/2}, \quad (67)$$

where the star denotes the epoch of horizon exit, $k = a_*H_*$. The spectral index specifying the slight scale dependence is given by^{14),15)}

$$n_{\delta\sigma} - 1 \equiv \frac{d \ln \mathcal{P}_{\delta\sigma}}{d \ln k} = 2\eta_{\sigma\sigma} - 2\epsilon_H, \quad (68)$$

where $\eta_{\sigma\sigma} \equiv U_{\sigma\sigma}/(3H^2)$.

Here we assume that the curvaton potential $U[\sigma]$ is given by the following quadratic form

$$U[\sigma] = \frac{1}{2}m_\sigma^2(\sigma - v)^2, \quad (69)$$

which has a global minimum at $\sigma = v$. The reason we have shifted the global minimum of the potential from the origin to $\sigma = v$ is as follows. In this model, as seen in §2 and §3, the dilatons φ_i initially exist near the origin, along with the potential $V[\varphi_i]$ in Eq. (3), and then rapidly evolve in the inflationary stage. Hence, the initial field amplitude of the field variables f and \tilde{f} is also small, and then increases in proportion to t^2 in the inflationary stage. Because the curvaton considered here corresponds to the scalar field with the lighter eigenmass, m_- , it is conjectured that the initial amplitude of the curvaton is small and then increases. In this case, it follows from the discussion in §4.3 that $|U_{\sigma\sigma}|/H^2 \simeq m_\sigma^2/H^2 \ll 1$, and hence the relation (65) is satisfied. The spectral index of the primordial curvature perturbation is estimated as 0.99 ± 0.04 by using the first year WMAP data only,¹¹⁾ where the uncertainty represents the 68% confidence interval. For example, in the case (v) in Table I appearing in §4.3, $\epsilon_H = 1/\omega = 2.4 \times 10^{-2}$ and $\eta_{\sigma\sigma} \simeq m_\sigma^2/(3H^2) = 2.8 \times 10^{-5}$. Hence, it follows from Eq. (68) that $n_{\delta\sigma} = 1 + 2\eta_{\sigma\sigma} - 2\epsilon_H = 0.98$, which is consistent with the above observational constraint of WMAP.

The curvaton remains over damped until the Hubble parameter falls below the curvaton mass m_σ . The curvaton then starts to oscillate about its vacuum value, taken to be $\sigma = v$ in the curvaton potential $U[\sigma]$ in Eq. (69). Here we suppose that the oscillation starts during

the radiation-dominated era after the reheating epoch following inflation. It follows from Eq. (67) that the spectrum of the fractional field perturbation at this stage is given by

$$\mathcal{P}_{\delta\sigma/\bar{\sigma}}^{1/2} = \frac{1}{2\pi} \frac{H_*}{\bar{\sigma}_*}, \quad (70)$$

where we have defined $\bar{\sigma} \equiv |\sigma - v|$. Moreover, the energy density in the oscillating field is given by¹⁵⁾

$$\rho_\sigma(\mathbf{x}, t) = m_\sigma^2 \tilde{\sigma}^2(\mathbf{x}, t), \quad (71)$$

where $\tilde{\sigma}(\mathbf{x}, t)$ is the amplitude of the oscillation. The perturbation in ρ_σ depends on the curvaton field perturbation through both a linear and a quadratic term. Assuming that the linear term dominates, however, we obtain

$$\frac{\delta\rho_\sigma}{\rho_\sigma} \approx 2 \left(\frac{\delta\sigma}{\bar{\sigma}} \right)_*, \quad (72)$$

where we have used $\delta\sigma = \delta\bar{\sigma}$.

5.3. Generation of the curvature perturbation

In the previous subsection, we considered the epoch just after the Hubble parameter falls below the curvaton mass, and the curvaton oscillation starts. Once the curvaton starts to oscillate, the energy density becomes a mixture of that of the curvaton and radiation. At this point, the pressure perturbation corresponding to this mixture becomes non-adiabatic, and hence, according to Eq. (58), the generation of the curvature perturbation begins. It ends when the pressure perturbation again becomes adiabatic, which is during the epoch of curvaton matter domination, or the epoch of curvaton decay, whichever is earlier. Moreover, at the stage at which the curvaton oscillation starts, it is assumed that the dominant portion of the energy density comes from the radiation. The curvaton, however, is assumed to be fairly long-lived, while decaying before nucleosynthesis. As long as the decay rate of the curvaton Γ_σ is negligible, i.e., $\Gamma_\sigma \ll H$, we obtain $\rho_\sigma \propto a^{-3}(t)$ and $\rho_r \propto a^{-4}(t)$, leading to $\rho_\sigma/\rho_r \propto a(t)$, where ρ_σ is the energy density of the curvaton σ , and ρ_r is that of the radiation. It is this increase in the relative curvaton energy density which generates the curvature perturbation.

In order to analyze the generation of the curvature perturbation, it is convenient to separately consider the curvature perturbations ζ_r and ζ_σ on, respectively, slices of uniform radiation density and curvaton density. These curvature perturbations on super-horizon scales are separately conserved,²⁰⁾ as the radiation and the curvaton are perfectly non-interacting

fluids. The curvature perturbations are given by²⁰⁾

$$\zeta = -H \frac{\delta\rho}{\dot{\rho}} = -H \frac{\delta\rho_r + \delta\rho_\sigma}{\dot{\rho}_r + \dot{\rho}_\sigma}, \quad (73)$$

$$\zeta_r = -H \frac{\delta\rho_r}{\dot{\rho}_r} = \frac{1}{4} \frac{\delta\rho_r}{\rho_r}, \quad (74)$$

$$\zeta_\sigma = -H \frac{\delta\rho_\sigma}{\dot{\rho}_\sigma} = \frac{1}{3} \frac{\delta\rho_\sigma}{\rho_\sigma}, \quad (75)$$

where the density perturbations are defined on the flat slicing of spacetime ($\psi = 0$). It follows from Eqs. (73)–(75) that the total curvature perturbation ζ can be written as

$$\zeta = (1 - r)\zeta_r + r\zeta_\sigma, \quad (76)$$

where the relative contribution of the curvaton to the total curvature is given by

$$r = \frac{3\rho_\sigma}{4\rho_r + 3\rho_\sigma}. \quad (77)$$

As is usually done in the curvaton scenario, here we assume that the curvature perturbation in the radiation produced at the end of inflation is negligible, i.e., $\zeta_r \approx 0$. Furthermore, we assume that the curvaton decays instantaneously at the time that $H = H_{\text{dec}} = \Gamma_\sigma$, where H_{dec} is the Hubble parameter in the decay epoch. In this case, it follows from Eq. (76) that the curvature perturbation during the decay epoch is given by

$$\zeta \approx r_{\text{dec}}\zeta_\sigma \quad (78)$$

$$\approx \frac{2}{3} r_{\text{dec}} \left(\frac{\delta\sigma}{\bar{\sigma}} \right)_*, \quad (79)$$

where the approximate equality in Eq. (79) follows from Eqs. (72) and (75). Here, r_{dec} is the relative contribution of the curvaton to the total curvature during the decay epoch, i.e., when $H = H_{\text{dec}}$. Consequently, it follows from Eqs. (70) and (79) that the spectrum of the curvature perturbation in the curvaton scenario is given by

$$\mathcal{P}_\zeta^{1/2} \approx \frac{2}{3} r_{\text{dec}} \mathcal{P}_{\delta\sigma/\bar{\sigma}}^{1/2} \quad (80)$$

$$\approx \frac{r_{\text{dec}}}{3\pi} \frac{H_*}{\bar{\sigma}_*}. \quad (81)$$

The amplitude of the curvature perturbation required by the WMAP measurement¹¹⁾ is estimated as

$$\mathcal{P}_\zeta^{1/2}(\text{WMAP}) \simeq 5.2 \times 10^{-5}. \quad (82)$$

In the case that the curvaton dominates the energy density before it decays (that is, $r_{\text{dec}} = 1$), it follows from Eq. (80) that the result in Eq. (82) implies the following amplitude of the spectrum of the fractional field perturbation of the curvaton:

$$\mathcal{P}_{\delta\sigma/\bar{\sigma}}^{1/2} \simeq 7.7 \times 10^{-5}. \quad (83)$$

Thus, in the case that the curvaton dominates the energy density before it decays, if the spectrum amplitude of the fractional field perturbation of the curvaton is approximately 7.7×10^{-5} (i.e., it follows from Eq. (70) that $H_*/\bar{\sigma}_* \simeq 4.8 \times 10^{-4}$), then the curvature perturbation with the amplitude suggested by observations obtained from WMAP can be generated.

Finally, we note that the arguments given in this section employ analysis in the Einstein frame. It is not obvious whether the same property exists in the Jordan frame, which is the conformal frame considered in this model. It is shown in Appendix C, however, that the power spectrum of the curvature perturbation in the Einstein frame exactly coincides with that in the Jordan frame. Thus, the results of the above arguments obtained from analysis in the Einstein frame, are identical to those in the Jordan frame.

§6. Conclusion

In the present paper, we have shown that the curvaton scenario is realized in a theory with two dilatons coupled to the scalar curvature.³⁾ This theory has been considered in order to simultaneously solve the hierarchy problem between gravity and particle physics mass scales and the small cosmological constant or the dark energy problem. We have found that when both coupling constants between two dilatons and the scalar curvature are much smaller than unity, power-law inflation in which the power-law exponent is much larger than unity is realized in this model, and that when the difference between the values of the two coupling constants is much smaller than these values themselves [in other words, the two dilatons have an approximate $O(2)$ symmetric coupling], there exists a scalar field corresponding to the curvaton whose mass is much smaller than the Hubble parameter in the inflationary stage in this theory. This is realized without introducing any other scalar field that may play the role of the curvaton. Furthermore, we have investigated the simple version of the curvaton scenario in which the curvaton has a quadratic potential, and we demonstrated that a curvature perturbation with a sufficiently large amplitude and a nearly scale-invariant spectrum suggested by observations obtained from WMAP can be generated.

Finally, we note that in the present model, $\varphi_r = \sqrt{\varphi_1^2 + \varphi_2^2} \approx \gamma(t)M$ exists around the origin of the potential V in Eq. (3) in the inflationary stage; i.e., the case $\gamma(t) \ll 1$ during

inflation is considered. After sufficient inflationary expansion, the amplitude of φ_r becomes large, and then φ_r approaches the first minimum of the potential. After the oscillation epoch, φ_r remains around the first potential minimum. Finally, the dilatons decay into radiation through coupling to other lighter fields, and then the universe is reheated. By contrast, in the scenario proposed in Ref. 3), the following case is considered: In the inflationary stage, φ_r goes over many local maxima of the potential and then changes for many periods; i.e., there exist the relations $\gamma(t) \gtrsim 1$ and $\gamma(t_f) - \gamma(t_i) \gg 1$, where t_f and t_i are the time at the end and beginning of inflation, respectively. Thus, the situation for the evolution of φ_r during inflation considered in the present model is different from that in the scenario proposed in Ref. 3).

Acknowledgements

The work of K.B. was partially supported by the Monbukagakusho 21st century COE Program ‘‘Towards a New Basic Science; Depth and Synthesis’’ and was also supported by a Grant-in-Aid provided by the Japan Society for the Promotion of Science.

Appendix A

— Eigenmasses of f and \tilde{f} —

In this appendix, following the discussion in §4, we first derive the fluctuation equation in terms of $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\tilde{f} = \epsilon_1\varphi_1^2 - \epsilon_2\varphi_2^2$ in order to derive these mass values in the inflationary stage. Next, we investigate these eigenmasses in order to show that an extremely large hierarchy of the mass values of f and \tilde{f} in the inflationary stage can be realized.

A.1. Background quantities

To begin with, as the preliminary stage for investigating the fluctuation equation in terms of f and \tilde{f} , we consider the terms on the right-hand side of Eqs. (10) and (11). From $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\varphi_r^2 = \varphi_1^2 + \varphi_2^2$, we find

$$\varphi_1^2 = \frac{f - \epsilon_2\varphi_r^2}{\epsilon_1 - \epsilon_2}, \quad \varphi_2^2 = \frac{-f + \epsilon_1\varphi_r^2}{\epsilon_1 - \epsilon_2}. \quad (\text{A}\cdot 1)$$

Then, using Eq. (A.1), we obtain

$$\frac{\epsilon_1^2\varphi_1^2 + \epsilon_2^2\varphi_2^2}{f} = (\epsilon_1 + \epsilon_2) - \epsilon_1\epsilon_2\frac{\varphi_r^2}{f}, \quad (\text{A}\cdot 2)$$

$$\frac{\varphi_1\varphi_2}{f} = \pm \frac{1}{\epsilon_1 - \epsilon_2} \sqrt{\left(-1 + \epsilon_1\frac{\varphi_r^2}{f}\right) \left(1 - \epsilon_2\frac{\varphi_r^2}{f}\right)}, \quad (\text{A}\cdot 3)$$

$$\tilde{f} = \epsilon_1 \varphi_1^2 - \epsilon_2 \varphi_2^2 = \frac{f}{\epsilon_1 - \epsilon_2} \left[(\epsilon_1 + \epsilon_2) - 2\epsilon_1 \epsilon_2 \frac{\varphi_r^2}{f} \right]. \quad (\text{A}\cdot 4)$$

Furthermore, from $\dot{f} = 2(\epsilon_1 \varphi_1 \dot{\varphi}_1 + \epsilon_2 \varphi_2 \dot{\varphi}_2)$ and $L = \varphi_1 \dot{\varphi}_2 - \varphi_2 \dot{\varphi}_1$, we find

$$\dot{\varphi}_1 = \frac{\varphi_1 \dot{f} - 2\epsilon_2 \varphi_2 L}{2f}, \quad \dot{\varphi}_2 = \frac{\varphi_2 \dot{f} + 2\epsilon_1 \varphi_1 L}{2f}. \quad (\text{A}\cdot 5)$$

Using Eq. (A·5), we obtain

$$\epsilon_1 \dot{\varphi}_1^2 + \epsilon_2 \dot{\varphi}_2^2 = \frac{1}{4f} \left(\dot{f}^2 + 4\epsilon_1 \epsilon_2 L^2 \right), \quad (\text{A}\cdot 6)$$

$$\epsilon_1 \dot{\varphi}_1^2 - \epsilon_2 \dot{\varphi}_2^2 = \frac{1}{4f} \left[\frac{\tilde{f}}{f} \left(\dot{f}^2 - 4\epsilon_1 \epsilon_2 L^2 \right) - 8\epsilon_1 \epsilon_2 \dot{f} L \left(\frac{\varphi_1 \varphi_2}{f} \right) \right], \quad (\text{A}\cdot 7)$$

$$\begin{aligned} \dot{\varphi}_1^2 + \dot{\varphi}_2^2 &= \dot{\varphi}_r^2 + \frac{L^2}{\varphi_r^2} \\ &= \frac{1}{4f^2} \left[\varphi_r^2 \dot{f}^2 + 4(\epsilon_1 - \epsilon_2) \varphi_1 \varphi_2 \dot{f} L + 4(\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2) L^2 \right]. \end{aligned} \quad (\text{A}\cdot 8)$$

It follows from Eq. (A·8) that the quadric equation in terms of L reads

$$\begin{aligned} \left[\left(\frac{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}{f} \right) \left(\frac{\varphi_r^2}{f} \right) - 1 \right] L^2 + (\epsilon_1 - \epsilon_2) \left(\frac{\varphi_r^2}{f} \right) \left(\frac{\varphi_1 \varphi_2}{f} \right) \dot{f} L \\ + \frac{1}{4} \left(\frac{\varphi_r^2}{f} \right)^2 \dot{f}^2 - \varphi_r^2 \dot{\varphi}_r^2 = 0. \end{aligned} \quad (\text{A}\cdot 9)$$

The solution of Eq. (A·9) is given by

$$L = \left[2(\epsilon_1 - \epsilon_2) \left(\frac{\varphi_1 \varphi_2}{f} \right) \right]^{-1} \left[- \left(\frac{\varphi_r^2}{f} \right) \dot{f} \pm 2\varphi_r \dot{\varphi}_r \right], \quad (\text{A}\cdot 10)$$

where in deriving this expression we have used Eq. (A·3). Thus, from Eqs. (A·2)–(A·4), (A·6)–(A·8), and (A·10), we see that all the terms on the right-hand side of Eqs. (10) and (11) can be represented in terms of f and φ_r .

Next, we consider approximate expressions of A , B , and C . We again assume $\varphi_1^2 \approx \varphi_2^2$ and $\dot{\varphi}_1^2 \approx \dot{\varphi}_2^2$, as in the case that we approximately determined ω in terms of ϵ_i in §3. Applying these approximate relations to Eq. (20) and using Eqs. (24), (25) and (28), we find

$$A \approx \frac{32}{3} \left(\frac{\beta}{\alpha} \right)^2 A, \quad (\text{A}\cdot 11)$$

where in deriving this approximate expression we have employed the relation $\epsilon_i \ll 1$, that is, $\alpha/\beta \gg 1$, and hence $\omega \gg 1$. In the same way, from Eqs. (21) and (A.11) we find

$$C^2 + \frac{B^2}{C^2} \approx \frac{32\beta}{3\alpha} \Lambda. \quad (\text{A}\cdot 12)$$

Furthermore, applying one of the above approximate relations, $\varphi_1^2 \approx \varphi_2^2$, to φ_r^2/f and using the ansatz (15), we find

$$\frac{\varphi_r^2}{f} = \frac{C^2}{A} \approx \frac{2}{\alpha}. \quad (\text{A}\cdot 13)$$

Thus, from Eqs. (A.11) and (A.13), we find

$$C \approx \sqrt{\frac{64\beta^2}{3\alpha^3}} \Lambda, \quad (\text{A}\cdot 14)$$

where we have taken the positive value of C .

On the other hand, substituting the ansatz (15) into the solution of the quadric equation in terms of L , (A.10), we find

$$B = \pm 2C^2 \left[\left(-1 + \epsilon_1 \frac{C^2}{A} \right) \left(1 - \epsilon_2 \frac{C^2}{A} \right) \right]^{-1/2} \quad (\text{A}\cdot 15)$$

$$\approx \pm \frac{128\beta^2}{3\alpha^3} \sqrt{\frac{\alpha^2}{2\beta - \alpha^2}} \Lambda, \quad (\text{A}\cdot 16)$$

where the second approximate equality follows from Eqs. (A.11) and (A.13).

A.2. Linearized equations

In this subsection, we discuss the fluctuation equation in terms of $f = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ and $\tilde{f} = \epsilon_1\varphi_1^2 - \epsilon_2\varphi_2^2$ in order to derive these mass values in the inflationary stage. With the ansatz

$$\begin{aligned} f &= f_0 + \delta f, & |\delta f| &\ll |f_0|, \\ \tilde{f} &= \tilde{f}_0 + \delta \tilde{f}, & |\delta \tilde{f}| &\ll |\tilde{f}_0|, \end{aligned} \quad (\text{A}\cdot 17)$$

where f_0 and \tilde{f}_0 are the zeroth-order quantities, which satisfy Eqs. (10) and (11), respectively, we obtain the linearized equations

$$\begin{pmatrix} \delta \ddot{f} + 3H\delta \dot{f} \\ \delta \ddot{\tilde{f}} + 3H\delta \dot{\tilde{f}} \end{pmatrix} = -\mathcal{M}^2 \begin{pmatrix} \delta f \\ \delta \tilde{f} \end{pmatrix}, \quad (\text{A}\cdot 18)$$

where we have retained the terms up to first order in δf and $\delta \tilde{f}$. Moreover, we have omitted the quantities in terms of $\dot{\delta f}$ and $\dot{\delta \tilde{f}}$ derived from the right-hand side of Eqs. (10) and (11). This is because, as we see from the ansatz (15) and Eq. (A.4), both f and \tilde{f} are proportional to t^2 , and thus, in the large t limit we have $|\dot{\delta f}| \ll \delta f$ and $|\dot{\delta \tilde{f}}| \ll \delta \tilde{f}$. Incidentally, the reason we have chosen the field variable \tilde{f} as the partner of the coupling f is that \tilde{f} is a simple quantity satisfying the conditions that it have the same dimensions as f and that it be expressed as a linear combination of φ_1^2 and φ_2^2 .

In deriving the linearized equations (A.18), we have taken into account the following point: From Eqs. (A.2)–(A.4), (A.6)–(A.8), and (A.10), we see that all the terms on the right-hand side of Eqs. (10) and (11) can be represented in terms of f and φ_r , as noted in the previous subsection. Furthermore, from Eq. (A.4), we find $\varphi_r^2 = [(\epsilon_1 + \epsilon_2)f - (\epsilon_1 - \epsilon_2)\tilde{f}] / (2\epsilon_1\epsilon_2)$. Hence, the fluctuation $\delta\varphi_r$ induced by δf and $\delta \tilde{f}$ can be represented as follows:

$$\delta\varphi_r = \frac{1}{4\epsilon_1\epsilon_2\varphi_r} \left[(\epsilon_1 + \epsilon_2)\delta f - (\epsilon_1 - \epsilon_2)\delta \tilde{f} \right]. \quad (\text{A.19})$$

Thus, all the fluctuations derived from the right-hand side of Eqs. (10) and (11) can be represented in terms of δf and $\delta \tilde{f}$ by using Eqs. (A.2)–(A.4), (A.6)–(A.8), (A.10) and (A.19).

From the above, we find that \mathcal{M}^2 is given by

$$\mathcal{M}^2 \equiv \begin{pmatrix} \mathcal{M}^2_{11} & \mathcal{M}^2_{12} \\ \mathcal{M}^2_{21} & \mathcal{M}^2_{22} \end{pmatrix}, \quad (\text{A.20})$$

with

$$\begin{aligned} \mathcal{M}^2_{11} &= -4F^{-2}(2F-1)Q\frac{1}{f}V \\ &\quad - 2F^{-1} \left[-1 + \frac{\epsilon_1 + \epsilon_2}{\epsilon_1\epsilon_2} \left(\frac{1}{4}\frac{f}{\varphi_r^2} + U \right) - \frac{6Q}{F} \right] \frac{1}{\varphi_r} V' \\ &\quad + F^{-1} \frac{\epsilon_1 + \epsilon_2}{2\epsilon_1\epsilon_2} \frac{f}{\varphi_r^2} V'' \\ &\quad - F^{-1} \left\{ \frac{Q}{F} \frac{\dot{\varphi}_1^2 + \dot{\varphi}_2^2}{f} + \frac{\epsilon_1 + \epsilon_2}{\epsilon_1\epsilon_2} U \frac{L^2}{\varphi_r^4} - \frac{1 + 6(\epsilon_1 + \epsilon_2)}{2F} J \right. \\ &\quad \left. + \frac{Q}{\epsilon_1\epsilon_2} \frac{I}{\varphi_r^2} \left[Q^2 \frac{I}{f} - Q \frac{f}{f} \pm (\epsilon_1 + \epsilon_2) \frac{\dot{\varphi}_r}{\varphi_r} \right] \right\} \\ &\approx -\frac{3\alpha(2\beta - \alpha^2)}{8\beta^2} \frac{V}{\Lambda t^2} + \frac{3\sqrt{3}\alpha^{3/2}(\alpha^2 - 4\beta)}{16\beta(\alpha^2 - \beta)} \frac{V'}{\sqrt{\Lambda t}} + \frac{\alpha^2}{2(\alpha^2 - \beta)} V'' \end{aligned} \quad (\text{A.21})$$

$$+ \frac{C_{11}}{\beta(\alpha^2 - \beta)(2\beta - \alpha^2)} \frac{1}{t^2}, \quad (\text{A}\cdot 22)$$

$$\begin{aligned} \mathcal{M}_{12}^2 = & -4F^{-2}(\epsilon_1 - \epsilon_2) \frac{1}{f} V \\ & - 2F^{-1}(\epsilon_1 - \epsilon_2) \left[-\frac{1}{\epsilon_1 \epsilon_2} \left(\frac{1}{4} \frac{f}{\varphi_r^2} + U \right) + \frac{6}{F} \right] \frac{1}{\varphi_r} V' \\ & - F^{-1} \frac{\epsilon_1 - \epsilon_2}{2\epsilon_1 \epsilon_2} \frac{f}{\varphi_r^2} V'' \\ & + F^{-1}(\epsilon_1 - \epsilon_2) \left[\frac{1}{F} \frac{\dot{\varphi}_1^2 + \dot{\varphi}_2^2}{f} + \frac{U}{\epsilon_1 \epsilon_2} \frac{L^2}{\varphi_r^4} + \frac{3}{F} J \right. \\ & \quad \left. + \frac{Q}{\epsilon_1 \epsilon_2} \frac{I}{\varphi_r^2} \left(Q \frac{I}{f} - \frac{\dot{f}}{f} \pm \frac{\dot{\varphi}_r}{\varphi_r} \right) \right] \end{aligned} \quad (\text{A}\cdot 23)$$

$$\begin{aligned} \approx & \pm \sqrt{2\beta - \alpha^2} \left\{ -\frac{3\alpha^2}{8\beta^2} \frac{V}{\Lambda t^2} + \frac{\sqrt{3}\alpha^{3/2}(\alpha^2 + 8\beta)}{16\alpha\beta(\alpha^2 - \beta)} \frac{V'}{\sqrt{\Lambda}t} - \frac{\alpha}{2(\alpha^2 - \beta)} V'' \right. \\ & \left. + \frac{C_{12}}{\beta(\alpha^2 - \beta)(2\beta - \alpha^2)} \frac{1}{t^2} \right\}, \end{aligned} \quad (\text{A}\cdot 24)$$

$$\begin{aligned} \mathcal{M}_{21}^2 = & 4F^{-2} \frac{Q}{\epsilon_1 - \epsilon_2} [(\epsilon_1 + \epsilon_2) + 24\epsilon_1 \epsilon_2] \frac{1}{f} V \\ & - \frac{F^{-1}}{\epsilon_1 - \epsilon_2} \left\{ Q \left(-\frac{12Q}{F} + \frac{\epsilon_1 + \epsilon_2}{2\epsilon_1 \epsilon_2} \frac{f}{\varphi_r^2} \right) + 48\epsilon_1 \epsilon_2 \left[\left(1 + \frac{6Q}{F} \right) P - Q \frac{\varphi_r^2}{f} \right] \right. \\ & \quad \left. + 12(\epsilon_1 + \epsilon_2) \left(2Q - P \frac{f}{\varphi_r^2} \right) + \frac{(\epsilon_1 + \epsilon_2)(\epsilon_1 - \epsilon_2)}{\epsilon_1 \epsilon_2} W \right\} \frac{1}{\varphi_r} V' \\ & - F^{-1} \frac{\epsilon_1 + \epsilon_2}{\epsilon_1 - \epsilon_2} \left(-\frac{Q}{2\epsilon_1 \epsilon_2} + 12P \right) \frac{f}{\varphi_r^2} V'' \\ & + \frac{1}{\epsilon_1 - \epsilon_2} \frac{1}{f^2} \left[Q \left(f^2 - 4\epsilon_1 \epsilon_2 L^2 \right) + (4\epsilon_1 \epsilon_2 P - Q^2) f I \right] \\ & - F^{-1} \left\{ \frac{Q}{(\epsilon_1 - \epsilon_2)F} [(\epsilon_1 + \epsilon_2) + 24\epsilon_1 \epsilon_2] \left(\frac{\dot{\varphi}_1^2 + \dot{\varphi}_2^2}{f} + 3J \right) \right. \\ & \quad \left. + W \left(3J + \frac{\epsilon_1 + \epsilon_2}{2\epsilon_1 \epsilon_2} \frac{L^2}{\varphi_r^4} \right) \right\} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\epsilon_1 - \epsilon_2} \left\{ \frac{\dot{f}}{f} - \left[Q + \frac{\epsilon_1 - \epsilon_2}{2\epsilon_1\epsilon_2} \frac{W}{F} \left(\frac{f}{\varphi_r^2} + 12\epsilon_1\epsilon_2 \right) \right] \frac{I}{f} \right\} \\
& \quad \times \left[Q^2 \frac{I}{f} - Q \frac{\dot{f}}{f} \pm (\epsilon_1 + \epsilon_2) \frac{\dot{\varphi}_r}{\varphi_r} \right] \tag{A.25}
\end{aligned}$$

$$\begin{aligned}
& \approx \pm \sqrt{2\beta - \alpha^2} \left\{ \frac{3\alpha^2}{8\beta^2} \frac{V}{\Lambda t^2} - \frac{9\sqrt{3}\alpha^{5/2}}{16\beta(\alpha^2 - \beta)} \frac{V'}{\sqrt{\Lambda}t} + \frac{\alpha}{2(\alpha^2 - \beta)} V'' \right\} \\
& \quad + \frac{C_{21}}{(\pm\sqrt{2\beta - \alpha^2})\beta(\alpha^2 - \beta)} \frac{1}{t^2}, \tag{A.26}
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{22}^2 &= -4F^{-1} \left[(\epsilon_1 + \epsilon_2) - 6(\epsilon_1 - \epsilon_2) \frac{W}{F} \right] \frac{1}{f} V \\
& + F^{-1} \left\{ 2 \left[1 - \frac{6Q}{F} + 6 \left(2Q - P \frac{f}{\varphi_r^2} \right) + 144\epsilon_1\epsilon_2 \frac{P}{F} \right] \right. \\
& \quad \left. + \frac{1}{2\epsilon_1\epsilon_2} \left[2(\epsilon_1 - \epsilon_2)W + Q \frac{f}{\varphi_r^2} \right] \right\} \frac{1}{\varphi_r} V' \\
& - F^{-1} \left(\frac{Q}{2\epsilon_1\epsilon_2} - 12P \right) \frac{f}{\varphi_r^2} V'' \\
& + \frac{1}{2f^2} \left(-\dot{f}^2 + 4\epsilon_1\epsilon_2 L^2 + 2Q\dot{f}I \right) \\
& + F^{-1} \left\{ \left[(\epsilon_1 + \epsilon_2) - 6(\epsilon_1 - \epsilon_2) \frac{W}{F} \right] \left(\frac{\dot{\varphi}_1^2 + \dot{\varphi}_2^2}{f} + 3J \right) + \frac{\epsilon_1 - \epsilon_2}{2\epsilon_1\epsilon_2} W \frac{L^2}{\varphi_r^4} \right\} \\
& - \left\{ \frac{\dot{f}}{f} - \left[Q + \frac{\epsilon_1 - \epsilon_2}{2\epsilon_1\epsilon_2} \frac{W}{F} \left(\frac{f}{\varphi_r^2} + 12\epsilon_1\epsilon_2 \right) \right] \frac{I}{f} \right\} \left(Q \frac{I}{f} - \frac{\dot{f}}{f} \pm \frac{\dot{\varphi}_r}{\varphi_r} \right) \tag{A.27}
\end{aligned}$$

$$\begin{aligned}
& \approx -\frac{3\alpha^3}{8\beta^2} \frac{V}{\Lambda t^2} + \frac{\sqrt{3}\alpha^{3/2}(14\beta - 5\alpha^2)}{16\beta(\alpha^2 - \beta)} \frac{V'}{\sqrt{\Lambda}t} - \frac{2\beta - \alpha^2}{2(\alpha^2 - \beta)} V'' \\
& \quad + \frac{C_{22}}{\beta(\alpha^2 - \beta)(2\beta - \alpha^2)} \frac{1}{t^2}, \tag{A.28}
\end{aligned}$$

where

$$F = 1 + 12 \left[(\epsilon_1 + \epsilon_2) - \epsilon_1\epsilon_2 \frac{\varphi_r^2}{f} \right] = 1 + 12U \approx 1 + 12\frac{\beta}{\alpha}, \tag{A.29}$$

$$Q = (\epsilon_1 + \epsilon_2) - 2\epsilon_1\epsilon_2 \frac{\varphi_r^2}{f} \approx \frac{2\beta - \alpha^2}{\alpha}, \quad (\text{A}\cdot 30)$$

$$U = (\epsilon_1 + \epsilon_2) - \epsilon_1\epsilon_2 \frac{\varphi_r^2}{f} \approx \frac{\beta}{\alpha}, \quad (\text{A}\cdot 31)$$

$$J = \frac{1}{f^2} \left(f^2 + 4\epsilon_1\epsilon_2 L^2 \right) \approx 4 \left(\frac{7\alpha^2 - 6\beta}{2\beta - \alpha^2} \right) \frac{1}{t^2}, \quad (\text{A}\cdot 32)$$

$$I = -\frac{1}{2P} \left[-\left(\frac{\varphi_r^2}{f} \right) f \pm 2\varphi_r \dot{\varphi}_r \right] \approx \frac{128}{3} \frac{\beta^2}{\alpha(2\beta - \alpha^2)} \Lambda t, \quad (\text{A}\cdot 33)$$

$$P = -\left(1 - \epsilon_1 \frac{\varphi_r^2}{f} \right) \left(1 - \epsilon_2 \frac{\varphi_r^2}{f} \right) \approx \frac{2\beta - \alpha^2}{\alpha^2}, \quad (\text{A}\cdot 34)$$

$$W = (\epsilon_1 - \epsilon_2) + \frac{\epsilon_1 + \epsilon_2}{\epsilon_1 - \epsilon_2} Q \approx \pm 2\sqrt{2\beta - \alpha^2}, \quad (\text{A}\cdot 35)$$

$$C_{11} = 7\alpha^4\beta - 16\beta^3 - 2\alpha^2\beta^2 - \alpha^6 - \frac{96\beta(\alpha^2 - \beta)^2(2\beta - \alpha^2)}{\alpha}, \quad (\text{A}\cdot 36)$$

$$C_{12} = 14\alpha\beta^2 - \alpha^3\beta - \alpha^5 + \frac{48\beta(\alpha^2 - \beta)[2\alpha(\alpha^2 - \beta) - 3\beta(7\alpha^2 - 6\beta)]}{\alpha}, \quad (\text{A}\cdot 37)$$

$$C_{21} = \alpha^5 - 8\alpha\beta^2 - 5\alpha^3\beta + 12\beta(-7\alpha^4 + \alpha^2\beta + 10\beta^2), \quad (\text{A}\cdot 38)$$

$$C_{22} = \alpha \left[\alpha(14\beta^2 - \alpha^4 - \alpha^2\beta) + 12\beta(7\alpha^4 - 9\alpha^2\beta - 2\beta^2) \right]. \quad (\text{A}\cdot 39)$$

Here, the expression for F given in (A·29) follows from Eqs. (12) and (A·2).

In deriving the last approximate equality in each of the equations (A·22), (A·24), (A·26), and (A·28)–(A·35), we have used Eqs. (24), (25), (A·11), (A·12), (A·14), and (A·16). Moreover, in the last approximate equality in each of the equations (A·22), (A·24), (A·26), and (A·28), we have retained all the leading-order terms and some of the sub-leading order terms in ϵ_i and omitted the rest of the sub-leading order terms, which are explicitly unimportant. Furthermore, in deriving the last approximate equality in (A·33), we have taken the negative sign, so that I has a finite value. If we take the positive sign, under the approximation in (A·13), we have $I \approx 0$. Since it follows from Eqs. (A·10), (A·33), and (A·34) that the angular momentum is given by $L = -(\pm P^{1/2}) I$, and because here we consider the case $L \neq 0$, we have taken the negative sign so that the angular momentum can have a finite value. Hence, in deriving the approximate expressions (A·22), (A·24), (A·26) and (A·28), we have taken the negative sign in the last term on the right-hand side of each of Eqs. (A·21), (A·23), (A·25) and (A·27), because the sign of these terms is determined by the angular momentum

L , and L is proportional to I , as shown above.

We finally emphasize that all the terms in each of the components of \mathcal{M}^2 can be expressed in terms of the background quantities $f = At^2$, $L = Bt$ and $\varphi_r = Ct$ in the inflationary stage, and hence can be approximately represented in terms of α and β by using the expressions (A·11), (A·12), (A·14) and (A·16), as shown in expressions (A·22), (A·24), (A·26) and (A·28).

A.3. Mass diagonalization

Next, in order to show that an extremely large hierarchy of the mass values of f and \tilde{f} in the inflationary stage can be realized, we investigate these eigenmasses by solving the characteristic equation of \mathcal{M}^2 ,

$$\det(\mathcal{M}^2 - \lambda E) = 0 \iff \lambda^2 - \text{Tr} \mathcal{M}^2 \lambda + \det \mathcal{M}^2 = 0, \quad (\text{A}\cdot 40)$$

where λ is the eigenvalue of \mathcal{M}^2 and E is the unit matrix. The solutions of Eq. (A·40) are equivalent to the diagonal components of the diagonalized form of \mathcal{M}^2 , namely, the square of the eigenmasses of f and \tilde{f} . The solution of Eq. (A·40) is given by

$$\lambda_{\pm} = \frac{\text{Tr} \mathcal{M}^2}{2} \left[1 \pm \sqrt{1 - \frac{4 \det \mathcal{M}^2}{(\text{Tr} \mathcal{M}^2)^2}} \right]. \quad (\text{A}\cdot 41)$$

In this subsection, in order to evaluate the solutions (A·41), we derive approximate expressions of $\text{Tr} \mathcal{M}^2$ and $\det \mathcal{M}^2$.

From Eq. (3) and the expressions (A·22), (A·24), (A·26), and (A·28), we can obtain approximate expressions for $\text{Tr} \mathcal{M}^2$ and $\det \mathcal{M}^2$. For convenience in describing these expressions, we define $\text{Tr} \mathcal{M}^2 \equiv (\text{Tr} \mathcal{M}^2)^{(0)} + (\text{Tr} \mathcal{M}^2)^{(1)}$ and $\det \mathcal{M}^2 \equiv (\det \mathcal{M}^2)^{(0)} + (\det \mathcal{M}^2)^{(1)} + (\det \mathcal{M}^2)^{(2)}$, where $(\text{Tr} \mathcal{M}^2)^{(0)}$ and $(\det \mathcal{M}^2)^{(0)}$ are the parts independent of the sine functions, while $(\text{Tr} \mathcal{M}^2)^{(1)}$ and $(\det \mathcal{M}^2)^{(1)}$ are the parts proportional to $\sin[\varphi_r/M + \arctan(X_1/X_2)]$ and $\sin[\varphi_r/M + \arctan(Y_1/Y_2)]$, respectively, and $(\det \mathcal{M}^2)^{(2)}$ is the part proportional to $\sin[2\varphi_r/M + \arctan(Z_1/Z_2)]$. [X_1 , X_2 , Y_1 , Y_2 , Z_1 and Z_2 are given in Eqs. (A·49)–(A·54).] We then obtain the approximate expressions for $\text{Tr} \mathcal{M}^2$ and $\det \mathcal{M}^2$ as

$$\begin{aligned} \text{Tr} \mathcal{M}^2 &= \mathcal{M}_{11}^2 + \mathcal{M}_{22}^2 \\ &\equiv (\text{Tr} \mathcal{M}^2)^{(0)} + (\text{Tr} \mathcal{M}^2)^{(1)}, \end{aligned} \quad (\text{A}\cdot 42)$$

$$(\text{Tr} \mathcal{M}^2)^{(0)} \approx \left[-\frac{3\alpha}{4\beta} + \frac{C_{11} + C_{22}}{\beta(\alpha^2 - \beta)(2\beta - \alpha^2)} \right] \frac{1}{t^2}, \quad (\text{A}\cdot 43)$$

$$(\text{Tr} \mathcal{M}^2)^{(1)} \approx \sqrt{X_1^2 + X_2^2} \sin\left(\frac{\varphi_r}{M} + \arctan \frac{X_1}{X_2}\right) \frac{1}{t^2}, \quad (\text{A}\cdot 44)$$

and

$$\begin{aligned}\det \mathcal{M}^2 &= \mathcal{M}_{11}^2 \mathcal{M}_{22}^2 - \mathcal{M}_{12}^2 \mathcal{M}_{21}^2 \\ &\equiv (\det \mathcal{M}^2)^{(0)} + (\det \mathcal{M}^2)^{(1)} + (\det \mathcal{M}^2)^{(2)},\end{aligned}\tag{A.45}$$

$$\begin{aligned}(\det \mathcal{M}^2)^{(0)} &\approx \left\{ \frac{3\alpha}{8\beta^3(\alpha^2 - \beta)} \left[-\frac{\alpha^2}{2\beta - \alpha^2} C_{11} - C_{22} + \alpha(C_{21} - C_{12}) \right] \right. \\ &\quad + \frac{C_{11}C_{22} - (2\beta - \alpha^2)C_{12}C_{21}}{\beta^2(\alpha^2 - \beta)^2(2\beta - \alpha^2)^2} \\ &\quad \left. + \frac{3\alpha^3}{64\beta^4} \left[9\alpha(2\beta - \alpha^2) + \frac{\alpha^6 - 11\alpha^2\beta^2 + 2\alpha^4\beta + 8\beta^3}{64(\alpha^2 - \beta)^2} \frac{1}{\mathcal{R}^2} \right] \right\} \frac{1}{t^4},\end{aligned}\tag{A.46}$$

$$(\det \mathcal{M}^2)^{(1)} \approx \sqrt{Y_1^2 + Y_2^2} \sin \left(\frac{\varphi_r}{M} + \arctan \frac{Y_1}{Y_2} \right) \frac{1}{t^4},\tag{A.47}$$

$$(\det \mathcal{M}^2)^{(2)} \approx \sqrt{Z_1^2 + Z_2^2} \sin \left(2\frac{\varphi_r}{M} + \arctan \frac{Z_1}{Z_2} \right) \frac{1}{t^4},\tag{A.48}$$

where

$$X_1 = -\frac{3\alpha}{4\beta} \left(1 + \frac{\alpha}{48\beta} \frac{1}{\mathcal{R}^2} \right),\tag{A.49}$$

$$X_2 = \frac{\sqrt{3}\alpha^{5/2}}{64\beta^2} \frac{1}{\mathcal{R}},\tag{A.50}$$

$$\begin{aligned}Y_1 &= \frac{3\alpha}{8\beta^3(\alpha^2 - \beta)} \left[-\frac{\alpha^2}{2\beta - \alpha^2} C_{11} - C_{22} + \alpha(C_{21} - C_{12}) \right] + \frac{9\alpha^4(2\beta - \alpha^2)}{16\beta^4} \\ &\quad - \frac{\alpha^2}{128\beta^3(\alpha^2 - \beta)} \left\{ \frac{1}{\alpha^2 - \beta} \left[-C_{11} + \frac{\alpha^2}{2\beta - \alpha^2} C_{22} + \alpha(C_{21} - C_{12}) \right] \right. \\ &\quad \left. + \frac{3\alpha(2\beta^2 - \alpha^4)}{4\beta} \right\} \frac{1}{\mathcal{R}^2},\end{aligned}\tag{A.51}$$

$$\begin{aligned}Y_2 &= -\frac{\sqrt{3}\alpha^{5/2}}{128\beta^3(\alpha^2 - \beta)} \left\{ \frac{1}{\alpha^2 - \beta} \left[\frac{(14\beta - 5\alpha^2)C_{11} + 3(\alpha^2 - 4\beta)C_{22}}{2\beta - \alpha^2} + 9\alpha C_{12} \right. \right. \\ &\quad \left. \left. - \frac{(\alpha^2 + 8\beta)C_{21}}{\alpha} \right] + \frac{3(12\alpha^3\beta - 22\alpha\beta^2 + \alpha^5)}{4\beta} \right\} \frac{1}{\mathcal{R}},\end{aligned}\tag{A.52}$$

$$Z_1 = \frac{3\alpha^3}{64\beta^4} \left[3\alpha(2\beta - \alpha^2) + \frac{7\alpha^6 - 5\alpha^2\beta^2 - 10\alpha^4\beta + 8\beta^3}{64(\alpha^2 - \beta)^2} \frac{1}{\mathcal{R}^2} \right], \quad (\text{A}\cdot 53)$$

$$Z_2 = \frac{\sqrt{3}\alpha^{5/2}}{1024\beta^4} \left[-\frac{3(12\alpha^3\beta - 22\alpha\beta^2 + \alpha^5)}{\alpha^2 - \beta} + \frac{\alpha^2}{4} \frac{1}{\mathcal{R}^2} \right] \frac{1}{\mathcal{R}}. \quad (\text{A}\cdot 54)$$

Here we have considered the case $V_0 \approx \Lambda \approx M^4$. Moreover, from Eq. (41), we have used the relation

$$M \approx \frac{H}{\mathcal{R}} = \frac{\omega}{\mathcal{R}} \frac{1}{t} \approx \frac{\alpha}{8\beta} \frac{1}{\mathcal{R}} \frac{1}{t}, \quad (\text{A}\cdot 55)$$

where the last approximate equality follows from Eq. (28).

Appendix B

— Power-Law Inflation in the Einstein Frame —

In this appendix, we show that power-law inflation can be realized in this model not only in the Jordan frame, as shown in §3, but also in the Einstein frame.

B.1. Action

To begin with, we make the following conformal transformation of the action in Eq. (1):

$$g_{\mu\nu} \rightarrow \hat{g}_{\mu\nu} = \Omega^2 g_{\mu\nu}, \quad (\text{B}\cdot 1)$$

with

$$\Omega = \sqrt{2f}. \quad (\text{B}\cdot 2)$$

Here, the hat denotes quantities in the Einstein frame. Moreover, $f(\varphi_i) = \epsilon_1\varphi_1^2 + \epsilon_2\varphi_2^2$ in Eq. (2) is the coupling between the dilaton and the scalar curvature. The action in the Einstein frame is then given by^(6),21)

$$S_E = \int d^4\hat{x} \sqrt{-\hat{g}} \left\{ -\frac{1}{2}\hat{R} + \frac{1}{2} \sum_i \left[\frac{3}{2} \left(\frac{f_{\varphi_i}}{f} \right)^2 + \frac{1}{2f} \right] \hat{g}^{\mu\nu} \partial_\mu \varphi_i \partial_\nu \varphi_i \right. \\ \left. + 6\epsilon_1\epsilon_2 \frac{\varphi_1\varphi_2}{f^2} \hat{g}^{\mu\nu} \partial_\mu \varphi_1 \partial_\nu \varphi_2 - \hat{V}[\varphi_i] + \hat{\mathcal{L}}_m \right\}, \quad (\text{B}\cdot 3)$$

with

$$\hat{V}[\varphi_i] = \frac{V[\varphi_i]}{4f^2}. \quad (\text{B}\cdot 4)$$

Here, the subscript φ_i denotes partial differentiation with respect to φ_i . Moreover, in deriving the expression for the third term on the right-hand side of Eq. (B-3), namely, the cross term, we have used the symmetry between μ and ν . Because there exists this cross term in the action in Eq. (B-3), we introduce new scalar fields defined as

$$\partial_\mu \tilde{\varphi}_1 \equiv \frac{1}{\sqrt{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}} (\epsilon_1 \varphi_1 \partial_\mu \varphi_1 + \epsilon_2 \varphi_2 \partial_\mu \varphi_2), \quad (\text{B}\cdot 5)$$

$$\partial_\mu \tilde{\varphi}_2 \equiv \frac{1}{\sqrt{\epsilon_1^2 \varphi_1^2 + \epsilon_2^2 \varphi_2^2}} (-\epsilon_2 \varphi_2 \partial_\mu \varphi_1 + \epsilon_1 \varphi_1 \partial_\mu \varphi_2), \quad (\text{B}\cdot 6)$$

so that the cross term can be removed. Then the action in Eq. (B-3) can be rewritten in the form

$$S_E = \int d^4 \hat{x} \sqrt{-\hat{g}} \left\{ -\frac{1}{2} \hat{R} + \frac{1}{2} \sum_i \left[\frac{3}{2} \left(\frac{f_{\varphi_i}}{f} \right)^2 + \frac{1}{2f} + (-1)^{1+i} 6 \left(\frac{\epsilon_2 \varphi_2}{f} \right)^2 \right] \hat{g}^{\mu\nu} \partial_\mu \tilde{\varphi}_i \partial_\nu \tilde{\varphi}_i - \hat{V}[\varphi_i] + \hat{\mathcal{L}}_m \right\}, \quad (\text{B}\cdot 7)$$

where i in the third term in the brackets on the right-hand side of Eq. (B-7) corresponds to the subscript of φ_i ($i = 1, 2$).

Furthermore, we introduce new scalar fields defined as

$$\frac{d\hat{\varphi}_i}{d\tilde{\varphi}_i} \equiv \sqrt{D_E(\varphi_i)}, \quad (\text{B}\cdot 8)$$

with

$$D_E(\varphi_i) = \frac{3}{2} \left(\frac{f_{\varphi_i}}{f} \right)^2 + \frac{1}{2f} + (-1)^{1+i} 6 \left(\frac{\epsilon_2 \varphi_2}{f} \right)^2, \quad (> 0) \quad (\text{B}\cdot 9)$$

so that the action in Eq. (B-7) can be rewritten in the following canonical form:

$$S_E = \int d^4 \hat{x} \sqrt{-\hat{g}} \left[-\frac{1}{2} \hat{R} + \frac{1}{2} \hat{g}^{\mu\nu} \partial_\mu \hat{\varphi}_i \partial_\nu \hat{\varphi}_i - \hat{V}[\varphi_i] + \hat{\mathcal{L}}_m \right]. \quad (\text{B}\cdot 10)$$

It is conjectured that if the potential \hat{V} is written in terms of $\hat{\varphi}_1$ and $\hat{\varphi}_2$, the form is complicated.

B.2. Field equations

Field equations can be derived by taking variations of the action in Eq. (B-10) with respect to the metric $\hat{g}_{\mu\nu}$ and the dilatons $\hat{\varphi}_i$ as follows:

$$\hat{R}_{\mu\nu} - \frac{1}{2} \hat{g}_{\mu\nu} \hat{R} = \hat{T}_{\mu\nu}^{(m)} + \hat{T}_{\mu\nu}^{(\varphi_i)}, \quad (\text{B}\cdot 11)$$

with

$$\hat{T}_{\mu\nu}^{(\varphi_i)} = \partial_\mu \hat{\varphi}_i \partial_\nu \hat{\varphi}_i - \hat{g}_{\mu\nu} \left[\frac{1}{2} \hat{g}^{\mu\nu} \partial_\mu \hat{\varphi}_i \partial_\nu \hat{\varphi}_i - \hat{V}[\varphi_i] \right], \quad (\text{B}\cdot 12)$$

and

$$\hat{\square} \hat{\varphi}_i = -\frac{\partial \hat{V}}{\partial \hat{\varphi}_i}, \quad (\text{B}\cdot 13)$$

where $\hat{\square} \equiv \hat{g}^{\mu\nu} \hat{\nabla}_\mu \hat{\nabla}_\nu$.

We now assume spatially flat Friedmann-Robertson-Walker (FRW) spacetime with the metric

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = dt^2 - a^2(t) d\mathbf{x}^2, \quad (\text{B}\cdot 14)$$

to which we apply the conformal transformation in Eq. (B.1) as follows:

$$d\hat{s}^2 = \hat{g}_{\mu\nu} dx^\mu dx^\nu = \Omega^2 ds^2 = \Omega^2 dt^2 - \Omega^2 a^2(t) d\mathbf{x}^2. \quad (\text{B}\cdot 15)$$

We write the above line element in the form

$$d\hat{s}^2 = d\hat{t}^2 - \hat{a}^2(\hat{t}) d\mathbf{x}^2, \quad (\text{B}\cdot 16)$$

through which \hat{t} is defined as the cosmic time in the Einstein frame, and $\hat{a}(\hat{t})$ as the scale factor. From Eqs. (B.15) and (B.16), we find

$$d\hat{t} = \Omega dt, \quad (\text{B}\cdot 17)$$

$$\hat{a} = \Omega a. \quad (\text{B}\cdot 18)$$

In the FRW metric (B.16), the background gravitational field equations read

$$\hat{H}^2 = \left(\frac{1}{\hat{a}} \frac{d\hat{a}}{d\hat{t}} \right)^2 = \frac{1}{3} \left\{ \hat{T}_{00}^{(m)} + \frac{1}{2} \left[\left(\frac{d\hat{\varphi}_1}{d\hat{t}} \right)^2 + \left(\frac{d\hat{\varphi}_2}{d\hat{t}} \right)^2 \right] + \hat{V} \right\}, \quad (\text{B}\cdot 19)$$

$$\frac{d\hat{H}}{d\hat{t}} = -\frac{1}{6} \left\{ -\hat{T}^{(m)} + 4\hat{T}_{00}^{(m)} + 3 \left[\left(\frac{d\hat{\varphi}_1}{d\hat{t}} \right)^2 + \left(\frac{d\hat{\varphi}_2}{d\hat{t}} \right)^2 \right] \right\}. \quad (\text{B}\cdot 20)$$

Moreover, the equations of motion for the background homogeneous scalar fields read

$$\frac{d^2 \hat{\varphi}_i}{d\hat{t}^2} + 3\hat{H} \frac{d\hat{\varphi}_i}{d\hat{t}} = -\frac{\partial \hat{V}}{\partial \hat{\varphi}_i}. \quad (\text{B}\cdot 21)$$

B.3. Power-law inflation

Next, we show that power-law inflation can be realized in the Einstein frame. Substituting Eq. (B·2) with $f = At^2$ in the ansatz (15) in the Jordan frame into Eq. (B·17), we find

$$d\hat{t} = \sqrt{2A}tdt. \quad (\text{B}\cdot 22)$$

Integrating Eq. (B·22), we obtain

$$\hat{t} = \sqrt{\frac{A}{2}}t^2, \quad (\text{B}\cdot 23)$$

where we have chosen the constant of integration so that $\hat{t} = 0$ at $t = 0$.

Here we seek solutions with the following ansatz:

$$\hat{a} \propto \hat{t}^{\hat{\omega}}. \quad (\text{B}\cdot 24)$$

We now assume that in the inflationary stage, the cosmic energy density is dominated by the dilatons, and hence $\hat{T}_{\mu\nu}^{(m)}$ is negligible. Substituting the ansatz (B·24) into Eqs. (B·19) and (B·20) and using Eq. (B·4) with $f = At^2$, we obtain

$$\frac{\hat{\omega}^2}{\hat{t}^2} = \frac{1}{6} \left[\left(\frac{d\hat{\varphi}_1}{d\hat{t}} \right)^2 + \left(\frac{d\hat{\varphi}_2}{d\hat{t}} \right)^2 \right] + \frac{V}{24A} \frac{1}{\hat{t}^2}, \quad (\text{B}\cdot 25)$$

$$-\frac{\hat{\omega}}{\hat{t}^2} = -\frac{1}{2} \left[\left(\frac{d\hat{\varphi}_1}{d\hat{t}} \right)^2 + \left(\frac{d\hat{\varphi}_2}{d\hat{t}} \right)^2 \right]. \quad (\text{B}\cdot 26)$$

Eliminating $\left[d\hat{\varphi}_1/d\hat{t} \right]^2 + \left[d\hat{\varphi}_2/d\hat{t} \right]^2$ from Eqs. (B·25) and (B·26), we obtain

$$\hat{\omega}^2 - \frac{1}{3}\hat{\omega} - \frac{V}{24A} = 0. \quad (\text{B}\cdot 27)$$

The solution of Eq. (B·27) is given by

$$\hat{\omega} = \frac{1}{6} \left(1 + \sqrt{1 + \frac{3V}{2A}} \right) \quad (\text{B}\cdot 28)$$

$$\approx \frac{1}{6} \left(1 + \sqrt{1 + 9\omega^2} \right), \quad (\text{B}\cdot 29)$$

where we have taken the positive solution of $\hat{\omega}$. Moreover, in deriving the approximate equality in Eq. (B·29), we have ignored variation of the potential, and hence replaced V

by its average value Λ (as done in §3 in deriving the solutions of power-law inflation in the Jordan frame) and used the relation $\Lambda/A \approx 3/(32) (\alpha/\beta)^2 \approx 6\omega^2$, where the first approximate equality follows from Eq. (A·11), and in deriving the last approximate equality we have used Eq. (28). Hence, it follows from Eq. (B·29) that if the power-law exponent ω in the Jordan frame is much larger than unity, the power-law exponent $\hat{\omega}$ in the Einstein frame can also be much larger than unity. Because power-law inflation with $\omega \gg 1$ can be realized in the Jordan frame, as shown in §3 and §4.2, power-law inflation can also be realized in the Einstein frame.

Finally, we estimate the number of e -folds during inflation in the Einstein frame. As in the Jordan frame, this number is defined as⁹⁾

$$\hat{N} \equiv \int_{\hat{t}_i}^{\hat{t}_f} \hat{H} d\hat{t} \quad (\text{B}\cdot\text{30})$$

$$= \hat{\omega} \ln \left(\frac{\hat{t}_f}{\hat{t}_i} \right), \quad (\text{B}\cdot\text{31})$$

where in deriving Eq. (B·31), we have used $\hat{H} = \hat{\omega}/\hat{t}$. Applying Eq. (B·23) to Eq. (B·31), we obtain

$$\hat{N} = 2\hat{\omega} \ln \left(\frac{t_f}{t_i} \right) \quad (\text{B}\cdot\text{32})$$

$$\approx 2\hat{\omega} \ln \left[\frac{\pi}{\gamma(t_i)} \right]. \quad (\text{B}\cdot\text{33})$$

Here, in deriving Eq. (B·33), we have used Eqs. (54) and (55). For example, in the case (v) in Table I, it follows from $\omega = 4.1 \times 10^1$ and Eq. (B·29) that $\hat{\omega} = 2.1 \times 10^1$. Furthermore, it follows from this value of $\hat{\omega}$, the relation $\gamma(t_i) \simeq 3.5 \times 10^{-4}$ and Eq. (B·33) that $\hat{N} = 3.7 \times 10^2 > 70$. Thus, the expansion of the universe during the power-law inflation in the Einstein frame is large enough to solve both the horizon and flatness problems. Thus our result in the Einstein frame is the same as that in the Jordan frame, considered in §4.3.

Appendix C

— Equivalence of the Power Spectra of the Curvature Perturbations in the Jordan and Einstein Frames —

In this appendix, we show that the power spectra of the curvature perturbations in the Jordan and Einstein frames are the same, following the outline given in Ref. 21). Here we note that in this appendix we consider the case in which there exists a scalar field coupled to the scalar curvature, as shown in Eq. (C·1).

C.1. Curvature perturbations in the Jordan frame

We first consider the power spectrum of curvature perturbations in the Jordan frame. To begin with, we consider the action

$$\mathcal{S}_J = \int d^4x \sqrt{-g} \left[-\frac{1}{2} \mathcal{F}(\phi) R + \frac{1}{2} (\nabla\phi)^2 - \mathcal{V}[\phi] \right], \quad (\text{C}\cdot 1)$$

where $\mathcal{F}(\phi)$ is the coupling between the scalar field ϕ and the scalar curvature R , and $\mathcal{V}[\phi]$ is the potential of the scalar field; both $\mathcal{F}(\phi)$ and $\mathcal{V}[\phi]$ are arbitrary functions of the scalar field ϕ .

We consider a general perturbed metric for scalar perturbations

$$ds^2 = (1 + 2\Phi) dt^2 - 2a(t) B_{p,i} dx^i dt - a^2(t) \left[(1 - 2\psi) \delta_{ij} + 2E_{p,i,j} \right] dx^i dx^j, \quad (\text{C}\cdot 2)$$

where a comma denotes a flat-space coordinate derivative. Here, it is convenient to introduce the comoving curvature perturbation \mathcal{R} defined as

$$\mathcal{R} \equiv -\psi - \frac{H}{\dot{\phi}} \delta\phi, \quad (\text{C}\cdot 3)$$

where $\delta\phi$ is the perturbation of the field ϕ . It follows from the action in Eq. (C.1) that the equation of motion is given by²²⁾

$$\frac{1}{a^3 Q_S} \left(a^3 Q_S \dot{\mathcal{R}} \right)^\bullet + \frac{k^2}{a^2} \mathcal{R} = 0, \quad (\text{C}\cdot 4)$$

with

$$Q_S = \frac{\dot{\phi}^2 + 3\dot{\mathcal{F}}^2 / (2\mathcal{F})}{\left[H + \dot{\mathcal{F}} / (2\mathcal{F}) \right]^2}. \quad (\text{C}\cdot 5)$$

If we ignore the contribution of the decaying mode, the curvature perturbation is conserved in the large-scale limit ($k \rightarrow 0$). We here introduce the new variables $z = a\sqrt{Q_S}$ and $u = a\mathcal{R}$, so that Eq. (C.4) can be rewritten in the form

$$u'' + \left(k^2 - \frac{z''}{z} \right) u = 0, \quad (\text{C}\cdot 6)$$

where the prime denotes differentiation with respect to the conformal time $\eta = \int dt/a(t)$. Here, the gravitational term z''/z can be written as

$$\frac{z''}{z} = (aH)^2 \left[(1 + \delta_S)(2 + \delta_S + \varepsilon) + \frac{\delta'_S}{aH} \right], \quad (\text{C}\cdot 7)$$

where

$$\varepsilon = \frac{\dot{H}}{H^2}, \quad (\text{C}\cdot 8)$$

$$\delta_S = \frac{\dot{Q}_S}{2HQ_S}. \quad (\text{C}\cdot 9)$$

In the context of slow-roll inflation, it is a good approximation to ignore the variations of ε and δ_S . In the case of power-law inflation, the conformal time is given by $\eta = -1/[(1+\varepsilon)aH]$. From this expression of η and Eq. (C.7), we obtain

$$\frac{z''}{z} = \frac{\gamma_S}{\eta^2}, \quad (\text{C}\cdot 10)$$

with

$$\gamma_S = \frac{(1 + \delta_S)(2 + \delta_S + \varepsilon)}{(1 + \varepsilon)^2}. \quad (\text{C}\cdot 11)$$

The solution of Eq. (C.6) is then given by

$$u = \frac{\sqrt{\pi|\eta|}}{2} [c_1 H_{\nu_S}^{(1)}(k|\eta|) + c_2 H_{\nu_S}^{(2)}(k|\eta|)], \quad (\text{C}\cdot 12)$$

where

$$\nu_S \equiv \sqrt{\gamma_S + \frac{1}{4}}. \quad (\text{C}\cdot 13)$$

Here $H_{\nu_S}^{(1)}$ and $H_{\nu_S}^{(2)}$ are ν_S th-order Hankel functions of types 1 and 2, respectively. We choose the coefficients to be $c_1 = 0$ and $c_2 = 1$, so that positive frequency solutions in the Minkowski vacuum can be recovered in the asymptotic past. Then, using the relation $H_{\nu_S}^{(2)}(k|\eta|) \rightarrow (i/\pi)\Gamma(\nu_S)(k|\eta|/2)^{-\nu_S}$ for long wavelength perturbations ($k \rightarrow 0$), we find that the power spectrum of the comoving curvature perturbation $\mathcal{P}_{\mathcal{R}} \equiv [k^3/(2\pi^2)]|\mathcal{R}|^2$ is given by

$$\mathcal{P}_{\mathcal{R}} = \frac{1}{Q_S} \left(\frac{H}{2\pi}\right)^2 \left(\frac{1}{aH|\eta|}\right)^2 \left(\frac{\Gamma(\nu_S)}{\Gamma(3/2)}\right)^2 \left(\frac{k|\eta|}{2}\right)^{3-2\nu_S}. \quad (\text{C}\cdot 14)$$

Furthermore, the spectral index $n_S \equiv 1 + d \ln \mathcal{P}_{\mathcal{R}} / (d \ln k)$ is given by

$$n_S - 1 = 3 - 2\nu_S = 3 - \sqrt{4\gamma_S + 1}, \quad (\text{C}\cdot 15)$$

where the last equality follows from the relation (C.13).

C.2. Curvature perturbations in the Einstein frame

Next, in order to show that the power spectrum of the curvature perturbations in the Jordan frame is identical to that in the Einstein frame, we consider the power spectrum of curvature perturbations in the Einstein frame.

To begin with, we apply the following conformal transformation to the action in Eq. (C·1):

$$g_{\mu\nu} \rightarrow \hat{g}_{\mu\nu} = \Omega_S g_{\mu\nu}, \quad (\text{C}\cdot 16)$$

with

$$\Omega_S = \mathcal{F}. \quad (\text{C}\cdot 17)$$

The action in the Einstein frame is then given by^(6),21)

$$\mathcal{S}_E = \int d^4\hat{x} \sqrt{-\hat{g}} \left\{ -\frac{1}{2}\hat{R} + \frac{1}{2} \left[\frac{3}{2} \left(\frac{\mathcal{F}_\phi}{\mathcal{F}} \right)^2 + \frac{1}{\mathcal{F}} \right] (\hat{\nabla}\phi)^2 - \hat{V}[\phi] \right\}, \quad (\text{C}\cdot 18)$$

with

$$\hat{V}[\phi] = \frac{V[\phi]}{\mathcal{F}^2}. \quad (\text{C}\cdot 19)$$

Here, the subscript ϕ denotes partial differentiation with respect to ϕ . Furthermore, we introduce the new scalar fields

$$\frac{d\hat{\phi}}{d\phi} \equiv \sqrt{\mathcal{D}_E(\phi)}, \quad (\text{C}\cdot 20)$$

with

$$\mathcal{D}_E(\phi) = \frac{3}{2} \left(\frac{\mathcal{F}_\phi}{\mathcal{F}} \right)^2 + \frac{1}{\mathcal{F}}, \quad (> 0) \quad (\text{C}\cdot 21)$$

so that the action in Eq. (C·18) can be rewritten in the following canonical form:

$$\mathcal{S}_E = \int d^4\hat{x} \sqrt{-\hat{g}} \left[-\frac{1}{2}\hat{R} + \frac{1}{2} (\hat{\nabla}\hat{\phi})^2 - \hat{V}[\phi] \right]. \quad (\text{C}\cdot 22)$$

Now we consider a perturbed metric in the Einstein frame,

$$d\hat{s}^2 = \Omega_S ds^2 \quad (\text{C}\cdot 23)$$

$$\begin{aligned} &= \left(1 + 2\hat{\Phi} \right) dt^2 - 2\hat{a}(\hat{t}) \hat{B}_{\text{p},i} d\hat{x}^i dt \\ &\quad - \hat{a}^2(\hat{t}) \left[\left(1 - 2\hat{\psi} \right) \delta_{ij} + 2\hat{E}_{\text{p},i,j} \right] d\hat{x}^i d\hat{x}^j, \end{aligned} \quad (\text{C}\cdot 24)$$

and decompose the conformal factor into background and perturbed parts as

$$\Omega_S(\mathbf{x}, t) = \bar{\Omega}_S(t) \left(1 + \frac{\delta\Omega_S(\mathbf{x}, t)}{\bar{\Omega}_S(t)} \right). \quad (\text{C}\cdot\text{25})$$

In what follows we drop the bar on $\bar{\Omega}_S(t)$. From the above, we obtain the following relations:

$$\begin{aligned} \hat{a} &= a\sqrt{\Omega_S}, \quad d\hat{t} = \sqrt{\Omega_S}dt, \quad \hat{H} = \frac{1}{\sqrt{\Omega_S}} \left(H + \frac{\dot{\Omega}_S}{2\Omega_S} \right), \\ \hat{\Phi} &= \Phi + \frac{\delta\Omega_S}{2\Omega_S}, \quad \hat{\psi} = \psi + \frac{\delta\Omega_S}{2\Omega_S}. \end{aligned} \quad (\text{C}\cdot\text{26})$$

Using these relations, it is shown that curvature perturbations in the Einstein frame exactly coincide with those in the Jordan frame as follow:

$$\hat{\mathcal{R}} \equiv -\hat{\psi} - \frac{\hat{H}}{d\hat{\phi}/(d\hat{t})} \delta\hat{\phi} \quad (\text{C}\cdot\text{27})$$

$$= -\psi - \frac{H}{\dot{\phi}} \delta\phi = \mathcal{R}. \quad (\text{C}\cdot\text{28})$$

Hence, from Eq. (C·28) we find

$$\hat{\mathcal{P}}_{\mathcal{R}} = \mathcal{P}_{\mathcal{R}}. \quad (\text{C}\cdot\text{29})$$

Furthermore, we introduce the following quantities:

$$\begin{aligned} \hat{\gamma}_S &= \frac{(1 + \hat{\delta}_S)(2 + \hat{\delta}_S + \hat{\varepsilon})}{(1 + \hat{\varepsilon})^2}, \quad \hat{\varepsilon} = \frac{d\hat{H}/(d\hat{t})}{\hat{H}^2}, \\ \hat{\delta}_S &= \frac{d\hat{Q}_S/(d\hat{t})}{2\hat{H}\hat{Q}_S}, \quad \hat{Q}_S = \left[\frac{d\hat{\phi}/(d\hat{t})}{\hat{H}} \right]^2 = \frac{Q_S}{\mathcal{F}}. \end{aligned} \quad (\text{C}\cdot\text{30})$$

The spectral index of scalar perturbations in the Einstein frame is then given by

$$\hat{n}_S - 1 = 3 - \sqrt{4\hat{\gamma}_S + 1}. \quad (\text{C}\cdot\text{31})$$

Moreover, the two quantities $\hat{\varepsilon}$ and $\hat{\delta}_S$ can be expressed as follows:

$$\hat{\varepsilon} = \frac{\varepsilon - \vartheta}{1 + \vartheta} + \frac{\dot{\vartheta}}{H(1 + \vartheta)^2}, \quad (\text{C}\cdot\text{32})$$

$$\hat{\delta}_S = \frac{\delta - \vartheta}{1 + \vartheta}, \quad (\text{C}\cdot\text{33})$$

with

$$\vartheta = \frac{\dot{\mathcal{F}}}{2H\mathcal{F}}. \quad (\text{C}\cdot\text{34})$$

When the variation of ϑ is negligible ($\dot{\vartheta} \simeq 0$), which is the case in the context of slow-roll inflation, it can be shown that $\hat{\gamma}_S = \gamma_S$. Hence, the spectral index in Eq. (C·31) in the Einstein frame coincides with that in the Jordan frame: $\hat{n}_S = n_S$.

Finally, we note that in the above discussion, we have considered comoving curvature perturbations \mathcal{R} and then shown that the power spectrum of such perturbations in the Einstein frame exactly coincides with that in the Jordan frame. Contrastingly, in §5 we considered curvature perturbations on uniform-density hypersurfaces ζ . Comoving curvature perturbations and curvature perturbations on uniform-density hypersurfaces ζ , however, are identical on superhorizon scales.²³⁾ Thus, the results of §5 obtained from analysis in the Einstein frame are identical to those in the Jordan frame.

References

- 1) S. Perlmutter et al., *Astrophys. J.* **517** (1999), 565.
B. P. Schmidt et al., *Astrophys. J.* **507** (1998), 46.
A.G. Riess et al., *Astron. J.* **116** (1998), 1009.
- 2) S. Weinberg, *Rev. Mod. Phys.* **61** (1989), 1.
- 3) M. Yoshimura, *Phys. Lett. B* **608** (2005), 183.
- 4) P. A. M. Dirac, *Nature* **139** (1937), 323.
- 5) C. Brans and R. H. Dicke, *Phys. Rev.* **124** (1961), 925.
S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, New York, 1972).
- 6) Y. Fujii and K. Maeda, *The Scalar-Tensor Theory of Gravitation* (Cambridge University Press, Cambridge, United Kingdom, 2003).
K. Maeda, *Phys. Rev. D* **39** (1989), 3159.
- 7) D. Kazanas, *Astrophys. J.* **241** (1980), L59.
A. A. Starobinsky, *Phys. Lett. B* **91** (1980), 99.
A. H. Guth, *Phys. Rev. D* **23** (1981), 347.
K. Sato, *Mon. Not. R. Astron. Soc.* **195** (1981), 467.
- 8) A. D. Linde, *Particle Physics and Inflationary Cosmology* (Harwood, Chur, Switzerland, 1990).
K. A. Olive, *Phys. Rep.* **190** (1990), 307.
D. H. Lyth and A. Riotto, *Phys. Rep.* **314** (1999), 1.

- 9) E. W. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley, Redwood City, California, 1990).
- 10) C. L. Bennett et al., *Astrophys. J., Suppl. Ser.* **148** (2003), 1.
- 11) D. N. Spergel et al., *Astrophys. J., Suppl. Ser.* **148** (2003), 175.
- 12) T. Moroi and T. Takahashi, *Phys. Lett. B* **522** (2001), 215; *Phys. Rev. D* **66** (2002), 063501.
- 13) K. Enqvist and M. S. Sloth, *Nucl. Phys. B* **626** (2002), 395.
- 14) D. H. Lyth and D. Wands, *Phys. Lett. B* **524** (2002), 5.
- 15) D. H. Lyth, C. Ungarelli and D. Wands, *Phys. Rev. D* **67** (2003), 023503.
- 16) A. Zee, *Phys. Rev. Lett.* **42** (1979), 417.
- 17) D. La and P. J. Steinhardt, *Phys. Rev. Lett.* **62** (1989), 376.
- 18) C. Mathiazhagan and V.B. Johri, *Class. Quantum Grav.* **1** (1984), L29.
- 19) J. M. Bardeen, *Phys. Rev. D* **22** (1980), 1882.
 H. Kodama and M. Sasaki, *Prog. Theor. Phys. Suppl. No. 78* (1984), 1.
 V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, *Phys. Rep.* **215** (1992), 203.
- 20) D. Wands, K. A. Malik, D. H. Lyth and A. R. Liddle, *Phys. Rev. D* **62** (2000), 043527.
- 21) S. Tsujikawa and B. Gumjudpai, *Phys. Rev. D* **69** (2004), 123523.
- 22) J. Hwang, *Phys. Rev. D* **53** (1996), 762.
 J. Hwang and H. Noh, *Phys. Rev. D* **54** (1996), 1460.
 H. Noh and J. Hwang, *Phys. Lett. B* **515** (2001), 231.
- 23) A. Riotto, hep-ph/0210162.




Fig. 1. Estimates of $\Xi_1 = \text{Tr } \tilde{\mathcal{M}}^2/M^2$ in the relation (46). The solid curve is for the case $\Upsilon = 1.5 \times 10^4$, and the dotted curve is for the case $\Upsilon = 3.5 \times 10^2$. The left panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($0 \leq j \leq 89$), where j is an integer, while the right panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 89$).




Fig. 2. Estimates of $\Xi_2 = \det \tilde{\mathcal{M}}^2/M^2$ in the relation (47). The solid curve is for the case $\Upsilon = 1.5 \times 10^4$, and the dotted curve is for the case $\Upsilon = 3.5 \times 10^2$. The left panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($0 \leq j \leq 89$), where j is an integer, while the right panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 89$). In the case $\Upsilon = 1.5 \times 10^4$, for $\epsilon_2 \leq 2.9 \times 10^{-3}$ and $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\Xi_2 < 0$. Similarly, in the case $\Upsilon = 3.5 \times 10^2$, for $\epsilon_2 \leq 6.9 \times 10^{-3}$ and $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\Xi_2 < 0$.




Fig. 3. Estimates of $\tilde{\Theta}$ in the relation (48). The solid curve is for the case $\Upsilon = 1.5 \times 10^4$, and the dotted curve is for the case $\Upsilon = 3.5 \times 10^2$. The left panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($0 \leq j \leq 89$), where j is an integer, while the right panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 89$). In the case $\Upsilon = 1.5 \times 10^4$, for $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\tilde{\Theta} < 0$. Similarly, in the case $\Upsilon = 3.5 \times 10^2$, for $\epsilon_2 \geq 4.6 \times 10^{-2}$, we have $\tilde{\Theta} < 0$.




Fig. 4. Estimates of $\langle \lambda_+ \rangle / M^2$ and $\langle \lambda_- \rangle / M^2$ in Eqs. (49) and (50). The solid curves are for the case $\Upsilon = 1.5 \times 10^4$, and the dotted curves are for the case $\Upsilon = 3.5 \times 10^2$. Thick curves represent $\langle \lambda_+ \rangle / M^2$, and the thin curves represent $\langle \lambda_- \rangle / M^2$. The left panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($0 \leq j \leq 89$), where j is an integer, while the right panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 89$). In both the cases $\Upsilon = 1.5 \times 10^4$ and $\Upsilon = 3.5 \times 10^2$, for $\epsilon_2 \leq 2.4 \times 10^{-3}$ and $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\langle \lambda_- \rangle / M^2 < 0$.




Fig. 5. Estimates of $\langle \lambda_- / \lambda_+ \rangle$ in Eq. (51). The solid curve is for the case $\mathcal{T} = 1.5 \times 10^4$, and the dotted curve is for the case $\mathcal{T} = 3.5 \times 10^2$. In this figure, the former and latter coincide. The left panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-4}$, $\epsilon_2 = 1.0 \times 10^{-3} + 1.0 \times 10^{-4}j$ ($0 \leq j \leq 89$), where j is an integer, while the right panel describes the case $\epsilon_1 = \epsilon_2 + 1.0 \times 10^{-3}$, $\epsilon_2 = 1.0 \times 10^{-2} + 1.0 \times 10^{-3}j$ ($0 \leq j \leq 89$). In both the cases $\mathcal{T} = 1.5 \times 10^4$ and $\mathcal{T} = 3.5 \times 10^2$, for $\epsilon_2 \leq 2.4 \times 10^{-3}$ and $\epsilon_2 \geq 4.5 \times 10^{-2}$, we have $\langle \lambda_- / \lambda_+ \rangle < 0$.

Title

Firstname FAMILYNAME

Name and Address of your affiliation

Write your ABSTRACT here.

§1. Section Title

Start your paper from here.

References

1)