## Spin Correlations in Top-Quark Pair Production  $\mathrm{at}\,\,e^+e^-\,$  Colliders

Stephen Parke [†](#page-9-0) and Yael Shadmi [∗](#page-9-0) Theoretical Physics Department

Fermi National Accelerator Laboratory P.O. Box 500, Batavia, IL 60510 (June 21, 1996)

## Abstract

We show that top-quark pairs are produced in an essentially unique spin configuration in polarized  $e^+e^-$  colliders at all energies above the threshold region. Since the directions of the electroweak decay products of polarized top-quarks are strongly correlated to the top-quark spin axis, this unique spin configuration leads to a distinctive topology for top-quark pair events which can be used to constrain anomalous couplings to the top-quark. A significant interference effect between the longitudinal and transverse W-bosons in the decay of polarized top-quarks is also discussed. These results are obtained at leading order in perturbation theory but radiative corrections are expected to be small.

Sincethe top-quark, with mass in the range of  $175 \text{ GeV}$  [[1,2\]](#page-9-0), decays electroweakly before hadronizing[[3\]](#page-9-0), there are significant angular correlations between the decay products of the top-quark and the spin of the top-quark. Therefore, if top-quark pairs are produced with large polarizations, there will be sizable angular correlations between the decay products and the spin orientation of each top-quark, as well as between the decay products of the top-quark and the decay products of the top anti-quark. These angular correlations depend sensitively on the top-quark couplings to the  $Z$  and the photon, and to the  $W$  and  $b$ -quark. Many authors have proposed to use the angular information in top-quark events produced at  $e^+e^-$  colliders to constrain deviations from Standard-Model couplings [\[4-8](#page-7-0)]. In most of these studies, the top-quark spin is decomposed in the helicity basis, i.e. along the direction of motion of the quark. Recently, Mahlon and Parke [\[9\]](#page-9-0) have shown that this decomposition in the helicity basis is far from the optimal decomposition for top-quarks produced at the Tevatron. In this paper we address the issue of what is the optimal decomposition of the topquark spins for  $e^+e^-$  colliders. Here, we only consider top-quark production at energies that are above the threshold region[[10](#page-9-0)]. We give the differential cross-section for top-quark pair production for a generic spin basis. We have found that there is a decomposition for which the top-quark pairs are produced in an essentially unique spin configuration at polarized  $e^+e^-$  colliders. We call this spin basis the "Off-diagonal" basis because the contribution from like-spin pairs vanishes to leading order in perturbation theory. Finally, we discuss the angular correlations between the decay products of the top-quarks, and point out a significant interference effect between the longitudinal and transverse W-bosons in the decays of the polarized top-quarks.

The generic spin basis we will consider in this paper is shown in Figure [1](#page-10-0). This choice is motivated by two features of the Standard Model prediction for top-quark pair production at  $e^+e^-$  colliders at leading-order in perturbation theory; the transverse top-quark polarization is zero, and CP is conserved. Therefore, we have defined our general basis so that the spins of the top quark and anti-quark are in the production plane, so there is no transverse polarization, and the spin four-vectors are back-to-back in the zero momentum frame, so <span id="page-2-0"></span>that states with opposite spins are CP eigenstates.

The top-quark spin states are defined in the top-quark rest-frame, where we decompose the top spin along the direction  $\hat{s}_t$ , which makes an angle  $\xi$  with the anti-top momentum in the clockwise direction. Similarly, the top anti-quark spin states are defined in the anti-top rest-frame, along the direction  $\hat{s}_{\bar{t}}$ , which makes the same angle  $\xi$  with the top momentum also in the clockwise direction. Thus, the state  $t_1\bar{t}_1(t_1\bar{t}_\perp)$  refers to a top with spin in the  $+\hat{s}_t$  $(-\hat{s}_t)$  direction in the top rest-frame, and an anti-top with spin  $+\hat{s}_{\bar{t}}(-\hat{s}_{\bar{t}})$  in the anti-top rest-frame.

Using this generic spin basis, the Standard Model leading-order polarized cross-sections for top-quark pair production at center-of-mass energy  $\sqrt{s}$ , top-quark speed  $\beta$  and top-quark scattering angle  $\theta^*$ , are given by

$$
\frac{d\sigma}{d\cos\theta^*} (e_L^- e_R^+ \to t_\uparrow \bar{t}_\uparrow) = \frac{d\sigma}{d\cos\theta^*} (e_L^- e_R^+ \to t_\downarrow \bar{t}_\downarrow)
$$

$$
= \left(\frac{3\pi\alpha^2}{2s}\beta\right) |A_{LR}\cos\xi - B_{LR}\sin\xi|^2,
$$

$$
\frac{d\sigma}{d\cos\theta^*} \left( e^-_L e^+_R \to t_\uparrow \,\bar{t}_\downarrow \, or \, t_\downarrow \,\bar{t}_\uparrow \right) = \left( \frac{3\pi\alpha^2}{2s} \beta \right) \, |A_{LR}\sin\xi \, + \, B_{LR}\cos\xi \, \pm \, D_{LR}|^2. \tag{1}
$$

Here  $\alpha$  is the QED fine structure constant and the quantities A, B and D depend on the kinematic variables  $\beta$  and cos  $\theta^*$ , and on the fermion couplings to the photon and Z-boson, in the following way

$$
A_{LR} = [(f_{LL} + f_{LR}) \sqrt{1 - \beta^2} \sin \theta^*]/2 ,
$$
  
\n
$$
B_{LR} = [f_{LL} (\cos \theta^* + \beta) + f_{LR} (\cos \theta^* - \beta)]/2 ,
$$
  
\n
$$
D_{LR} = [f_{LL} (1 + \beta \cos \theta^*) + f_{LR} (1 - \beta \cos \theta^*)]/2 ,
$$
\n(2)

with

$$
f_{IJ} = Q_{\gamma}(e)Q_{\gamma}(t) + Q_Z^I(e)Q_Z^J(t)\Big(\frac{1}{\sin^2 \theta_W}\Big)\Big(\frac{s}{(s - M_Z^2) + iM_Z\Gamma_Z}\Big) ,\qquad (3)
$$

where  $M_Z$  is the Z mass,  $\Gamma_Z$  is the Z width, and  $I, J \in (L, R)$ . The electron couplings are given by

$$
Q_{\gamma}(e) = -1, \quad Q_Z^L(e) = \frac{2\sin^2\theta_W - 1}{2\cos\theta_W}, \quad Q_Z^R(e) = \frac{\sin^2\theta_W}{\cos\theta_W}, \tag{4}
$$

and the top-quark couplings are given by

$$
Q_{\gamma}(t) = \frac{2}{3}, \quad Q_Z^L(t) = \frac{3 - 4\sin^2\theta_W}{6\cos\theta_W}, \quad Q_Z^R(t) = \frac{-2\sin^2\theta_W}{3\cos\theta_W}.
$$
 (5)

The angle  $\theta_W$  is the Weinberg angle. In the limit  $s \gg M_Z^2$  the products of fermion couplings for top-quark production are

$$
f_{LL} = -1.19, \quad f_{LR} = -0.434, \quad f_{RR} = -0.868, \quad f_{RL} = -0.217 \tag{6}
$$

where, as throughout this paper, we use  $\sin^2 \theta_W = 0.232$ .

Thecross-sections for  $e_R^- e_L^+$  may be obtained from eqns. ([1-3](#page-2-0)) by interchanging L and R as well as  $\uparrow$  and  $\downarrow$ . These cross-sections can be conveniently derived using the spinor helicity method for massive fermions described in [\[9](#page-9-0)].

Our generic basis reproduces the familiar helicity basis for the special case

$$
\cos \xi = \pm 1 \tag{7}
$$

for which the top-quark spin is defined along its direction of motion. Substituting eqn. (7) in our general polarized cross-section expressions([1\)](#page-2-0), we recover the well known helicity cross-sections

$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_L \bar{t}_L \right) = \frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_R \bar{t}_R \right)
$$

$$
= \left( \frac{3\pi\alpha^2}{8s} \beta \right) |f_{LL} + f_{LR}|^2 \left( 1 - \beta^2 \right) \sin^2\theta^* ,
$$

$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_R \bar{t}_L \text{ or } t_L \bar{t}_R \right) = \left( \frac{3\pi\alpha^2}{8s} \beta \right) |f_{LL}(1 \mp \beta) + f_{LR}(1 \pm \beta)|^2 \left( 1 \mp \cos\theta^* \right)^2 . \tag{8}
$$

Another useful basis is the "Beamline basis" [\[9](#page-9-0)], in which the top-quark spin axis is the positron direction in the top rest-frame, and the top anti-quark spin axis is the electron direction in the anti-top rest-frame. In terms of the spin angle  $\xi$ , this corresponds to

$$
\cos \xi = \frac{\cos \theta^* + \beta}{1 + \beta \cos \theta^*} \ . \tag{9}
$$

<span id="page-4-0"></span>The polarized cross-sections in this basis are

$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\uparrow \, \bar{t}_\uparrow \right) = \frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\downarrow \, \bar{t}_\downarrow \right)
$$
\n
$$
= \left( \frac{3\pi\alpha^2}{2s} \beta \right) |f_{LR}|^2 \frac{\beta^2 (1-\beta^2) \sin^2\theta^*}{(1+\beta\cos\theta^*)^2},
$$
\n
$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\downarrow \, \bar{t}_\uparrow \right) = \left( \frac{3\pi\alpha^2}{2s} \beta \right) |f_{LR}|^2 \frac{\beta^4 \sin^4\theta^*}{(1+\beta\cos\theta^*)^2},
$$
\n
$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\uparrow \, \bar{t}_\downarrow \right) = \left( \frac{3\pi\alpha^2}{2s} \beta \right) \left| f_{LL}(1+\beta\cos\theta^*) + f_{LR} \frac{(1-\beta^2)}{(1+\beta\cos\theta^*)} \right|^2. \tag{10}
$$

Note that in this basis, three out of the four polarized cross-sections are proportional to  $|f_{LR}|^2$  which is much smaller  $|f_{LL}|^2$ . These three components are further suppressed by at least two powers of  $\beta$ . The remaining component,  $t_{\uparrow} \bar{t}_{\downarrow}$ , will therefore account for most of the total cross-section. Hence the top-quark spin is strongly correlated with the positron spin direction determined in the top-quark rest-frame.

The production threshold for top-quarks of mass 175 GeV is far above the Z-boson pole. In this region, the Z width is negligible and we can take the factors  $f_{IJ}$  as real. With real  $f_{IJ}$ 's one can choose a basis in which the  $t_{\uparrow}\bar{t}_{\uparrow}$  and  $t_{\downarrow}\bar{t}_{\downarrow}$  components vanish identically, see eqn.([1](#page-2-0)). In this basis, which we refer to as the "Off-diagonal basis", the spin angle  $\xi$  is given by

$$
\tan \xi = \frac{(f_{LL} + f_{LR})\sqrt{1 - \beta^2} \sin \theta^*}{f_{LL}(\cos \theta^* + \beta) + f_{LR}(\cos \theta^* - \beta)}
$$
(11)

and the polarized cross-sections are

$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\uparrow \, \bar{t}_\uparrow \right) = \frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\downarrow \, \bar{t}_\downarrow \right) = 0 \,,
$$
\n
$$
\frac{d\sigma}{d\cos\theta^*} \left( e_L^- e_R^+ \to t_\uparrow \, \bar{t}_\downarrow \, \text{or} \, t_\downarrow \, \bar{t}_\uparrow \right) = \left( \frac{3\pi\alpha^2}{8s} \beta \right) \left[ f_{LL}(1 + \beta\cos\theta^*) + f_{LR}(1 - \beta\cos\theta^*) \right]
$$
\n
$$
\pm \sqrt{(f_{LL}(1 + \beta\cos\theta^*) + f_{LR}(1 - \beta\cos\theta^*))^2 - 4f_{LL}f_{LR}\beta^2\sin^2\theta^*} \Bigg]^2. \tag{12}
$$

For  $|f_{LL}| \gg |f_{LR}|$  only the  $t_{\uparrow}t_{\downarrow}$  component is substantially different from zero, and to leading

order in  $\frac{f_{LR}}{f_{LL}}$ , the  $t\downarrow \bar{t}_{\uparrow}$  component is given by the same expression as for the Beamline basis<sup>1</sup>. In general there are two "Off-diagonal" bases for fermion pair production, one for  $e^-_L e^+_R$ R scattering and the other for  $e^-_R e^+_L$  $L_t^+$ , since in general  $\frac{f_{LR}}{f_{LL}} \neq \frac{f_{RL}}{f_{RR}}$  $\frac{f_{RL}}{f_{RR}}$ . However for top-quark production the two ratios are approximately equal in both sign and magnitude, so that the two "Off-diagonal" bases are almost identical. In the rest of this paper we will only use the Off-diagonal basis for  $e^-_L e^+_R$  defined by eqn. [\(11](#page-4-0)) even when discussing  $e^-_R e^+_L$  $\frac{1}{L}$  scattering.

To illustrate the different spin bases we now consider top-quark pair production, at a 400 GeV  $e^+e^-$  collider. We take the top-quark mass to be 175 GeV ( $\beta \sim 0.5$ ). Fig. [2](#page-11-0) shows the dependence of the spin direction angle,  $\xi$ , on the scattering angle,  $\theta^*$ , for the helicity, Beamline and Off-diagonal bases. The Beamline basis lies close to the Off-diagonal basis for all scattering angles. For  $\cos \theta^*$  near zero there is a marked difference between the helicity basis and the other two bases. Note that as  $\beta \to 1$ , both the Beamline and Off-diagonal bases approach the helicity basis. Therefore, for lepton colliders with center-of-mass energies greater than 1 TeV, all three bases are equivalent.

The cross-sections for producing  $t\bar{t}$  pairs of definite helicities are shown in Fig. [3](#page-12-0) for both  $e^-_L e^+_R$  and  $e^-_R e^+_L$  $L<sup>+</sup>$  scattering. While the dominant components of the signal are  $t_L\bar{t}_R$ for a left-handed electron and  $t_R\bar{t}_L$  for a right-handed electron, other spin components make up more than 40% of the total cross-section. In the Off-diagonal basis, in contrast, only one spin component is appreciably non-zero for all values of the scattering angle;  $t_{\uparrow}$   $\bar{t}_{\downarrow}$  for  $e^-_L$   $e^+_R$  and  $t_\downarrow$   $\bar{t}_\uparrow$  for  $e^-_R$   $e^+_L$  $_L^+$ , see Fig. [4.](#page-13-0) All other components are more than two and a half

<sup>1</sup>Thepolarized cross-section formulae ([1\)](#page-2-0) are also valid for top-quark pair production in  $q\bar{q}$ scattering, with an appropriate change of couplings. The Off-diagonal basis for  $q\bar{q}$ -scattering is given by eqn. [\(11\)](#page-4-0) with  $f_{LL} = f_{LR} = f_{RL} = f_{RR}$ , for which  $\tan \xi = \sqrt{1 - \beta^2} \tan \theta^*$  with

$$
\frac{d\sigma}{d\cos\theta^*} \left( q_L^- \, \bar{q}_R^+ \to t_\uparrow \, \bar{t}_\uparrow \, \text{or} \, \, t_\downarrow \, \bar{t}_\downarrow \right) = 0
$$

and

$$
\frac{d\sigma}{d\cos\theta^*} \left( q_L^- \, \bar{q}_R^+ \to t_\uparrow \, \bar{t}_\downarrow \, \text{or} \, t_\downarrow \, \bar{t}_\uparrow \right) = \left( \frac{\pi \alpha_s^2}{9s} \beta \, \right) \left[ 1 \pm \sqrt{1 - \beta^2 \sin^2\theta^*} \, \right]^2 \, .
$$

orders of magnitude smaller than these dominant contributions. Thus, when defined in the Off-diagonal basis, the spins of the top quark and anti-quark produced in  $e^-_L e^+_R$  and  $e^-_R e^+_L$ L scattering are essentially determined. This continues to hold for top-quark pair production at higher energies. In Fig. [5](#page-14-0) we show, for  $e^-_L e^+_R$  collisions, the fraction of top-quark events in the  $t_1\bar{t}_\downarrow$  configuration, defined in the Off-diagonal basis, as a function of the top-quark speed  $\beta$ . Also shown is the fraction of top-quark pairs in the  $t_L\bar{t}_R$  helicity configuration. The Off-diagonal basis gives a very clean  $t_{\uparrow}\bar{t}_{\downarrow}$  spin state for all values of  $\beta$  in these collisions. Similarly, the spin state  $t\downarrow \bar{t}_{\uparrow}$  dominates  $e^{-}_{R}e^{+}_{L}$  $<sub>L</sub><sup>+</sup>$  collisions at all energies.</sub>

We do not show here cross-section plots for the Beamline basis, since they are almost identical to those of Fig. [4,](#page-13-0) except that the non-dominant contributions are now at the 1% level for a 400 GeV collider. For both the Beamline basis and the Off-diagonal basis, the contribution of higher order corrections to the non-dominant components is expected to increase their total contribution to the few percent level.

Since the top-quark pairs are produced in an unique spin configuration, and the electroweak decay products of polarized top-quarks are strongly correlated to the spin axis, the top-quark events at  $e^+e^-$  collider have a very distinctive topology. Deviations from this topology would signal anomalous couplings. In the Standard Model, the predominant decay mode of the top-quark is  $t \to bW^+$ , with the  $W^+$  decaying either hadronically or leptonically. For definiteness we consider here the decay  $t \to b W^+ \to b e^+ \nu$ . The differential decay width of a polarized top-quark depends non-trivially on three angles. The first is the angle,  $\chi_w^t$ , between the top-quark spin and the direction of motion of the W-boson in the top-quark rest-frame. Next is the angle between the direction of motion of the b-quark and the positron in the W-boson rest-frame. We call this angle  $\pi - \chi_e^w$ . Finally, in the top-quark rest-frame, we have the azimuthal angle, Φ, between the positron direction of motion and the top-quark spin around the direction of motion of the W-boson.

The differential polarized top-quark decay distribution in terms of these three angles is given by

<span id="page-7-0"></span>
$$
\frac{1}{\Gamma_T} \frac{d^3 \Gamma}{d \cos \chi_w^t \ d \cos \chi_e^w \ d\Phi} = \frac{3}{16\pi \ (m_t^2 + 2m_W^2)} \left[ m_t^2 (1 + \cos \chi_w^t) \sin^2 \chi_e^w \right. \\
\left. + m_W^2 (1 - \cos \chi_w^t) (1 - \cos \chi_e^w)^2 + 2m_t m_W (1 - \cos \chi_e^w) \sin \chi_e^w \sin \chi_w^t \cos \Phi \right],\tag{13}
$$

where  $m_t$  is the top-quark mass,  $m_W$  is the W mass, and  $\Gamma_T$  is the total decay width (we neglect the b-quark mass). The first and second terms in (13) give the contributions of longitudinal and transverse W-bosons respectively. The interference term, given by the third term in (13), does not contribute to the total width, but its effects on the angular distribution of the top-quark decay products are sizable. Fig. [6](#page-15-0) shows contour plots of the differential angular decay distribution in the  $\chi_e^w - \chi_w^t$  plane<sup>2</sup>, after integrating over the azimuthal angle  $\Phi$ . Fig. [7](#page-16-0) shows analogous contours integrated over  $\Phi$  for positive (solid lines) and for negative (dashed lines) values of  $\cos \Phi$  separately. The pronounced difference between these is related to the size of the interference term, which can be seen from the Φ-distribution

$$
\frac{1}{\Gamma_T} \frac{d\Gamma}{d\Phi} = \frac{1}{2\pi} \left[ 1 + \frac{3\pi^2 m_t m_W}{16(m_t^2 + 2m_W^2)} \cos \Phi \right]. \tag{14}
$$

For a 175 GeV top-quark the coefficient in front of the cosine term has a value equal to 0.59, therefore the maximum and minimum values of this distribution are approximately 4 to 1.

There are also significant correlations of the angle between the top-quark spin and the momentum of the *i*-th decay product,  $\chi_i^t$ , measured in the top-quark rest-frame. The differential decay rate of the top-quark is given by

$$
\frac{1}{\Gamma_T} \frac{d\Gamma}{d\cos\chi_i^t} = \frac{1}{2} \left[ 1 + \alpha_i \cos\chi_i^t \right],\tag{15}
$$

where  $\alpha_b = -0.41$ ,  $\alpha_{\nu} = -0.31$  and  $\alpha_{e^+} = 1$ , for  $m_t = 175$  GeV, see ref. [\[11\]](#page-9-0).

In summary, we have presented simple analytic expressions for the polarized cross-section for top-quark pair production in polarized  $e^+e^-$  colliders. For a particular choice of axes,

<sup>&</sup>lt;sup>2</sup>We take  $M_W = 80$  GeV.

the Off-diagonal basis, not only do the like-spin contributions vanish, but one spin configuration dominates the total cross-section. In this configuration, the top-quark spin is strongly correlated with the positron spin direction determined in the top-quark rest-frame. The subsequent electroweak decays of the top-quark pair give decay products whose angular distributions are highly correlated with the parent top-quark spin. Top-quark pair events thus have a distinctive topology. This topology is sensitive to the top-quark couplings to the Z-boson and to the photon, which determine the orientation and the size of the topquark and top anti-quark polarizations, as well as to the top-quark couplings to the W and the b-quark, which determine its decay distributions. Angular correlations in top-quark events may therefore be used to constrain deviations from the Standard Model. We have also shown that the interference between the *longitudinal* and *transverse W*-bosons has a significant impact on the angular distribution of the top-quark decay products, and thus will provide additional means for testing the Standard Model predictions for top-quark decays.

## ACKNOWLEDGMENTS

The Fermi National Accelerator Laboratory is operated by Universities Research Association, Inc., under contract DE-AC02-76CHO3000 with the U.S. Department of Energy.

## REFERENCES

- <span id="page-9-0"></span>† Electronic address: parke@fnal.gov
- <sup>∗</sup> Electronic address: yael@fnth06.fnal.gov
- [1] F. Abe, et. al., Phys. Rev. Lett. 74, 2626 (1995).
- [2] S. Abachi, et. al., Phys. Rev. Lett. 74, 2632 (1995).
- [3] I. Bigi, Y. Dokshitzer, V. Khoze, J. Kühn, and P. Zerwas, Phys. Lett. **181B**, 157 (1986).
- [4] M.E. Peskin, in Physics and Experiments at Linear Colliders, R. Orava, P. Eeorla and M. Nordberg, eds. (World Scientific, 1992).
- [5] G.A. Ladinsky and C.P. Yuan, Phys. Rev. D49, 4415 (1994).
- [6] T.L. Barklow and C.R. Schmidt, in DPF '94: The Albuquerque Meeting, S. Seidel, ed. (World Scientific, Singapore, 1995).
- [7] C.R. Schmidt, preprint SCIPP-95-14, [hep-ph/9504434.](http://arxiv.org/abs/hep-ph/9504434)
- [8] G.L. Kane, J. Pumplin, and W. Repko, Phys. Rev. Lett. 41, 1689 (1978); M. Anselmino, P. Kroll, and B. Pire, Phys. Lett. B167, 113 (1986); G.L. Kane, G.A. Ladinsky, and C.–P. Yuan, Phys. Rev. D45, 124 (1992); D. Atwood and A. Soni, Phys. Rev. D45, 2405 (1992); C.–P. Yuan, Phys. Rev. D45, 782 (1992); W. Bernreuther, J.P. Ma, and T. Schröder, Phys. Lett. **B297**, 318 (1992); W. Bernreuther, O. Nachtmann, P. Overmann, and T. Schröder, Nucl. Phys. **B388** 53, (1992); erratum **B406**, 516 (1993); T. Arens and L.M. Seghal, Nucl. Phys. B393, 46 (1993); F. Cuypers and S.D. Rindani, Phys. Lett. **B343**, 333 (1995); P. Poulose, S.D. Rindani, Phys. Lett. **B349**, 379 (1995);
- [9] G. Mahlon and S. Parke, Phys. Rev 53, 4886 (1996).
- [10] For top polarization results near threshold see, for example: R. Harlander, M. Jezabek, J.H. Kühn, and T. Teubner, Phys. Lett. **B346**, 137 (1995); R. Harlander, M. Jeżabek, J.H. Kühn and M. Peter, preprint TTP-95-48, [hep-ph/9604328](http://arxiv.org/abs/hep-ph/9604328), and references therein.
- [11] M. Je $\dot{z}$ abek and J.H. Kühn, Phys. Lett. **B329**, 317 (1994).



<span id="page-10-0"></span>

FIG. 1. The scattering process in the center-of-mass frame (a), in the top-quark rest-frame (b) and in the top anti-quark rest-frame (c).  $s_t$  ( $s_{\bar{t}}$ ) is the top (anti-top) spin axis.

<span id="page-11-0"></span>

FIG. 2. The dependence of the spin angle,  $\xi$ , on the scattering angle,  $\theta^*$ , for the helicity, Beamline and Off-diagonal (defined for  $e^-_L\ e^+_R$  $R$ <sup>+</sup> scattering) bases for a 175 GeV top-quark produced by a 400 GeV  $e^+e^-$  collider.

<span id="page-12-0"></span>

FIG. 3. The differential cross-sections for producing top-quark pairs at a 400 GeV  $e^+e^$ collider in the following helicity configurations:  $t_L\bar{t}_R$  (LR),  $t_R\bar{t}_L$  (RL), and the sum of  $t_L\bar{t}_L$  and  $t_R\bar{t}_R$  (LL+RR), for left-handed and right-handed electron beams.

<span id="page-13-0"></span>

FIG. 4. The differential cross-sections for producing top-quark pairs at a 400 GeV  $e^+e^$ collider in the following spin configurations in the Off-diagonal basis (defined for  $e^-_L$   $e^+_R$  $R$  scattering):  $t_{\uparrow}\bar{t}_{\downarrow}$  (UD),  $t_{\downarrow}\bar{t}_{\uparrow}$  (DU), and the sum of  $t_{\uparrow}\bar{t}_{\uparrow}$  and  $t_{\downarrow}\bar{t}_{\downarrow}$  (UU+DD), for left-handed and right-handed electron beams.

<span id="page-14-0"></span>

FIG. 5. The fraction of top-quark pairs in the dominant spin configuration in the Off-diagonal basis and in the helicity basis, as a function of the top-quark speed,  $\beta$ , in  $e^-_L e^+_R$  $R$ <sup>+</sup> scattering. The solid line gives  $\sigma(e^-_L e^+_R \to t_1 \bar{t}_\downarrow)/\sigma_T$ , defined in the Off-diagonal basis. The dot-dashed line gives  $\sigma(e^-_L e^+_R \to t_L \bar{t}_R)/\sigma_T$  in the helicity basis. Here  $\sigma_T$  is the total cross-section for  $e^-_L e^+_R \to t\bar{t}$ .

<span id="page-15-0"></span>

FIG. 6. Contours of the top-quark decay distribution, eqn. [\(13\)](#page-7-0), integrated over all Φ, in the  $\chi_e^w - \chi_w^t$  plane.

<span id="page-16-0"></span>

FIG. 7. Contours of the top-quark decay distribution, eqn. [\(13\)](#page-7-0), integrated over  $\Phi$  for  $\cos \Phi > 0$ (solid), and for  $\cos \Phi < 0$  (dashed) in the  $\chi_e^w - \chi_w^t$  plane. The solid and dashed curves join continuously at the edges of the plot, where the interference term is zero.