

NORDITA-96/40 P

hep-ph/9606472

June 28, 1996

Generating heavy quarkonia in a perturbative QCD cascade

Per Ernström and Leif Lönnblad

NORDITA

Blegdamsvej 17, DK-2100 Copenhagen Ø

per@nordita.dk, leif@nordita.dk

Abstract

We present an implementation of heavy quarkonium production within a perturbative QCD cascade based on the Color Dipole Cascade model. We consider the processes most relevant in the context of the ψ' surplus at the Tevatron; $g \rightarrow \psi'$ and $c \rightarrow \psi'$ in the color-singlet model and $g \rightarrow \psi'$ through the color-octet mechanism. Our implementation is, however, easily extendible to other quarkonia and other production mechanisms.

Where comparison is possible we find good agreement with analytical calculations.

We present some suggestions for measurements at the Tevatron that would be sensitive to the shape of the fragmentation functions. Our calculations indicate that such measurements could be used to test the color-octet mechanism solution to the ψ' surplus at the Tevatron.

1 Introduction

Recent Tevatron data [1, 2] on prompt charmonium (ψ and ψ') production at large p_T has attracted a lot of attention.

At large p_T , charmonium production is believed to be dominated by parton fragmentation [3], and the Tevatron data [1, 2] does indeed show a $1/p_T^5$ behavior consistent with the predictions of a fragmentation process.

The large c quark mass m_c make, however, predictions possible also for the magnitude of the cross section. Production of charmonium states can be factorized into the hard QCD production of a heavy $c\bar{c}$ pair within a region $1/m_c$, followed by the nonperturbative formation of the charmonium state. In [4], a general factorization scheme was developed to all orders in the relative velocity v of the c and \bar{c} quark in the charmonium state. For S-wave states, treated to leading order in v^2 , this factorization scheme reduces to the color singlet model [5]; the $c\bar{c}$ pair must be produced in a color singlet state with the same spin and angular momentum as the charmonium state. The subsequent nonperturbative formation of the charmonium state can be described by a single process independent parameter. Fixing this parameter from measurements on the electro-magnetic decay of the ψ and ψ' , predictions can be made.

A color singlet model calculation to LO in α_s of large p_T ψ' production fall, however, more than an order of magnitude below the Tevatron data [1, 2, 6, 7]. Although this is the largest deviation from LO color singlet model predictions seen, it should be noted that large deviations have been observed also at low p_T [8, 9].

A calculation of prompt ψ production [6, 7, 10] give a similar result, but in this case contributions from P-wave charmonium states decaying to ψ obscure the interpretation of the data.

Braaten *et al.* [11] have proposed that the ψ' surplus at the Tevatron could be explained by a v suppressed production mechanism, usually referred to as the color octet mechanism. The $c\bar{c}$ pair is produced in a color octet 3S_1 state, and forms the ψ' through soft gluon emission or absorption. The new nonperturbative parameter giving the “probability” for the color octet $c\bar{c}$ pair to form the ψ' is suppressed by v^4 as compared to the color singlet mechanism. The hard gluon fragmentation production of a color octet $c\bar{c}$ pair is, on the other hand, of lower order in α_s as compared to the production of a 3S_1 color singlet $c\bar{c}$ pair.

In addition, the color octet fragmentation process is enhanced by the so called trigger bias effect. The fragmentation contribution to the charmonium production cross section can be written as a convolution

$$d\sigma_{\mathcal{O}}(p) = \sum_i \int_0^1 dz d\sigma_i(p/z, \mu) D_{i \rightarrow \mathcal{O}}(z, \mu), \quad (1)$$

of parton production cross sections $d\sigma_i$ and fragmentation functions $D_{i \rightarrow \mathcal{O}}$, describing the fragmentation of a parton i into a charmonium state \mathcal{O} . Since the parton production cross sections fall steeply with the parton transverse momentum p_T/z , fragmentation processes where the charmonium state takes a large fraction z of the parton momentum are favored. In a strict LO calculation the color octet fragmentation function is proportional to $\delta(1-z)$. When leading logs are re-summed, fragmentation functions are softened, but the fact remains that in comparison with other production mechanisms the color octet mechanism is characterized by large z fragmentation.

The arguments in favor of the color octet solution are of a qualitative nature, given a new nonperturbative parameter one can obviously fit to data. It is therefore crucial to find independent means to check the hypotheses by Braaten *et al.*

One way would be to look for other processes where the same color octet

production mechanism could be important, see eg. [12]. Alternatively one might look for characteristics of the color octet production mechanism at the Tevatron. It has eg. been noted that ψ' produced through the color octet mechanism are dominantly transversely polarized [13].

Here we will make use of the fact that a ψ' produced in a fragmentation process will be part of a jet. We investigate the possibilities to extract more information on the fragmentation process by detecting the associated jet. At a superficial level one can identify the jet momentum with the parton momentum and thus get a measure of the fraction z of the parton momentum taken by the ψ' . As noted above, large z fragmentation is a characteristic of the color octet production mechanism.

In [14] it was proposed that trigger bias enhanced higher order corrections might explain the ψ' surplus within the color singlet model. It was shown explicitly that trigger bias enhanced higher order corrections to light quark fragmentation into η_c dominate over the LO result. For ψ and ψ' higher order calculations are however lacking. If this proposal is correct one would expect the fragmentation to be intermediate in hardness between the soft LO color singlet mechanisms and the very hard color octet mechanism.

In order to make a quantitative analysis it is necessary to implement the fragmentation processes in a perturbative QCD cascade, allowing the use of measurable variables such as $z_{\text{jet}} = p_{T\psi'}/p_{T\text{jet}}$.

In this paper we present an implementation of the LO color octet and singlet production mechanisms relevant for ψ' production, but the extension to other mechanisms and other quarkonia is trivial.

In section 2 we describe the implementation of fragmentation production of quarkonium states in a leading-log QCD shower generator based on the Dipole Cascade Model (DCM), first in general and then more specifically for the

LO color octet and singlet production mechanisms relevant for ψ' production.

In section 3 we give some results of our simulations and discuss the possibilities to distinguish between different fragmentation mechanisms based on the shape of fragmentation functions.

Our conclusions are presented in section 4, which is followed by an appendix describing the analytical approximations used to verify our implementation.

2 The Implementation

The fragmentation production of quarkonia may be viewed as a two step process: 1. the *hard* fragmentation into a color singlet or octet $c\bar{c}$ -pair, and 2. the *nonperturbative* formation of the quarkonium state.

Here we implement both these steps in conjunction in a hard QCD shower generator, by including splitting functions not into $c\bar{c}$ -pairs but directly into quarkonium states. Thus, the nonperturbative formation of the quarkonium state is implemented at the perturbative level, before normal hadronization sets in. This is motivated by the fact that all hard interactions are already included in the splitting functions. The $c\bar{c}$ -pairs are thus effectively non-interacting until the hadronization onset scale. Finally, according to BBL factorization [4], the “probability” that a color octet or singlet $c\bar{c}$ -pair form a quarkonium state is given by process independent parameters (matrix elements). BBL factorization is however fully inclusive and does not describe how the $c\bar{c}$ -pair effects the hadronization of the remaining partons. In the color singlet case, the $c\bar{c}$ -pair is color disconnected from the other partons in the event and it is certainly reasonable to let the other partons hadronize independently of the $c\bar{c}$ -pair. The color octet case is more troublesome. In our implementation we enforce the radiation of a soft gluon that inherits

the color connection of the $c\bar{c}$ -pair to the other partons. That is, we simply implement splitting into the quarkonium state plus a soft gluon. The hadronization is done using the stringfragmentation model in JETSET which is infrared safe w.r.t. soft gluons, suggesting insensitivity to the exact modelling of these soft interactions. We will, however, return to this nontrivial question in a forthcoming publication.

In this paper we present an implementation of heavy quarkonium production in the Ariadne program [15] which implements the Dipole Cascade Model (DCM) [16, 17] for QCD cascades.

The DCM differs from conventional parton shower models in that gluon emission is treated as dipole radiation from color dipoles between partons. In eg. e^+e^- annihilation, the first gluon, g_1 , is emitted from the dipole between the initial $q\bar{q}$ pair. An emission of a second softer gluon can then be described in terms of radiation from two independent dipoles, one between the q and g_1 and one between g_1 and \bar{q} . Further gluon emissions are given by three independent dipoles etc.

One problem with this model is that only gluon radiation is modelled and the process of splitting a gluon into a $q\bar{q}$ pair is not included in a natural way. In [18] it was shown that this process can be included without leaving the dipole picture by taking the ordinary splitting kernel for $g \rightarrow q\bar{q}$ and dividing it in two equal parts describing the contribution from each connecting dipole to the splitting separately. This procedure works very well and describes correctly the amount of secondary heavy quarks produced at LEP [19] although some theoretical questions have been raised [20].

The process of $g \rightarrow \mathcal{O} + g$ will here be treated in the same way as the $g \rightarrow q\bar{q}$ one. The splitting kernel $d_{g \rightarrow \mathcal{O}}(z, S)$, which is related to the fragmentation

function through

$$D_{g \rightarrow \mathcal{O}}(z, \mu) = \int_0^{\mu^2} \frac{dS}{S} d_{g \rightarrow \mathcal{O}}(z, S) \quad (2)$$

is hence divided into two equal parts assigned to each of the connecting dipoles. Assuming a dipole between a quark and a gluon in its center of mass system, with the gluon along the positive z -axis, z is then the fraction of the total positive light-cone momenta of the dipole carried by the quarkonium. The partons in the DCM is always on mass shell, therefore, S is no longer the mass square of the split gluon, but is still the mass square of the decay products. Four-momentum is conserved by “borrowing” momentum from the other parton in the dipole – this is a valid approach in the limit of strongly ordered emissions.

For practical reasons, the generations of quarkonium production is done in the variables $m_{\perp}^2 \equiv m_{\mathcal{O}}^2 + p_{\perp}^2 = zS$ and $y = \ln \frac{zW^2}{S}$, the transverse mass and rapidity of the quarkonium in the dipoles center of mass, W is the total mass of the emitting dipole.

For eg. a q - g dipole there are now several competing processes that can occur; a gluon can be emitted according to the standard DCM, the initial gluon can be split into a $q\bar{q}$ pair of different flavors and it can be split into a quarkonium plus gluons according to several different mechanisms. This competition is as usual governed by the Sudakov formfactor mechanism using the transverse momentum squared as ordering variable. Hence the probability $P_i(k_{\perp}^2)$ of the process i to occur at a scale k_{\perp}^2 is given by

$$P_i(k_{\perp}^2) = d_i(k_{\perp}^2) e^{-\int_{k_{\perp}^2}^{k_{\perp}^2, \max} \sum_j d_j(p_{\perp}^2) dp_{\perp}^2} \quad (3)$$

where $d_i(k_{\perp}^2)$ is the splitting kernel with the dependence on rapidity integrated out.

It is now straight forward to implement basically any mechanism for fragmentation production of any heavy quarkonium. In this paper we discuss three mechanisms for ψ' production:

- Color-Octet mechanism $g \rightarrow c\bar{c}(^3S_1^{(8)}) \rightarrow \psi' \pm$ soft gluons. A perturbative transition of a gluon into a collinear $c\bar{c}$ pair in a color octet 3S_1 state, followed by a nonperturbative transition into ψ' involving emission and/or absorption of soft gluons. As discussed above we implement this process as $g \rightarrow \psi' + g$, with one soft final state gluon. Since the gluons are soft, the splitting kernel is essentially a delta-function,

$$d_{g \rightarrow \psi'}(z, S) = \frac{\pi\alpha_s \langle 0 | O_8^{\psi'}(^3S_1) | 0 \rangle}{8m_c} \delta(1-z) \delta(S - 4m_c^2). \quad (4)$$

In the implementation we give for practical reasons the splitting kernel a small width, but we do not in this paper study possible effects of the physical $\mathcal{O}(v^2)$ width [21]. Since the characteristic velocity v of the c -quarks in the charmonium state is not that small ($v^2 \approx 0.2$), the physical width could effect the event structure and through trigger bias enhancement also the normalization of the large p_T quarkonium production. The Color-Octet matrix element $\langle 0 | O_8^{\psi'}(^3S_1) | 0 \rangle$ was taken to be 0.0042 GeV^3 as given by the fit in [11] to CDF data on large p_T ψ' production.

- Color-Singlet mechanism $g \rightarrow c\bar{c}(^3S_1^{(8)}) + 2g \rightarrow \psi' + 2g$ is theoretically simpler since the hard splitting color disconnects the $c\bar{c}$ -pair from the rest of the event. The implementation is complicated, however, due to the final 3-particle phase space. For simplicity we treat also this process as $g \rightarrow \psi' + g$. The error introduced by this simplification, which will be investigated in a future publication, may be assumed to be small since two gluons nearby in phase space are usually well approximated

with one within the framework of Lund string fragmentation. The color-singlet gluon fragmentation function is given as a double integral in [22]. Performing one of the integrals we find an analytical expression for the fictitious $g \rightarrow \psi' + g$ splitting kernel. The color singlet parameter $|R(0)|^2$ relevant also for the last production mechanisms is taken to be 0.231 GeV^3 .

- Color-Singlet mechanism for $c \rightarrow \psi' + c$. Here the whole splitting kernel, given in [23], is assigned to the only dipole connected to the c quark. The scale is the running α_S is, as in the other cases, taken to be m_\perp^2 .

3 What comes out

The ARIADNE program is interfaced to PYTHIA and JETSET [24, 25] for handling of the hard sub-process and hadronization respectively. We use this to generate ψ' production events at the Tevatron. In fig. 1 we see the resulting $p_{T\psi'}$ spectrum for the different production mechanisms. Using the same parameters as in [11], we reproduce approximately their result where the Octet mechanism totally dominates the cross section for high $p_{T\psi'}$, and we also reproduce the measured $p_{T\psi' \rightarrow \mu^+ \mu^-}$ spectrum [1].

The difference between the Octet and Singlet mechanisms is larger in our case than in [6, 11] even after taking into account the small variations in the parameters used. The generated spectra from ARIADNE does, however, agree with the analytical approximation in the appendix at large p_T . At small p_T we do not expect these to agree because there the simple assumption in the analytical calculation that the gluon spectrum behaving like $1/p_T^5$ breaks down. Also effects of initial-state QCD radiation, implemented through the so-called Soft Radiation Model [26] in ARIADNE, are absent in the analytical




Figure 1: *The transverse momentum distribution of $\psi' \rightarrow \mu^+ \mu^-$ production at the Tevatron for different production mechanisms compared to CDF data [1]. Full line is the color octet mechanism, dotted line is the sum of the color singlet mechanisms, and the long-dashed and short-dashed lines are the gluon and c -quark initiated singlet mechanisms respectively. The histograms were generated with ARIADNE and the smooth lines are from the analytical approximation in the Appendix.*

calculation. In addition, ARIADNE implements a more complete treatment of the parton evolution, in contrast to the analytical calculations, which, like earlier numerical calculations, uses the approximative though asymptotically correct AP evolution. The full evolution equation and a discussion of the approximation made in the use of the AP evolution equation can be found in [6].

In fig. 2 we show the generated z -spectrum for $20 < p_{T\psi'} < 25$ GeV, where z is now $p_{T\psi'}/\hat{p}_T$ and \hat{p}_T is the transverse momentum in the hard subprocess. It is clear that the z -spectrum is much harder for the octet mechanism as expected. This is also seen in fig. 3, where the average z is shown as a function of $p_{T\psi'}$. Here we also find a good agreement with the




Figure 2: The z -distribution for $20 < p_{T,\psi'} < 25$ GeV as generated by ARIADNE for different production mechanisms.




Figure 3: The average z as a function of $p_{T,\psi'}$ for different production mechanisms. The smooth lines are from the analytical approximation in the appendix and the (due to low statistics) broken lines were generated with ARIADNE.




Figure 4: *The z distribution for $20 < p_{T\psi'} < 25$ GeV as generated by ARIADNE using the color octet mechanism. Full line is the generated spectrum, long-dashed line is after fragmentation using a cone radius of 1 to define the associated jet, and short-dashed is the same with the addition of the underlying event according to the multiple scattering model in PYTHIA.*

analytical approximation at large $p_{T\psi'}$.

With the availability of ψ' production in an event generator we can now also study hadronization effects. To study eg. the z -spectrum experimentally one could sum up the transverse energy $E_{T\text{jet}}$ in a cone in the pseudorapidity–azimuth plane around the detected ψ' , defining $z_{\text{jet}} = p_{T\psi'}/E_{T\text{jet}}$. In fig. 4 we compare the z -spectrum for such a procedure with the generated spectrum in fig. 2 for the octet mechanism. We note that just adding hadronization, a smearing of the jet is introduced and some of the jet energy falls outside the cone, which makes the spectrum considerably harder. Adding also multiple interactions to model the underlying event [27] in PYTHIA makes it softer again. In fig. 5 we see that the differences in hardness between the octet and singlet mechanisms remains after all hadronization effects although the




Figure 5: *The z distribution for $20 < p_{T\psi'} < 25$ GeV as generated with ARIADNE with different production mechanisms, after hadronization and inclusion of multiple interactions.*

shapes are somewhat distorted.

4 Conclusions

We have implemented different quarkonium production mechanisms in the ARIADNE event generator. Besides giving a more accurate treatment of the initial- and final-state parton evolution than recent analytical approximations, such a generator is an essential tool for understanding experimental corrections and hadronization effects in case one wants to study non-inclusive aspects of high p_T onium production at e.g. the Tevatron.

In interpreting the results it is important to keep in mind the status of the implementations of the different production mechanisms:

- The implementation of the color singlet $c \rightarrow \psi' + c$ fragmentation is unproblematic and trustworthy.

- The color singlet $g \rightarrow \psi' + gg$ fragmentation has been implemented as $g \rightarrow \psi' + g$. The error introduced by this simplification is believed to be small, and in the future the full $g \rightarrow \psi' + gg$ splitting will be implemented.
- The color octet $g \rightarrow c\bar{c}(^3S_1^{(8)}) \rightarrow \psi' \pm$ soft gluons, has been implemented as $g \rightarrow \psi' +$ one soft gluon, and the effects of the physical $\mathcal{O}(v^2)$ width of the fragmentation function has not been investigated. In the future the sensitivity of the results to the modelling of the soft interactions will be studied.

The production mechanisms we have implemented here are, after a simple change of normalization relevant also for ψ production. In this case it is necessary, however, to take into account also contributions from P-wave states decaying to ψ . It is therefore important to implement also P-wave production mechanisms.

There are also other experiments where the fragmentation production of quarkonium states is important. One interesting example is Z^0 decays at LEP. Here the situation is quite different as compared to large p_T production. There are more quarks than gluons available for fragmentation, and there is no trigger bias enhancement. Consequently other production mechanisms might be dominant and measurements would be more sensitive to the low z shape of fragmentation functions.

We have shown that it might be possible to measure the z -distribution of the fragmentation function at the Tevatron and from that learn about the production mechanism.

This result is of course dependent on our very crude approximation of the experimental environment. But with the implementation presented here, the determination of optimal choices of jet definitions and experimental cuts is

clearly facilitated and it does not seem unlikely that the z spectrum, if measured at the Tevatron, could be used to better understand the quarkonium production mechanism.

Acknowledgement. We are grateful for discussions with Paul Hoyer, Michelangelo Mangano and Mikko Vänttinen.

Appendix

Here we present a very simple calculation that exhibits the gross features of charmonium production at large p_T .

We start out from a rapidity integrated version of eq. (1)

$$\frac{d\sigma_{\psi'}(p_T)}{dp_T} = \sum_i \int_0^1 dz \frac{d\sigma_i}{dp_{iT}}(p_T/z, \mu) D_{g \rightarrow \psi'}(z, \mu). \quad (5)$$

We choose the factorization scale as $\mu = p_T/z$, and note that large logs $\log(4m_c^2/\mu^2)$ can be resummed by AP evolution [28] of the fragmentation functions. We calculate the parton production cross sections as convolutions of hard subprocess cross sections ($\propto p_T^{-4}$) and parton distributions, and find that the parton production cross sections fall roughly as p_T^{-5} . Consequently we take $d\sigma_i/dp_{iT} \approx c_i/p_{iT}^5$ in eq. (5) and find

$$\frac{d\sigma_{\psi'}}{dp_T} \approx \sum_i \int_0^1 dz \frac{c_i}{p_T^5} z^5 D_{i \rightarrow \psi'}(z, p_{iT}/z) \approx \sum_i \frac{c_i}{p_T^5} D_{i \rightarrow \psi'}^{(6)}(p_T). \quad (6)$$

Here $D_{i \rightarrow \psi'}^{(6)}(\mu) \equiv \int_0^1 dz z^5 D_{i \rightarrow \psi'}(z, \mu)$ is the sixth moment of the fragmentation function. In eq. (6) we have neglected the evolution of the fragmentation functions between $\mu = p_T$ and p_T/z , implicitly assuming that the dominant contributions to the integral in eq. (6) come from large values of z .

We find $c_g \approx 7 \cdot 10^9 \text{GeV}^4 \text{nb}$, and $c_c \approx 4 \cdot 10^7 \text{GeV}^4 \text{nb}$, to LO in α_s . The gluon production cross section is two orders of magnitude larger than the quark production cross section. Within the color singlet model to LO in α_s , the c quark fragmentation function is, however, much larger than the gluon fragmentation function. We keep therefore both gluon and c quark fragmentation in eq. (6).

In [6] mixing effects were taken into account by including NLO corrections to the hard scattering charm production cross section. Here as well as in our

QCD shower generator implementation, we include mixing explicitly and use the LO charm production cross section to avoid double counting.

For moments ($n > 1$) the AP evolution equations can be solved analytically. The solution has the form:

$$D_{g \rightarrow \psi'}^{(n)}(\mu) = \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{d_g^{(n)}} \left((1 - \epsilon^{(n)}) D_{g \rightarrow \psi'}^{(n)}(\mu_0) - \delta^{(n)} D_{c \rightarrow \psi'}^{(n)}(\mu_0) \right) + \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{d_c^{(n)}} \left(\epsilon^{(n)} D_{g \rightarrow \psi'}^{(n)}(\mu_0) + \delta^{(n)} D_{c \rightarrow \psi'}^{(n)}(\mu_0) \right) \quad (7)$$

$$D_{c \rightarrow \psi'}^{(n)}(\mu) = \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{d_c^{(n)}} \left((1 - \epsilon^{(n)}) D_{c \rightarrow \psi'}^{(n)}(\mu_0) + \rho^{(n)} D_{g \rightarrow \psi'}^{(n)}(\mu_0) \right) + \left(\frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{d_g^{(n)}} \left(\epsilon^{(n)} D_{c \rightarrow \psi'}^{(n)}(\mu_0) - \rho^{(n)} D_{g \rightarrow \psi'}^{(n)}(\mu_0) \right), \quad (8)$$

with $\rho^{(n)} \delta^{(n)} = \epsilon^{(n)}(1 - \epsilon^{(n)})$. For the moments n relevant here, the mixing parameters $\rho^{(n)}$, $\delta^{(n)}$ and $\epsilon^{(n)}$ are all small, and we may put $\rho^{(n)}$ and $\epsilon^{(n)}$ to zero. Within the color singlet model the $\delta^{(n)} D_{c \rightarrow \psi}^{(n)}$ terms in eq. (7) are important however since as noted above the c quark fragmentation function is much larger than the gluon fragmentation function. For simplicity we choose the initial scales for charm and gluon fragmentation and the mixing onset scale all to $2m_c$ and ignore flavor thresholds (refinements are simple). Combining eq. (6),(7) and (8) we find

$$\frac{d\sigma_{\psi'}}{dp_T} \approx \frac{c_g}{p_T^5} \left(\frac{\alpha_s(p_T)}{\alpha_s(2m_c)} \right)^{d_g^{(6)}} \left(D_{g \rightarrow \psi'}^{(6)}(2m_c) - \delta^{(6)} D_{c \rightarrow \psi'}^{(6)}(2m_c) \right) + \frac{c_c + \delta^{(6)} c_g}{p_T^5} \left(\frac{\alpha_s(p_T)}{\alpha_s(2m_c)} \right)^{d_c^{(6)}} D_{c \rightarrow \psi'}^{(6)}(2m_c). \quad (9)$$

According to the hypotheses by Braaten *et al.* [11] the color octet mechanism gluon fragmentation completely dominates ψ' production. We may then of course ignore charm fragmentation all together in eq. (9).

Similarly we can calculate the mean value of z in the convolution in eq. (5). Superficially identifying z with $p_{T\psi'}/p_{T\text{jet}}$, this gives us a hint of the possibilities to use the form of fragmentation functions to distinguish between different production mechanisms. More importantly, it can be used to check our QCD cascade implementation of quarkonium production.

For convenience we give numerical values for the moment evolution parameters $\delta^{(n)}$, $d_g^{(n)}$, $d_c^{(n)}$ for 4 (5) quark flavors, as well as the values we have used for the moments of the color singlet fragmentation functions $D_{c\rightarrow\psi'}^{(n)}$ and $D_{g\rightarrow\psi'}^{(n)}$:

n	5	6	7
$\delta^{(n)}$	0.030 (0.028)	0.023 (0.022)	0.018 (0.017)
$d_g^{(n)}$	2.18 (2.46)	2.46 (2.76)	2.68 (3.00)
$d_c^{(n)}$	0.97 (1.05)	1.08 (1.17)	1.17 (1.27)
$D_{c\rightarrow\psi'}^{(n)}(2m_c)$	$1.2 \cdot 10^{-5}$	$9.7 \cdot 10^{-6}$	$7.7 \cdot 10^{-6}$
$D_{g\rightarrow\psi'}^{(n)}(2m_c)$	$1.8 \cdot 10^{-7}$	$1.3 \cdot 10^{-7}$	$1.1 \cdot 10^{-7}$

For the moments of the color octet gluon fragmentation function we have used the value $D_{g\rightarrow\psi'}^{(n)}(2m_c) \approx 4.2 \cdot 10^{-5}$, independent of the moment n .

References

- [1] CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **69** (1992) 3704;
CDF Collaboration, V. Papadimitriou *et al.*, Fermilab-Conf-95/128-E.
- [2] D0 Collaboration, S. Abachi *et al.*, Fermilab-Conf-95/205-E and
Fermilab-Conf-95/206-E.
- [3] E. Braaten and T.C. Yuan, *Phys. Rev. Lett.* **71** (1993) 1673.
- [4] G.T. Bodwin, E. Braaten and G.P. Lepage, *Phys. Rev.* **51** (1995) 1125.

- [5] R. Baier and R. Rückl, *Z. Phys.* **C19** (1983) 251, and references therein.
- [6] E. Braaten, M.A. Doncheski, S. Fleming and M. Mangano, *Phys. Lett.* **B333** (1994) 548;
- [7] D.P. Roy and K. Sridhar, *Phys. Lett.* **B339** (1994) 141.
- [8] G. A. Schuler, CERN-TH.7170/94, hep-ph/9403387, to appear in *Phys. Rep.*
- [9] M. Vanttinen, P. Hoyer, S.J. Brodsky and W.-K. Tang, *Phys. Rev.* **D51** (1995) 3332.
- [10] M. Cacciari and M. Greco, *Phys. Rev. Lett.* **73** (1994) 1586;
- [11] E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74** (1995) 3327.
- [12] E. Braaten and Y.-Q. Chen, *Phys. Rev. Lett.* **76** (1996) 730.
- [13] P. Cho and M.B. Wise, *Phys. Lett.* **B346** (1995) 129;
M. Beneke and I.Z. Rothstein, *Phys. Lett.* **B372** (1996) 157.
- [14] P. Ernström, P. Hoyer, M. Vanttinen, NORDITA - 96/3 P,
hep-ph/9601333, to appear in *Z. Phys. C*.
- [15] L. Lönnblad, *Comput. Phys. Commun.* **71** (1992) 15. The implementation of quarkonium production will be available as of release 4.09 of the ARIADNE program.
- [16] G. Gustafson, *Phys. Lett.* **B175** (1986) 453.
- [17] G. Gustafson, U. Pettersson, *Nucl. Phys.* **B306** (1988) 746.
- [18] B. Andersson, *et al.*, *Nucl. Phys.* **B339** (1990) 393.
- [19] OPAL Collaboration, R. Akers *et al.*, *Phys. Lett.* **B353** (1995) 163.
- [20] M.H. Seymour, *Nucl. Phys.* **B436** (1995) 163.
- [21] E. Braaten, S. Fleming and T. C. Yuan, To appear in *Ann. Rev. Nucl. Part. Sci.* hep-ph/9602374.

- [22] E. Braaten and T.C. Yuan, *Phys. Rev.* **D52** (1995) 6627,
hep-ph/9507398.
- [23] E. Braaten, K. Cheung and T.C. Yuan *Phys. Rev.* **D48** (1993) 4230.
- [24] T. Sjöstrand, CERN-TH.7112/93, December 1993.
- [25] T. Sjöstrand, *Comput. Phys. Commun.* **82** (1994) 74.
- [26] B. Andersson *et al.*, *Z. Phys.* **C43** (1989) 625.
- [27] T. Sjöstrand, M. van Zijl, *Phys. Rev.* **D36** (1987) 2019.
- [28] G. Altarelli and G. Parisi, *Nucl. Phys.* **B126** (1977) 298.