

J/ψ Production: Tevatron and Fixed-Target Collisions

A. Petrelli^{a*}

^aArgonne National Laboratory
9700 S. Cass Avenue, Argonne, IL 60439 USA

In this talk I show the results of a fit of the NRQCD matrix elements to the CDF data for direct J/ψ production, by including the radiative corrections to the $gg \rightarrow {}^3S_1^{[1]}g$ channel and the effect of the k_T -smearing. Furthermore I perform the NLO NRQCD analysis of J/ψ production in fixed-target proton-nucleon collisions and I fit the colour-octet matrix elements to the available experimental data. The results are compared to the Tevatron ones.

1. INTRODUCTION

The J/ψ production cross-section within the NRCQD factorization theory [1] is given by the expression:

$$d\sigma^{J/\psi} = \sum_n \langle 0 | \mathcal{O}^{J/\psi}[n] | 0 \rangle d\hat{\sigma}(c\bar{c}[n]) \quad (1)$$

where $n = {}^{2S+1}L_J^{[1,8]}$. The relevant long-distance matrix elements up to order v^4 are $\langle \mathcal{O}_8^{J/\psi}({}^3S_1) \rangle$ and the linear combination $\Delta_8^{J/\psi}(k) = \langle \mathcal{O}_8^{J/\psi}({}^1S_0) \rangle + k \langle \mathcal{O}_8^{J/\psi}({}^3P_0) \rangle / m^2$. The phenomenological consistency of the NRQCD factorization formalism rests upon the universality of the long-distance matrix elements. The phenomenological determination of the non-perturbative matrix elements relies on the accuracy in the computation of the short distance kernels. In this talk I consider the Tevatron VS fixed-target universality issue. First I perform a fit of $\langle \mathcal{O}_8^{J/\psi}({}^3S_1) \rangle$ and $\Delta_8^{J/\psi}$ (3.5) matrix elements to the Tevatron CDF data [2] by considering two possible deviation from the standard fits [3–5], namely a) the $O(\alpha_s^4)$ colour-singlet contribution and b) the effect of the k_T -smearing. I successively perform a fit of the MEs to a wide fixed-target data sample based upon a NLO QCD analysis.

2. THE COLOUR-SINGLET $O(\alpha_s^4)$ CONTRIBUTION

The equation (1) is usually interpreted as a double expansion in the strong coupling α_s and the velocity v . In the J/ψ p_T -differential cross-section a third expansion parameter has to be considered, and specifically $1/p_T$ [6]. In the triple-expansion paradigm It is straightforward to realize that the process $ij \rightarrow {}^3S_1^{[1]}kl$ is a priori large. The scaling of its partonic cross-section is in fact $O(\alpha_s^4 v^0 / p_T^2)$ as compared to $O(\alpha_s^3 v^4 / p_T^6)$ of the C-even colour-octet configurations (${}^1S_0^{[8]}$ and ${}^3P_J^{[8]}$). In this section I will show the effect of the channels $ij \rightarrow {}^3S_1^{[1]}kl$ in the extraction of the colour-octet matrix elements at the Tevatron. The details of the calculation will appear in a forthcoming paper [7]. In the present document I just confine myself to draw the general lines of the computation. The three $O(\alpha_s^4)$ channels I am going to consider are:

$$q\bar{q} \rightarrow {}^3S_1^{[1]}gg, \quad (2)$$

$$qg \rightarrow {}^3S_1^{[1]}qg, \quad (3)$$

$$gg \rightarrow {}^3S_1^{[1]}gg. \quad (4)$$

The one-loop colour-singlet channel is presently unknown for hadroproduction of J/ψ . It is only available for J/ψ photoproduction [8] and annihilation into light hadrons [9].

Nevertheless, scaling arguments show that the virtual channel gives a subleading contribution

*Talk given at the QCD99 Euroconference, Montpellier, France, July 1999. Report No. ANL-HEP-CP-99-107.




Figure 1. Different channels contributing to the J/ψ production at the Tevatron. The NLO ${}^3S_1^{[1]}$ channel is given by the dashed curves ($s_{\min} = M_{J/\psi}^2$ (lower dash) and $s_{\min} = M_{J/\psi}^2/20$ (upper dash)). The fit of the colour-octet MEs to data are performed by considering the upper dashed curve. The resulting fitted curves are shown (${}^3S_1^{[8]}$ (dots) and $\Delta_8^{J/\psi}(3.5)$ (dotdash)).

at high p_T , being $O(1/p_T^8)$ its fall-off (same as the born, which is already known to be negligible). The tree-level QED-like diagrams in the channel (4) are also suppressed at high p_T , but they have been included to double check the global gauge invariance of the process. The omission of the abelian diagrams would generate a $1/p_T^2$ -suppressed gauge dependence. To evaluate the amplitudes relative to the processes (2)-(4) I make use of the covariant projection technique [11,12]. The evaluation of the channels (2) and (3) is straightforward. The process (4) demands the helicity amplitude formalism:

$$\begin{aligned} \mathcal{M}(P^\epsilon, k_1^{h_1}, k_2^{h_2}, k_3^{h_3}, k_4^{h_4}) = \\ \frac{\delta_{ij}}{\sqrt{N}} \text{Tr} \{ (\not{P} + M) \gamma^\alpha \mathcal{A}_{ij}^{\mu_1 \mu_2 \mu_3 \mu_4} \} \times \\ \mathcal{E}_\alpha^\epsilon(P) \epsilon_{\mu_1}^{h_1}(k_1) \epsilon_{\mu_2}^{h_2}(k_2) \epsilon_{\mu_3}^{h_3}(k_3) \epsilon_{\mu_4}^{h_4}(k_4) \end{aligned} \quad (5)$$

I use the Calcult collaboration representation for

the external gluons [13]

$$\begin{aligned} \not{\epsilon}^\pm(k, p, q) = \frac{1}{[8(k \cdot p)(k \cdot q)(p \cdot q)]^{1/2}} \times \\ [\not{k} \not{p} \not{q} (1 \mp \gamma_5) + \not{q} \not{p} \not{k} (1 \pm \gamma_5) - 2(p \cdot q) \not{k}] \end{aligned} \quad (6)$$

	No NLO Sing	NLO Sing
$\langle \mathcal{O}_8^{J/\psi}({}^3S_1) \rangle$	1.4 ± 0.26	1.5 ± 0.26
$\Delta_8^{J/\psi}(3.5)$	12.5 ± 2.8	9.6 ± 2.8

Table 1

NLO colour-singlet effect in the colour-octet MEs extraction at the Tevatron. The values are expressed in units of 10^{-2} GeV^3 . In the first column is reported the standard fit. The second one shows the fit obtained by considering NLO ${}^3S_1^{[1]}$ contribution with a democratic cut on any jet pair $s_{ij} > s_{\min} = M_{J/\psi}^2/20$. The latter effect lowers the value of $\Delta_8^{J/\psi}(3.5)$ but leaves $\langle \mathcal{O}_8^{J/\psi}({}^3S_1) \rangle$ essentially unchanged.

which corresponds to the choice of a light-like axial gauge, being k the momentum of the given gluon and p, q the two light-like reference momenta. There are three independent helicity configurations, namely $(+, +, -, -)$, $(+, +, +, -)$, $(+, +, +, +)$. In particular the all-plus configuration turns out to be zero. Each helicity amplitude is expanded in terms of the six colour structures represented by the six traces $\text{tr}(\lambda^{a_1} \lambda^{a_2} \lambda^{a_3} \lambda^{a_4})$. When the square is performed the C-parity symmetry allows the overall factorization of the colour.

Being the virtual-emission channel missing, in performing the numerical analysis one is forced to put phenomenological cuts to avoid collinear and soft regions of the phase space. In particular: a) azimuth-pseudorapidity separation cut between the two final jets, b) p_T cut on the two final jets. Another way to proceed is to put a minimum invariant mass cut on any jet pair: $s_{ij} = (k_i + k_j)^2 > s_{\min}$. I choose the latter way. The




Figure 2. Different contributions to J/ψ production at the Tevatron. Dots: ${}^3S_1^{[8]}$. Dashes: ${}^1S_0^{[8]} + {}^3P_J^{[8]}$. Dotdash: NLO ${}^3S_1^{[1]}$. The effect of k_T -smearing is included for three different values of $\langle k_T \rangle$ ($\langle k_T \rangle = 0, 1, 1.5$ GeV). The results are given for three different sets of pdfs. The NLO ${}^3S_1^{[1]}$ effect is only included in the $\langle k_T \rangle = 0$ case.

		MRS(R2)	MRS(A)	CTEQ4M
$\Delta_8^{J/\psi}(3.5)$	$\langle k_T \rangle = 0$	9.6 ± 2.8	19.7 ± 3.7	11.9 ± 2.8
	$\langle k_T \rangle = 1$	7.7 ± 2.0	14.8 ± 2.7	8.6 ± 2.1
	$\langle k_T \rangle = 1.5$	4.1 ± 1.4	8.4 ± 1.9	4.5 ± 1.5
$\langle \mathcal{O}_8^{J/\psi}({}^3S_1) \rangle$	$\langle k_T \rangle = 0$	1.5 ± 0.26	1.5 ± 0.26	1.5 ± 0.26
	$\langle k_T \rangle = 1$	1.6 ± 0.26	1.7 ± 0.26	1.5 ± 0.22
	$\langle k_T \rangle = 1.5$	1.7 ± 0.19	1.9 ± 0.23	1.7 ± 0.19

Table 2

Effects of intrinsic transverse momentum in the colour-octet MEs fit in J/ψ production at the Tevatron. Fits are performed for three different values of $\langle k_T \rangle$ and three pdfs. Values in units of 10^{-2} GeV³.




Figure 3. Fits of the matrix element $\Delta_8^{J/\psi}(6.4)$ to the fixed-target proton-nucleon collisions data. Fits are performed for three sets of pdfs. The value of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ is taken from the above Tevatron fits (for any correspondent pdf) with $\langle k_T \rangle = 0$. (The value of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ is only slightly affected by the intrinsic k_T anyway. The indirect impact of the Tevatron k_T -smearing in the extraction of $\Delta_8^{J/\psi}(6.4)$ is not appreciable).

figure (1) shows the cut dependence of the process for $M_\psi^2/20 < s_{\min} < M_\psi^2$. The NLO colour-singlet contribution is given by the dashed lines for $s_{\min} = M_\psi^2$ (lower dashes) and $s_{\min} = M_\psi^2/20$ (upper dashes). The colour-singlet matrix element² is set to $\langle \mathcal{O}_1^{J/\psi}(^3S_1) \rangle = 1.2 \text{ GeV}^3$. As p_T increases, the cut sensitivity becomes milder and milder, like expected. The fit of the colour-octet MEs is performed by assuming $s_{\min} = M_\psi^2/20$ and compared to the standard fit (that is without $O(\alpha_s^4)$ correction). The results are summarized in the table (1). The NLO colour-singlet corrections lower the value of the matrix element $\Delta_8^{J/\psi}(3.5)$. The VEV $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ instead is nailed by the high p_T tail of the data distribution and is quite insensitive to the low p_T effects. At high p_T the $^3S_1^{[8]}$ channel develops large collinear logarithms $(\alpha_s \log p_T/2m)^n$ which make the fixed order cross section unreliable. The leading logarithms are resummed by using the standard DGLAP equation for the fragmentation function of the gluon into J/ψ . The accuracy of the cross section in the whole Tevatron p_T -range

is achieved by matching the fixed order to the fragmentation cross-section, according to the following equation:

$$\frac{d\sigma}{dp_T^2}(^3S_1^{[8]}) = \frac{d\sigma^{\text{FXD}}}{dp_T^2} - \frac{d\sigma^{\text{ASY}}}{dp_T^2} + \frac{d\sigma^{\text{FRG}}}{dp_T^2} \quad (7)$$

being

$$\frac{d\sigma^{\text{FRG}}}{dp_T^2} = \frac{d\hat{\sigma}_g}{dp_{T,g}^2} \otimes D_{g \rightarrow \psi} \quad (8)$$

$$\frac{d\sigma^{\text{ASY}}}{dp_T^2} = \left. \frac{d\sigma^{\text{FXD}}}{dp_T^2} \right|_{p_T \gg m}, \quad (9)$$

where the meaning of the symbols is transparent. From the table (1) can be deduced that the inclusion of the $O(\alpha_s^4)$ $^3S_1^{[1]}$ contribution in the J/ψ production at the Tevatron does not affect in a dramatic way the extraction of the colour-octet MEs.

3. THE EFFECT OF THE INTRINSIC k_T

The effect of the intrinsic transverse momentum of the partons in the J/ψ differential cross-section at the Tevatron is phenomenologically implemented by performing a gaussian smear-

²For the the colour-singlet operator I use the original normalization defined in the ref. [1]

	MRS(R2)	MRS(A)	CTEQ4M
$\Delta_8^{J/\psi}(6.4)$	1.0	1.8	1.1

Table 3

Values of the fitted $\Delta_8^{J/\psi}(6.4)$ at fixed-target collisions for different pdfs. In units of 10^{-2} GeV^3 .

ing of the p_T -distributions. The smearing is implemented channel by channel for three values of $\langle k_T \rangle = \langle k_T^2 \rangle^{1/2}$, namely $\langle k_T \rangle = 0, 1, 1.5 \text{ GeV}$ and for three pdf parameterizations (CTEQ4M, MRS(A) and MRS(R2)). The complete NLO calculation of colour-octet channels [10] shows that the Sudakov effect is likely confined below the 2 GeV p_T -region. The region we are analysing is therefore free of Sudakov effects. The figure (2) synthesizes the results of the Tevatron fits with both k_T -smearing and colour-singlet radiative corrections. Since the k_T -smearing essentially attacks the p_T -slope, the colour-octet C-even channels are stronger affected than the flatter $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ distribution, which is instead only slightly sensitive to the transverse momentum of the partons. The basic effect of the k_T -kick is to tilt clockwise the dashed curves in figure (2) and eventually lower the fitted value of the matrix element $\Delta_8^{J/\psi}(3.5)$. The obtained fits are also summarized in the table (2). The lack of accuracy of the NLO colour-singlet cross section at low p_T (due again to the fact that the one-loop channel is still unknown) might affect the shape at intermediate p_T once the k_T -smearing is turned on. That is why the NLO colour-singlet channel is only present in the $k_T = 0$ case.

4. FIXED-TARGET

In this section I perform the fit of NRQCD MEs to a compilation of fixed-target data by using the NLO QCD cross sections evaluated in the reference [10]. A comprehensive LO analysis can be found in the ref [15]. The references of the experimental data can be found in the papers [14,15]. Note that the quoted experiments do not distinguish the direct J/ψ from the ones coming from

the ψ' and χ_J feed-down. Let me fix the VEVs relative to the feed-down first. For the χ_J feed-down I choose $\langle \mathcal{O}_1^{x_0}(^3P_0) \rangle / m^2 = 4.4 \times 10^{-2} \text{ GeV}^3$ and $\langle \mathcal{O}_8^{x_0}(^3S_1) \rangle = 3.2 \times 10^{-3} \text{ GeV}^3$. For the ψ' : $\langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle = 4.4 \times 10^{-3} \text{ GeV}^3$ and $\Delta_8^{\psi'}(6.4) = 2.0 \times 10^{-3} \text{ GeV}^3$ (the latter number is a result of an independent fit that will be shown somewhere; the pdf-dependence of the ψ' VEVs is not considered here since its effect on the J/ψ cross section is negligible). Once the MEs relative to the χ_J and ψ' feed-down have been fixed, I focus on the direct component of J/ψ production. The cross section for direct J/ψ production according to the NRQCD factorization formalism is expressed by the formula:

$$\begin{aligned} \sigma(J/\psi) = & \quad (10) \\ & \hat{\sigma}(^3S_1^{[1]}) \frac{\langle \mathcal{O}_1^\psi(^3S_1) \rangle}{m^5} + \hat{\sigma}(^3S_1^{[8]}) \frac{\langle \mathcal{O}_8^\psi(^3S_1) \rangle}{m^5} + \\ & \hat{\sigma}(^1S_0^{[8]}) \frac{\langle \mathcal{O}_8^\psi(^1S_0) \rangle}{m^5} + \hat{\sigma}(^3P_J^{[8]}) \frac{\langle \mathcal{O}_8^\psi(^3P_0) \rangle}{m^7} \end{aligned}$$

which is accurate up to order four in the velocity expansion. The second line of the previous equation can be rewritten as

$$\sigma = \frac{\hat{\sigma}(^1S_0^{[8]})}{m^5} \left(\langle \mathcal{O}_8^\psi(^1S_0) \rangle + k(E_{\text{beam}}) \frac{\langle \mathcal{O}_8^\psi(^3P_0) \rangle}{m^2} \right)$$

The coefficient $k(E_{\text{beam}})$ is independent of E_{beam} at LO ($k(E_{\text{beam}}) = 7$ at LO) and mildly dependent on E_{beam} at NLO: its average value is around 6.4 ($k(100 \text{ GeV}) = 6.6$, $k(1500 \text{ GeV}) = 6.3$). In fixed-target collisions the matrix element $\Delta_8^{J/\psi}(k)$ appears in a linear combination which is different from the Tevatron one. On the other hand at fixed-target it is not possible to fit simultaneously $\Delta_8^{J/\psi}(6.4)$ and $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ since the all channels have essentially the same shape in E_{beam} . Therefore –following the procedure adopted in the ref [15]– I use the value of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ fitted at Tevatron and extract $\Delta_8^{J/\psi}(6.4)$ from the fixed-target data. In particular I pick the values of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ obtained from the Tevatron at $k_T = 0$ (we have seen that $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ is not sensitive to the intrinsic k_T anyway) and I fit $\Delta_8^{J/\psi}(6.4)$ for three different

pdfs. The fitted curves are shown in the figure (3). The table (3) reports the obtained values for $\Delta_8^{J/\psi}(6.4)$. The NLO QCD corrections lower by about a factor of two the fitted value of $\Delta_8^{J/\psi}(6.4)$ at fixed-target.

5. CONCLUSIONS

Both the radiative corrections to the $^3S_1^{[1]}$ channel and the k_T -kick lower the value of $\Delta_8^{J/\psi}(3.5)$ extracted at the Tevatron. The previous effects vice versa don't have a significant impact on the determination of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$. On the other hand the value of $\Delta_8^{J/\psi}(6.4)$ obtained by fitting the fixed-target data is still sensibly lower than $\Delta_8^{J/\psi}(3.5)$. If one believes that the NRQCD MEs are positive then $\Delta_8^{J/\psi}(6.4) > \Delta_8^{J/\psi}(3.5)$ should hold. Probably the gap would be partially bridged by the inclusion of the $O(\alpha_s^4)$ radiative corrections also for the colour-octet channels the Tevatron. The large theoretical uncertainties in the evaluation of the charmonium total cross-section certainly affect the reliability of the MEs extracted from fixed-target experiments. Even if one does not rely in the current understanding of the mechanisms of charmonium production at low- p_T , the reduction of the Tevatron colour-octet MEs is still welcome in the Hera-Tevatron universality perspective [16].

6. ACKNOWLEDGEMENTS

The results presented here have been obtained with the collaboration of F. Maltoni and M. L. Mangano. This work is supported in part by the U.S. Department of Energy, High Energy Physics Division, Contract W-31-109-Eng-38.

REFERENCES

1. G.T. Bodwin, E. Braaten and G.P. Lepage, Phys. Rev. **D51** (1995) 1125, erratum *ibid.* **D55** (1997) 5853.
2. CDF Collaboration, F. Abe et al., Phys. Rev. Lett. **79** (1997) 578.
3. P. Cho and A.K. Leibovich, Phys. Rev. **D53** (1996) 150; Phys. Rev. **D53** (1996) 6203.
4. M. Cacciari, M. Greco, M.L. Mangano and A. Petrelli, Phys. Lett. **B356** (1995) 560.
5. M. Beneke and M. Krämer, Phys. Rev. **D55** (1997) 5269.
6. E. Braaten and T.C. Yuan, Phys. Rev. Lett. **71** (1993) 1673.
7. F. Maltoni and A. Petrelli, *embryonal*.
8. M. Krämer, Nucl. Phys. **B458** (1996) 3.
9. P. B. MacKenzie and G. P. Lepage, Phys. Rev. Lett. **47** (1981) 1244.
10. A. Petrelli, M. Cacciari, M. Greco, F. Maltoni and M.L. Mangano, Nucl. Phys. **B514** (1998) 245.
11. J.H. Kühn, J. Kaplan and E.G.O. Safiani, Nucl. Phys. **B157** (1979) 125.
12. B. Guberina, J.H. Kühn, R.D. Peccei and R. Rückl, Nucl. Phys. **B174** (1980) 317.
13. P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, Phys. Lett. **B105** (1981) 215.
14. G.A. Schuler, CERN-TH.7170 (1994), [hep-ph/9403387](#), to appear in Phys. Rep. C.
15. M. Beneke and I.Z. Rothstein, Phys. Rev. **D54** (1996) 2005, erratum *ibid.* **B389** (1996) 769.
16. M. Cacciari and M. Krämer, Phys. Rev. Lett. **76** (1996) 4128; P. Ko, J. Lee and H. S. Song, Phys. Rev. **D54** (1996) 4312; B. A. Kniehl and G. Kramer, Eur.Phys.J. **C6** (1999) 493.