

Fractional Quantum Mechanics and Lévy Path Integrals

Nikolai Laskin*

Isotrace Laboratory, University of Toronto
60 St. George Street, Toronto, ON M5S 1A7
Canada

Abstract

The fractional quantum and statistical mechanics have been developed via new path integrals approach.

PACS number(s): 03.65.-w, 05.30.-d, 05.40. Fb, 03.65. Db

From a functional integration point of view the quantum mechanics in Feynman's path integrals treatment [1], [2] is the theory of functionals on measure generated by the Brownian motion-Wiener stochastic process [3], [4]. The Wiener process is the Markov, Gaussian or normal stochastic process the statistical description of which is given by well-known diffusion equation [4].

The natural generalization of the Brownian motion is the Lévy flights (or Lévy motion). The mathematical foundation of this generalization is the theory of stable probability distributions developed by Lévy [5]. The most fundamental property of the Lévy distributions is the stability under addition, following from the generalized central limit theorem valid for Lévy distributions. So, from the probability theory point of view the stable probability law is the generalization of well-known Gaussian law.

Lévy flights is widely used to model a variety processes such as anomalous diffusion [6], turbulence [7], chaotic dynamics [8], plasma physics [9],

*E-mail: nlaskin@rocketmail.com

financial dynamics [10], biology and physiology[11]. The most recent studies are related to fractional Lévy motion [12] which is the generalization of the fractional Brownian motion [6].

This paper deals with new path integrals over measures induced by the stochastic process of Lévy flights. I have generalized the Feynman and Wiener path integrals and developed the new fractional quantum mechanics (fQM) and fractional statistical mechanics (fSM). The fQM and fSM include the Feynman quantum and statistical mechanics in similar way as the Lévy process (Lévy flights) generalizes the Wiener process (Brownian motion).

In this paper I have developed the foundations of fQM and fSM in their functional formulations. As the next step we should think about the concrete fQM problems such as, for example, fractional harmonic oscillator, fractional hydrogen atom, fractional scattering theory, relativistic fractional quantum mechanics, fractional quantum gauge field theory etc.

1. The Feynman quantum mechanics deals with Feynman's quantum mechanical functionals $K_F(x_b t_b | x_a t_a)$ [1]

$$K_F(x_b t_b | x_a t_a) = \int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}_{Feynman} x(\tau) \cdot \exp\left\{\frac{i}{\hbar} \int_{t_a}^{t_b} d\tau V(x(\tau))\right\}, \quad (1)$$

where $V(x(\tau))$ is the potential energy as a functional of a particle path $x(\tau)$ and the Feynman functional measure is defined as

$$\int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}_{Feynman} x(\tau) \dots = \lim_{N \rightarrow \infty} \int dx_1 \dots dx_{N-1} \left(\frac{2\pi i \hbar \varepsilon}{m}\right)^{-N/2} \times \quad (2)$$

$$\times \prod_{j=1}^N \exp\left\{\frac{im}{2\hbar \varepsilon} (x_j - x_{j-1})^2\right\} \dots,$$

here m is the mass of the quantum mechanical particle, \hbar is the Planck's constant, $x_0 = x_a$, $x_N = x_b$ and $\varepsilon = (t_b - t_a)/N$.

I propose the fractional quantum mechanics based on the new path integral

$$K_L(x_b t_b | x_a t_a) = \int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}_{Laskin} x(\tau) \cdot \exp\left\{\frac{i}{\hbar} \int_{t_a}^{t_b} d\tau V(x(\tau))\right\}, \quad (3)$$

where $V(x(\tau))$ is the potential energy as a functional of the Lévy particle path and the Laskin path integral measure is defined as

$$\begin{aligned} & \int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}_{Laskin} x(\tau) \dots = \\ & = \lim_{N \rightarrow \infty} \int dx_1 \dots dx_{N-1} \left(\frac{iD_\alpha \varepsilon}{\hbar}\right)^{-N/\alpha} \cdot \prod_{j=1}^N L_\alpha \left\{ \left(\frac{\hbar}{iD_\alpha \varepsilon}\right)^{1/\alpha} |x_j - x_{j-1}| \right\} \dots, \end{aligned} \quad (4)$$

where D_α is the generalized "diffusion coefficient", \hbar denotes the Planck's constant, $x_0 = x_a$, $x_N = x_b$, $\varepsilon = (t_b - t_a)/N$, and the Lévy function L_α is expressed in terms of Fox's H functions[13], [14], [15]

$$\begin{aligned} & \left(\frac{D_\alpha t}{\hbar}\right)^{-1/\alpha} L_\alpha \left\{ \left(\frac{\hbar}{D_\alpha t}\right)^{1/\alpha} |x| \right\} = \\ & = \frac{\pi}{\alpha |x|} H_{2,2}^{1,1} \left[\left(\frac{\hbar}{D_\alpha t}\right)^{1/\alpha} |x| \mid \begin{matrix} (1, 1/\alpha), (1, 1/2) \\ (1, 1), (1, 1/2) \end{matrix} \right], \end{aligned} \quad (5)$$

here α is the Lévy index, $0 < \alpha \leq 2$.

The Eqs (3)-(5) define the fractional quantum mechanics via Laskin path integral.

The fractional free particle propagator $K_L^{(0)}(x_b t_b | x_a t_a)$ has the form

$$K_L^{(0)}(x_b t_b | x_a t_a) = \int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}_{Laskin} x(\tau) \cdot 1 = \quad (6)$$

$$= \left(\frac{iD_\alpha(t_b - t_a)}{\hbar} \right)^{-1/\alpha} L_\alpha \left\{ \left(\frac{\hbar}{iD_\alpha(t_b - t_a)} \right)^{1/\alpha} |x_b - x_a| \right\}.$$

It is known that at $\alpha = 2$ the Lévy distribution is transformed to Gaussian and the Lévy flights process is transformed to the process of the Brownian motion. The Eq.(6) in accordance with definition given by the Eq.(5) and the properties of the Fox's function $H_{2,2}^{1,1}$ at $\alpha = 2$ (see, [14], [15]) is transformed to Feynman free particle propagator[1]

$$K_F^{(0)}(x_b t_b | x_a t_a) = \left(\frac{2\pi i \hbar (t_b - t_a)}{m} \right)^{-1/2} \cdot \exp \left\{ \frac{im(x_b - x_a)^2}{2\hbar(t_b - t_a)} \right\}. \quad (7)$$

In terms of Fourier integral (momentum representation) the propagator $K_L^{(0)}(x_b t_b | x_a t_a)$ is written as follows

$$K_L^{(0)}(x_b t_b | x_a t_a) = \frac{1}{2\pi \hbar} \int dp \cdot \exp \left\{ i \frac{p(x_b - x_a)}{\hbar} - i \frac{D_\alpha(t_b - t_a) |p|^\alpha}{\hbar} \right\}. \quad (8)$$

We see that the energy E_p of the fractional quantum mechanical particle with the momentum p is given by

$$E_p = D_\alpha |p|^\alpha. \quad (9)$$

The comparison of the Eq.(7) and the Eq.(8) allows to conclude that at $\alpha = 2$ we should put $D_2 = 1/2m$. Then the Eq.(9) is transformed to $E_p = p^2/2m$.

Using the Eq.(9) we can define the *Laskin functional measure in the momentum representation* by the following way

$$\begin{aligned} & \int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}_{Laskin} x(\tau) \dots = \\ & = \lim_{N \rightarrow \infty} \int dx_1 \dots dx_{N-1} \frac{1}{(2\pi \hbar)^N} \int dp_1 \dots dp_N \cdot \exp \left\{ i \frac{p_1(x_1 - x_a)}{\hbar} - i \frac{D_\alpha \varepsilon |p_1|^\alpha}{\hbar} \right\} \times \dots \end{aligned} \quad (10)$$

$$\times \exp \left\{ i \frac{p_N(x_b - x_{N-1})}{\hbar} - i \frac{D_\alpha \varepsilon |p_N|^\alpha}{\hbar} \right\} \dots$$

here $\varepsilon = (t_b - t_a)/N$.

Taking into account the Eq.(8) it is easy to check on the consistency condition

$$K_L^{(0)}(x_b t_b | x_a t_a) = \int dx' K_L^{(0)}(x_b t_b | x' t') \cdot K_L^{(0)}(x' t' | x_a t_a). \quad (11)$$

The fractional free particle propagator is governed by the fractional Schrödinger equation

$$i\hbar \frac{\partial}{\partial t_b} K_L^{(0)}(x_b t_b | x_a t_a) = -D_\alpha (\hbar \nabla_b)^\alpha K_L^{(0)}(x_b t_b | x_a t_a), \quad (12)$$

where the Riesz quantum fractional derivative $(\hbar \nabla)^\alpha$ is defined through its Fourier transform (see, for example, [16])

$$(\hbar \nabla)^\alpha \dots = -\frac{1}{2\pi\hbar} \int dp e^{i\frac{px}{\hbar}} |p|^\alpha \dots \quad (13)$$

The propagator $K_L(x_b t_b | x_a t_a)$ describes the evolution of the fractional quantum-mechanical system

$$\psi_f(x_b, t_b) = \int dx_a K_L(x_b t_b | x_a t_a) \cdot \psi_i(x_a, t_a), \quad (14)$$

where $\psi_i(x_a, t_a)$ is the fractional wave function of initial (at the $t = t_a$) state and $\psi_f(x_b, t_b)$ is the fractional wave function of final (at the $t = t_b$) state.

Taking into account the Eqs.(3), (14) and the definition (13) we can derive the fractional Schrödinger equation for the wave function of quantum mechanical system

$$i\hbar \frac{\partial \psi}{\partial t} = -D_\alpha (\hbar \nabla)^\alpha \psi + V(x)\psi. \quad (15)$$

It is easy to see that the fractional (or Levy) free particle wave function has the form

$$\psi(x, t) = \frac{1}{2\pi\hbar} \int dp \cdot \exp \left\{ i \frac{px}{\hbar} - i \frac{D_\alpha t |p|^\alpha}{\hbar} \right\}. \quad (16)$$

Thus, the Eqs.(3), (4) and (14), (15) are the basic equations of new fQM. They should be analyzed for the particular fractional quantum systems.

2. In order to develop the fractional statistical mechanics let us go in the previous consideration from imaginary time to "inverse temperature" $\beta = 1/k_B T$, where k_B is the Boltzmann's constant and T is the temperature, $it \rightarrow \beta$. In the framework of traditional functional approach to the statistical mechanics we deal with functionals over the Wiener measure[1], [4], [17]

$$\rho_W(x, \beta|x_0) = \int_{x(0)=x_0}^{x(\beta)=x} \mathcal{D}_{Wiener} x(u) \cdot \exp \left\{ - \int_0^\beta du V(x(u)) \right\}, \quad (17)$$

where $\rho_W(x, \beta|x_0)$ is the density matrix of the statistical system in the external field $V(x)$ and the Wiener measure is given by

$$\begin{aligned} \int_{x(0)=x_0}^{x(\beta)=x} \mathcal{D}_{Wiener} x(u) \dots &= \lim_{N \rightarrow \infty} \int dx_1 \dots dx_{N-1} \left(\frac{2\pi\hbar^2\zeta}{m} \right)^{-N/2} \times \\ &\times \prod_{j=1}^N \exp \left\{ - \frac{m}{2\hbar^2\zeta} (x_j - x_{j-1})^2 \right\} \dots, \end{aligned} \quad (18)$$

here $\zeta = \beta/N$ and $x_N = x$.

The fSM deals with Lévy or fractional density matrix $\rho_L(x, \beta|x_0)$ which is given by

$$\rho_L(x, \beta|x_0) = \int_{x(0)=x_0}^{x(\beta)=x} \mathcal{D}_{Lévy} x(u) \cdot \exp \left\{ - \int_0^\beta du V(x(u)) \right\}, \quad (19)$$

where the Lévy functional measure is defined as follows

$$\int_{x(0)=x_0}^{x(\beta)=x} \mathcal{D}_{Lévy} x(u) \dots = \lim_{N \rightarrow \infty} \int dx_1 \dots dx_{N-1} (D_\alpha \varsigma)^{-N/\alpha} \times \quad (20)$$

$$\prod_{j=1}^N L_\alpha \left\{ \frac{|x_j - x_{j-1}|}{(D_\alpha \varsigma)^{1/\alpha}} \right\} \dots,$$

here $\varsigma = \beta/N$, $x_N = x$ and the Lévy function L_α is given by the Eq.(5).

It was shown by Lévy [18] (and later by Doob [19]) that the Eq.(20) defines the functional measure in the space \mathfrak{D} of left (or right) continuous functions having only discontinuities of the first kind.

In Fourier space the fractional density matrix $\rho_L^{(0)}(x, \beta|x_0)$ of a free particle ($V = 0$) has the form

$$\rho_L^{(0)}(x, \beta|x_0) = \frac{1}{2\pi\hbar} \int dp \exp \left\{ i \frac{p(x - x_0)}{\hbar} - \beta D_\alpha |p|^\alpha \right\}. \quad (21)$$

When $\alpha = 2$ and $D_2 = 1/2m$ the Eq.(21) gives the well-known density matrix for one-dimensional free particle [17]

$$\rho_W^{(0)}(x, \beta|x_0) = \left(\frac{m}{2\pi\hbar^2\beta} \right)^{1/2} \exp \left\{ -\frac{m}{2\hbar^2\beta} (x - x_0)^2 \right\}. \quad (22)$$

The density matrix $\rho_L(x, \beta|x_0)$ obeys the fractional differential equation

$$-\frac{\partial \rho_L(x, \beta|x_0)}{\partial \beta} = -D_\alpha (\hbar \nabla_x)^\alpha \rho_L(x, \beta|x_0) + V(x) \rho_L(x, \beta|x_0). \quad (23)$$

The Eqs. (19) - (21) and (23) are the basic equations of new fSM in its functional formulation.

3. I have developed the new path integral approach to fQM and fSM. My approach is based on the new functional measures generated by the stochastic process of the Lévy flights. The *Laskin* and *Lévy* path integrals are the generalization of the *Feynman* and *Wiener* path integrals respectively.

It will be interesting to find the physical systems and objects the quantum dynamics (statistical mechanics) of which is governed by the developed fQM (fSM).

References

- [1] R. P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York, 1965).
- [2] C. Grosche and F. Steiner, *Handbook of Feynman Path Integrals* (Springer-Verlag, Berlin Heidelberg, 1998).
- [3] N. Wiener, Proc. Lond. Math. Soc. **22**, 454 (1924).
- [4] M. Kac, *Probability and Related Topics in Physical Sciences* (Interscience, New York, 1959), Chap. IV.
- [5] P. Lévy, *Théorie de l'Addition des Variables Aléatoires* (Gauthier-Villars, Paris, 1937).
- [6] B.B. Mandelbrot and J.W. van Ness, SIAM (Soc. Ind. Appl. Math.) Rev. **10**, 422 (1968).
- [7] J. Klafter, A. Blumen and M.F. Shlesinger, Phys. Rev. **A35**, 3081 (1987).
- [8] G.M. Zaslavsky, Physica **D76**, 110 (1994).
- [9] G. Zimbaro, P. Veltri, G. Basile and S. Principato, Phys. Plasmas **2** (7), 2653 (1995).
- [10] R.N. Mantega and H.E. Stanley, Nature, **376**, 46 (1995).
- [11] B.J. West and W. Deering, Phys. Rep. **246**, 1 (1994).
- [12] A. V. Chechkin, V. Yu. Gonchar, cond-mat/9903400.
- [13] C. Fox, Trans. Am. Math. Soc. **98**, 395 (1961).
- [14] A.M. Mathai and R.K. Saxena, *The H-function with Applications in Statistics and Other Disciplines* (Wiley Eastern, New Delhi, 1978).
- [15] B.J. West and P. Grigolini, Phys. Rev. **E55**, 99 (1997).
- [16] K.B. Oldham and J. Spanier, *The Fractional Calculus* (Academic Press, New York, 1974).

- [17] R.P. Feynman, *Statistical Mechanics* (Benjamin. Reading, Mass. 1972).
- [18] P. Lévy, *Ann. Pisa*, **3**, 337 (1934).
- [19] J.L. Doob, *Stochastic processes* (Wiley, New York 1953).