

On Ghost Structure of Vacuum Superstring Field Theory

Kazuki Ohmori¹

*Department of Physics, Faculty of Science, University of Tokyo
Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan*

Abstract

After discussing the general form of the kinetic operator \hat{Q} around the tachyon vacuum, we determine the specific form of the pure-ghost kinetic operator \hat{Q} by requiring the twist invariance of the action. We obtain a novel result that the Grassmann-even piece Q_{even} of \hat{Q} must act differently on GSO(+) sector and on GSO(-) sector to preserve the twist invariance, and show that this structure is crucial for gauge invariance of the action. With this choice of \hat{Q} , we construct a solution in an approximation scheme which is conjectured to correspond to a non-BPS D9-brane. We consider both 0-picture cubic and Berkovits' non-polynomial superstring field theories for the NS sector.

¹E-mail: ohmori@hep-th.phys.s.u-tokyo.ac.jp

1 Introduction

For the past one and a half years vacuum string field theory (VSFT) [1, 2] has been studied intensively. The distinguishing feature of this theory is that the kinetic operator is made purely out of ghost operators so that its equation of motion allows us to have solutions of matter-ghost factorized form. Since in bosonic string theory there is essentially only one kind of D-brane, it has been assumed that D-brane solutions in VSFT have the factorized form $\Psi = \Psi_m \otimes \Psi_g$ and that its ghost part Ψ_g is common to all Dp -branes [3]. This assumption has successfully been tested by showing that such solutions correctly reproduce the ratios of D-brane tensions, in which only the matter part of the solutions was needed. As a matter of fact, many encouraging results have been obtained from the analysis of the matter sector: not only the ratios of tensions [3, 4, 5, 6, 7] and various D-brane configurations [8, 4] but also the overall D25-brane tension [9] and the fluctuation spectrum around the brane solution [4, 10, 11, 12, 13, 14, 9]. On the other hand, it has not been discussed whether the solutions of the ghost equation of motion proposed in [10, 15, 16, 17] are suited for the description of the D-brane or not. Anyway, in section 3 of the present paper we will give arguments based on the assumption that the twisted ghost sliver state introduced in [15] really describes the universal ghost solution for bosonic D-branes.

In the type II superstring case, there are two kinds of D-branes: stable BPS D-branes and unstable non-BPS ones. Because of the qualitative difference between these two families, even if all D-brane solutions in vacuum superstring field theory have the matter-ghost factorized form and each of the families has its universal ghost solutions, there is no apparent reason to assume that the two ghost solutions should also agree with each other. In case they are really different, we will be forced to consider the full matter+ghost system even for the calculation of ratios of tensions between a BPS D-brane and a non-BPS D-brane. As another issue peculiar to the superstring case, there is a question of how the spacetime supersymmetry is restored around the tachyon vacuum: After the non-supersymmetric non-BPS D-branes or brane–antibrane systems have completely decayed through the tachyon condensation, it is believed that there remains the true vacuum for the type II closed superstrings without any D-branes, where $d = 10, \mathcal{N} = 2$ spacetime supersymmetry should exist. Considering that the exact tachyon vacuum solution has not been obtained so far and that it seems difficult to investigate the supersymmetric structure in the level truncation scheme, the only way to proceed is to construct superstring field theory around the tachyon vacuum directly. Although we have no idea up to now how this phenomenon of supersymmetry restoration

should be described in terms of *open* string degrees of freedom, if we are to take the vacuum string field theory proposal seriously this is a problem of the kind that should be resolved within this theory. For this purpose, it is obvious that we must reveal the complete structure of the theory including both Neveu-Schwarz (NS) and Ramond (R) sectors.² As a first step toward this goal, we consider in this paper the NS sector only. The above two interesting problems have motivated us to study in detail the ghost structure of superstring field theory around the tachyon vacuum.

The study of ghost kinetic operator in vacuum superstring field theory was initiated by Aref'eva *et. al.* [20, 21]. Those authors argued that the kinetic operator \widehat{Q} around the tachyon vacuum may contain a Grassmann-even operator Q_{even} that mixes the GSO(\pm) sectors of the string field, and proposed two possible candidates for the pure-ghost kinetic operator \widehat{Q} where Q_{even} was represented by linear combinations of $\gamma(i)$ and $\gamma(-i)$. However, they have not shown on what principle they determined the precise form of \widehat{Q} . In the present paper we adopt as a guiding principle the ‘twist invariance’ of the action, which exists in the original (bosonic, 0-picture cubic super and Berkovits’ super) string field theories on D-branes, reinforced with the gauge invariance. In fact, if the tachyon vacuum solution is represented by a twist-even string field configuration as has been the case in the level truncation calculations, this twist symmetry should survive in vacuum string field theory action. As a result, we will put forward a candidate different from theirs, and show that our choice gives rise to consistent gauge invariant actions.

This paper is organized as follows. In section 2 we fix the form of the ghost kinetic operator \widehat{Q} from the requirement of the twist invariance of the cubic action, and then examine its properties. In section 3 we try to solve the equations of motion, but we find a solution only in an approximation scheme. In section 4 we extend the results obtained in the previous sections for the cubic theory to the non-polynomial superstring field theory by Berkovits. In section 5 we summarize our results and have discussion on further research. In Appendices we expose the technical details about the inner derivation formula for Q_{even} and the twist invariance of our action.

²Spacetime supersymmetry transformations for open string fields containing both GSO(\pm) sectors have been discussed by Yoneya [18] in the context of Witten’s cubic open superstring field theory [19].

2 Proposal for the (Super)ghost Kinetic Operator: Cubic Theory

In sections 2 and 3, we consider the cubic vacuum superstring field theory [20, 21]. We first argue that a GSO(\pm)-mixing Grassmann-even sector Q_{even} generally arises in the kinetic operator \widehat{Q} around the tachyon vacuum, following [20], and that this is in fact necessary in order to have the expected structure of vacuum superstring field theory action. We then use the twist symmetry to determine the form of purely ghostly kinetic operator \widehat{Q} , and propose that this operator describes the open superstrings around one of the tachyon vacua.

2.1 Structure of the kinetic operator around the tachyon vacuum

Let us begin with the cubic action for the Neveu-Schwarz sector string field \widehat{A} in the background of a non-BPS D-brane [20]:

$$S = -\frac{1}{2g_o^2} \text{Tr} \left[\frac{1}{2} \langle\langle \widehat{Y}_{-2} | \widehat{A}, \widehat{Q}_B \widehat{A} \rangle\rangle + \frac{1}{3} \langle\langle \widehat{Y}_{-2} | \widehat{A}, \widehat{A} * \widehat{A} \rangle\rangle \right], \quad (2.1)$$

where the *internal Chan-Paton* structure is

$$\begin{aligned} \widehat{Q}_B &= Q_B \otimes \sigma_3, \\ \widehat{Y}_{-2} &= Y(i)Y(-i) \otimes \sigma_3 \quad (Y(z) = c\partial\xi e^{-2\phi}(z)), \\ \widehat{A} &= A_+ \otimes \sigma_3 + A_- \otimes i\sigma_2, \end{aligned} \quad (2.2)$$

and the trace Tr is taken over the space of these 2×2 matrices (σ_i 's are the Pauli matrices). The GSO(+) sector string field A_+ is Grassmann-odd and consists of states with integer weights, while the GSO(-) field A_- is Grassmann-even and has half-odd-integer weights. Both of them have ghost number 1 and picture number 0. The bracket $\langle\langle Y_{-2} | \dots \rangle\rangle$ is defined in terms of the correlation functions in the world-sheet CFT as³

$$\langle\langle Y_{-2} | A_1, A_2 \rangle\rangle = \left\langle Y(i)Y(-i) f_1^{(2)} \circ A_1(0) f_2^{(2)} \circ A_2(0) \right\rangle_{\text{UHP}}, \quad (2.3)$$

$$\langle\langle Y_{-2} | A_1, A_2 * A_3 \rangle\rangle = \left\langle Y(i)Y(-i) f_1^{(3)} \circ A_1(0) f_2^{(3)} \circ A_2(0) f_3^{(3)} \circ A_3(0) \right\rangle_{\text{UHP}}, \quad (2.4)$$

³Strictly speaking, the action of conformal transformations $f_k^{(n)} \circ A(0) = (f_k^{(n)'(0)})^h A(f_k^{(n)}(0))$ is not well-defined for vertex operators of non-integer weights h unless we fix the phase ambiguity. We will mention it when it is needed.

where

$$f_k^{(n)}(z) = h^{-1} \left(e^{2\pi i \frac{k-1}{n}} h(z)^{\frac{2}{n}} \right), \quad (2.5)$$

$$h(z) = \frac{1+iz}{1-iz}, \quad h^{-1}(z) = -i \frac{z-1}{z+1},$$

and in particular,

$$f_1^{(2)}(z) = z, \quad f_2^{(2)}(z) = h^{-1}(e^{\pi i} h(z)) = -\frac{1}{z} \equiv I(z). \quad (2.6)$$

Note that since the inverse picture-changing operator $Y(z)$ is a primary field of conformal weight 0, it is not affected by conformal transformations that keep the open string midpoint $\pm i$ fixed.

By taking the trace over the internal Chan-Paton matrices in advance, the action (2.1) can be rewritten explicitly in terms of A_{\pm} as

$$S = -\frac{1}{g_o^2} \left[\frac{1}{2} \langle\langle Y_{-2} | A_+, Q_B A_+ \rangle\rangle + \frac{1}{2} \langle\langle Y_{-2} | A_-, Q_B A_- \rangle\rangle \right. \\ \left. + \frac{1}{3} \langle\langle Y_{-2} | A_+, A_+ * A_+ \rangle\rangle - \langle\langle Y_{-2} | A_+, A_- * A_- \rangle\rangle \right]. \quad (2.7)$$

From this expression, one can immediately see that this action is invariant under the sign-flip of the GSO(-) sector string field: $A_- \rightarrow -A_-$. This symmetry guarantees that the effective potential for the ‘real’⁴ tachyon field living in the GSO(-) sector takes the left-right symmetric double-well form. Now, postulating the solution \widehat{A}_0 corresponding to one of the doubly-degenerate tachyon vacua, we expand the string field \widehat{A} around it as $\widehat{A} = \widehat{A}_0 + \widehat{a}$. Then part of the action quadratic in the fluctuation fields $\widehat{a} = a_+ \otimes \sigma_3 + a_- \otimes i\sigma_2$ is found to be

$$-g_o^2 S_{\text{quad}} = \frac{1}{2} \langle\langle Y_{-2} | a_+, (Q_B a_+ + A_{0+} * a_+ + a_+ * A_{0+} - A_{0-} * a_- - a_- * A_{0-}) \rangle\rangle \quad (2.8) \\ + \frac{1}{2} \langle\langle Y_{-2} | a_-, (Q_B a_- + A_{0+} * a_- - a_- * A_{0+} - A_{0-} * a_+ + a_+ * A_{0-}) \rangle\rangle,$$

where we have used the cyclicity relation of the 3-string vertex:

$$\langle\langle Y_{-2} | A_1, A_2 * A_3 \rangle\rangle = e^{2\pi i h_{A_3}} \langle\langle Y_{-2} | A_3, A_1 * A_2 \rangle\rangle \quad (2.9)$$

with h_{A_3} denoting the conformal weight of A_3 . That is to say, if the vertex operator A_3 to be moved is in the GSO(-) sector an additional minus sign arises.⁵ This is due to the

⁴Remember that in 0-picture cubic superstring field theory there also exists an ‘auxiliary’ tachyon field in the GSO(+) sector [22].

⁵I’d like to thank I. Kishimoto for reminding me of this fact.

2π -rotation $\mathcal{R}_{2\pi}(z) \equiv h^{-1}(e^{2\pi i}h(z))$ of the unit disk, which provides a conformal factor $[\mathcal{R}'_{2\pi}(0)]^{h_{A_3}} = e^{2\pi i h_{A_3}}$ [23, 22]. Note that (2.8) contains cross-terms among a_+ and a_- through the help of A_{0-} . In the matrix notation (2.1), the same quadratic action is also written as

$$-g_o^2 S_{\text{quad}} = \frac{1}{4} \text{Tr} \langle \langle \widehat{Y}_{-2} | \widehat{a}, (\widehat{Q}_B \widehat{a} + \widehat{A}_0 * \widehat{a} + \widehat{a} * \widehat{A}_0) \rangle \rangle, \quad (2.10)$$

where we have used the fact that the hatted fields satisfy the cyclicity relation without any sign factor because the $-$ sign for the $\text{GSO}(-)$ field appearing in (2.9) is compensated for by one more sign arising from $i\sigma_2 \widehat{Y}_{-2} = -\widehat{Y}_{-2} i\sigma_2$. Here we define the new kinetic operator around the tachyon vacuum as

$$\widehat{Q}\widehat{\Phi} \equiv \widehat{Q}_B \widehat{\Phi} + \widehat{A}_0 * \widehat{\Phi} - (-1)^{\#\text{gh}(\widehat{\Phi})} \widehat{\Phi} * \widehat{A}_0, \quad (2.11)$$

where $\#\text{gh}(\widehat{\Phi})$ denotes the ghost number of $\widehat{\Phi}$. To understand why we have adopted the ghost number grading instead of the Grassmannality, recall that the internal Chan-Paton factors have originally been introduced in such a way that the $\text{GSO}(\pm)$ string fields with different Grassmannalities obey the same algebraic relations [23]. Note that if we restrict ourselves to the $\text{GSO}(+)$ states, these two gradings agree with each other: $(-1)^{\#\text{gh}(\widehat{\Phi}_+)} = (-1)^{|\widehat{\Phi}_+|}$.⁶ By using \widehat{Q} defined in (2.11), eq.(2.10) can further be rewritten as

$$-g_o^2 S_{\text{quad}} = \frac{1}{2} \langle \langle Y_{-2} | a_+, \frac{1}{2} \text{Tr}(\widehat{Q}\widehat{a}) \rangle \rangle + \frac{1}{2} \langle \langle Y_{-2} | a_-, \frac{1}{2} \text{Tr}(\sigma_1 \widehat{Q}\widehat{a}) \rangle \rangle. \quad (2.12)$$

Comparing eqs.(2.8) and (2.12), we notice that $\text{Tr}(\widehat{Q}i\sigma_2)a_-$ and $\text{Tr}(\sigma_1 \widehat{Q}\sigma_3)a_+$ (as well as $\text{Tr}(\widehat{Q}\sigma_3)a_+$ and $\text{Tr}(\sigma_1 \widehat{Q}i\sigma_2)a_-$) must be non-zero in general, because the tachyon vacuum solution \widehat{A}_0 contains non-zero $\text{GSO}(-)$ components A_{0-} and there is no reason that $-A_{0-} * a_- - a_- * A_{0-}$ and $-A_{0-} * a_+ + a_+ * A_{0-}$ should vanish in (2.8). This can be achieved by letting \widehat{Q} have the following internal Chan-Paton structure [20]:

$$\widehat{Q} = Q_{\text{odd}} \otimes \sigma_3 - Q_{\text{even}} \otimes i\sigma_2, \quad (2.13)$$

where Q_{odd} and Q_{even} are Grassmann-odd and Grassmann-even operators respectively. Explicit actions of $Q_{\text{odd}}, Q_{\text{even}}$ can be found by comparing the both sides of eq.(2.11). When \widehat{Q} acts on a string field \widehat{X} of *odd ghost number* of the form

$$\widehat{X} = X_+ \otimes \sigma_3 + X_- \otimes i\sigma_2, \quad (2.14)$$

⁶Generically a relation $(-1)^{\#\text{gh}(\widehat{\Phi})}(-1)^{|\widehat{\Phi}|}(-1)^{\text{GSO}(\widehat{\Phi})} = 1$ holds among the ghost number $\#\text{gh}(\widehat{\Phi})$, Grassmannality $|\widehat{\Phi}|$ and GSO parity $\text{GSO}(\widehat{\Phi})$ of $\widehat{\Phi}$.

we find

$$\begin{aligned}
Q_{\text{odd}}X_+ &= Q_B X_+ + A_{0+} * X_+ + X_+ * A_{0+}, \\
Q_{\text{odd}}X_- &= Q_B X_- + A_{0+} * X_- - X_- * A_{0+}, \\
Q_{\text{even}}X_+ &= -A_{0-} * X_+ + X_+ * A_{0-}, \\
Q_{\text{even}}X_- &= -A_{0-} * X_- - X_- * A_{0-},
\end{aligned} \tag{2.15}$$

where the GSO(+) component X_+ is Grassmann-odd and the GSO(-) component X_- is Grassmann-even. On the other hand, when \widehat{Q} acts on a string field \widehat{Y} of *even ghost number* having the form

$$\widehat{Y} = Y_+ \otimes \mathbf{1} + Y_- \otimes \sigma_1, \tag{2.16}$$

we obtain

$$\begin{aligned}
Q_{\text{odd}}Y_+ &= Q_B Y_+ + A_{0+} * Y_+ - Y_+ * A_{0+}, \\
Q_{\text{odd}}Y_- &= Q_B Y_- + A_{0+} * Y_- + Y_- * A_{0+}, \\
Q_{\text{even}}Y_+ &= -A_{0-} * Y_+ + Y_+ * A_{0-}, \\
Q_{\text{even}}Y_- &= -A_{0-} * Y_- - Y_- * A_{0-},
\end{aligned} \tag{2.17}$$

where Y_+ is Grassmann-even and Y_- is Grassmann-odd. These two sets (2.15), (2.17) of equations can be written in a unified manner as

$$Q_{\text{odd}}a = Q_B a + A_{0+} * a - (-1)^{|a|} a * A_{0+}, \tag{2.18}$$

$$Q_{\text{even}}a = -A_{0-} * a + (-1)^{\text{GSO}(a)} a * A_{0-}, \tag{2.19}$$

for a of any ghost number and of any GSO parity, where $|a|$ denotes the Grassmannality of a ($|a| = 0/1 \pmod{2}$ if a is Grassmann-even/odd) and $\text{GSO}(a)$ represents the GSO parity of a ($(-1)^{\text{GSO}(a)} = \pm 1$ if a is in the GSO(\pm) sector). The difference in the Chan-Paton structures for even and odd ghost number string fields, (2.14) and (2.16),⁷ comes from the consistency of the $*$ -product: For example, $*$ -multiplication of two string fields both having ghost number 1 must give rise to a structure appropriate for a string field of ghost number 2, and the set of all ghost number 0 string fields must form a closed subalgebra under the $*$ -product.

Here we argue that non-zero Q_{even} is necessary to have a sensible vacuum superstring field theory. To this end, suppose that we are given $Q_{\text{even}} = 0$, so $\widehat{Q} = Q_{\text{odd}} \otimes \sigma_3$. Since this \widehat{Q} has the same structure as $\widehat{Q}_B = Q_B \otimes \sigma_3$, the action expanded around the tachyon

⁷The same assignment of the Chan-Paton matrices to string fields had been proposed in [24].

vacuum would again take the same form as the original one (2.7) with Q_B replaced by Q_{odd} and A_{\pm} denoting the fluctuations around the tachyon vacuum. Then, for the same reason as the one mentioned above, the ‘string field theory around the tachyon vacuum’ would have a \mathbb{Z}_2 -reflection symmetry under $a_- \rightarrow -a_-$, which means that if we have a solution $a_+ \otimes \sigma_3 + a_- \otimes i\sigma_2$ then we find one more solution $a_+ \otimes \sigma_3 - a_- \otimes i\sigma_2$ with the same energy density. However, we do not expect such a degeneracy of solutions to exist in vacuum superstring field theory (Figure 1). There remains a possibility that any




Figure 1: A schematic picture of the tachyon potential. No reflection symmetry is expected around the tachyon vacuum ($a = 0$).

relevant solutions in this theory, such as D-branes, consist only of GSO(+) components so that we can avoid having a pair of degenerate solutions, but we do not believe that this is the case. Next we show that Q_{even} plays the rôle of removing this unwanted degeneracy. Making use of the new kinetic operator \widehat{Q} defined in (2.11) or (2.18)–(2.19), we can write the cubic superstring field theory action around the tachyon vacuum as

$$\begin{aligned}
S &= -\frac{1}{g_o^2} \left[\frac{1}{2} \langle\langle Y_{-2} | a_+, Q_{\text{odd}} a_+ \rangle\rangle + \frac{1}{2} \langle\langle Y_{-2} | a_-, Q_{\text{odd}} a_- \rangle\rangle \right. \\
&\quad + \frac{1}{2} \langle\langle Y_{-2} | a_+, Q_{\text{even}} a_- \rangle\rangle + \frac{1}{2} \langle\langle Y_{-2} | a_-, Q_{\text{even}} a_+ \rangle\rangle \\
&\quad \left. + \frac{1}{3} \langle\langle Y_{-2} | a_+, a_+ * a_+ \rangle\rangle - \langle\langle Y_{-2} | a_+, a_- * a_- \rangle\rangle \right] \quad (2.20)
\end{aligned}$$

$$= -\frac{1}{2g_o^2} \text{Tr} \left[\frac{1}{2} \langle\langle \widehat{Y}_{-2} | \widehat{a}, \widehat{Q} \widehat{a} \rangle\rangle + \frac{1}{3} \langle\langle \widehat{Y}_{-2} | \widehat{a}, \widehat{a} * \widehat{a} \rangle\rangle \right], \quad (2.21)$$

where we have omitted a constant term. In the second line of eq.(2.20) the GSO(−) string field a_- enters the action linearly, so that the above-mentioned \mathbb{Z}_2 symmetry is absent in

this action.⁸ At this stage the kinetic operator \widehat{Q} is regular and is not considered to be pure ghost. In the next subsection we will try to determine the form of the purely ghostly kinetic operator $\widehat{\mathcal{Q}}$ which is supposed to arise after a suitable singular field redefinition.

2.2 Ghost kinetic operator of cubic vacuum superstring field theory

Let us briefly review the argument of Gaiotto, Rastelli, Sen and Zwiebach [15] about the origin of a pure-ghost kinetic operator of bosonic vacuum string field theory. First assume that a regular representative Q of an equivalence class of kinetic operators around the tachyon vacuum takes the following form

$$Q = \int_{-\pi}^{\pi} d\sigma a_c(\sigma)c(\sigma) + \sum_r \int_{-\pi}^{\pi} d\sigma a_r(\sigma)O_r(\sigma), \quad (2.22)$$

where $a_{c,r}$ are functions of σ and O_r 's are local operators of ghost number 1 with conformal weights higher than that of c . Then consider performing a reparametrization of the open string coordinate: $\sigma \rightarrow f(\sigma)$, which keeps the open string midpoint $\pm\pi/2$ fixed and is symmetric about it. While this operation does not change the $*$ -product, it induces a transformation on the operator (2.22) as

$$Q \longrightarrow \mathcal{Q} = \int_{-\pi}^{\pi} d\sigma a_c(\sigma)(f'(\sigma))^{-1}c(f(\sigma)) + \sum_r \int_{-\pi}^{\pi} d\sigma a_r(\sigma)(f'(\sigma))^{h_r}O_r(f(\sigma)). \quad (2.23)$$

If we choose $f(\sigma)$ such that $f'(\sigma) \simeq (\sigma \mp \frac{\pi}{2})^2 + \varepsilon_r^2$ near $\sigma = \pm\pi/2$ with small ε_r , the integrand of the first term becomes large around $\sigma = \pm\pi/2$ and, in the limit $\varepsilon_r \rightarrow 0$, all other contributions can be neglected. In this way we have obtained simple but singular constituents of pure-ghost kinetic operator: $\varepsilon_r^{-1}c(i)$ and $\varepsilon_r^{-1}c(-i)$ in the upper half plane coordinate. The relative coefficient between these two terms will be fixed by requiring that the kinetic operator \mathcal{Q} preserve the twist invariance of the action.

It is known that the bosonic cubic open string field theory action

$$S(\Phi) = -\frac{1}{g_o^2} \left[\frac{1}{2} \langle \Phi, Q_B \Phi \rangle + \frac{1}{3} \langle \Phi, \Phi * \Phi \rangle \right] \quad (2.24)$$

has a twist symmetry [25, 26]. On an arbitrary L_0^{tot} -eigenstate $|\Phi\rangle$ the twist operator Ω acts as

$$\Omega|\Phi\rangle = (-1)^{h_\Phi+1}|\Phi\rangle, \quad (2.25)$$

⁸This argument relies on the very fundamental assumption that in vacuum superstring field theory we do not perform the GSO-projection on the open string field, as in the case of the open superstring theory on non-BPS D-branes.

where h_Φ is the L_0^{tot} -eigenvalue of $|\Phi\rangle$. It can be shown that the action (2.24) is twist-invariant, $S(\Omega\Phi) = S(\Phi)$ [26]. Thanks to this property, we could restrict the string field to be twist-even in computing the tachyon potential [27]. The proof of the twist-invariance of the action uses the fact that the usual BRST operator Q_B commutes with the twist operator Ω :

$$\Omega(Q_B|\Phi\rangle) = Q_B(\Omega|\Phi\rangle). \quad (2.26)$$

This property holds because Q_B is the zero-mode of the BRST current j_B so that it does not change the L_0^{tot} -eigenvalue of the state. Let us now turn to bosonic vacuum string field theory whose action is given by

$$S_V(\Psi) = -\kappa_0 \left[\frac{1}{2} \langle \Psi, \mathcal{Q}\Psi \rangle + \frac{1}{3} \langle \Psi, \Psi * \Psi \rangle \right], \quad (2.27)$$

where, according to the arguments of the last paragraph, the kinetic operator \mathcal{Q} is some linear combination of $c(i)$ and $c(-i)$. Since the original action (2.24) has the twist symmetry and the tachyon vacuum solution is believed to be represented by a twist-even configuration [27], it is natural to assume that the VSFT action (2.27) also has twist symmetry. For this action to be twist invariant, \mathcal{Q} must commute with Ω :

$$\Omega(\mathcal{Q}|\Psi\rangle) = \mathcal{Q}(\Omega|\Psi\rangle). \quad (2.28)$$

Since we have

$$\Omega(c_n|\Psi\rangle) = (-1)^{(h_\Psi - n) + 1} (c_n|\Psi\rangle) = (-1)^{-n} c_n(\Omega|\Psi\rangle),$$

\mathcal{Q} satisfies the twist-invariance condition (2.28) if \mathcal{Q} consists of even modes c_{2n} only. This requirement uniquely fixes the relative normalization as [15]

$$\begin{aligned} \mathcal{Q} = \mathcal{Q}_{\text{GRSZ}} &\equiv \frac{1}{2i} (c(i) - c(-i)) \\ &= c_0 - (c_2 + c_{-2}) + (c_4 + c_{-4}) - \dots, \end{aligned} \quad (2.29)$$

where an overall normalization constant has been absorbed into the definition of the string field. This kinetic operator was shown [15, 28] to agree with the one found in [10] by requiring that the Siegel gauge solution should solve the equations of motion in the full gauge-unfixed field configuration space.

Now we turn to the superstring case, where there are two negative-dimensional operators c and γ . Suppose that after a reparametrization of the string coordinate implemented by a function f , the kinetic operator is written as

$$\begin{aligned} \widehat{\mathcal{Q}} &= \int_{-\pi}^{\pi} d\sigma a_c(\sigma) [f'(\sigma)]^{-1} c(f(\sigma)) \otimes \sigma_3 \\ &\quad + \int_{-\pi}^{\pi} d\sigma a_\gamma(\sigma) [f'(\sigma)]^{-\frac{1}{2}} \gamma(f(\sigma)) \otimes i\sigma_2 + \dots \end{aligned} \quad (2.30)$$

Let us postulate a function f which around $\sigma = \frac{\pi}{2}$ behaves as

$$[f'(\sigma)]^{-\frac{1}{2}} \sim \frac{1}{\varepsilon_r} \delta\left(\sigma - \frac{\pi}{2}\right) \quad \text{and} \quad [f'(\sigma)]^{-1} \sim \frac{1}{\varepsilon_r^2} \delta\left(\sigma - \frac{\pi}{2}\right)$$

in the singular limit $\varepsilon_r \rightarrow 0$. If we take such f that behaves similarly near $\sigma = -\frac{\pi}{2}$ and is regular everywhere except at $\sigma = \pm\frac{\pi}{2}$, then $\widehat{\mathcal{Q}}$ (2.30) in the limit $\varepsilon_r \rightarrow 0$ is dominated by

$$\begin{aligned} \widehat{\mathcal{Q}} &= \frac{1}{\varepsilon_r^2} \left(a_c\left(\frac{\pi}{2}\right) c\left(\frac{\pi}{2}\right) + a_c\left(-\frac{\pi}{2}\right) c\left(-\frac{\pi}{2}\right) \right) \otimes \sigma_3 \\ &\quad + \frac{1}{\varepsilon_r} \left(a_\gamma\left(\frac{\pi}{2}\right) \gamma\left(\frac{\pi}{2}\right) + a_\gamma\left(-\frac{\pi}{2}\right) \gamma\left(-\frac{\pi}{2}\right) \right) \otimes i\sigma_2, \end{aligned} \quad (2.31)$$

where we have used $f\left(\pm\frac{\pi}{2}\right) = \pm\frac{\pi}{2}$. We then require $\widehat{\mathcal{Q}}$ to preserve the twist invariance of the action, by which the form of $\widehat{\mathcal{Q}}$ can further be restricted without knowing the precise values of $a_{c,\gamma}\left(\pm\frac{\pi}{2}\right)$. In 0-picture cubic superstring field theory, it has been shown in [22] that the action (2.1) is invariant under the twist operation

$$\Omega|A\rangle = \begin{cases} (-1)^{h_A+1}|A\rangle & \text{for GSO(+)} \text{ states } (h_A \in \mathbb{Z}) \\ (-1)^{h_A+\frac{1}{2}}|A\rangle & \text{for GSO(-)} \text{ states } (h_A \in \mathbb{Z} + \frac{1}{2}). \end{cases} \quad (2.32)$$

Since $\mathcal{Q}_{\text{GRSZ}}$ (2.29) preserves the twist eigenvalues on both GSO(\pm) sectors and is Grassmann-odd, the odd part Q_{odd} of the kinetic operator (2.13), after the singular reparametrization, becomes

$$Q_{\text{odd}} \longrightarrow \mathcal{Q}_{\text{odd}} = \frac{1}{2i\varepsilon_r^2} (c(i) - c(-i)) \quad (\varepsilon_r \rightarrow 0), \quad (2.33)$$

where we have made a finite rescaling of ε_r for convenience. On the other hand, since $\gamma(z)$ has half-odd-integer modes in the NS sector and mixes the GSO(\pm) sectors, its twist property becomes much more complicated. For example, let us consider a GSO(+) state $|A_+\rangle$ with L_0^{tot} -eigenvalue h_+ . From eq.(2.32), we have

$$\Omega|A_+\rangle = (-1)^{h_++1}|A_+\rangle. \quad (2.34)$$

When γ_r ($r \in \mathbb{Z} + \frac{1}{2}$) acts on $|A_+\rangle$, the resulting state $\gamma|A_+\rangle$ is in the GSO(-) sector and hence its twist eigenvalue must be evaluated as a GSO(-) state:

$$\Omega(\gamma_r|A_+\rangle) = (-1)^{(h_+-r)+\frac{1}{2}}(\gamma_r|A_+\rangle). \quad (2.35)$$

Combining eqs.(2.34) and (2.35), we find the following relation:

$$\Omega(\gamma_r|A_+\rangle) = (-1)^{-r-\frac{1}{2}}\gamma_r(\Omega|A_+\rangle). \quad (2.36)$$




Figure 2: (a) The action of γ_r on GSO(+) twist-even states. γ_r with $r \in 2\mathbb{Z} - \frac{1}{2}$ preserve the twist eigenvalues (indicated by solid arrows), whereas the wrong ones with $r \in 2\mathbb{Z} + \frac{1}{2}$ reverse the twist eigenvalues (dotted arrows). (b) The action of γ_r on GSO(-) twist-even states. Now it is γ_r with $r \in 2\mathbb{Z} + \frac{1}{2}$ that preserve the twist.

Thus we conclude that γ_r acting on a GSO(+) state $|A_+\rangle$ commutes with the twist

$$\Omega(\gamma_r|A_+\rangle) = \gamma_r(\Omega|A_+\rangle) \quad (2.37)$$

when $r \in 2\mathbb{Z} - \frac{1}{2}$. This argument is visualized in Figure 2(a). Since we find

$$\frac{1}{2i}(\gamma(i) - \gamma(-i)) = \frac{1}{2i} \left(\sum_{r \in \mathbb{Z} + \frac{1}{2}} \frac{\gamma_r}{i^{r - \frac{1}{2}}} - \sum_{r \in \mathbb{Z} + \frac{1}{2}} \frac{\gamma_r}{(-i)^{r - \frac{1}{2}}} \right) = \sum_{n \in \mathbb{Z}} (-1)^n \gamma_{-\frac{1}{2} + 2n}, \quad (2.38)$$

we identify a candidate for the twist-preserving kinetic operator as

$$Q_{\text{even}} \longrightarrow Q_{\text{even}}^{\text{GSO}(+)} = \frac{q_1}{2i\varepsilon_r} (\gamma(i) - \gamma(-i)) \quad (\varepsilon_r \rightarrow 0), \quad (2.39)$$

where q_1 is a finite real constant. However, it turns out that this kinetic operator, when acting on a GSO(-) state, does *not* preserve the twist eigenvalue. To see this, consider a GSO(-) state $|A_-\rangle$ with L_0^{tot} -eigenvalue h_- . From the relation

$$\Omega(\gamma_r|A_-\rangle) = (-1)^{(h_- - r) + 1} (\gamma_r|A_-\rangle) = (-1)^{-r + \frac{1}{2}} \gamma_r(\Omega|A_-\rangle), \quad (2.40)$$

γ_r commutes with Ω if $r \in 2\mathbb{Z} + \frac{1}{2}$, rather than $r \in 2\mathbb{Z} - \frac{1}{2}$ (see Figure 2(b)). Therefore, the twist-preserving kinetic operator acting on a $\text{GSO}(-)$ state should take the form

$$\begin{aligned} \mathcal{Q}_{\text{even}} \longrightarrow \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} &= \frac{q_2}{2\varepsilon_r} (\gamma(i) + \gamma(-i)) \\ &= \frac{q_2}{\varepsilon_r} \sum_{n \in \mathbb{Z}} (-1)^n \gamma_{\frac{1}{2}+2n} \end{aligned} \quad (2.41)$$

in the $\varepsilon_r \rightarrow 0$ limit, where q_2 is another constant.

Our proposal that the kinetic operator $\mathcal{Q}_{\text{even}}$ takes different forms (2.39), (2.41) depending on the GSO parity of the states on which $\mathcal{Q}_{\text{even}}$ acts may seem strange, but such a behavior is in fact necessary for the construction of gauge-invariant actions: To show the gauge invariance of the action, we need the hermiticity relation for $\widehat{\mathcal{Q}} = \mathcal{Q}_{\text{odd}} \otimes \sigma_3 - \mathcal{Q}_{\text{even}} \otimes i\sigma_2$,⁹

$$\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{A}, \widehat{B} \rangle\rangle = -(-1)^{\#\text{gh}(\widehat{A})} \langle\langle \widehat{Y}_{-2} | \widehat{A}, \widehat{\mathcal{Q}} \widehat{B} \rangle\rangle. \quad (2.42)$$

Given the internal Chan-Paton structure

$$\begin{aligned} \widehat{A} &= A_+ \otimes \sigma_3 + A_- \otimes i\sigma_2 && \text{for } \#\text{gh}(\widehat{A}) \text{ odd} \\ \widehat{A} &= A_+ \otimes \mathbf{1} + A_- \otimes \sigma_1 && \text{for } \#\text{gh}(\widehat{A}) \text{ even,} \end{aligned} \quad (2.43)$$

(similarly for \widehat{B}) and concentrating on the $\mathcal{Q}_{\text{even}}$ part, eq.(2.42) is rewritten as

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}} A_+, B_- \rangle\rangle = -\langle\langle Y_{-2} | A_+, \mathcal{Q}_{\text{even}} B_- \rangle\rangle, \quad (2.44)$$

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}} A_-, B_+ \rangle\rangle = \langle\langle Y_{-2} | A_-, \mathcal{Q}_{\text{even}} B_+ \rangle\rangle. \quad (2.45)$$

Here, let us closely look at the 2-point vertex (BPZ inner product)

$$\langle\langle Y_{-2} | A_1, A_2 \rangle\rangle = \langle Y(i)Y(-i)A_1(0)I \circ A_2(0) \rangle_{\text{UHP}}.$$

The inversion $I(z) = -\frac{1}{z} = h^{-1}(-h(z))$ can be written in the following two ways: $\mathcal{R}_\pi(z) \equiv h^{-1}(e^{\pi i}h(z))$ and $\mathcal{R}_{-\pi}(z) \equiv h^{-1}(e^{-\pi i}h(z))$. Recalling that the 2π -rotation $\mathcal{R}_{2\pi}(z) = h^{-1}(e^{2\pi i}h(z)) \simeq z$ acts non-trivially on $\text{GSO}(-)$ states with half-integer weights, we must define the action of the inversion $I(z) = -1/z$ on A_2 more precisely. We adopt the convention

$$\begin{aligned} \langle\langle Y_{-2} | A_1, A_2 \rangle\rangle &\equiv \langle Y(i)Y(-i) A_1(0) \mathcal{R}_\pi \circ A_2(0) \rangle_{\text{UHP}} \\ &= [\mathcal{R}'_\pi(0)]^{h_{A_2}} \langle Y(i)Y(-i) A_1(0) A_2(\mathcal{R}_\pi(0)) \rangle_{\text{UHP}}, \end{aligned} \quad (2.46)$$

⁹Since eq.(2.42) is derived from the derivation property (2.68) of $\widehat{\mathcal{Q}}$ and the fact that $\widehat{\mathcal{Q}}$ annihilates $\langle \mathcal{I} | Y_{-2}$, self-consistency also requires eq.(2.42) to hold.

where

$$[\mathcal{R}'_{\pi}(z)]^h \equiv z^{-2h}, \quad [\mathcal{R}'_{-\pi}(z)]^h \equiv e^{-2\pi ih} z^{-2h} = (-1)^{2h} z^{-2h}, \quad (2.47)$$

and we have assumed A_2 to be a primary field in (2.46). The above definition (2.47) is consistent with the composition laws

$$\mathcal{R}_{\pi} \circ \mathcal{R}_{\pi}(z) = \mathcal{R}_{2\pi}(z), \quad \mathcal{R}_{\pi} \circ \mathcal{R}_{-\pi}(z) = \mathcal{R}_{-\pi} \circ \mathcal{R}_{\pi}(z) = z,$$

and $[\mathcal{R}'_{2\pi}(z)]^h = e^{2\pi ih}$. Defined this way, the inner product (2.46) has been made well-defined for GSO(-) states as well. Now let us see eq.(2.45) in detail. The left-hand side of (2.45) can be written as

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} A_-, B_+ \rangle\rangle = \langle Y(i)Y(-i)A_-(0) \mathcal{Q}_{\text{even}}^{\text{GSO}(-)}(\mathcal{R}_{\pi} \circ B_+(0)) \rangle_{\text{UHP}}, \quad (2.48)$$

while the right-hand side of (2.45) is

$$\langle\langle Y_{-2} | A_-, \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} B_+ \rangle\rangle = \langle\langle Y(i)Y(-i)A_-(0)(\mathcal{R}_{\pi} \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(+)}) (\mathcal{R}_{\pi} \circ B_+(0)) \rangle\rangle_{\text{UHP}}. \quad (2.49)$$

For these two expressions to agree with each other, we must have

$$\mathcal{R}_{\pi} \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} = \mathcal{Q}_{\text{even}}^{\text{GSO}(-)}, \quad (2.50)$$

but this equation *cannot be satisfied if we stick to the case* $\mathcal{Q}_{\text{even}}^{\text{GSO}(+)} = \mathcal{Q}_{\text{even}}^{\text{GSO}(-)}$ because neither of $\gamma(\pm i), \gamma(i) \pm \gamma(-i)$ is self-conjugate under the inversion $\mathcal{R}_{\pm\pi}$.¹⁰ Thus we conclude that in order for $\widehat{\mathcal{Q}}$ to satisfy the hermiticity relation (2.45) $\mathcal{Q}_{\text{even}}$ must inevitably take different forms on GSO(\pm) sectors. In fact, we find from (2.39)

$$\begin{aligned} \mathcal{R}_{\pi} \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} &= \frac{q_1}{2i\varepsilon_r} \mathcal{R}_{\pi} \circ (\gamma(i) - \gamma(-i)) \\ &= \frac{q_1}{2i\varepsilon_r} \left((\mathcal{R}'_{\pi}(i))^{-\frac{1}{2}} \gamma(\mathcal{R}_{\pi}(i)) - (\mathcal{R}'_{\pi}(-i))^{-\frac{1}{2}} \gamma(\mathcal{R}_{\pi}(-i)) \right) \\ &= \frac{q_1}{2\varepsilon_r} (\gamma(i) + \gamma(-i)) \end{aligned} \quad (2.51)$$

because $(\mathcal{R}'_{\pi}(\pm i))^{-\frac{1}{2}} = \pm i$ due to the definition (2.47). Thus, we see that the hermiticity condition (2.50) is satisfied by our choice (2.39) and (2.41) of kinetic operator if we set $q_1 = q_2$. With this choice, one can verify that eq.(2.44) also holds true.

As shown above, the ratio q_1/q_2 of the finite normalization constants of $\mathcal{Q}_{\text{even}}^{\text{GSO}(\pm)}$ has been fixed by requiring the hermiticity condition. Then, how can we determine the value of q_1 itself? This question seems difficult to answer because it requires the

¹⁰Generically, operators of half-integer weights satisfy $\mathcal{R}_{\pi} \circ \mathcal{R}_{\pi} \circ \mathcal{O} = -\mathcal{O}$ so that it seems impossible to construct operators which are real and self-conjugate under \mathcal{R}_{π} .

detailed information about the reparametrization, by which the kinetic operator around the tachyon vacuum has been brought to the simple purely ghost form, and the functions $a_{c,\gamma}(\sigma)$ appearing in (2.30). We only note here that the sign of q_1 , which corresponds to the relative sign between the normalization constant of \mathcal{Q}_{odd} and that of $\mathcal{Q}_{\text{even}}$, is related to the choice of the tachyon vacuum around which vacuum superstring field theory is constructed. This fact can immediately be seen from the definition (2.19) of $\mathcal{Q}_{\text{even}}$ whose sign is flipped under $A_{0-} \rightarrow -A_{0-}$. Since two degenerate tachyon vacua are considered to be physically equivalent, we may take q_1 to be positive without loss of generality.

To summarize, we have seen that the twist invariance condition (2.37) combined with the hermiticity condition (2.42) points to the choice

$$\begin{aligned}\mathcal{Q}_{\text{even}}^{\text{GSO}(+)} &= \frac{q_1}{2i\varepsilon_r}(\gamma(i) - \gamma(-i)) \\ \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} &= \frac{q_1}{2\varepsilon_r}(\gamma(i) + \gamma(-i)),\end{aligned}\tag{2.52}$$

or collectively

$$\mathcal{Q}_{\text{even}}|\psi\rangle = q_1 \frac{1-i}{4\varepsilon_r} (1 - i(-1)^{\text{GSO}(\psi)}) (\gamma(i) - (-1)^{\text{GSO}(\psi)}\gamma(-i)) |\psi\rangle.\tag{2.53}$$

The next step is to check whether the above choice of $\mathcal{Q}_{\text{even}}$ together with \mathcal{Q}_{odd} given in (2.33) satisfies the axioms imposed on the kinetic operator of superstring field theory. Some of the proofs given below overlap with the ones presented in [20].

Nilpotency of $\widehat{\mathcal{Q}}$

To get a nilpotent kinetic operator $\widehat{\mathcal{Q}}$, it turns out that we must add to \mathcal{Q}_{odd} a non-leading term in ε_r as¹¹ [20]

$$\mathcal{Q}_{\text{odd}} = \frac{1}{2i\varepsilon_r^2}(c(i) - c(-i)) + \frac{q_1^2}{2} \oint \frac{dz}{2\pi i} b\gamma^2(z).\tag{2.54}$$

Here $Q_{\text{ABC}} \equiv \oint \frac{dz}{2\pi i} b\gamma^2(z)$ is part of the BRST charge Q_B , and was considered in [29]. Recently, Q_{ABC} was used to propose the pregeometrical formulation of Berkovits' superstring field theory [30]. Since Q_{ABC} is the zero mode of a weight 1 primary $b\gamma^2$, it manifestly preserves the twist eigenvalues. Since we have

$$\widehat{\mathcal{Q}}\widehat{\mathcal{Q}}|\widehat{A}\rangle = \{(\mathcal{Q}_{\text{odd}}^2 - \mathcal{Q}_{\text{even}}^2) \otimes \mathbf{1} - [\mathcal{Q}_{\text{odd}}, \mathcal{Q}_{\text{even}}] \otimes \sigma_1\} |\widehat{A}\rangle,\tag{2.55}$$

¹¹The choice of the second term in (2.54) is not unique. For example, $\oint \frac{dz}{2\pi i} f(z)b\gamma^2(z)$ will do if a scalar function $f(z)$ is regular in an annular region around $|z|=1$ and satisfies $f(\pm i)=1$. Or, we can add to \mathcal{Q}_{odd} more terms which anticommute with $c(\pm i)$ and Q_{ABC} and are nilpotent themselves. Here we consider the simplest choice Q_{ABC} .

we must show both $(\mathcal{Q}_{\text{odd}}^2 - \mathcal{Q}_{\text{even}}^2)|\widehat{A}\rangle = 0$ and

$$[\mathcal{Q}_{\text{odd}}, \mathcal{Q}_{\text{even}}]|A_{\pm}\rangle = (\mathcal{Q}_{\text{odd}}\mathcal{Q}_{\text{even}}^{\text{GSO}(\pm)} - \mathcal{Q}_{\text{even}}^{\text{GSO}(\pm)}\mathcal{Q}_{\text{odd}})|A_{\pm}\rangle = 0. \quad (2.56)$$

The latter holds because \mathcal{Q}_{odd} (2.54) contains no β field and \mathcal{Q}_{odd} does not change the GSO parity, as indicated in (2.56).¹² The former one can be shown as follows:

$$\mathcal{Q}_{\text{odd}}^2|A_{\pm}\rangle = \frac{q_1^2}{4i\varepsilon_r^2} \left\{ \oint \frac{dz}{2\pi i} b\gamma^2(z), c(i) - c(-i) \right\} |A_{\pm}\rangle = \frac{q_1^2}{4i\varepsilon_r^2} (\gamma(i)^2 - \gamma(-i)^2) |A_{\pm}\rangle, \quad (2.57)$$

and, from eqs.(2.52),

$$\mathcal{Q}_{\text{even}}^2|A_{\pm}\rangle = \mathcal{Q}_{\text{even}}^{\text{GSO}(\mp)}\mathcal{Q}_{\text{even}}^{\text{GSO}(\pm)}|A_{\pm}\rangle = \frac{q_1^2}{4i\varepsilon_r^2} (\gamma(i)^2 - \gamma(-i)^2) |A_{\pm}\rangle, \quad (2.58)$$

where $|A_{+/-}\rangle$ denote any states in the GSO(+/-) sectors respectively, and we have used the fact that $\mathcal{Q}_{\text{even}}$ reverses the GSO parity of the states. From (2.57) and (2.58), it follows that $(\mathcal{Q}_{\text{odd}}^2 - \mathcal{Q}_{\text{even}}^2)|\widehat{A}\rangle = 0$. This completes the proof that $\widehat{\mathcal{Q}} = \mathcal{Q}_{\text{odd}} \otimes \sigma_3 - \mathcal{Q}_{\text{even}} \otimes i\sigma_2$ with \mathcal{Q}_{odd} given in (2.54) and $\mathcal{Q}_{\text{even}}$ in (2.52) is nilpotent.

$\langle \mathcal{I} | \widehat{\mathcal{Q}} = 0$

Given that the identity string field $\langle \mathcal{I} |$ is defined as

$$\langle \mathcal{I} | \varphi \rangle = \langle f_1^{(1)} \circ \varphi(0) \rangle_{\text{UHP}} \quad (2.59)$$

with $f_1^{(1)}(z) = h^{-1}(h(z)^2) = \frac{2z}{1-z^2}$ for an arbitrary Fock space state $|\varphi\rangle = \varphi(0)|0\rangle$, both $\langle \mathcal{I} | c(\pm i)$ and $\langle \mathcal{I} | \gamma(\pm i)$ contain divergences because the conformal factors $(f_1^{(1)'(\pm i)})^h$ diverge for $h < 0$. However, as pointed out in [15, 20], they can be regularized by the following prescription. If we make the replacements

$$\begin{aligned} c(i) &\longrightarrow c_{\epsilon}(i) = \frac{1}{2} (e^{-i\epsilon} c(i e^{i\epsilon}) + e^{i\epsilon} c(i e^{-i\epsilon})), \\ c(-i) &\longrightarrow c_{\epsilon}(-i) = \frac{1}{2} (e^{-i\epsilon} c(-i e^{i\epsilon}) + e^{i\epsilon} c(-i e^{-i\epsilon})), \\ \gamma(i) &\longrightarrow \gamma_{\epsilon}(i) = \frac{1}{e^{-\frac{\pi i}{4}} - e^{\frac{\pi i}{4}}} \left(e^{-\frac{\pi i}{4} - \frac{i\epsilon}{2}} \gamma(i e^{i\epsilon}) - e^{\frac{\pi i}{4} + \frac{i\epsilon}{2}} \gamma(i e^{-i\epsilon}) \right), \\ \gamma(-i) &\longrightarrow \gamma_{\epsilon}(-i) = \frac{1}{e^{-\frac{\pi i}{4}} - e^{\frac{\pi i}{4}}} \left(e^{-\frac{\pi i}{4} - \frac{i\epsilon}{2}} \gamma(-i e^{i\epsilon}) - e^{\frac{\pi i}{4} + \frac{i\epsilon}{2}} \gamma(-i e^{-i\epsilon}) \right), \end{aligned} \quad (2.60)$$

¹²Note that, if we had implemented the grading in (2.53) by the Grassmannality or the ghost number of the states $|\psi\rangle$, then $\mathcal{Q}_{\text{even}}$ would not commute with \mathcal{Q}_{odd} because \mathcal{Q}_{odd} itself is Grassmann-odd and has ghost number 1. So the grading had to be implemented by the GSO parity for $\widehat{\mathcal{Q}}$ to be nilpotent.

in \widehat{Q} , then all of $c_\epsilon(\pm i), \gamma_\epsilon(\pm i)$ turn out to annihilate $\langle \mathcal{I} |$, while in the $\epsilon \rightarrow 0$ limit they naively reduce to the original midpoint insertions. Here we give a slightly different proof than in [20] that $\gamma_\epsilon(i)$ kills the identity. Let us consider

$$\gamma_{(a,b)}(i) = \frac{1}{a+b} (a\gamma(i e^{i\epsilon}) + b\gamma(i e^{-i\epsilon})) \quad (2.61)$$

with ϵ -dependent constants a and b , and see when $\langle \mathcal{I} | \gamma_{(a,b)}(i) | \varphi \rangle$ vanishes. Note that in the $\epsilon \rightarrow 0$ limit $\gamma_{(a,b)}(i)$ reduces to $\gamma(i)$ irrespective of the values of a and b . From the definition (2.59) of the identity, we have

$$\begin{aligned} \langle \mathcal{I} | \gamma_{(a,b)}(i) | \varphi \rangle &= \frac{a}{a+b} \left\langle \left(f_1^{(1)'}(i e^{i\epsilon}) \right)^{-\frac{1}{2}} \gamma(f_1^{(1)}(i e^{i\epsilon})) f_1^{(1)} \circ \varphi(0) \right\rangle_{\text{UHP}} \\ &+ \frac{b}{a+b} \left\langle \left(f_1^{(1)'}(i e^{-i\epsilon}) \right)^{-\frac{1}{2}} \gamma(f_1^{(1)}(i e^{-i\epsilon})) f_1^{(1)} \circ \varphi(0) \right\rangle_{\text{UHP}}. \end{aligned} \quad (2.62)$$

Using the relations

$$\begin{aligned} f_1^{(1)}(i e^{-i\epsilon}) &= \frac{2i e^{-i\epsilon}}{1 + e^{-2i\epsilon}} = \frac{2i e^{i\epsilon}}{e^{2i\epsilon} + 1} = f_1^{(1)}(i e^{i\epsilon}), \\ f_1^{(1)'}(i e^{-i\epsilon}) &= \frac{2(1+z^2)}{(1-z^2)^2} \Big|_{z=i e^{-i\epsilon}} = \frac{2(1 - e^{-2i\epsilon})}{(1 + e^{-2i\epsilon})^2} = e^{2i\epsilon} \frac{2(e^{2i\epsilon} - 1)}{(e^{2i\epsilon} + 1)^2} \\ &= e^{\pi i + 2i\epsilon} \frac{2(1 - e^{2i\epsilon})}{(1 + e^{2i\epsilon})^2} = e^{\pi i + 2i\epsilon} f_1^{(1)'}(i e^{i\epsilon}), \end{aligned}$$

where we have conventionally defined $-1 = e^{\pi i}$, we rewrite eq.(2.62) as

$$\langle \mathcal{I} | \gamma_{(a,b)}(i) | \varphi \rangle = \frac{1}{a+b} \left(f_1^{(1)'}(i e^{i\epsilon}) \right)^{-\frac{1}{2}} \left(a + b e^{-\frac{\pi i}{2} - i\epsilon} \right) \left\langle \gamma(f_1^{(1)}(i e^{i\epsilon})) f_1^{(1)} \circ \varphi(0) \right\rangle_{\text{UHP}}.$$

This expression vanishes when

$$\frac{b}{a} = -e^{\frac{\pi i}{2} + i\epsilon}$$

is satisfied. So, by choosing

$$a = e^{-\frac{\pi i}{4} - \frac{i\epsilon}{2}}, \quad b = -e^{\frac{\pi i}{4} + \frac{i\epsilon}{2}},$$

$\gamma_{(a,b)}(i)$ (2.61) essentially reproduces $\gamma_\epsilon(i)$ of (2.60). In the same way we can prove $\langle \mathcal{I} | \gamma_\epsilon(-i) = 0$, $\langle \mathcal{I} | c_\epsilon(\pm i) = 0$ as well. Furthermore, Q_{ABC} also annihilates $\langle \mathcal{I} |$ because

$$\langle \mathcal{I} | Q_{\text{ABC}} | \varphi \rangle = \left\langle f_1^{(1)} \circ \left(\oint \frac{dz}{2\pi i} b \gamma^2(z) \varphi(0) \right) \right\rangle_{\text{UHP}} = \left\langle \oint \frac{dz'}{2\pi i} b \gamma^2(z') \left(f_1^{(1)} \circ \varphi(0) \right) \right\rangle_{\text{UHP}}$$

is shown to vanish by the contour deformation argument. Therefore, if we define \mathcal{Q}_{odd} and $\mathcal{Q}_{\text{even}}^{\text{GSO}(\pm)}$ as the $\epsilon \rightarrow 0$ limit of

$$\begin{aligned}\mathcal{Q}_{\text{odd},\epsilon} &= \frac{1}{2i\epsilon_r^2} (c_\epsilon(i) - c_\epsilon(-i)) + \frac{q_1^2}{2} \oint \frac{dz}{2\pi i} b\gamma^2(z), \\ \mathcal{Q}_{\text{even},\epsilon}^{\text{GSO}(+)} &= \frac{q_1}{2i\epsilon_r} (\gamma_\epsilon(i) - \gamma_\epsilon(-i)), \\ \mathcal{Q}_{\text{even},\epsilon}^{\text{GSO}(-)} &= \frac{q_1}{2\epsilon_r} (\gamma_\epsilon(i) + \gamma_\epsilon(-i)),\end{aligned}$$

respectively, then we obtain an operator $\widehat{\mathcal{Q}}$ which annihilates the identity.

To give still another argument that $\mathcal{Q}_{\text{even}}$ kills $|\mathcal{I}\rangle$, we notice that the action of $\mathcal{Q}_{\text{even}}$ on a state $|\psi\rangle$ can be expressed¹³ as an inner derivation,¹⁴

$$\begin{aligned}\mathcal{Q}_{\text{even}}|\psi\rangle &= \lim_{\epsilon \rightarrow 0} (|\Sigma_\epsilon * \psi\rangle - (-1)^{\text{GSO}(\psi)} |\psi * \Sigma_\epsilon\rangle), \\ |\Sigma_\epsilon\rangle &= \Gamma_\epsilon |\mathcal{I}\rangle, \\ \Gamma_\epsilon &= q_1 \frac{1-i}{4\epsilon_r} (\gamma(i e^{i\epsilon}) + i\gamma(-i e^{-i\epsilon})).\end{aligned}\tag{2.63}$$

As shown in Appendix A, by considering the inner product $\langle \varphi | \mathcal{Q}_{\text{even}} | \psi \rangle$ with a Fock space state $\langle \varphi |$ we actually recover the previous expression (2.53)

$$\langle \varphi | \mathcal{Q}_{\text{even}} | \psi \rangle = q_1 \frac{1-i}{4\epsilon_r} (1 - i(-1)^{\text{GSO}(\psi)}) \langle \varphi | (\gamma(i) - (-1)^{\text{GSO}(\psi)} \gamma(-i)) | \psi \rangle.\tag{2.64}$$

From the expression (2.63), it is obvious that $\mathcal{Q}_{\text{even}}$ annihilates the identity $|\mathcal{I}\rangle$. Substituting $|\psi\rangle = |\mathcal{I}\rangle$ and $(-1)^{\text{GSO}(\mathcal{I})} = +1$, one finds

$$\mathcal{Q}_{\text{even}}|\mathcal{I}\rangle = \lim_{\epsilon \rightarrow 0} (|\Sigma_\epsilon * \mathcal{I}\rangle - |\mathcal{I} * \Sigma_\epsilon\rangle) = \lim_{\epsilon \rightarrow 0} (|\Sigma_\epsilon\rangle - |\Sigma_\epsilon\rangle) = 0.$$

Derivation property of $\widehat{\mathcal{Q}}$

It is known [15] that $\mathcal{Q}_{\text{GRSZ}} = \frac{1}{2i}(c(i) - c(-i))$ is a graded derivation of the $*$ -algebra because $\mathcal{Q}_{\text{GRSZ}}$ can be written as

$$\begin{aligned}\mathcal{Q}_{\text{GRSZ}} &= \sum_{n=0}^{\infty} (-1)^n \mathcal{C}_{2n}, \\ \mathcal{C}_0 &= c_0, \quad \mathcal{C}_n = c_n + (-1)^n c_{-n} \quad \text{for } n \neq 0,\end{aligned}$$

¹³As was shown in [15], almost the same is true of \mathcal{Q}_{odd} : see eq.(2.78).

¹⁴Note that this expression has the same structure as the original definition (2.19) of $\mathcal{Q}_{\text{even}}$. Since the $\epsilon \rightarrow 0$ limit of Σ_ϵ is not well-defined, $\mathcal{Q}_{\text{even}}$ itself may not be considered as an inner derivation.

and each \mathcal{C}_n obeys the Leibniz rule graded by the Grassmannality [31, 1]. The derivation property of Q_{ABC} , which is the zero-mode of a primary field of conformal weight 1, is proven by the contour deformation argument [31]. Taking the internal Chan-Paton factors into account, $\widehat{\mathcal{Q}}_{\text{odd}} = \mathcal{Q}_{\text{odd}} \otimes \sigma_3$ satisfies

$$\widehat{\mathcal{Q}}_{\text{odd}}(\widehat{A} * \widehat{B}) = (\widehat{\mathcal{Q}}_{\text{odd}}\widehat{A}) * \widehat{B} + (-1)^{\#_{\text{gh}}(\widehat{A})}\widehat{A} * (\widehat{\mathcal{Q}}_{\text{odd}}\widehat{B}), \quad (2.65)$$

where the internal Chan-Paton structure of \widehat{A} and \widehat{B} is given by (2.43). For the case of $\mathcal{Q}_{\text{even}}$, we will make use of the expression (2.63). Let us consider $\mathcal{Q}_{\text{even}}$ acting on the $*$ -product $A * B$ of two states A and B . From the property of the GSO parity that $(-1)^{\text{GSO}(A*B)} = (-1)^{\text{GSO}(A)}(-1)^{\text{GSO}(B)}$ one obtains

$$\begin{aligned} \mathcal{Q}_{\text{even}}|A * B\rangle &= |\Sigma_\epsilon * A * B\rangle - (-1)^{\text{GSO}(A*B)}|A * B * \Sigma_\epsilon\rangle \\ &= (|\Sigma_\epsilon * A\rangle - (-1)^{\text{GSO}(A)}|A * \Sigma_\epsilon\rangle) * |B\rangle \\ &\quad + (-1)^{\text{GSO}(A)}|A\rangle * (|\Sigma_\epsilon * B\rangle - (-1)^{\text{GSO}(B)}|B * \Sigma_\epsilon\rangle) \\ &= |(\mathcal{Q}_{\text{even}}A) * B\rangle + (-1)^{\text{GSO}(A)}|A * (\mathcal{Q}_{\text{even}}B)\rangle, \end{aligned}$$

where we have omitted the symbol $\lim_{\epsilon \rightarrow 0}$. Attaching the Chan-Paton factors to A and B , and then multiplying $i\sigma_2$ from the left, we have for $\widehat{\mathcal{Q}}_{\text{even}} = \mathcal{Q}_{\text{even}} \otimes i\sigma_2$

$$\widehat{\mathcal{Q}}_{\text{even}}|\widehat{A} * \widehat{B}\rangle = |(\widehat{\mathcal{Q}}_{\text{even}}\widehat{A}) * \widehat{B}\rangle + (-1)^{\text{GSO}(\widehat{A})}i\sigma_2|\widehat{A} * (\mathcal{Q}_{\text{even}}\widehat{B})\rangle. \quad (2.66)$$

When $i\sigma_2$ passes \widehat{A} , we find from (2.43) a rule

$$\begin{aligned} i\sigma_2 \cdot \widehat{A} &= -(-1)^{\text{GSO}(\widehat{A})}\widehat{A} \cdot i\sigma_2 && \text{for } \#_{\text{gh}}(\widehat{A}) \text{ odd,} \\ i\sigma_2 \cdot \widehat{A} &= (-1)^{\text{GSO}(\widehat{A})}\widehat{A} \cdot i\sigma_2 && \text{for } \#_{\text{gh}}(\widehat{A}) \text{ even,} \end{aligned}$$

which can be written collectively in the form

$$i\sigma_2 \cdot \widehat{A} = (-1)^{\#_{\text{gh}}(\widehat{A})}(-1)^{\text{GSO}(\widehat{A})}\widehat{A} \cdot i\sigma_2.$$

Applying it to eq.(2.66), we eventually find

$$\widehat{\mathcal{Q}}_{\text{even}}(\widehat{A} * \widehat{B}) = (\widehat{\mathcal{Q}}_{\text{even}}\widehat{A}) * \widehat{B} + (-1)^{\#_{\text{gh}}(\widehat{A})}\widehat{A} * (\widehat{\mathcal{Q}}_{\text{even}}\widehat{B}). \quad (2.67)$$

Since both $\widehat{\mathcal{Q}}_{\text{odd}}$ and $\widehat{\mathcal{Q}}_{\text{even}}$ obey the same Leibniz rule (2.65), (2.67) with the ghost number grading, so does $\widehat{\mathcal{Q}} = \widehat{\mathcal{Q}}_{\text{odd}} - \widehat{\mathcal{Q}}_{\text{even}}$:

$$\widehat{\mathcal{Q}}(\widehat{A} * \widehat{B}) = (\widehat{\mathcal{Q}}\widehat{A}) * \widehat{B} + (-1)^{\#_{\text{gh}}(\widehat{A})}\widehat{A} * (\widehat{\mathcal{Q}}\widehat{B}). \quad (2.68)$$

Hermiticity of $\widehat{\mathcal{Q}}$ in the presence of Y_{-2}

As mentioned before, in the proof of gauge invariance of the action (2.83) we are going to use the hermiticity relations

$$\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{A}, \widehat{B} \rangle\rangle = -(-1)^{\#\text{gh}(\widehat{A})} \langle\langle \widehat{Y}_{-2} | \widehat{A}, \widehat{\mathcal{Q}} \widehat{B} \rangle\rangle, \quad (2.69)$$

or more generally

$$\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{A}_1, \widehat{A}_2 * \dots * \widehat{A}_n \rangle\rangle = -(-1)^{\#\text{gh}(\widehat{A}_1)} \langle\langle \widehat{Y}_{-2} | \widehat{A}_1, \widehat{\mathcal{Q}}(\widehat{A}_2 * \dots * \widehat{A}_n) \rangle\rangle. \quad (2.70)$$

To show them we decompose $\widehat{\mathcal{Q}}, \widehat{A}, \widehat{B}$ into their GSO components. For the first one (2.69) explicit forms are

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{odd}} A_+, B_+ \rangle\rangle = -(-1)^{|A_+|} \langle\langle Y_{-2} | A_+, \mathcal{Q}_{\text{odd}} B_+ \rangle\rangle, \quad (2.71)$$

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{odd}} A_-, B_- \rangle\rangle = -(-1)^{|A_-|} \langle\langle Y_{-2} | A_-, \mathcal{Q}_{\text{odd}} B_- \rangle\rangle, \quad (2.72)$$

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}} A_+, B_- \rangle\rangle = -\langle\langle Y_{-2} | A_+, \mathcal{Q}_{\text{even}} B_- \rangle\rangle, \quad (2.73)$$

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}} A_-, B_+ \rangle\rangle = \langle\langle Y_{-2} | A_-, \mathcal{Q}_{\text{even}} B_+ \rangle\rangle. \quad (2.74)$$

(2.73), (2.74) have already been proved below eq.(2.45). We can show that relations (2.71), (2.72) are satisfied by $\mathcal{Q}_{\text{GRSZ}}$ from the precise definition of the 2-vertex,

$$\langle\langle Y_{-2} | A, B \rangle\rangle = \langle Y(i)Y(-i)A(0)\mathcal{R}_\pi \circ B(0) \rangle_{\text{UHP}}, \quad (2.75)$$

and the conformal transformation of $\mathcal{Q}_{\text{GRSZ}}$ under \mathcal{R}_π ,

$$\mathcal{R}_\pi \circ \mathcal{Q}_{\text{GRSZ}} = -\mathcal{Q}_{\text{GRSZ}}.$$

For Q_{ABC} , eqs.(2.71), (2.72) are equivalent to the statement

$$\left\langle Y(i)Y(-i) \oint_C \frac{dz}{2\pi i} b\gamma^2(z)(A_\pm(0) \mathcal{R}_\pi \circ B_\pm(0)) \right\rangle_{\text{UHP}} = 0, \quad (2.76)$$

where C is an integration contour encircling 0 and $\mathcal{R}_\pi(0) = \infty$, but not $\pm i$. By deforming the contour this equation is restated as

$$[Q_{\text{ABC}}, Y_{-2}] \equiv \left(\oint_{C(i)} \frac{dz}{2\pi i} b\gamma^2(z)Y(i) \right) Y(-i) + \left(\oint_{C(-i)} \frac{dz}{2\pi i} b\gamma^2(z)Y(-i) \right) Y(i) = 0, \quad (2.77)$$

where $C(\pm i)$ are small contours encircling $\pm i$, respectively. This equality holds because the operator product between $b\gamma^2 = b\eta\partial\eta e^{2\phi}$ and $Y = c\partial\xi e^{-2\phi}$ is non-singular. Thus we have shown that $\widehat{\mathcal{Q}}$ satisfies the hermiticity relation (2.69).

In the more general case (2.70), we make use of the inner derivation formulas: eq.(2.63) for $\mathcal{Q}_{\text{even}}$ and

$$\begin{aligned}\mathcal{Q}_{\text{GRSZ}}|\psi\rangle &= \lim_{\epsilon \rightarrow 0} (|S_\epsilon * \psi\rangle - (-1)^{|\psi|} |\psi * S_\epsilon\rangle), \\ |S_\epsilon\rangle &= \frac{1}{4i} (e^{-i\epsilon} c(i e^{i\epsilon}) - e^{i\epsilon} c(-i e^{-i\epsilon})) |\mathcal{I}\rangle,\end{aligned}\tag{2.78}$$

for $\mathcal{Q}_{\text{GRSZ}}$ [15, 32]. We find

$$\begin{aligned}\langle\langle Y_{-2} | \mathcal{Q}_{\text{GRSZ}} A_{1+}, (\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle &= \lim_{\epsilon \rightarrow 0} \left(\langle\langle Y_{-2} | S_\epsilon * A_{1+}, (\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle \right. \\ &\quad \left. - (-1)^{|A_{1+}|} \langle\langle Y_{-2} | A_{1+} * S_\epsilon, (\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle \right) \\ &= -(-1)^{|A_{1+}|} \lim_{\epsilon \rightarrow 0} \left(\langle\langle Y_{-2} | A_{1+}, S_\epsilon * (\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle \right. \\ &\quad \left. - (-1)^{|\widehat{A}_2 * \dots * \widehat{A}_n|} \langle\langle Y_{-2} | A_{1+}, (\widehat{A}_2 * \dots * \widehat{A}_n)_+ * S_\epsilon \rangle\rangle \right) \\ &= -(-1)^{|A_{1+}|} \langle\langle Y_{-2} | A_{1+}, \mathcal{Q}_{\text{GRSZ}}(\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle,\end{aligned}\tag{2.79}$$

and similarly

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{GRSZ}} A_{1-}, (\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle = -(-1)^{|A_{1-}|} \langle\langle Y_{-2} | A_{1-}, \mathcal{Q}_{\text{GRSZ}}(\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle,\tag{2.80}$$

where we have used the associativity of the $*$ -product and cyclicity for a GSO(+/-) state S_ϵ , and $(\widehat{A}_2 * \dots * \widehat{A}_n)_{+/-}$ denote the GSO(+/-) components of $\widehat{A}_2 * \dots * \widehat{A}_n$. Note that $(-1)^{|A_{1+}|}$ and $(-1)^{|A_2 * \dots * A_n|}$ agree with each other since the whole insertion inside the correlator must be Grassmann-odd to give a non-vanishing value. In the same way as above, we get

$$\begin{aligned}\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}} A_{1+}, (\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle &= \lim_{\epsilon \rightarrow 0} \left(\langle\langle Y_{-2} | \Sigma_\epsilon * A_{1+}, (\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle \right. \\ &\quad \left. - (-1)^{\text{GSO}(A_{1+})} \langle\langle Y_{-2} | A_{1+} * \Sigma_\epsilon, (\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle \right) \\ &= -\lim_{\epsilon \rightarrow 0} \left(\langle\langle Y_{-2} | A_{1+}, \Sigma_\epsilon * (\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle - (-1) \langle\langle Y_{-2} | A_{1+}, (\widehat{A}_2 * \dots * \widehat{A}_n)_- * \Sigma_\epsilon \rangle\rangle \right) \\ &= -\langle\langle Y_{-2} | A_{1+}, \mathcal{Q}_{\text{even}}(\widehat{A}_2 * \dots * \widehat{A}_n)_- \rangle\rangle,\end{aligned}\tag{2.81}$$

and

$$\langle\langle Y_{-2} | \mathcal{Q}_{\text{even}} A_{1-}, (\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle = \langle\langle Y_{-2} | A_{1-}, \mathcal{Q}_{\text{even}}(\widehat{A}_2 * \dots * \widehat{A}_n)_+ \rangle\rangle.\tag{2.82}$$

In the second equality of (2.81), an additional sign factor (-1) has arisen because Σ_ϵ lives in the GSO(-) sector (see eq.(2.9)). The proof of (2.70) for \mathcal{Q}_{ABC} is essentially the same as in the previous case (2.76). Collecting eqs.(2.79)–(2.82) in the matrix form gives eq.(2.70).

2.3 Gauge invariant cubic action of vacuum superstring field theory

In this subsection we will show that the following cubic action

$$S_V = -\frac{\kappa_0}{2} \text{Tr} \left[\frac{1}{2} \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{Q}} \widehat{\mathcal{A}} \rangle\rangle + \frac{1}{3} \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle \right] \quad (2.83)$$

with $\widehat{\mathcal{Q}}$ given above (eqs.(2.54), (2.52)) is invariant under the gauge transformation

$$\delta \widehat{\mathcal{A}} = \widehat{\mathcal{Q}} \widehat{\Lambda} + \widehat{\mathcal{A}} * \widehat{\Lambda} - \widehat{\Lambda} * \widehat{\mathcal{A}}, \quad (2.84)$$

where $\widehat{\Lambda}$ is an infinitesimal gauge parameter of ghost number 0 and picture number 0, whose internal Chan-Paton structure is

$$\widehat{\Lambda} = \Lambda_+ \otimes \mathbf{1} + \Lambda_- \otimes \sigma_1. \quad (2.85)$$

The gauge variation of the action linear in $\widehat{\Lambda}$ is

$$\begin{aligned} \delta S_V &= -\frac{\kappa_0}{2} \text{Tr} \left[\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{Q}}(\delta \widehat{\mathcal{A}}) \rangle\rangle + \langle\langle \widehat{Y}_{-2} | \delta \widehat{\mathcal{A}}, \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle \right] \\ &= -\frac{\kappa_0}{2} \text{Tr} \left[\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{Q}}^2 \widehat{\Lambda} \rangle\rangle + \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{Q}}(\widehat{\mathcal{A}} * \widehat{\Lambda}) \rangle\rangle - \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{Q}}(\widehat{\Lambda} * \widehat{\mathcal{A}}) \rangle\rangle \right. \\ &\quad \left. + \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{\Lambda}, \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle + \langle\langle \widehat{Y}_{-2} | (\widehat{\mathcal{A}} * \widehat{\Lambda}), \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle - \langle\langle \widehat{Y}_{-2} | (\widehat{\Lambda} * \widehat{\mathcal{A}}), \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle \right], \end{aligned} \quad (2.86)$$

where we have used the cyclicity of the vertices and the hermiticity (2.69) of $\widehat{\mathcal{Q}}$. Making use of the cyclicity and the associativity of the $*$ -product, the last two terms cancel each other. The first term vanishes as well thanks to the nilpotency of $\widehat{\mathcal{Q}}$. Using the cyclicity, the ‘partial integration’ (2.70) and the derivation property (2.68) of $\widehat{\mathcal{Q}}$, the second and the third terms are written as

$$\begin{aligned} &\text{Tr} \left(\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{\Lambda}, \widehat{\mathcal{A}} * \widehat{\Lambda} \rangle\rangle - \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{\Lambda}, \widehat{\Lambda} * \widehat{\mathcal{A}} \rangle\rangle \right) \\ &= \text{Tr} \left(\langle\langle \widehat{Y}_{-2} | \widehat{\Lambda}, \widehat{\mathcal{Q}} \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle - \langle\langle \widehat{Y}_{-2} | \widehat{\Lambda}, \widehat{\mathcal{A}} * \widehat{\mathcal{Q}} \widehat{\mathcal{A}} \rangle\rangle \right) \\ &= \text{Tr} \langle\langle \widehat{Y}_{-2} | \widehat{\Lambda}, \widehat{\mathcal{Q}}(\widehat{\mathcal{A}} * \widehat{\mathcal{A}}) \rangle\rangle = -\text{Tr} \langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{Q}} \widehat{\Lambda}, \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle\rangle, \end{aligned}$$

which precisely cancels the fourth term. This completes the proof of gauge invariance. Therefore the action (2.83) defines a consistent gauge theory of a string field, at least at the classical level. Moreover, the kinetic operator $\widehat{\mathcal{Q}}$, which governs the perturbative nature of the string field around $\widehat{\mathcal{A}} = 0$, has the following properties:

1. $\widehat{\mathcal{Q}}$ is made purely out of ghost operators;
2. $\widehat{\mathcal{Q}}$ has vanishing cohomology, so that there are no perturbative physical open string states¹⁵ around the vacuum $\widehat{\mathcal{A}} = 0$;
3. $\widehat{\mathcal{Q}}$ contains non-zero $\mathcal{Q}_{\text{even}}$ so that the unwanted \mathbb{Z}_2 reflection symmetry is broken, as it should be;
4. $\widehat{\mathcal{Q}}$ has been constructed in such a way that the action S_V (2.83) is invariant under the twist transformation (2.32).

The property 1 is obvious from the explicit form (2.54), (2.52) of $\widehat{\mathcal{Q}}$. To show the property 2, suppose that we have a state $|\widehat{\mathcal{A}}\rangle$ which is annihilated by $\widehat{\mathcal{Q}}$. Then $|\widehat{\mathcal{A}}\rangle$ itself can be written as

$$|\widehat{\mathcal{A}}\rangle = \{\widehat{\mathcal{Q}}, \varepsilon_r^2 \widehat{b}_0\} |\widehat{\mathcal{A}}\rangle = \widehat{\mathcal{Q}}(\varepsilon_r^2 \widehat{b}_0 |\widehat{\mathcal{A}}\rangle),$$

where $\widehat{b}_0 = b_0 \otimes \sigma_3$. Since any $\widehat{\mathcal{Q}}$ -closed state $|\widehat{\mathcal{A}}\rangle$ has, at least formally, been expressed as a $\widehat{\mathcal{Q}}$ -exact form, it follows that $\widehat{\mathcal{Q}}$ has vanishing cohomology. Twist invariance of the action is explicitly shown in Appendix B. From the properties mentioned above, it seems that the action (2.83) is well suited for the description of cubic superstring field theory around one of the tachyon vacua. Although this action is gauge-invariant and is well-defined even for finite ε_r , if we are to consider that the vacuum superstring field theory action (2.83) is derived from the original one (2.1) through the tachyon condensation followed by a field redefinition, the parameter ε_r should be taken to zero.

3 Construction of a Spacetime-Independent Solution: ε_r -Expansion

Now that we have determined the precise form of the cubic vacuum superstring field theory action, we set about constructing solutions in this theory. Varying the action (2.83), we obtain the following equations of motion:

$$\mathcal{F}(\widehat{\mathcal{A}}) \equiv \widehat{\mathcal{Q}}\widehat{\mathcal{A}} + \widehat{\mathcal{A}} * \widehat{\mathcal{A}} = 0 \tag{3.1}$$

or, in the GSO-component form $\mathcal{F}(\widehat{\mathcal{A}}) = \mathcal{F}_+ \otimes \mathbf{1} + \mathcal{F}_- \otimes \sigma_1$,

$$\mathcal{F}_+ \equiv \mathcal{Q}_{\text{odd}}\mathcal{A}_+ + \mathcal{A}_+ * \mathcal{A}_+ + \mathcal{Q}_{\text{even}}\mathcal{A}_- - \mathcal{A}_- * \mathcal{A}_- = 0, \tag{3.2}$$

$$\mathcal{F}_- \equiv \mathcal{Q}_{\text{odd}}\mathcal{A}_- + \mathcal{Q}_{\text{even}}\mathcal{A}_+ + \mathcal{A}_+ * \mathcal{A}_- - \mathcal{A}_- * \mathcal{A}_+ = 0. \tag{3.3}$$

¹⁵We are ignoring the effect of the non-trivial kernel of Y_{-2} on the physical open string spectrum [33].

It has been discussed in the literature [12, 11, 13, 9] that, even if the inner product $\langle \widehat{\varphi} | \mathcal{F}(\widehat{\mathcal{A}}) \rangle$ with any Fock space state $\langle \widehat{\varphi} |$ vanishes, it does *not* follow that $\langle \widehat{X} | \mathcal{F}(\widehat{\mathcal{A}}) \rangle$ vanishes for more general states $\langle \widehat{X} |$. However, since it seems that what matters in this argument is the normalization of the solution $\widehat{\mathcal{A}}$ and we are not in a position to deal with it in detail, we consider the simplest case where we require

$$\langle \widehat{\varphi} | \mathcal{F}(\widehat{\mathcal{A}}) \rangle = \langle \widehat{\varphi} | \widehat{\mathcal{Q}}\widehat{\mathcal{A}} + \widehat{\mathcal{A}} * \widehat{\mathcal{A}} \rangle = 0 \quad (3.4)$$

to hold for any Fock space state $\langle \widehat{\varphi} |$ of ghost number 1 and *picture number* -2 , instead of inserting the double-step inverse picture changing operator Y_{-2} . In the component form, it becomes

$$\langle \varphi_+ | \mathcal{F}_+ \rangle = \langle \varphi_- | \mathcal{F}_- \rangle = 0, \quad (3.5)$$

where φ_+, φ_- denote the GSO(+)-, GSO(-)-components of $\widehat{\varphi} = \varphi_+ \otimes \sigma_3 + \varphi_- \otimes i\sigma_2$, respectively.

The fact that we must handle two quite different types of operators $\mathcal{Q}_{\text{GRSZ}} = \frac{1}{2i}(c(i) - c(-i))$, $Q_{\text{ABC}} = \oint \frac{dz}{2\pi i} b\gamma^2(z)$ at the same time makes it difficult to find solutions of the equations of motion. Now let us recall that in the ‘realistic’ vacuum superstring field theory the parameter ε_r has to be taken to zero and that $\mathcal{Q}_{\text{GRSZ}}$ and Q_{ABC} enter \mathcal{Q}_{odd} in different orders in ε_r . Multiplied by ε_r^4 , the equations of motion (3.5) can be written as

$$\begin{aligned} \langle \varphi_+ | \left(\mathcal{Q}_{\text{GRSZ}} + \frac{q_1^2 \varepsilon_r^2}{2} Q_{\text{ABC}} \right) (\varepsilon_r^2 \mathcal{A}_+) + (\varepsilon_r^2 \mathcal{A}_+) * (\varepsilon_r^2 \mathcal{A}_+) \\ + \varepsilon_r \Gamma(\varepsilon_r^2 \mathcal{A}_-) - (\varepsilon_r^2 \mathcal{A}_-) * (\varepsilon_r^2 \mathcal{A}_-) \rangle = 0, \end{aligned} \quad (3.6)$$

$$\begin{aligned} \langle \varphi_- | \left(\mathcal{Q}_{\text{GRSZ}} + \frac{q_1^2 \varepsilon_r^2}{2} Q_{\text{ABC}} \right) (\varepsilon_r^2 \mathcal{A}_-) + \varepsilon_r \Gamma(\varepsilon_r^2 \mathcal{A}_+) \\ + (\varepsilon_r^2 \mathcal{A}_+) * (\varepsilon_r^2 \mathcal{A}_-) - (\varepsilon_r^2 \mathcal{A}_-) * (\varepsilon_r^2 \mathcal{A}_+) \rangle = 0, \end{aligned} \quad (3.7)$$

where we have introduced the notation $\Gamma = \varepsilon_r \mathcal{Q}_{\text{even}}$ which is finite in the $\varepsilon_r \rightarrow 0$ limit. If we assume that $\mathcal{A}'_{\pm} \equiv \varepsilon_r^2 \mathcal{A}_{\pm}$ can be expanded in a power series in ε_r as

$$\mathcal{A}'_+ = \mathcal{A}_+^{(0)} + \varepsilon_r \mathcal{A}_+^{(1)} + \varepsilon_r^2 \mathcal{A}_+^{(2)} + \dots, \quad (3.8)$$

$$\mathcal{A}'_- = \mathcal{A}_-^{(0)} + \varepsilon_r \mathcal{A}_-^{(1)} + \varepsilon_r^2 \mathcal{A}_-^{(2)} + \dots, \quad (3.9)$$

then we can try to solve the equations of motion order by order in ε_r . In the $\varepsilon_r \rightarrow 0$ limit we can expect that the full solutions \mathcal{A}'_{\pm} are well approximated by terms of lowest orders. At order $(\varepsilon_r)^0$, we have

$$\langle \varphi_+ | (\mathcal{Q}_{\text{GRSZ}} \mathcal{A}_+^{(0)} + \mathcal{A}_+^{(0)} * \mathcal{A}_+^{(0)} - \mathcal{A}_-^{(0)} * \mathcal{A}_-^{(0)}) \rangle = 0, \quad (3.10)$$

$$\langle \varphi_- | (\mathcal{Q}_{\text{GRSZ}} \mathcal{A}_-^{(0)} + \mathcal{A}_+^{(0)} * \mathcal{A}_-^{(0)} - \mathcal{A}_-^{(0)} * \mathcal{A}_+^{(0)}) \rangle = 0. \quad (3.11)$$

These equations admit a solution $\mathcal{A}_-^{(0)} = 0$ with $\mathcal{A}_+^{(0)} \neq 0$. Then we are left with

$$\langle \varphi_+ | (\mathcal{Q}_{\text{GRSZ}} \mathcal{A}_+^{(0)} + \mathcal{A}_+^{(0)} * \mathcal{A}_+^{(0)}) \rangle = 0, \quad (3.12)$$

which has exactly the same form as the equation of motion of bosonic VSFT. It is known that this equation is solved by the ‘ bc -twisted sliver state’ [15] of ghost number 1,

$$|\mathcal{A}_+^{(0)}\rangle = \mathcal{N}_+^{(0)} |\Xi'\rangle, \quad (3.13)$$

$$\langle \Xi' | \varphi \rangle \equiv \lim_{n \rightarrow \infty} \langle f^{(n)} \circ \varphi'(0) \rangle'_{\text{UHP}} = \lim_{n \rightarrow \infty} \kappa^{(n)} \langle \tilde{f}^{(n)} \circ \varphi'(0) \rangle'_{\text{UHP}}, \quad (3.14)$$

where the conformal maps $f^{(n)}, \tilde{f}^{(n)}$ are defined by

$$\begin{aligned} f^{(n)}(z) &= h^{-1}(h(z)^{2/n}), \\ \tilde{f}^{(n)}(z) &= M_n \circ f^{(n)}(z) = \frac{n}{2} h^{-1}(h(z)^{2/n}), \quad \left(M_n(z) = \frac{n}{2} z \right), \end{aligned} \quad (3.15)$$

and $|\varphi\rangle = \varphi'(0)|0'\rangle$ with $|0'\rangle$ denoting the $SL(2, \mathbb{R})$ -invariant vacuum of the twisted conformal field theory CFT' and $\langle \dots \rangle'_{\text{UHP}}$ represents the correlation function on the upper half plane in CFT': See [15] for more detail. Two expressions in (3.14) agree with each other thanks to the $SL(2, \mathbb{R})$ -invariance of the correlation function, but we have put a multiplicative factor $\kappa^{(n)}$ which might arise due to the conformal anomaly because CFT' has a non-vanishing total central charge $c' = 24$. The advantage of defining the sliver as the $n \rightarrow \infty$ limit of finite- n wedge states is that we can explicitly compute the $*$ -product using the generalized gluing and resmoothing theorem, even when some operators are inserted on the sliver. The normalization factor $\mathcal{N}_+^{(0)}$ in front of Ξ' is determined by the equation of motion (3.12) as

$$\mathcal{N}_+^{(0)} = -\frac{1}{e^{2c'K} K^{(3)} |f_1^{(3)'}(i)|^{1/4} |(h \circ \tilde{f}^{(\infty)})'(i)|^{-1/4} |h \circ \tilde{f}^{(\infty)}(i) - h \circ \tilde{f}^{(\infty)}(-i)|^{1/4}}, \quad (3.16)$$

though its precise expression is not important. A constant $K^{(3)}$ arises when we relate correlation functions in the untwisted CFT and those in the twisted CFT' [15, 34]:

$$\begin{aligned} &\langle f_1^{(3)} \circ \Phi_1(0) f_2^{(3)} \circ \Phi_2(0) f_3^{(3)} \circ \Phi_3(0) \rangle_{\text{UHP}} \\ &= K^{(3)} \langle f_1^{(3)} \circ \Phi_1'(0) f_2^{(3)} \circ \Phi_2'(0) f_3^{(3)} \circ \Phi_3'(0) e^{\frac{1}{2}\rho(i)} e^{\frac{1}{2}\tilde{\rho}(i)} \rangle'_{\text{UHP}}, \end{aligned} \quad (3.17)$$

where ρ is the bosonized ghost field defined as $c(z) = e^{\rho(z)}, b(z) = e^{-\rho(z)}$ with the OPE $\rho(z)\rho(0) \sim \log z$, whereas another constant K arises when we use the gluing theorem in a conformal field theory with a non-vanishing central charge c [35]:

$$\begin{aligned} &\sum_r \langle f_1 \circ \Phi_{r_1}(0) \dots f_n \circ \Phi_{r_n}(0) f \circ \Phi_r(0) \rangle_{\mathcal{D}_1} \langle g_1 \circ \Phi_{s_1}(0) \dots g_m \circ \Phi_{s_m}(0) g \circ \Phi_r^c(0) \rangle_{\mathcal{D}_2} \\ &= e^{cK} \langle F_1 \circ f_1 \circ \Phi_{r_1}(0) \dots F_n \circ f_n \circ \Phi_{r_n}(0) \widehat{F}_2 \circ g_1 \circ \Phi_{s_1}(0) \dots \widehat{F}_2 \circ g_m \circ \Phi_{s_m}(0) \rangle_{\mathcal{D}}. \end{aligned} \quad (3.18)$$

At order $(\varepsilon_r)^1$, the equations of motion become

$$\langle \varphi_+ | (\mathcal{Q}_{\text{GRSZ}} \mathcal{A}_+^{(1)} + \mathcal{A}_+^{(0)} * \mathcal{A}_+^{(1)} + \mathcal{A}_+^{(1)} * \mathcal{A}_+^{(0)}) \rangle = 0, \quad (3.19)$$

$$\langle \varphi_- | (\mathcal{Q}_{\text{GRSZ}} \mathcal{A}_-^{(1)} + \Gamma \mathcal{A}_+^{(0)} + \mathcal{A}_+^{(0)} * \mathcal{A}_-^{(1)} - \mathcal{A}_-^{(1)} * \mathcal{A}_+^{(0)}) \rangle = 0, \quad (3.20)$$

where we have already used the result $\mathcal{A}_-^{(0)} = 0$ at the zeroth order. First we note that eq.(3.19) is solved by $\mathcal{A}_+^{(1)} = 0$. In fact, (3.12) + $\varepsilon_r \times$ (3.19) gives

$$\langle \varphi_+ | \left(\mathcal{Q}_{\text{GRSZ}} (\mathcal{A}_+^{(0)} + \varepsilon_r \mathcal{A}_+^{(1)}) + (\mathcal{A}_+^{(0)} + \varepsilon_r \mathcal{A}_+^{(1)}) * (\mathcal{A}_+^{(0)} + \varepsilon_r \mathcal{A}_+^{(1)}) \right) \rangle = 0$$

up to the ε_r^2 -term. Since $\mathcal{A}_+^{(0)}$ and $\mathcal{A}_+^{(0)} + \varepsilon_r \mathcal{A}_+^{(1)}$ satisfy the same equation, $\mathcal{A}_+^{(1)}$ should be equal to zero. Next we compute each term in the left hand side of eq.(3.20). We begin by showing that

$$\langle \varphi_- | \mathcal{Q}_{\text{GRSZ}} | \psi_- \rangle = \langle \psi_- | \mathcal{Q}_{\text{GRSZ}} | \varphi_- \rangle, \quad (3.21)$$

$$\langle \varphi_- | \Gamma^{\text{GSO}(+)} | \psi_+ \rangle = \langle \psi_+ | \Gamma^{\text{GSO}(-)} | \varphi_- \rangle, \quad (3.22)$$

hold for any Fock space states φ_- and ψ_{\pm} , with the subscripts \pm denoting their GSO parities. The left hand side of (3.21) can be handled as

$$\begin{aligned} \langle \varphi_- | \mathcal{Q}_{\text{GRSZ}} | \psi_- \rangle &= \frac{1}{2i} \langle \varphi_-(0) \mathcal{R}_\pi \circ (c(i) - c(-i)) \mathcal{R}_\pi \circ \psi_-(0) \rangle_{\text{UHP}} \\ &= \frac{1}{2i} \langle \psi_-(0) (\mathcal{R}_\pi \circ \mathcal{R}_{-\pi}) \circ (c(i) - c(-i)) \mathcal{R}_{-\pi} \circ \varphi_-(0) \rangle_{\text{UHP}} \\ &= \frac{1}{2i} \langle \psi_-(0) \mathcal{R}_\pi \circ (-c(i) + c(-i)) (-\mathcal{R}_\pi \circ \varphi_-(0)) \rangle_{\text{UHP}} \\ &= \langle \psi_- | \mathcal{Q}_{\text{GRSZ}} | \varphi_- \rangle, \end{aligned}$$

where we used the $SL(2, \mathbb{R})$ -invariance of the correlator under $\mathcal{R}_{-\pi}$, and the fact that the vertex operators φ_-, ψ_- associated with the ghost number 1, $\text{GSO}(-)$ states are Grassmann-even. In the same way, we find the left hand side of eq.(3.22) to be

$$\begin{aligned} \langle \varphi_- | \Gamma^{\text{GSO}(+)} | \psi_+ \rangle &= \frac{q_1}{2i} \langle \varphi_-(0) \mathcal{R}_\pi \circ (\gamma(i) - \gamma(-i)) \mathcal{R}_\pi \circ \psi_+(0) \rangle_{\text{UHP}} \\ &= \frac{q_1}{2i} \langle \psi_+(0) (\mathcal{R}_\pi \circ \mathcal{R}_{-\pi}) \circ (\gamma(i) - \gamma(-i)) \mathcal{R}_{-\pi} \circ \varphi_-(0) \rangle_{\text{UHP}} \\ &= \frac{q_1}{2i} \langle \psi_+(0) \mathcal{R}_\pi \circ ((-i)\gamma(i) - i\gamma(-i)) (-\mathcal{R}_\pi \circ \varphi_-(0)) \rangle_{\text{UHP}} \\ &= \langle \psi_+(0) \mathcal{R}_\pi \circ \left(\frac{q_1}{2} (\gamma(i) + \gamma(-i)) \right) \mathcal{R}_\pi \circ \varphi_-(0) \rangle_{\text{UHP}} \\ &= \langle \psi_+ | \Gamma^{\text{GSO}(-)} | \varphi_- \rangle. \end{aligned}$$

Although $\mathcal{A}_+^{(0)}$ and $\mathcal{A}_-^{(1)}$ are not Fock space states, we should consider them to satisfy eqs.(3.21), (3.22) with ψ_\pm replaced by \mathcal{A}_\pm . Now, we make an ansatz for the solution $\langle \mathcal{A}_-^{(1)} |$ as

$$\langle \mathcal{A}_-^{(1)} | \varphi \rangle = \lim_{n \rightarrow \infty} \left\langle \oint_{f^{(n)}(\mathcal{C})} \frac{dz}{2\pi i} s(z) b'(z) \gamma(z) f^{(n)} \circ \varphi'(0) \right\rangle'_{\text{UHP}}, \quad (3.23)$$

or conveniently

$$\langle \mathcal{A}_-^{(1)} | = \langle \Xi' | \oint_{\mathcal{C}} \frac{d\xi}{2\pi i} s(\xi) b' \gamma(\xi). \quad (3.24)$$

Here $s(z)$ is a globally defined holomorphic field of conformal weight $1/2$ such that $dz s(z) b' \gamma(z)$ transforms as a 1-form in CFT'. $s(z)$ is required to be holomorphic everywhere, except at the puncture $z = 0$, in the global complex z -plane which is obtained as a double-cover of the upper half plane for interacting open strings. For $s(z)$ to be regular at infinity, $\lim_{z \rightarrow \infty} z s(z)$ must be finite. Since γ field in this representation is periodic in the NS sector, $s(z)$ must also be periodic under $z \rightarrow e^{2\pi i} z$ for the integral to be well-defined. Furthermore, $s(z)$ must be a Grassmann-even quantity because the GSO(-) state $\langle \mathcal{A}_-^{(1)} |$ of ghost number 1 should be Grassmann-even. (Note that Ξ' itself is Grassmann-odd.) The integration contour \mathcal{C} goes along the double cover of the ‘open string’ $|\xi| = 1$ counterclockwise. More precisely, anticipating the insertions at the open string midpoint $z = i$ and its mirror image $z = -i$, we define the contour \mathcal{C} to be the one indicated in Figure 3. That is to say, the points $\pm i$ lie *outside* the contour, where




Figure 3: The closed contour \mathcal{C} along the open string $|\xi| = 1$.

we are referring to the left-side region of an oriented closed contour, when we walk along it, as the ‘inside’ of the contour. The overall normalization of $\mathcal{A}_-^{(1)}$ has been absorbed in

the definition of s . At last we compute

$$\begin{aligned} \langle \varphi_- | \mathcal{Q}_{\text{GRSZ}} | \mathcal{A}_-^{(1)} \rangle &= \langle \mathcal{A}_-^{(1)} | \mathcal{Q}_{\text{GRSZ}} | \varphi_- \rangle \\ &= \lim_{n \rightarrow \infty} \kappa^{(n)} \left\langle \oint_{\tilde{f}^{(n)}(\mathcal{C})} \frac{dz}{2\pi i} s(z) b' \gamma(z) \frac{1}{2} \left(c'(\tilde{f}^{(n)}(i)) + c'(\tilde{f}^{(n)}(-i)) \right) \tilde{f}^{(n)} \circ \varphi'_-(0) \right\rangle'_{\text{UHP}}, \end{aligned}$$

where we have used eqs.(3.21), (3.23) and (3.14). In CFT' $\mathcal{Q}_{\text{GRSZ}}$ is written as $\frac{1}{2}(c'(i) + c'(-i))$ and c' has conformal weight 0. Since $sb'\gamma(z)$ is holomorphic everywhere except at the origin, the integration contour $\tilde{f}^{(n)}(\mathcal{C})$ can be deformed in the way indicated in Figure 4. We call the new contour $\mathcal{C}'_{(n)}$. In the limit $n \rightarrow \infty$, both $\tilde{f}^{(n)}(i) = ni/2$ and




Figure 4: Deformation of the integration contour from $\tilde{f}^{(n)}(\mathcal{C})$ to $\mathcal{C}'_{(n)}$.

$\tilde{f}^{(n)}(-i) = -ni/2$ go to the same point at infinity and $\mathcal{C}'_{(\infty)}$ becomes a small contour encircling the infinity in the clockwise direction. Picking up the pole coming from the operator product of b' with c' , we obtain

$$\langle \varphi_- | \mathcal{Q}_{\text{GRSZ}} | \mathcal{A}_-^{(1)} \rangle = -\kappa^{(\infty)} s(\infty) \langle \gamma(\infty) \tilde{f}^{(\infty)} \circ \varphi'_-(0) \rangle'_{\text{UHP}}. \quad (3.25)$$

The next one to calculate is

$$\langle \varphi_- | \Gamma | \mathcal{A}_+^{(0)} \rangle = \langle \mathcal{A}_+^{(0)} | \Gamma^{\text{GSO}(-)} | \varphi_- \rangle = \frac{q_1}{2} \mathcal{N}_+^{(0)} \langle \Xi' | (\gamma(i) + \gamma(-i)) | \varphi_- \rangle.$$

This is easily evaluated to give

$$\langle \varphi_- | \Gamma | \mathcal{A}_+^{(0)} \rangle = \frac{q_1}{2} \mathcal{N}_+^{(0)} \kappa^{(\infty)} \left([\tilde{f}^{(\infty)'}(i)]^{-\frac{1}{2}} + [\tilde{f}^{(\infty)'(-i)}]^{-\frac{1}{2}} \right) \langle \gamma(\infty) \tilde{f}^{(\infty)} \circ \varphi'_-(0) \rangle'_{\text{UHP}}. \quad (3.26)$$

We then turn to the third term in (3.20). Inserting the complete set of states $\mathbf{1} = \sum_r |\Phi_r\rangle\langle\Phi_r^c|$, we have

$$\begin{aligned} \langle\varphi_-|\mathcal{A}_+^{(0)} * \mathcal{A}_-^{(1)}\rangle &= \mathcal{N}_+^{(0)} \sum_{r,s} \langle\varphi_-| \left(|\Phi_r\rangle * \left(\mathcal{R}_{-\pi} \circ \oint_{\mathcal{C}} \frac{d\xi}{2\pi i} s(\xi) b' \gamma(\xi) \right) |\Phi_s\rangle \right) \langle\Phi_r^c|\Xi'\rangle \langle\Phi_s^c|\Xi'\rangle \\ &= \mathcal{N}_+^{(0)} \sum_{r,s} \left\langle f_1^{(3)} \circ \varphi_-(0) f_2^{(3)} \circ \Phi_r(0) f_3^{(3)} \circ \left[\mathcal{R}_{-\pi} \circ \left(\oint_{\mathcal{C}} \frac{d\xi}{2\pi i} s(\xi) \xi b(\xi) \gamma(\xi) \right) \Phi_s(0) \right] \right\rangle_{\text{UHP}} \\ &\quad \times \lim_{n \rightarrow \infty} \langle f^{(n)} \circ \Phi_r^{c'}(0) \rangle'_{\text{UHP}} \langle f^{(n)} \circ \Phi_s^{c'}(0) \rangle'_{\text{UHP}}, \end{aligned} \quad (3.27)$$

where we have used

$$|\mathcal{A}_-^{(1)}\rangle = \mathcal{R}_{-\pi} \circ \left(\oint_{\mathcal{C}} \frac{d\xi}{2\pi i} s(\xi) b' \gamma(\xi) \right) |\Xi'\rangle.$$

After rewriting the CFT correlator in (3.27) as a CFT' correlator using eq.(3.17), we can express (3.27) as a single correlator by making use of the gluing theorem (3.18) twice. From the formulas given in [32] we obtain the result

$$\begin{aligned} \langle\varphi_-|\mathcal{A}_+^{(0)} * \mathcal{A}_-^{(1)}\rangle &= \mathcal{N}_+^{(0)} K^{(3)} |f_1^{(3)'}(i)|^{1/4} e^{2c'K} \\ &\quad \times \lim_{n \rightarrow \infty} \kappa^{(n)} \left\langle \oint_{\mathcal{C}'_{(2n-1)}} \frac{dz}{2\pi i} s(z) b' \gamma(z) \tilde{f}^{(2n-1)} \circ \left(e^{\frac{1}{2}\rho(i)} e^{\frac{1}{2}\rho(-i)} \varphi'_-(0) \right) \right\rangle'_{\text{UHP}}, \end{aligned} \quad (3.28)$$

where the integration contour, after the deformation shown in Figure 5(a), has become the one $\mathcal{C}'_{(2n-1)}$ previously defined. In the $n \rightarrow \infty$ limit $e^{\frac{1}{2}\rho(\tilde{f}^{(2n-1)}(i))}$ and $e^{\frac{1}{2}\rho(\tilde{f}^{(2n-1)}(-i))}$




Figure 5: We obtain the contours $\bar{\mathcal{C}}_{(2n-1)}, \bar{\bar{\mathcal{C}}}_{(2n-1)}$ just after applying the gluing theorem. They can be deformed into $\mathcal{C}'_{(2n-1)}$.

come close to each other so that their product can be replaced by the leading term $e^\rho = c'$

in the OPE. This fact can be made more transparent by moving from the upper half plane to the unit disk representation by the conformal map $h(z)$. Performing the contour integration, we finally reach

$$\begin{aligned} \langle \varphi_- | \mathcal{A}_+^{(0)} * \mathcal{A}_-^{(1)} \rangle &= -\mathcal{N}_+^{(0)} K^{(3)} e^{2c'K} \kappa^{(\infty)} |f_1^{(3)'}(i)|^{1/4} |(h \circ \tilde{f}^{(\infty)})'(i)|^{-1/4} \\ &\times |h \circ \tilde{f}^{(\infty)}(i) - h \circ \tilde{f}^{(\infty)}(-i)|^{1/4} s(\infty) \langle \gamma(\infty) \tilde{f}^{(\infty)} \circ \varphi'_-(0) \rangle'_{\text{UHP}}. \end{aligned} \quad (3.29)$$

The last term in eq.(3.20) is found to be

$$\begin{aligned} -\langle \varphi_- | \mathcal{A}_-^{(1)} * \mathcal{A}_+^{(0)} \rangle &= -\mathcal{N}_+^{(0)} \sum_{r,s} \lim_{n \rightarrow \infty} \langle f^{(n)} \circ \Phi_r^{c'}(0) \rangle'_{\text{UHP}} \langle f^{(n)} \circ \Phi_s^{c'}(0) \rangle'_{\text{UHP}} \\ &\times \left\langle f_1^{(3)} \circ \varphi_-(0) f_2^{(3)} \circ \left(\mathcal{R}_{-\pi} \circ \oint_{\mathcal{C}} \frac{d\xi}{2\pi i} s(\xi) \xi b(\xi) \gamma(\xi) \right) f_2^{(3)} \circ \Phi_r(0) f_3^{(3)} \circ \Phi_s(0) \right\rangle_{\text{UHP}}. \end{aligned} \quad (3.30)$$

For $|\Phi_r\rangle \langle \Phi_r^c | \Xi' \rangle$ to be non-vanishing, Φ_r must share its properties with Ξ' . In particular, $\Phi_r(0)$ is Grassmann-odd. So when $f_2^{(3)} \circ \Phi_r(0)$ passes through $f_2^{(3)} \circ \mathcal{R}_{-\pi} \circ \oint_{\mathcal{C}} \frac{d\xi}{2\pi i} s(\xi) \xi b(\xi) \gamma(\xi)$ there arises a minus sign, which cancels the overall minus sign in front of the right hand side of (3.30). Comparing it with eq.(3.27), we find that $\langle \varphi_- | \mathcal{A}_+^{(0)} * \mathcal{A}_-^{(1)} \rangle$ and $-\langle \varphi_- | \mathcal{A}_-^{(1)} * \mathcal{A}_+^{(0)} \rangle$ give the same expression except for the integration contour. Calculating it further, we get the same expression as eq.(3.28), but with the contour replaced by $\bar{\mathcal{C}}_{(2n-1)}$ shown in Figure 5. This contour, however, can be deformed into $\mathcal{C}'_{(2n-1)}$ (Figure 5(b)) due to the holomorphicity, so $-\langle \varphi_- | \mathcal{A}_-^{(1)} * \mathcal{A}_+^{(0)} \rangle$ gives exactly the same result as $\langle \varphi_- | \mathcal{A}_+^{(0)} * \mathcal{A}_-^{(1)} \rangle$ (3.29). Substituting our results (3.25), (3.26), (3.29) into the equation of motion (3.20), we find that it is solved if we choose

$$s(\infty) \equiv s \left(z = \tilde{f}^{(\infty)}(i) = \infty \right) = -\frac{q_1}{2} \mathcal{N}_+^{(0)} \left([\tilde{f}^{(\infty)'}(i)]^{-1/2} + [\tilde{f}^{(\infty)'}(-i)]^{-1/2} \right), \quad (3.31)$$

where we have used the explicit form (3.16) of $\mathcal{N}_+^{(0)}$ to simplify the expression. Again, what is important is not the precise expression (3.31) for $s(\infty)$ but the fact that we could solve the equations of motion (3.19), (3.20) at order $(\varepsilon_r)^1$ by $\mathcal{A}_+^{(1)} = 0$ and $\mathcal{A}_-^{(1)}$ given in (3.23) with a suitable choice of a parameter $s(\infty)$.

Since the above $s(\infty)$ looks like a constant, one might suspect that this solution does not satisfy the regularity at infinity: $\lim_{z \rightarrow \infty} z s(z) < \infty$. However, $\tilde{f}^{(\infty)'}(z) = 1/(1+z^2)$ diverges when $z \rightarrow \pm i$ so that $s(\infty)$ is in fact vanishingly small. Regularizing the expression by $i \rightarrow i + \epsilon$ and taking the contribution from $\mathcal{N}_+^{(0)}$ into account, we find

$$\begin{aligned} \lim_{z \rightarrow \infty} z s(z) &\equiv \lim_{\epsilon \rightarrow 0} \tilde{f}^{(\infty)}(i + \epsilon) s(\tilde{f}^{(\infty)}(i + \epsilon)) \\ &\sim \lim_{\epsilon \rightarrow 0} \log \epsilon \left(\epsilon^{\frac{1}{12} - \frac{1}{4}} (\log \epsilon)^{\frac{1}{4}} \right) \epsilon^{\frac{1}{2}} \sim \lim_{\epsilon \rightarrow 0} \epsilon^{\frac{1}{3}} = 0, \end{aligned}$$

where we have used $\tilde{f}^{(\infty)}(i + \epsilon) = \tan^{-1}(i + \epsilon) \simeq \frac{1}{2i} \log \frac{i\epsilon}{2} \sim \log \epsilon$, $\tilde{f}^{(n)'}(i + \epsilon) \sim \epsilon^{\frac{2}{n}-1}$. Therefore the choice (3.31) of $s(\infty)$ does not violate the regularity condition. On the other hand, the equations of motion at order $(\epsilon_r)^2$ have not constrained the functional form of $s(z)$ except its value at the open string midpoint, as long as the chosen $s(z)$ satisfies the requirements mentioned below eq.(3.24). We cannot tell without solving the equations of motion at higher orders to what degree the form of s should be determined.

Our approximate solution $\hat{\mathcal{A}}' = \mathcal{A}_+^{(0)} \otimes \sigma_3 + \epsilon_r \mathcal{A}_-^{(1)} \otimes i\sigma_2$ given in (3.13), (3.23) has the following properties:

- At the lowest order $(\epsilon_r)^0$ the solution $\mathcal{A}_+^{(0)}$ takes the same form as the twisted sliver solution of bosonic vacuum string field theory, which is supposed to represent a spacetime-filling D25-brane;
- Both $\mathcal{A}_+^{(0)}$ and $\mathcal{A}_-^{(1)}$ have the matter-ghost split form and their matter part is simply the matter sliver [36, 3, 4, 37, 38].

These properties suggest that, if we can extend it to all orders in ϵ_r , the full solution should represent a spacetime-filling non-BPS D9-brane of type IIA superstring theory.¹⁶

4 Ghost Kinetic Operator and Solutions in Non-polynomial Vacuum Superstring Field Theory

In this section we consider Berkovits' non-polynomial superstring field theory around the tachyon vacuum. First of all, let us note that, if we expand the GSO-unprojected string field $\hat{\Phi}$ as $e^{\hat{\Phi}} = e^{\hat{\Phi}_0} * e^{\hat{\phi}}$ around a classical solution $\hat{\Phi}_0$ of the equation of motion $\hat{\eta}_0(e^{-\hat{\Phi}} \hat{Q}_B e^{\hat{\Phi}}) = 0$ derived from the Wess-Zumino-Witten-like action [39, 23]

$$S = \frac{1}{4g_o^2} \text{Tr} \left\langle \left\langle \left(e^{-\hat{\Phi}} \hat{Q}_B e^{\hat{\Phi}} \right) \left(e^{-\hat{\Phi}} \hat{\eta}_0 e^{\hat{\Phi}} \right) - \int_0^1 dt \left(e^{-t\hat{\Phi}} \partial_t e^{t\hat{\Phi}} \right) \left\{ \left(e^{-t\hat{\Phi}} \hat{Q}_B e^{t\hat{\Phi}} \right), \left(e^{-t\hat{\Phi}} \hat{\eta}_0 e^{t\hat{\Phi}} \right) \right\} \right\rangle \right\rangle, \quad (4.1)$$

then the action for the fluctuation field $\hat{\phi}$ takes the same form as the original one (4.1) with the BRST operator \hat{Q}_B replaced by another operator \hat{Q} given by

$$\hat{Q}\hat{X} = \hat{Q}_B\hat{X} + \hat{A}_0 * \hat{X} - (-1)^{\#\text{gh}(\hat{X})} \hat{X} * \hat{A}_0, \quad (4.2)$$

¹⁶We are assuming that we take the boundary conformal field theory associated with a non-BPS D9-brane of type IIA theory as the reference BCFT in whose state space the classical string field takes value.

where we have defined $\widehat{A}_0 = e^{-\widehat{\Phi}_0} \widehat{Q}_B e^{\widehat{\Phi}_0}$. This is a trivial extension of the results shown in [40, 38] for the GSO-projected theory. Since the above formula (4.2) for \widehat{Q} is very similar to the one (2.11) for the cubic case and both cubic and Berkovits' superstring field theories reproduce qualitatively the same tachyon potential of the double-well form [22, 23], the logic we presented in section 2 should be valid in this non-polynomial case as well: We need in \widehat{Q} the Grassmann-even sector $Q_{\text{even}} \otimes i\sigma_2$ as well as the Grassmann-odd piece $Q_{\text{odd}} \otimes \sigma_3$ to violate the \mathbb{Z}_2 reflection symmetry, and through a singular reparametrization \widehat{Q} should be dominated by the midpoint insertions of the lowest-dimensional local operators. Moreover, Berkovits' superstring field theory action (4.1) including both the GSO(\pm) sectors was shown to be invariant under the twist operation [23]

$$\Omega|\Phi\rangle = \begin{cases} (-1)^{h_\Phi+1}|\Phi\rangle & \text{for GSO(+)} \text{ states } (h_\Phi \in \mathbb{Z}) \\ (-1)^{h_\Phi+\frac{1}{2}}|\Phi\rangle & \text{for GSO(-)} \text{ states } (h_\Phi \in \mathbb{Z} + \frac{1}{2}). \end{cases}, \quad (4.3)$$

which is the same as the action of the twist operator (2.32) in the 0-picture cubic superstring field theory. Hence, if we require the kinetic operator \widehat{Q} around the tachyon vacuum to commute with the twist operator Ω , we will obtain the same \widehat{Q} as in the cubic case and the resulting action will become twist-invariant under (4.3). From these considerations, we propose that the non-polynomial vacuum superstring field theory action for the NS sector is given by

$$S = \frac{\kappa_0}{4} \text{Tr} \left\langle\left\langle \left(e^{-\widehat{\Phi}} \widehat{Q} e^{\widehat{\Phi}} \right) \left(e^{-\widehat{\Phi}} \widehat{\eta}_0 e^{\widehat{\Phi}} \right) - \int_0^1 dt \left(e^{-t\widehat{\Phi}} \partial_t e^{t\widehat{\Phi}} \right) \left\{ \left(e^{-t\widehat{\Phi}} \widehat{Q} e^{t\widehat{\Phi}} \right), \left(e^{-t\widehat{\Phi}} \widehat{\eta}_0 e^{t\widehat{\Phi}} \right) \right\} \right\rangle\right\rangle, \quad (4.4)$$

with

$$\begin{aligned} \widehat{Q} &= Q_{\text{odd}} \otimes \sigma_3 - Q_{\text{even}} \otimes i\sigma_2, \\ Q_{\text{odd}} &= \frac{1}{2i\varepsilon_r^2} (c(i) - c(-i)) + \frac{q_1^2}{2} \oint \frac{dz}{2\pi i} b\gamma^2(z), \\ Q_{\text{even}}^{\text{GSO}(+)} &= \frac{q_1}{2i\varepsilon_r} (\gamma(i) - \gamma(-i)), \quad Q_{\text{even}}^{\text{GSO}(-)} = \frac{q_1}{2\varepsilon_r} (\gamma(i) + \gamma(-i)). \end{aligned} \quad (4.5)$$

This \widehat{Q} is a nilpotent derivation of the $*$ -algebra, and can be regularized in such a way that \widehat{Q} annihilates the identity string field \mathcal{I} , as has been proven in subsection 2.2. In addition, \widehat{Q} anticommutes with $\widehat{\eta}_0 = \eta_0 \otimes \sigma_3$ because \widehat{Q} contains no factor of ξ_0 and $\widehat{\eta}_0$ does not change the GSO parity of the states. These properties guarantee that the action (4.4) is invariant under the infinitesimal gauge transformation

$$\delta(e^{\widehat{\Phi}}) = (\widehat{Q}\widehat{\Omega}) * e^{\widehat{\Phi}} + e^{\widehat{\Phi}} * (\widehat{\eta}_0 \widehat{\Lambda}). \quad (4.6)$$

where the gauge parameters $\widehat{\Omega}, \widehat{\Lambda}$ are of ghost number -1 . Actually we need not repeat the proof of gauge invariance here because the set $(\widehat{\Phi}, \widehat{\mathcal{Q}}, \widehat{\eta}_0)$ of the GSO-unprojected string field $\widehat{\Phi} = \Phi_+ \otimes \mathbf{1} + \Phi_- \otimes \sigma_1$ and the operators $\widehat{\mathcal{Q}}, \widehat{\eta}_0$ equipped with the internal Chan-Paton matrices satisfies exactly the same algebraic relations as that (Φ, Q_B, η_0) in the GSO-projected theory, so that the proof goes in the identical way to the one given in the GSO-projected theory (see [23]), as can be seen from the processes of the proof presented in subsection 2.3.

What we want to do next is to find classical solutions¹⁷ corresponding to D-branes. For this purpose, we will follow the strategy proposed in [34]. Let us consider the specific combination $\widehat{A}(\widehat{\Phi}) \equiv e^{-\widehat{\Phi}} \widehat{\mathcal{Q}} e^{\widehat{\Phi}}$ of the string field $\widehat{\Phi}$ of this theory. Since $\widehat{\Phi}$ has vanishing ghost and picture numbers, we find that

$$\widehat{A} \text{ has ghost number 1 and picture number 0.} \quad (4.7)$$

Written in terms of $\widehat{A}(\widehat{\Phi})$, the equation of motion following from the action (4.4) is

$$\widehat{\eta}_0(\widehat{A}) = 0. \quad (4.8)$$

In addition, $\widehat{A}(\widehat{\Phi})$ satisfies by definition

$$\widehat{\mathcal{Q}}\widehat{A} + \widehat{A} * \widehat{A} = 0, \quad (4.9)$$

because $\widehat{\mathcal{Q}}$ is a nilpotent derivation and annihilates $\mathcal{I} \otimes \mathbf{1} = e^{-\widehat{\Phi}} * e^{\widehat{\Phi}}$. On the other hand, in 0-picture cubic superstring field theory the *string field* $\widehat{\mathcal{A}}$ was defined to have

$$\text{ghost number 1 and picture number 0.} \quad (4.10)$$

Since this cubic theory was formulated within the “small” Hilbert space, $\widehat{\mathcal{A}}$ must not contain the zero mode of ξ . In other words, we have to impose

$$\widehat{\eta}_0(\widehat{\mathcal{A}}) = 0. \quad (4.11)$$

As given in (3.1), the equation of motion of the cubic theory was

$$\widehat{\mathcal{Q}}\widehat{\mathcal{A}} + \widehat{\mathcal{A}} * \widehat{\mathcal{A}} = 0. \quad (4.12)$$

Comparing eqs.(4.7)–(4.9) with eqs.(4.10)–(4.12), we find that the sets of equations we should solve in looking for classical solutions in these two theories coincide with each

¹⁷Recently, a general method of constructing exact solutions of the equation of motion of Berkovits’ string field theory was given in [41].

other.¹⁸ This fact can be used to find a solution $\widehat{\Phi}_0$ in the non-polynomial theory which ‘corresponds’ to a solution $\widehat{\mathcal{A}}_0$ in the cubic theory. Suppose that we are given a solution $\widehat{\mathcal{A}}_0$ in the cubic theory satisfying (4.10)–(4.12). From (4.10) and (4.12), it is *possible* that there exists a string field configuration $\widehat{\Phi}_0$ which satisfies

$$e^{-\widehat{\Phi}_0} \widehat{\mathcal{Q}} e^{\widehat{\Phi}_0} = \widehat{\mathcal{A}}_0. \quad (4.13)$$

If we can find such a $\widehat{\Phi}_0$, it then gives a solution in the non-polynomial theory because we have

$$\widehat{\eta}_0(e^{-\widehat{\Phi}_0} \widehat{\mathcal{Q}} e^{\widehat{\Phi}_0}) = \widehat{\eta}_0(\widehat{\mathcal{A}}_0) = 0,$$

where (4.11) was used. Furthermore, we expect that the solution $\widehat{\Phi}_0$ in the non-polynomial theory has the same physical interpretation as the corresponding one $\widehat{\mathcal{A}}_0$ in the cubic theory. In particular, the forms of the new kinetic operators around these solutions are, respectively,

$$\widehat{\mathcal{Q}}'_{\text{cubic}} \widehat{X} = \widehat{\mathcal{Q}} \widehat{X} + \widehat{\mathcal{A}}_0 * \widehat{X} - (-1)^{\#_{\text{gh}}(\widehat{X})} \widehat{X} * \widehat{\mathcal{A}}_0 \quad \text{in cubic theory}$$

and

$$\widehat{\mathcal{Q}}'_{\text{non-poly}} \widehat{X} = \widehat{\mathcal{Q}} \widehat{X} + \widehat{A}(\widehat{\Phi}_0) * \widehat{X} - (-1)^{\#_{\text{gh}}(\widehat{X})} \widehat{X} * \widehat{A}(\widehat{\Phi}_0) \quad \text{in non-polynomial theory.}$$

Since we have determined $\widehat{\Phi}_0$ such that $\widehat{\mathcal{A}}_0 = \widehat{A}(\widehat{\Phi}_0)$ is satisfied, these kinetic operators $\widehat{\mathcal{Q}}'_{\text{cubic}}, \widehat{\mathcal{Q}}'_{\text{non-poly}}$, each of which describes the perturbative physics around the corresponding solution, should agree with each other. This result strongly supports the claim that the solutions $\widehat{\mathcal{A}}_0$ and $\widehat{\Phi}_0$ of the two theories related through (4.13) share a common physical interpretation. Therefore, in order to find D-brane solutions $\widehat{\Phi}_0$ in non-polynomial vacuum superstring field theory, we have to

- (1) find solutions $\widehat{\mathcal{A}}_0$ in 0-picture cubic vacuum superstring field theory which can be interpreted as D-branes, and then
- (2) solve the equation (4.13) with respect to $\widehat{\Phi}_0$ for $\widehat{\mathcal{A}}_0$ just found above.

However, we have not yet found any explicit algorithm to solve the equation (4.13) for a given $\widehat{\mathcal{A}}_0$. This problem is left to future study.

¹⁸Since the picture changing operations have been completely eliminated from the theory in Berkovits’ formulation, the equation (4.9) should be considered by taking the inner product with a state of ghost number 0, picture number -1 *including* ξ_0 , with no picture changing operators inserted. Since we considered eq.(3.4) in section 3 instead of $\langle\langle \widehat{Y}_{-2} | \widehat{\varphi}, \mathcal{F}(\widehat{\mathcal{A}}) \rangle\rangle = 0$, the solution found there can be used here.

Finally we remark that, even if $\widehat{\mathcal{A}}_0$ can be written in the form $e^{-\widehat{\Phi}_0}\widehat{\mathcal{Q}}e^{\widehat{\Phi}_0}$, it does not immediately follow that this $\widehat{\mathcal{A}}_0$ is pure-gauge in the cubic theory. This is because in Berkovits' formulation we can seek an appropriate configuration $\widehat{\Phi}_0$ in the “large” Hilbert space including ξ_0 , whereas to make an assertion that $\widehat{\mathcal{A}}_0$ is pure-gauge in the cubic theory we must find a suitable gauge parameter $\widehat{\Lambda}$ satisfying $\widehat{\mathcal{A}}_0 = e^{-\widehat{\Lambda}}\widehat{\mathcal{Q}}e^{\widehat{\Lambda}}$ within the “small” Hilbert space. Conversely, let us suppose that we have a pure-gauge configuration $\widehat{\mathcal{A}}_0$ in the cubic theory. Then there exists a gauge parameter $\widehat{\Lambda}$ which has ghost number 0 and satisfies $\widehat{\mathcal{A}}_0 = e^{-\widehat{\Lambda}}\widehat{\mathcal{Q}}e^{\widehat{\Lambda}}$ and $\widehat{\eta}_0\widehat{\Lambda} = 0$. If we regard this $\widehat{\Lambda}$ as a string field in Berkovits' non-polynomial theory, such a configuration (*i.e.* annihilated by $\widehat{\eta}_0$) turns out to be pure-gauge, as shown in [34]. So the story is quite consistent in the sense that a pure-gauge configuration in one theory is mapped under (4.13) to some pure-gauge configuration in the other theory.¹⁹

5 Summary and Discussion

In this paper we have discussed the construction of vacuum superstring field theory itself and the brane solutions in it. We have first argued what form the kinetic operator $\widehat{\mathcal{Q}}$ around the tachyon vacuum should have in general, and seen that we need a Grassmann-even operator $Q_{\text{even}} \otimes i\sigma_2$ to have a structure expected of the vacuum superstring field theory action. We have then determined the form of the simple pure-ghost kinetic operator $\widehat{\mathcal{Q}}$ as unambiguously as possible by requiring that (i) the most singular parts of $\widehat{\mathcal{Q}}$ be made up of midpoint insertions of the lowest-dimensional operators, following [15], (ii) $\widehat{\mathcal{Q}}$ preserve the twist invariance of the string field theory action, and (iii) the action be gauge invariant. We have obtained a somewhat surprising result that the form of Q_{even} depends on the GSO parity of the state on which Q_{even} acts, but such a structure is indispensable for consistency. We have examined in detail the properties of $\widehat{\mathcal{Q}}$ determined this way, and explicitly shown that it satisfies all of the axioms which guarantee the gauge invariance of the action. We have observed from the known properties of the twist operation that $\widehat{\mathcal{Q}}$ should take the same form in both cubic and non-polynomial types of vacuum superstring field theory.

We have also tried to solve the equations of motion of cubic vacuum superstring field theory, but obtained only an approximate solution in an ε_r -expansion method. Finally

¹⁹More generally, we can show that two gauge-equivalent string field configurations $e^{\widehat{\Phi}_0}, h_1^{\widehat{\mathcal{Q}}}e^{\widehat{\Phi}_0}h_2^{\widehat{\eta}_0}$ in non-polynomial theory are mapped to two configurations $\widehat{\mathcal{A}}_0, (h_2^{\widehat{\eta}_0})^{-1}(\widehat{\mathcal{Q}} + \widehat{\mathcal{A}}_0)h_2^{\widehat{\eta}_0}$ which are also related through a formally valid gauge transformation in the cubic theory, where $h_1^{\widehat{\mathcal{Q}}}, h_2^{\widehat{\eta}_0}$ are gauge parameters annihilated by $\widehat{\mathcal{Q}}, \widehat{\eta}_0$, respectively.

we have suggested a way of generating the solution in the non-polynomial theory which physically corresponds to a given solution in the cubic theory, with its precise algorithm left to be clarified.

To completely understand the nature of superstring field theory around the tachyon vacuum, many things still remain to be done. At the level of the determination of the kinetic operator \widehat{Q} , aside from a proportionality constant we have not been able to uniquely fix the form of the non-leading part of Q_{odd} , which has been added *by hand* in order to make \widehat{Q} be nilpotent. It might be that, after successfully including the Ramond sector field in vacuum superstring field theory, the requirement of spacetime supersymmetry restored around the ‘type II closed string vacuum’ puts further constraints on the form of \widehat{Q} .

At the level of the construction of classical solutions, we have not obtained exact expressions even for the solution representing the most fundamental non-BPS D9-brane. Moreover, the expressions for our approximate solution are plagued with many vanishing or diverging factors, as is often the case with the singular representative of vacuum string field theory. In the bosonic case, such a ‘formal’ argument was partly justified by showing [15, 16] that the twisted sliver solution coincides with the Siegel gauge solution found in [10] in a less singular algebraic approach. So it is desirable that we could find a solution algebraically in this superstring case as well and compare it with ours. This might also give us insight into the construction of another tachyon vacuum or BPS D-brane solutions.

Acknowledgements

I am grateful to I. Kishimoto for many valuable discussions and comments. I’d like to thank T. Eguchi and Y. Matsuo for discussions, and A.A. Giryavets for E-mail correspondence. I also thank the members of Hokkaido University and the organizers of YITP workshop YITP-W-02-04 on “Quantum Field Theory 2002”, where I was given opportunities to have discussions on this research. This work is supported by JSPS Research Fellowships for Young Scientists.

Appendices

A Inner Derivation Formula for $\mathcal{Q}_{\text{even}}$

In this Appendix we show that the expression (2.63) reproduces eq.(2.64). Taking the inner product of (2.63) with a Fock space state $\langle\varphi|$ gives

$$\begin{aligned}
\langle\varphi|\mathcal{Q}_{\text{even}}|\psi\rangle &= \langle\varphi|(\Gamma_\epsilon\mathcal{I}) * \psi\rangle - (-1)^{\text{GSO}(\psi)}\langle\varphi|\psi * (\Gamma_\epsilon\mathcal{I})\rangle \tag{A.1} \\
&= \sum_r \langle\varphi|(\Gamma_\epsilon\Phi_r) * \psi\rangle\langle\Phi_r^c|\mathcal{I}\rangle - (-1)^{\text{GSO}(\psi)}\sum_r \langle\varphi|\psi * (\Gamma_\epsilon\Phi_r)\rangle\langle\Phi_r^c|\mathcal{I}\rangle \\
&= \sum_r (-1)^{\text{GSO}(\psi)} \left\langle f_1^{(3)} \circ \psi(0) f_2^{(3)} \circ \varphi(0) f_3^{(3)} \circ \Gamma_\epsilon f_3^{(3)} \circ \Phi_r(0) \right\rangle \langle f_1^{(1)} \circ \Phi_r^c(0) \rangle \\
&\quad - (-1)^{\text{GSO}(\psi)} \sum_r \left\langle f_1^{(3)} \circ \varphi(0) f_2^{(3)} \circ \psi(0) f_3^{(3)} \circ \Gamma_\epsilon f_3^{(3)} \circ \Phi_r(0) \right\rangle \langle f_1^{(1)} \circ \Phi_r^c(0) \rangle \\
&= (-1)^{\text{GSO}(\psi)} \left\langle F_1 \circ f_1^{(3)} \circ \psi(0) F_1 \circ f_2^{(3)} \circ \varphi(0) F_1 \circ f_3^{(3)} \circ \Gamma_\epsilon \right\rangle \\
&\quad - (-1)^{\text{GSO}(\psi)} \left\langle F_1 \circ f_1^{(3)} \circ \varphi(0) F_1 \circ f_2^{(3)} \circ \psi(0) F_1 \circ f_3^{(3)} \circ \Gamma_\epsilon \right\rangle,
\end{aligned}$$

where we have inserted the complete set of states, $\mathbf{1} = \sum_r |\Phi_r\rangle\langle\Phi_r^c|$, and we are omitting the symbol $\lim_{\epsilon\rightarrow 0}$. From the GGRT formulas given in [32] F 's are found to be

$$\begin{aligned}
F_1(z) &= h^{-1} \left(e^{\frac{\pi}{2}i} h(z)^{\frac{3}{2}} \right), \\
\widehat{F}_2(z) &= h^{-1} \left(e^{\frac{3\pi}{2}i} h(z)^{\frac{1}{2}} \right).
\end{aligned}$$

Calculating the compositions of the conformal maps carefully and using the $SL(2, \mathbb{R})$ -invariance of the correlators, we obtain

$$\begin{aligned}
\langle\varphi|\mathcal{Q}_{\text{even}}|\psi\rangle &= (-1)^{\text{GSO}(\psi)} \langle\varphi(0)\mathcal{R}_\pi \circ \Gamma_\epsilon \mathcal{R}_{-\pi} \circ \psi(0)\rangle \\
&\quad - (-1)^{\text{GSO}(\psi)} \langle\varphi(0)\mathcal{R}_{2\pi} \circ \Gamma_\epsilon \mathcal{R}_\pi \circ \psi(0)\rangle \\
&= \langle\varphi| (\Gamma_\epsilon - (-1)^{\text{GSO}(\psi)}\mathcal{R}_\pi \circ \Gamma_\epsilon) |\psi\rangle. \tag{A.2}
\end{aligned}$$

Now we can take the $\epsilon \rightarrow 0$ limit without encountering any singularities. From the definition (2.46), (2.47) of the action of \mathcal{R}_π we find

$$\begin{aligned}
\lim_{\epsilon\rightarrow 0} (\Gamma_\epsilon - (-1)^{\text{GSO}(\psi)}\mathcal{R}_\pi \circ \Gamma_\epsilon) &= q_1 \frac{1-i}{4\varepsilon_r} \left\{ \gamma(i) + i\gamma(-i) \right. \\
&\quad \left. - (-1)^{\text{GSO}(\psi)} (i\gamma(\mathcal{R}_\pi(i)) + i(-i)\gamma(\mathcal{R}_\pi(-i))) \right\} \\
&= q_1 \frac{1-i}{4\varepsilon_r} (1 - i(-1)^{\text{GSO}(\psi)}) (\gamma(i) - (-1)^{\text{GSO}(\psi)}\gamma(-i)), \tag{A.3}
\end{aligned}$$

which completes the proof of (2.64).

B Twist Invariance of the Cubic Action

Here we explicitly show that the cubic action (2.83) with our choice of $\widehat{\mathcal{Q}}$ is invariant under the twist transformation Ω (2.32). Since the cubic interaction term is not changed from the original one (2.1) at all, its twist invariance is shown in the same way as in [22]. Hence we will concentrate on the quadratic term. For \mathcal{Q}_{odd} we consider

$$\begin{aligned} \langle\langle Y_{-2} | \mathcal{O}_1, \mathcal{Q}_{\text{odd}} \mathcal{O}_2 \rangle\rangle &= \langle Y(i)Y(-i) (f_1^{(2)} \circ \mathcal{O}_1(0))(f_2^{(2)} \circ \mathcal{Q}_{\text{odd}})(f_2^{(2)} \circ \mathcal{O}_2(0)) \rangle_{\text{UHP}} \\ &= \langle Y(0)Y(\infty) (g_1^{(2)} \circ \mathcal{O}_1(0))(g_2^{(2)} \circ \mathcal{Q}_{\text{odd}})(g_2^{(2)} \circ \mathcal{O}_2(0)) \rangle_{\text{disk}}, \end{aligned} \quad (\text{B.1})$$

where $\langle \dots \rangle_{\text{disk}}$ denotes the correlation function on the unit disk, and

$$g_1^{(2)}(z) = h \circ f_1^{(2)}(z) = h(z), \quad g_2^{(2)}(z) = h \circ f_2^{(2)}(z) = e^{\pi i} h(z).$$

Note that for the expression (B.1) to be non-vanishing \mathcal{O}_1 and \mathcal{O}_2 must live in the same GSO sector. Defining $M(z) \equiv e^{i\pi} z$, we find the following relations for any primary vertex operator $\varphi(z)$:

$$\begin{aligned} g_1^{(2)} \circ M \circ \varphi(z) &= \tilde{I} \circ \mathcal{R}_{2\pi} \circ g_1^{(2)} \circ \varphi(z), \\ g_2^{(2)} \circ M \circ \varphi(z) &= \tilde{I} \circ g_2^{(2)} \circ \varphi(z), \end{aligned} \quad (\text{B.2})$$

where $\tilde{I}(z) \equiv 1/z$, $[\tilde{I}'(z)]^h \equiv e^{-i\pi h} z^{-2h}$, and $\mathcal{R}_{2\pi}(z) = e^{2\pi i} z$. If we assume that $\mathcal{O}_1, \mathcal{O}_2$ are primary fields of conformal weight h_1, h_2 respectively, we have

$$\begin{aligned} \mathcal{O}_1(0) &= e^{-i\pi h_1} M \circ \mathcal{O}_1(0), \\ \mathcal{O}_2(0) &= e^{-i\pi h_2} M \circ \mathcal{O}_2(0), \end{aligned} \quad (\text{B.3})$$

$$\begin{aligned} M \circ \mathcal{Q}_{\text{odd}} &= \frac{1}{2i\varepsilon_r^2} (M \circ c(i) - M \circ c(-i)) + \frac{q_1^2}{2} \oint \frac{dz}{2\pi i} M \circ (b\gamma^2(z)) \\ &= \frac{1}{2i\varepsilon_r^2} (e^{-i\pi} c(i e^{i\pi}) - e^{-i\pi} c(-i e^{i\pi})) + \frac{q_1^2}{2} \oint \frac{dz}{2\pi i} e^{\pi i} b\gamma^2(e^{\pi i} z) \\ &= \frac{1}{2i\varepsilon_r^2} (c(i) - c(-i)) + \frac{q_1^2}{2} \oint \frac{dz'}{2\pi i} b\gamma^2(z') = \mathcal{Q}_{\text{odd}}. \end{aligned} \quad (\text{B.4})$$

Then (B.1) can be rewritten as

$$\begin{aligned} \langle\langle Y_{-2} | \mathcal{O}_1, \mathcal{Q}_{\text{odd}} \mathcal{O}_2 \rangle\rangle &= e^{-i\pi(h_1+h_2)} \langle Y(0)Y(\infty) (g_1^{(2)} \circ M \circ \mathcal{O}_1(0))(g_2^{(2)} \circ M \circ \mathcal{Q}_{\text{odd}})(g_2^{(2)} \circ M \circ \mathcal{O}_2(0)) \rangle_{\text{disk}} \\ &= e^{-i\pi(h_1+h_2)} \langle \tilde{I} \circ \left(Y(0)Y(\infty) (\mathcal{R}_{2\pi} \circ g_1^{(2)} \circ \mathcal{O}_1(0))(g_2^{(2)} \circ \mathcal{Q}_{\text{odd}})(g_2^{(2)} \circ \mathcal{O}_2(0)) \right) \rangle_{\text{disk}} \\ &= (-1)^{h_1+h_2} (-1)^{2h_1} \langle Y(0)Y(\infty) g_1^{(2)} \circ \mathcal{O}_1(0) g_2^{(2)} \circ (\mathcal{Q}_{\text{odd}} \mathcal{O}_2(0)) \rangle_{\text{disk}} \\ &= (-1)^{h_1+h_2} (-1)^{2h_1} \langle\langle Y_{-2} | \mathcal{O}_1, \mathcal{Q}_{\text{odd}} \mathcal{O}_2 \rangle\rangle, \end{aligned} \quad (\text{B.5})$$

where we have used (B.2)–(B.4), the invariance of $Y(0)Y(\infty)$ under the conformal transformation \tilde{I} , the invariance of the disk correlator under the $SL(2, \mathbb{R})$ map $\tilde{I}(z)$, and $\mathcal{R}_{2\pi} \circ \mathcal{O}_1(0) = e^{2\pi i h_1} \mathcal{O}_1(0)$. $(-1)^{h_1+h_2}$ and $(-1)^{2h_1}$ are well-defined because $h_1 + h_2$ and $2h_1$ are always integers. Since eq.(B.5) says that $\langle\langle Y_{-2} | \mathcal{O}_1, \mathcal{Q}_{\text{odd}} \mathcal{O}_2 \rangle\rangle$ is equal to itself multiplied by $(-1)^{h_1+h_2}(-1)^{2h_1}$, this expression vanishes unless $(-1)^{h_1+h_2}(-1)^{2h_1} = 1$ holds. This condition is satisfied only when both \mathcal{O}_1 and \mathcal{O}_2 are twist-odd or twist-even. For example, consider a case $h_1 \in 2\mathbb{Z} + \frac{1}{2}$ (i.e. \mathcal{O}_1 is *twist-odd* and in the $\text{GSO}(-)$ sector). Since $(-1)^{2h_1} = -1$, h_2 must take value in $2\mathbb{Z} + \frac{1}{2}$ so that $(-1)^{h_1+h_2}$ will equal -1 . In other words, \mathcal{O}_2 must also be *twist-odd* and in the $\text{GSO}(-)$ sector. One can repeat similar arguments for other three cases to verify the above statement. To sum up, we have shown that twist-odd states enter $\langle\langle Y_{-2} | \mathcal{O}_1, \mathcal{Q}_{\text{odd}} \mathcal{O}_2 \rangle\rangle$ always *in pairs* (otherwise the expression vanishes), so that it is invariant under the twist operation Ω .

Now we turn to

$$\langle\langle Y_{-2} | \mathcal{O}_1^{(+)}, \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} \mathcal{O}_2^{(-)} \rangle\rangle = \langle Y(0)Y(\infty) g_1^{(2)} \circ \mathcal{O}_1^{(+)}(0) g_2^{(2)} \circ (\mathcal{Q}_{\text{even}}^{\text{GSO}(-)} \mathcal{O}_2^{(-)}(0)) \rangle_{\text{disk}}, \quad (\text{B.6})$$

where the superscripts (\pm) for $\mathcal{O}_1^{(\pm)}, \mathcal{O}_2^{(\pm)}$ denote their GSO parities. In a similar way to the above argument, we rewrite (B.6) as

$$\begin{aligned} \langle\langle Y_{-2} | \mathcal{O}_1^{(+)}, \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} \mathcal{O}_2^{(-)} \rangle\rangle &= e^{-i\pi(h_1+h_2-\frac{1}{2})} \langle Y(0)Y(\infty) (g_1^{(2)} \circ M \circ \mathcal{O}_1^{(+)}(0)) \\ &\quad \times (g_2^{(2)} \circ M \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(-)})(g_2^{(2)} \circ M \circ \mathcal{O}_2^{(-)}(0)) \rangle_{\text{disk}} \\ &= (-1)^{h_1+h_2-\frac{1}{2}} (-1)^{2h_1} \langle\langle Y_{-2} | \mathcal{O}_1^{(+)}, \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} \mathcal{O}_2^{(-)} \rangle\rangle, \end{aligned} \quad (\text{B.7})$$

where we have used

$$M \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} = \frac{q_1}{2\varepsilon_r} (e^{i\pi})^{-\frac{1}{2}} (\gamma(i e^{i\pi}) + \gamma(-i e^{i\pi})) = e^{-\frac{\pi i}{2}} \mathcal{Q}_{\text{even}}^{\text{GSO}(-)}.$$

Since $\mathcal{O}_1^{(+)}$ lives in the $\text{GSO}(+)$ sector, $(-1)^{2h_1} = 1$. If $\mathcal{O}_1^{(+)}$ is twist-odd, i.e. $h_1 \in 2\mathbb{Z}$, then $(-1)^{h_1+h_2-\frac{1}{2}} = 1$ requires $h_2 \in 2\mathbb{Z} + \frac{1}{2}$ which means that the $\text{GSO}(-)$ state $\mathcal{O}_2^{(-)}$ should be twist-odd. Conversely, if $\mathcal{O}_1^{(+)}$ is twist-even, $\mathcal{O}_2^{(-)}$ must be twist-even for $(-1)^{h_1+h_2-\frac{1}{2}} = 1$ to hold. We again conclude that twist-odd states appear in non-zero $\langle\langle Y_{-2} | \mathcal{O}_1^{(+)}, \mathcal{Q}_{\text{even}}^{\text{GSO}(-)} \mathcal{O}_2^{(-)} \rangle\rangle$ *pairwise* so that this expression is twist invariant. Finally we consider

$$\begin{aligned} \langle\langle Y_{-2} | \mathcal{O}_1^{(-)}, \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} \mathcal{O}_2^{(+)} \rangle\rangle &= \langle Y(0)Y(\infty) g_1^{(2)} \circ \mathcal{O}_1^{(-)}(0) g_2^{(2)} \circ (\mathcal{Q}_{\text{even}}^{\text{GSO}(+)} \mathcal{O}_2^{(+)}(0)) \rangle_{\text{disk}} \\ &= e^{-i\pi(h_1+h_2+\frac{1}{2})} \langle Y(0)Y(\infty) (g_1^{(2)} \circ M \circ \mathcal{O}_1^{(-)}(0)) \\ &\quad \times (g_2^{(2)} \circ M \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(+)})(g_2^{(2)} \circ M \circ \mathcal{O}_2^{(+)}(0)) \rangle_{\text{disk}} \\ &= (-1)^{h_1+h_2+\frac{1}{2}} (-1)^{2h_1} \langle\langle Y_{-2} | \mathcal{O}_1^{(-)}, \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} \mathcal{O}_2^{(+)} \rangle\rangle, \end{aligned} \quad (\text{B.8})$$

where we have used

$$M \circ \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} = \frac{q_1}{2i\epsilon_r} (e^{i\pi})^{-\frac{1}{2}} (\gamma(-i) - \gamma(i)) = -e^{-\frac{i\pi}{2}} \mathcal{Q}_{\text{even}}^{\text{GSO}(+)} \equiv e^{\frac{i\pi}{2}} \mathcal{Q}_{\text{even}}^{\text{GSO}(+)}.$$

The argument goes along the same line: If $h_1 \in 2\mathbb{Z} + \frac{1}{2}$ ($\mathcal{O}_1^{(-)}$ is twist-odd) then we must have $h_2 \in 2\mathbb{Z}$ ($\mathcal{O}_2^{(+)}$ is twist-odd) because of $(-1)^{2h_1} = -1$. Putting the above considerations together, we have completed the proof that the quadratic part $\text{Tr}\langle\langle \widehat{Y}_{-2} | \widehat{\mathcal{A}}, \widehat{\mathcal{Q}} \widehat{\mathcal{A}} \rangle\rangle$ of the action with our $\widehat{\mathcal{Q}}$ is twist invariant.

References

- [1] L. Rastelli, A. Sen and B. Zwiebach, “String Field Theory Around the Tachyon Vacuum,” *Adv. Theor. Math. Phys.* **5** (2002) 353-392 [hep-th/0012251].
- [2] L. Rastelli, A. Sen and B. Zwiebach, “Vacuum String Field Theory,” hep-th/0106010.
- [3] L. Rastelli, A. Sen and B. Zwiebach, “Classical Solutions in String Field Theory Around the Tachyon Vacuum,” *Adv. Theor. Math. Phys.* **5** (2002) 393-428 [hep-th/0102112].
- [4] L. Rastelli, A. Sen and B. Zwiebach, “Boundary CFT Construction of D-branes in Vacuum String Field Theory,” *JHEP* **0111** (2001) 045 [hep-th/0105168].
- [5] P. Mukhopadhyay, “Oscillator Representation of the BCFT Construction of D-branes in Vacuum String Field Theory,” *JHEP* **0112** (2001) 025 [hep-th/0110136].
- [6] K. Okuyama, “Ratio of Tensions from Vacuum String Field Theory,” *JHEP* **0203** (2002) 050 [hep-th/0201136].
- [7] P. Matlock, R.C. Rashkov, K.S. Viswanathan and Y. Yang, “Dp-brane Tension from Tachyons and B-field in Vacuum String Field Theory,” *Phys. Rev.* **D66** (2002) 026004 [hep-th/0203071].
- [8] L. Rastelli, A. Sen and B. Zwiebach, “Half-strings, Projectors, and Multiple D-branes in Vacuum String Field Theory,” *JHEP* **0111** (2001) 035 [hep-th/0105058].
- [9] Y. Okawa, “Open string states and D-brane tension from vacuum string field theory,” *JHEP* **0207** (2002) 003 [hep-th/0204012].

- [10] H. Hata and T. Kawano, “Open String States around a Classical Solution in Vacuum String Field Theory,” *JHEP* **0111** (2001) 038 [hep-th/0108150].
- [11] H. Hata and S. Moriyama, “Observables as Twist Anomaly in Vacuum String Field Theory,” *JHEP* **0201** (2002) 042 [hep-th/0111034];
H. Hata, S. Moriyama and S. Teraguchi, “Exact Results on Twist Anomaly,” *JHEP* **0202** (2002) 036 [hep-th/0201177];
H. Hata and S. Moriyama, “Reexamining Classical Solution and Tachyon Mode in Vacuum String Field Theory,” hep-th/0206208.
- [12] L. Rastelli, A. Sen and B. Zwiebach, “A Note on a Proposal for the Tachyon State in Vacuum String Field Theory,” *JHEP* **0202** (2002) 034 [hep-th/0111153].
- [13] R. Rashkov and K. S. Viswanathan, “A Note on the Tachyon State in Vacuum String Field Theory,” hep-th/0112202.
- [14] R. Rashkov and K.S. Viswanathan, “A Proposal for the Vector State in Vacuum String Field Theory,” hep-th/0201229.
- [15] D. Gaiotto, L. Rastelli, A. Sen and B. Zwiebach, “Ghost Structure and Closed Strings in Vacuum String Field Theory,” hep-th/0111129.
- [16] T. Okuda, “The Equality of Solutions in Vacuum String Field Theory,” hep-th/0201149.
- [17] R. Potting and J. Raeymaekers, “Ghost Sector of Vacuum String Field Theory and the Projection Equation,” *JHEP* **0206** (2002) 002 [hep-th/0204172].
- [18] T. Yoneya, “Spontaneously Broken Space-Time Supersymmetry in Open String Theory without GSO Projection,” *Nucl. Phys.* **B576** (2000) 219-240 [hep-th/9912255].
- [19] E. Witten, “Interacting Field Theory of Open Superstrings,” *Nucl. Phys.* **B276**(1986)291.
- [20] I.Y. Aref’eva, D.M. Belov and A.A. Giryavets, “Construction of the Vacuum String Field Theory on a non-BPS Brane,” *JHEP* **0209** (2002) 050 [hep-th/0201197].
- [21] I.Y. Aref’eva, A.A. Giryavets and A.S. Koshelev, “NS Ghost Slivers,” *Phys. Lett.* **B536** (2002) 138-146 [hep-th/0203227].

- [22] I.Y. Aref'eva, D.M. Belov, A.S. Koshelev and P.B. Medvedev, "Tachyon Condensation in Cubic Superstring Field Theory," *Nucl. Phys.* **B638** (2002) 3-20 [hep-th/0011117].
- [23] N. Berkovits, A. Sen and B. Zwiebach, "Tachyon Condensation in Superstring Field Theory," *Nucl. Phys.* **B587**(2000)147-178 [hep-th/0002211].
- [24] P.-J. De Smet, "Tachyon Condensation: Calculations in String Field Theory," hep-th/0109182.
- [25] M. R. Gaberdiel and B. Zwiebach, "Tensor Constructions of Open String Theories I: Foundations," *Nucl. Phys.* **B505** (1997) 569-624 [hep-th/9705038].
- [26] B. Zwiebach, "Komaba Lectures on String Field Theory and Tachyon Condensation," proceeding of the Workshop KOMABA 2000, <http://www-hep.phys.s.u-tokyo.ac.jp/etc/2000/komaba/Zwiebach.ps>
- [27] V.A. Kostelecký and S. Samuel, "On a Nonperturbative Vacuum for the Open Bosonic String," *Nucl. Phys.* **B336** (1990) 263;
 A. Sen and B. Zwiebach, "Tachyon Condensation in String Field Theory," *JHEP* **0003** (2000) 002 [hep-th/9912249];
 N. Moeller and W. Taylor, "Level truncation and the tachyon in open bosonic string field theory," *Nucl.Phys.* **B583**(2000)105-144 [hep-th/0002237].
- [28] K. Okuyama, "Ghost Kinetic Operator of Vacuum String Field Theory," *JHEP* **0201** (2002) 027 [hep-th/0201015].
- [29] J.N. Acosta, N. Berkovits and O. Chandia, "A Note on the Superstring BRST Operator," *Phys. Lett.* **B454** (1999) 247-248 [hep-th/9902178].
- [30] J. Klusoň, "Proposal for Background Independent Berkovits' Superstring Field Theory," *JHEP* **0107** (2001) 039 [hep-th/0106107];
 M. Sakaguchi, "Pregeometrical Formulation of Berkovits' Open RNS Superstring Field Theories," hep-th/0112135.
- [31] L. Rastelli and B. Zwiebach, "Tachyon Potentials, Star Products and Universality," *JHEP* **0109** (2001) 038, [hep-th/0006240].
- [32] I. Kishimoto and K. Ohmori, "CFT Description of Identity String Field: Toward Derivation of the VSFT Action," *JHEP* **0204** (2002) 036 [hep-th/0112169].

- [33] N. Berkovits, “Review of Open Superstring Field Theory,” hep-th/0105230.
- [34] K. Ohmori, “Comments on Solutions of Vacuum Superstring Field Theory,” *JHEP* **04** (2002) 059 [hep-th/0204138].
- [35] A. S. Schwarz and A. Sen, “Gluing Theorem, Star Product and Integration in Open String Field Theory in Arbitrary Background Fields,” *Int. J. Mod. Phys.* **A6** (1991) 5387–5407.
- [36] V. A. Kostelecký and R. Potting, “Analytical construction of a nonperturbative vacuum for the open bosonic string,” *Phys. Rev.* **D63**(2001)046007 [hep-th/0008252].
- [37] I.Y. Aref’eva, A.A. Giriyavets and P.B. Medvedev, “NS Matter Sliver,” *Phys. Lett.* **B532** (2002) 291-296 [hep-th/0112214].
- [38] M. Mariño and R. Schiappa, “Towards Vacuum Superstring Field Theory: The Supersliver,” hep-th/0112231.
- [39] N. Berkovits, “Super-Poincaré Invariant Superstring Field Theory,” *Nucl. Phys.* **B459** (1996) 439 [hep-th/9503099].
- [40] J. Klusoň, “Some Remarks About Berkovits’ Superstring Field Theory,” *JHEP* **0106** (2001) 045 [hep-th/0105319].
- [41] O. Lechtenfeld, A.D. Popov and S. Uhlmann, “Exact Solutions of Berkovits’ String Field Theory,” *Nucl. Phys.* **B637** (2002) 119-142 [hep-th/0204155].