

hep-th/0211011

SPIN-2002/31

ITF-2002/51

PUPT-2056

Open strings in the plane wave background I: Quantization and symmetries

Kostas Skenderis^{*} and Marika Taylor[†]

^{}Physics Department, Princeton University,
Princeton, NJ 08544, USA*

email:kostas@feynman.princeton.edu

*[†]Spinoza Institute, University of Utrecht,
Postbus 80.195, 3508 TD Utrecht, The Netherlands*

email:taylor@phys.uu.nl

ABSTRACT

We systematically investigate open strings in the plane wave background of type IIB string theory. We carefully analyze possible boundary conditions for open strings and find static as well as time-dependent branes. The branes fall into equivalence classes depending on whether they are related by the action of target space isometries. In particular static branes localized at the origin of transverse space and certain time-dependent branes fall into the same equivalence class. We analyze thoroughly the symmetries of all branes we discuss. Apart from symmetries descending from target space isometries, the worldsheet action being free admits a countably infinite number of other global worldsheet symmetries. We find that one can use such worldsheet symmetries to restore seemingly broken target space symmetries. In particular, we show that D-branes localized at arbitrary constant positions which were thought to be 1/4 supersymmetric in fact have sixteen supercharges whilst D-branes which were thought to be non-supersymmetric have eight supercharges. We discuss in detail the quantization in all cases.

Contents

1	Introduction and summary of results	2
2	Action and boundary conditions	6
2.1	Static D-branes	9
2.2	Symmetry-related D-branes	13
2.3	Time-dependent D-branes	15
3	Canonical Quantization	16
3.1	Mode expansions	17
4	Worldsheet symmetries and conserved charges	20
4.1	Closed strings	21
4.2	Open strings	25
5	Conclusions	37
A	Conventions	40
B	Symmetry currents and closed string superalgebra	41
C	Mode expansions for closed strings	42
D	Poisson-Dirac brackets for oscillators	43
D.1	D_- branes	44
D.2	D_+ branes	45
E	Kappa symmetry projections for time dependent branes	48

1 Introduction and summary of results

Understanding string theory on backgrounds with Ramond-Ramond (RR) fluxes is an important subject. Apart from its intrinsic importance, it has acquired a renewed interest due to the AdS/CFT duality. Understanding the gauge theory/gravity correspondence in the regime where the gauge theory is weakly coupled requires understanding string theory on such backgrounds. Despite considerable effort, however, our understanding remains somewhat limited.

Progress was made recently in [1, 2] where it was shown that closed string theory on a specific plane wave background with a RR flux is exactly solvable in the light cone gauge. This background is particularly interesting because it is obtained from the $AdS_5 \times S^5$ solution by a Penrose limit [3, 4]. The AdS/CFT correspondence then suggests a direct relation between gauge theory and string theory on this background, and indeed such a connection was proposed in [5].

Given the importance of string theory on RR backgrounds, it is of obvious interest to understand open strings and D-branes on such backgrounds. D-branes capture non-perturbative phenomena in string theory, and as such are bound to play an important role in the gravity/gauge theory correspondence. Moreover, much of our intuition about D-branes comes from flat space computations, and understanding in detail cases where the background is curved and contains RR fields may provide us with new insights.

Motivated by these issues, we start in this paper a systematic investigation of open strings and D-branes on the maximally supersymmetric plane wave background of type IIB supergravity [3], henceforth called “the plane wave background”. We analyze in this paper the possible D-branes, the quantization of open strings and the symmetries of the open string action. In a companion paper we will discuss the corresponding superalgebras and the spectra of these D-branes [6].

D-branes on the plane wave background have been investigated by various authors. Boundary states were constructed in [7, 8], D-branes in open string theory were studied in [9, 10], supersymmetric embeddings were classified in [11], supergravity solutions were constructed in [12], open-closed string duality was addressed in [8] and the duality with gauge theory was discussed in [13, 14, 11, 15]. Aspects of D-branes on this and other plane wave backgrounds were also discussed in [16, 17]. All these investigations yielded results consistent with each other. We extend these results in various directions and in particular find that certain D-branes are more supersymmetric than found previously. We will show that certain worldsheet symmetries can be used to restore seemingly broken target space

symmetries. The additional supersymmetries originate from such a mechanism.

Let us first review the existing results. Recall that in the plane wave background the eight directions transverse to the lightcone are divided into two sets of four. The target space supersymmetry in the lightcone gauge can be divided into dynamical supersymmetries Q^- and kinematical supersymmetries Q^+ . We shall call “dynamical” the supercharges that anticommute to the Hamiltonian (plus possibly other generators) and “kinematical” the supercharges that anticommute to P^+ . In the lightcone gauge the latter is just a c-number. The kinematical supercharges when evaluated on-shell are proportional to zero modes, but the dynamical ones depend on all modes. The half-supersymmetric D-branes are found to be of the type $(+, -, m, m \pm 2)$, where the notation indicates the worldvolume coordinates and m ($m \pm 2$) is the number of worldvolume coordinates in the first (second) set of four transverse directions. These D-branes preserve eight dynamical and eight kinematical supersymmetries when located at the origin of transverse space [9, 11], but only eight kinematical ones when located at an arbitrary constant position [11]. D-branes with different splitting of coordinates were found to break supersymmetry completely except for the case of the D1 brane which preserves eight dynamical supersymmetries irrespectively of its transverse position. Supersymmetric embeddings for D-branes that are not visible in open string theory in the light-cone gauge (because they wrap only one of the light-cone directions) were also discussed in [11].

The plane wave/gauge theory duality for the D5 $(+, -, 3, 1)$ case was discussed in [13], for the D7 $(+, -, 4, 2)$ in [14] and a uniform discussion for all D-branes was presented in [11]. In particular, it was proposed, and evidence was given, that all half supersymmetric D-branes are dual to defect conformal field theories. It was shown in [11] that the $(+, -, m, m \pm 2)$ probe branes located in the origin of transverse space originate from $AdS_{m+1} \times S^{m+1 \pm 2}$ embeddings in $AdS_5 \times S^5$ after the Penrose limit. The latter embeddings originate from orthogonal intersections $D(2m+1 \pm 2) \perp D3$ in the near-horizon limit. The defect conformal field theories involved in the gauge theory/plane wave duality are the theories of the $(2m + 2 \pm 2)$ -3 strings (i.e. the strings that stretch between the $D(2m + 1 \pm 2)$ and the $D3$ brane in the orthogonal intersection $D(2m + 1 \pm 2) \perp D3$).

In this paper we revisit the analysis of open string theory in the plane wave background. Such an analysis for (some) static D-branes was presented in [9, 10] and we will comment below on the relation of our results to theirs. The first step in our analysis is to determine the possible boundary conditions consistent with the variational problem. In addition, the Green-Schwarz action should be invariant under local κ -symmetry and the closed string

action constructed in [1] is indeed κ -symmetric. In order for the corresponding open string to be invariant under κ -symmetry one has to verify that the boundary terms that arise when checking the invariance of the action also vanish. Such a computation was carried out in [18] for flat space, and in [10] for the plane wave background. From the results of [10] one finds that when one imposes the fermionic lightcone gauge $\bar{\gamma}^+\theta = 0$ the boundary terms for all the branes we consider vanish. In [10] it was also shown that the boundary terms vanish for branes of type $(+, -, m, m \pm 2)$ localized at the origin of transverse space without imposing $\bar{\gamma}^+\theta = 0$. In our case this condition is always satisfied: the starting point of our analysis is the action where the kappa symmetry has already been gauge fixed, and we soon impose the lightcone and conformal gauges as well.

In our analysis we discuss both time-dependent as well as static branes. The static D-branes are the direct analogue of the ordinary flat space D-branes, i.e. they involve Neumann boundary conditions along the worldvolume of the brane and Dirichlet boundary conditions along the transverse directions. However, as reviewed above, the branes are also distinguished by the distribution of worldvolume coordinates in the transverse to the lightcone directions. It is convenient to introduce the notation D_- for the branes with worldvolume coordinates $(+, -, m, m \pm 2)$ and D_+ for all other $(+, -, m, n)$ branes (recall that in open string theory in the lightcone gauge, the lightcone directions are necessarily worldvolume coordinates). The reason for this terminology will become apparent in next section. Notice that there are only D_-p branes for $p = 3, 5, 7$.

The boundary conditions corresponding to static branes break some of the target space isometries. One may use the broken generators to generate new branes which we call here “symmetry-related branes”. These are equivalent to the original ones since the two are related by an isometric reparametrization of the target spacetime. In particular, we discuss in detail the branes obtained by translations and/or boosts of the static branes that are localized at the origin of transverse space. These symmetry-related branes satisfy time-dependent boundary conditions and in certain cases describe Penrose limits of giant gravitons and other rotating branes.

The boundary conditions are time dependent since in Brinkmann coordinates, which are the natural ones for string quantization, the corresponding Killing vectors depend on x^+ [3] and x^+ is identified with the worldsheet time τ in the light-cone gauge. As emphasized in [12], such branes are static in the Rosen coordinate system, that is, their transverse position is constant, since here the translations act as in flat space [4]. Notice that the symmetry-related branes that are located away from the origin are distinct from static branes located

at arbitrary constant position. (The latter in turn would translate to time-dependent branes in the Rosen coordinate system.) We also discuss more general time-dependent boundary conditions that are compatible with the variational problem.

We then proceed to quantize the open string with the boundary conditions just discussed. Since the model is free quantization is straightforward: one first expands in modes and then imposes the canonical commutation relations. The D_- and D_+ branes differ in their fermionic zero modes. For D_- branes these zero modes depend only on the worldsheet time coordinate τ , see (3.12), and can be obtained from the zero modes of the closed string fermions, see (C.2), upon identification. In the case of D_+ branes, however, the zero modes of the fermions depend on the worldsheet space coordinate σ , see (3.21), and there is no direct relation with the zero modes of the closed string.

The commutation relations for D_- branes were discussed in [9], and for type $D1$ -branes in [10]. Our results differ from those in these papers in numerical coefficients of the fermionic anticommutators, and in that the anticommutator of the zero modes for all D_+ branes is non-trivial. Note that our results have the correct flat space limit including coefficients.

The target space isometries that are compatible with the boundary conditions become symmetries of the open string action. We discuss in detail these symmetries and compute the corresponding Noether currents for all cases. In particular, part of the target space supersymmetries, which from now on will be called “closed string supersymmetries”, are compatible with the open string boundary conditions. Recall that for every Killing spinor of the background κ -symmetry implies that the closed string worldsheet action is invariant under a target space global supersymmetry. In the open string case, one has to take into account the boundary conditions. This was done in [10] and it was found that the allowed D -branes preserve exactly the target space supersymmetries found in the analysis of [11].

One of our main results is that one can restore some of the broken symmetries by using worldsheet symmetries. The worldsheet theory in the light cone gauge is quadratic in the fields. This implies that there are an infinite number of symmetries. The transformation rules are given by shifting each field by a parameter that satisfies the field equations. Such transformations leave the Lagrangian invariant up to total derivative terms. The transformations that are compatible with the boundary conditions are true symmetries of the theory. Expanding the parameter in a basis we find that there are a countably infinite number of such symmetries. The corresponding charges when evaluated on-shell are proportional to oscillators.

Worldsheet transformations that are not compatible with the boundary conditions play

a role as well. One may combine them with a seemingly broken target space symmetry to obtain a good symmetry. In this way we shall show that we can find eight additional dynamical supersymmetries in the case of D_- branes located away from the origin, and eight additional kinematical supersymmetries in the case of D1 branes. Thus these branes have sixteen supersymmetries but the additional supercharges do not originate from closed string supercharges.

One may understand why the branes possess additional supercharges as follows. In flat space, open string boundary conditions compatible with $2N$ target space supercharges lead to N fermionic oscillators at the k th mode. In the plane wave case the number of closed string supersymmetries preserved by our branes is in some cases less than 16, but there are still 8 fermionic oscillators at each mode level. These oscillators are spectrum generating and in all cases the spectrum automatically falls into supersymmetry representations. The corresponding supercharges, however, do not descend from target space superisometries, but are instead the new supercharges.

The paper is organized as follows. In the next section we discuss the variational problem for the open string in the plane wave background and possible consistent boundary conditions (D-branes). In section 3 we discuss the quantization of the open string for all cases considered in section 2 and in section 4 the corresponding symmetries and conserved charges. We conclude in section 5 with a discussion of our results. We include several appendices. In appendix A we give conventions and in appendices B and C we review results from closed strings in the plane wave background. In appendix D we discuss in detail the computation of the (anti)commutation relation for the oscillators. Appendix E contains a computation of the supersymmetry of certain time dependent D-branes from the D-brane worldvolume perspective.

2 Action and boundary conditions

In what follows we will consider open strings propagating in the maximally supersymmetric plane wave background [3] with Brinkmann metric

$$ds^2 = 2dx^+ dx^- + \sum_{I=1}^8 (dx^I dx^I - \mu^2 (x^I)^2 (dx^+)^2), \quad (2.1)$$

and RR flux

$$F_{+1234} = F_{+5678} = 4\mu. \quad (2.2)$$

It is useful to retain explicitly the mass parameter μ rather than scale it to one since the flat space limit will then be manifest¹.

The Green-Schwarz action in this background was constructed by Metsaev using supercoset techniques [1]. Just as for flat space, if one fixes the lightcone fermionic gauge the action becomes quadratic in fermions: the κ symmetry gauge fixed Green-Schwarz action takes the form

$$\begin{aligned}
S = T \int_{\Sigma} d^2\sigma & \left(-\frac{1}{2}\sqrt{-g}g^{ab}(2\partial_a x^+ \partial_b x^- - \mu^2 x_I^2 \partial_a x^+ \partial_b x^+ + \partial_a x^I \partial_b x^I) \right. \\
& -i\sqrt{-g}g^{ab} \partial_b x^+ (\bar{\theta}\bar{\gamma}^- \partial_a \theta + \theta\bar{\gamma}^- \partial_a \bar{\theta} + 2i\mu \partial_a x^+ \bar{\theta}\bar{\gamma}^- \Pi\theta) \\
& \left. +i\epsilon^{ab} \partial_a x^+ (\theta\bar{\gamma}^- \partial_b \theta + \bar{\theta}\bar{\gamma}^- \partial_b \bar{\theta}) \right). \tag{2.3}
\end{aligned}$$

In this expression g_{ab} is the worldsheet metric with (τ, σ) the worldsheet coordinates and $\epsilon^{01} = 1$. (x^+, x^-, x^I) are the bosonic coordinates of the target superspace whilst θ is a complex Weyl spinor satisfying the fermionic lightcone gauge condition $\bar{\gamma}^+ \theta = 0$. Here we use the spinor notation of [1], namely γ^μ and $\bar{\gamma}^\mu$ are the off-diagonal parts of the 32-dimensional gamma matrices Γ^μ . Our conventions are given in appendix A. Note that the complex Weyl spinor can be replaced by two real Majorana spinors such that

$$\theta = \frac{1}{\sqrt{2}}(\theta^1 + i\theta^2); \tag{2.4}$$

we will use both in what follows, according to convenience. The explicit appearance of $\Pi = \gamma^{1234}$ in the action reflects the fact that the RR flux background breaks the transverse symmetry to $SO(4) \times SO(4)'$. The overall normalization of the action T is taken to be the inverse of the length of the string (that is, 2π for the closed string and π for the open string).

We would now like to derive the field equations for the open strings. To do this one should consider variations of the action (2.3); these will include both bulk and boundary terms which may schematically be represented as

$$\delta S = T \int_{\Sigma} d^2\sigma \delta z^m F_m(z^n) + \int_{\partial\Sigma} d\sigma_a \delta z^m G_m^a(z^n), \tag{2.5}$$

where z^m extends over all the coordinates of the target superspace, which in this case are $(x^+, x^-, x^I, \theta^1, \theta^2)$, and the second integral is over the worldsheet boundary, $\partial\Sigma$. One should then choose boundary conditions and if necessary add boundary terms such that

¹We use the supergravity conventions given in appendix A of [11]. Compared with the conventions in [1] one should rescale F_5 by a factor of 2. We also note that the coordinates used here are related to those used in (6.12)-(6.13) of [11] by $x^+ \rightarrow \mu x^+, x^- \rightarrow -1/(2\mu)x^-, x^I \rightarrow x^I$.

the variational problem is well defined, i.e. so that $\delta S = 0$ implies the field equations. We discuss this procedure in detail here since we are interested in non-standard boundary conditions.

Usually the boundary conditions chosen will automatically set to zero $\delta z^m G_m^a$ for each m , and then $F_m = 0$ will give us the full complement of field equations. However, one may also wish to implement boundary conditions for which one or more component of $\delta z^m G_m^a$ is non-vanishing and is not cancelled by another component. In this case it would be necessary to add further boundary terms to the action such that under variations with these boundary conditions the total variation is zero on the boundary (for each m)². We will impose fermionic boundary conditions such that $\delta\theta^1 G_a^{\theta^1}$ cancels with $\delta\theta^2 G_a^{\theta^2}$; this relates θ^1 to θ^2 on the boundary and reduces the number of independent fermionic degrees of freedom propagating on the worldsheet. Note that locality requires that the boundary terms vanish separately at each boundary: only non-local conditions could implement cancellation between terms at different boundaries.

Before presenting the explicit forms for the variations of the action (2.3) let us comment that by starting with an action in which the kappa symmetry has already been fixed we will necessarily exclude boundary conditions (and hence D-branes) which are not compatible with this fermionic gauge choice. These include branes along one lightcone direction discussed in [11]. Furthermore, as mentioned in the introduction, it is important that the boundary conditions we consider here are, using the results of [10], compatible with the kappa symmetry of the original non gauge-fixed action given in [1].

So let us now consider the variation of the bulk action (2.3). Under x^- variations one obtains

$$\delta S = T \left(\int_{\Sigma} d^2\sigma \delta x^- \partial_a - \int_{\partial\Sigma} d\sigma_a \delta x^- \right) \sqrt{-g} g^{ab} \partial_b x^+, \quad (2.6)$$

whilst under x^I variations one finds

$$\begin{aligned} \delta S &= T \int_{\Sigma} d^2\sigma \left(\partial_a (\sqrt{-g} g^{ab} \partial_b x^I) + \mu^2 x^I \sqrt{-g} g^{ab} \partial_a x^+ \partial_b x^+ \right) \delta x^I \\ &\quad - T \int_{\partial\Sigma} d\sigma_a \sqrt{-g} g^{ab} \partial_b x^I \delta x^I. \end{aligned} \quad (2.7)$$

The x^+ variations give

$$\begin{aligned} \delta S &= T \left(\int_{\Sigma} d^2\sigma \delta x^+ \partial_a - \int_{\partial\Sigma} d\sigma_a \delta x^+ \right) \left(-i\epsilon^{ab} (\theta^1 \bar{\gamma}^- \partial_b \theta^1 - \theta^2 \bar{\gamma}^- \partial_b \theta^2) + \right. \\ &\quad \left. \sqrt{-g} g^{ab} [\partial_b x^- - \mu^2 x_I^2 \partial_b x^+ + i(\theta^1 \bar{\gamma}^- \partial_b \theta^1 + \theta^2 \bar{\gamma}^- \partial_b \theta^2 - 4\mu \partial_b x^+ \theta^1 \bar{\gamma}^- \Pi \theta^2)] \right). \end{aligned} \quad (2.8)$$

²One would also have to check that corresponding boundary terms added to the original action (before fixing kappa symmetry) ensure that the total action is κ symmetric with these boundary conditions.

The θ^1 variations give

$$\begin{aligned}\delta S &= T \int_{\Sigma} d^2\sigma \delta\theta^1 \left(-i\partial_a(\sqrt{g}g^{ab}\partial_b x^+)(\bar{\gamma}^-\theta^1) \right. \\ &\quad \left. -2i\sqrt{-g}g^{ab}\partial_b x^+(\bar{\gamma}^-\partial_a\theta^1 - \mu\partial_a x^+\bar{\gamma}^-\Pi\theta^2) + 2i\epsilon^{ab}\partial_a x^+\bar{\gamma}^-\partial_b\theta^1 \right) \\ &\quad + T \int_{\partial\Sigma} d\sigma_a \left(\sqrt{-g}g^{ab} + \epsilon^{ab} \right) i\partial_b x^+ \delta\theta^1 \bar{\gamma}^-\theta^1\end{aligned}\tag{2.9}$$

whilst the θ^2 variations give

$$\begin{aligned}\delta S &= T \int_{\Sigma} d^2\sigma \delta\theta^2 \left(-i\partial_a(\sqrt{g}g^{ab}\partial_b x^+)(\bar{\gamma}^-\theta^2) \right. \\ &\quad \left. -2i\sqrt{-g}g^{ab}\partial_b x^+(\bar{\gamma}^-\partial_a\theta^2 + \mu\partial_a x^+\bar{\gamma}^-\Pi\theta^1) - 2i\epsilon^{ab}\partial_a x^+\bar{\gamma}^-\partial_b\theta^1 \right) \\ &\quad + T \int_{\partial\Sigma} d\sigma_a \left(\sqrt{-g}g^{ab} - \epsilon^{ab} \right) i\partial_b x^+ \delta\theta^2 \bar{\gamma}^-\theta^2.\end{aligned}\tag{2.10}$$

Furthermore the worldsheet energy momentum tensor is

$$T_{ab} = -\frac{2}{T\sqrt{-g}} \frac{\delta S}{\delta g^{ab}} = L_{ab} - \frac{1}{2}(g^{cd}L_{cd})g_{ab};\tag{2.11}$$

$$\begin{aligned}L_{ab} &= 2\partial_{(a}x^+\partial_{b)}x^- - \mu^2 x^2 \partial_a x^+ \partial_b x^+ + \partial_a x^I \partial_b x^I \\ &\quad + 2i\partial_{(a}x^+(\theta^1\bar{\gamma}^-\partial_{b)}\theta^1 + \theta^2\bar{\gamma}^-\partial_{b)}\theta^2 - 2\mu\partial_{b)}x^+\theta^1\bar{\gamma}^-\Pi\theta^2),\end{aligned}\tag{2.12}$$

Here and in all that follows we (anti)symmetrise with unit strength. Notice that the stress energy tensor is traceless, i.e.

$$g^{ab}T_{ab} = 0.\tag{2.13}$$

The field equations obtained by varying the worldsheet metric (Virasoro constraints) amount to $T_{ab} = 0$.

In the following sections we will consider open strings with a variety of different boundary conditions correspondent to D-branes. We now describe in detail these boundary conditions.

2.1 Static D-branes

The simplest and most familiar choice of boundary conditions corresponds to static D-branes. For these, the bosonic boundary conditions are as follows. We first impose Neumann conditions on $(p-1)$ transverse coordinates and Dirichlet boundary conditions on the remaining transverse coordinates:

$$\begin{aligned}\partial_{\sigma}x^r| &= 0, & r &= 1, \dots, (p-1) \\ \partial_{\tau}x^{r'}| &= 0, & r' &= p, \dots, 8.\end{aligned}\tag{2.14}$$

Our index conventions are as follows: we use indices, $r, s, t = 1, \dots, (p-1)$, from the end of the latin alphabet to denote coordinates with Neumann boundary conditions, and primed indices of the same letters, $r', s', t' = p, \dots, 8$, to denote coordinates with Dirichlet boundary conditions. Latin indices, $i, j = 1, \dots, 4$, and $i', j' = 5, \dots, 8$, from the middle of the alphabet are used for the two sets of the transverse coordinates that transform among themselves under the $SO(4)$ and the $SO(4)'$, respectively.

We fix the boundary of the worldsheet to be at $\sigma = 0, \pi$ and denote by $A|$ the restriction of a quantity to the boundary. We also at this stage fix conformal gauge for the worldsheet metric $-g_{00} = g_{11} = 1$. Note that as usual we have assumed that all variations vanish at $\tau \rightarrow \pm\infty$ and so we need only consider the spatial boundary of the worldsheet. We also choose x^+ to satisfy a Neumann condition.

Now such boundary conditions remove all boundary terms in (2.6) and (2.7). The resulting field equation for x^+ together with the Neumann boundary condition forces us into bosonic lightcone gauge $x^+ = p^+\tau$. (Note that one can also include an overall integration constant x_0^+ but since it does not play a role we do not include it here.)

Appropriate fermion boundary conditions are

$$\theta^1| = \Omega\theta^2|, \quad (2.15)$$

at $\sigma = 0$ and $\sigma = \pi$. Ω is a real (constant) 16×16 dimensional matrix. Different matrices Ω can be used at $\sigma = 0$ and $\sigma = \pi$, as representing open string extending between distinct branes, but here we take Ω to be the same at both ends. Such a boundary condition necessarily reduces the number of independent fermion modes by one half and preserves at most one half of the supersymmetry. As previously advertised, the fermionic boundary condition (2.15) can cancel boundary terms between (2.9) and (2.10). That is, given the Neumann boundary condition on x^+ and since $\delta\theta^1 = \Omega\delta\theta^2$ on the boundary to preserve (2.15) we are left with a fermion boundary term proportional to

$$(\delta\theta^1\bar{\gamma}^-\theta^1 - \delta\theta^2\bar{\gamma}^-\theta^2)|, \quad (2.16)$$

which vanishes provided that Ω satisfies

$$\Omega^t\bar{\gamma}^-\Omega = \bar{\gamma}^-, \quad (2.17)$$

where Ω^t is the transpose of Ω . The fact that θ^1 and θ^2 have the same chirality implies that Ω should contain an even number of γ matrices, and using the lightcone condition $\gamma^+\theta^{1,2} = 0$ one finds that Ω may not contain γ^+ or γ^- . It then follows from (2.17) that

Ω should be an orthogonal matrix, $\Omega^t \Omega = 1$. Furthermore, the study of supersymmetry in later sections divides the discussion into Ω that satisfy one of the following two conditions

$$D_- : \quad \Omega \Pi \Omega \Pi = -1, \quad D_+ : \quad \Omega \Pi \Omega \Pi = 1. \quad (2.18)$$

We will call the branes that satisfy the former condition “ D_- branes”, and the ones that satisfy the latter condition “ D_+ branes”. As we will discuss in the next section, for a single D-brane the appropriate Ω is the product of gamma matrices $\gamma^{r'}$ where r' is transverse to the D-brane³.

After fixing bosonic lightcone gauge ($x^+ = p^+ \tau$) and conformal gauge ($-g_{00} = g_{11} = 1$) the action becomes

$$S = T \int d^2 \sigma \left(p^+ \partial_\tau x^- + \frac{1}{2} ((\partial_\tau x^I)^2 - (\partial_\sigma x^I)^2 - m^2 (x^I)^2) + i p^+ (\theta^1 \bar{\gamma}^- \partial_+ \theta^1 + \theta^2 \bar{\gamma}^- \partial_- \theta^2 - 2m \theta^1 \bar{\gamma}^- \Pi \theta^2) \right). \quad (2.19)$$

Here $\partial_\pm = \partial_\tau \pm \partial_\sigma$ and $m = \mu p^+$. The field equations for $(x^I, \theta^1, \theta^2)$ in bosonic lightcone gauge and conformal gauge ($-g_{00} = g_{11} = 1$) are then

$$\begin{aligned} (\partial_- \partial_+ + m^2) x^I &= 0; \\ (\partial_+ \theta^1 - m \Pi \theta^2) &= 0; \\ (\partial_- \theta^2 + m \Pi \theta^1) &= 0, \end{aligned} \quad (2.20)$$

to be solved together with the boundary conditions (2.14) and (2.15).

The Virasoro constraints in bosonic lightcone gauge and in conformal gauge for the metric are

$$\begin{aligned} T_{\sigma\tau} &= p^+ [\partial_\sigma x^- + i(\theta^1 \bar{\gamma}^- \partial_\sigma \theta^1 + \theta^2 \bar{\gamma}^- \partial_\sigma \theta^2)] + \partial_\tau x^I \partial_\sigma x^I = 0; \\ T_{\tau\tau} &= p^+ [\partial_\tau x^- + i(\theta^1 \bar{\gamma}^- \partial_\tau \theta^1 + \theta^2 \bar{\gamma}^- \partial_\tau \theta^2 - 2m \theta^1 \bar{\gamma}^- \Pi \theta^2)] \\ &\quad - \frac{1}{2} (m^2 (x^I)^2 - (\partial_\tau x^I)^2 - (\partial_\sigma x^I)^2) = 0. \end{aligned} \quad (2.21)$$

From (2.8), once we fix both bosonic lightcone gauge and conformal gauge, we get

$$\begin{aligned} \delta S &= T \int d^2 \sigma \delta x^+ (p^+)^{-1} (-\partial_\tau T_{\tau\tau} + \partial_\sigma T_{\sigma\tau} + \partial_\tau x^I (\partial_+ \partial_- x^I + m^2 x^I) \\ &\quad - 2i p^+ [\partial_\tau \theta^1 \bar{\gamma}^- (\partial_+ \theta^1 - m \Pi \theta^2) + \partial_\tau \theta^2 \bar{\gamma}^- (\partial_- \theta^2 + m \Pi \theta^1)]) \\ &\quad - T \int d\tau \delta x^+ (\partial_\sigma x^- + i(\theta^1 \bar{\gamma}^- \partial_\sigma \theta^1 + \theta^2 \bar{\gamma}^- \partial_\sigma \theta^2)). \end{aligned} \quad (2.22)$$

³Other choices of Ω are compatible with preserving closed string kinematical supersymmetries but not with dynamical supersymmetries. They will also generically break worldvolume covariance. We will not pursue such alternative boundary conditions here.

In the bulk term we have rewritten terms involving x^- using $(T_{\tau\tau}, T_{\tau\sigma})$ from (2.21) whilst in the boundary term we have already dropped terms which vanish because of the boundary conditions imposed on $(x^I, x^+, \theta^1, \theta^2)$. Now to make the x^+ variational problem well defined, the boundary term should vanish. This means that

$$(\partial_\sigma x^- + i(\theta^1 \bar{\gamma}^- \partial_\sigma \theta^1 + \theta^2 \bar{\gamma}^- \partial_\sigma \theta^2))| = 0 \quad (2.23)$$

at the boundary. This condition is automatically satisfied provided the Virasoro constraint $T_{\sigma\tau} = 0$ holds on the boundary. For both D_- and D_+ branes the fermion bilinear in (2.23) vanishes at the boundary⁴, so x^- satisfies a Neumann boundary condition.

Each term in the bulk variation involves either a Virasoro constraint or a field equation. Thus the bulk variation term vanishes automatically for all δx^+ provided that we impose the other field equations and the Virasoro constraints. One can understand this result as follows. Recall that given a conserved current \mathcal{G}^a that generates symmetry transformations $\delta\phi^A$, where ϕ^A are all the fields in the theory, its divergence is equal to

$$\partial_a \mathcal{G}^a = \delta\phi^A \frac{\delta S}{\delta\phi^A} \quad (2.24)$$

where it is understood that the parameter of the transformation has been removed from $\delta\phi^A$.⁵ In addition, in the presence of boundaries the spatial components of the current should vanish at the boundary. The derivation of the latter is elementary and will be presented in section 4.2.

Now let us apply this result to the current $T_{\alpha\tau}$ that generates worldsheet time translations,

$$\partial^\alpha T_{\alpha\tau} = \partial_\tau x^+ \frac{\delta S}{\delta x^+} + \partial_\tau x^I \frac{\delta S}{\delta x^I} + \partial_\tau \theta^1 \frac{\delta S}{\delta \theta^1} + \partial_\tau \theta^2 \frac{\delta S}{\delta \theta^2} + \partial_\tau x^- \frac{\delta S}{\delta x^-}. \quad (2.25)$$

In the light-cone gauge $\partial_\tau x^+ = p^+$, so

$$\frac{\delta S}{\delta x^+} = (p^+)^{-1} (\partial^\alpha T_{\alpha\tau} - \partial_\tau x^I \frac{\delta S}{\delta x^I} - \partial_\tau \theta^1 \frac{\delta S}{\delta \theta^1} - \partial_\tau \theta^2 \frac{\delta S}{\delta \theta^2} - \partial_\tau x^- \frac{\delta S}{\delta x^-}). \quad (2.26)$$

It follows that the x^+ field equation is implied by the Virasoro constraints and the other field equations (notice that in our case $\delta S/\delta x^-$ is identically equal to zero). One can readily check that (2.22) agrees with (2.26). In addition, $T_{\sigma\tau}$ should vanish at the boundary in order for the current $T_{a\tau}$ to lead to a conserved current, and this gives exactly (2.23).

⁴To show this notice that for D_- and D_+ branes the boundary condition (2.15) together with the field equation imply $|\partial_\sigma \theta^1| = -\Omega \partial_\sigma \theta^2|$ and $|\partial_\sigma \theta^1| = (-\Omega \partial_\sigma \theta^2 + 2m\Pi\theta^2)|$, respectively. In the latter case the vanishing of the bilinear in (2.23) follows upon using the fact that $\Omega\Pi$ is a symmetric matrix, for the Ω given in (3.20).

⁵Given a Lagrangian L that transforms under a symmetry as $\delta L = \partial^a k_a$, for some k_a (so that the action is invariant), the Noether current is given by $\mathcal{G}^a = \partial L / (\partial(\partial_a \phi^A)) \delta\phi^A - k^a$. Taking the divergence of this expression one gets (2.24).

2.2 Symmetry-related D-branes

Let G be the isometry group of the target spacetime. An isometry of the target spacetime may or may not leave invariant a given set of boundary conditions for open strings. Let H be the subgroup of G that respects the boundary conditions. These transformations descend to symmetries of the open string action and will be discussed in the next section. On the other hand, the action of the coset generators G/H results in new D-branes (i.e. open strings with a different boundary condition), which however may be considered as equivalent to the original ones since the two D-branes are related by an isometric reparametrization of the target spacetime. In this subsection we discuss the branes that are in the same equivalence class as the static branes; we refer to them as symmetry related branes.

The full set of (infinitesimal) bosonic isometries of the pp-wave background is as follows [3] (fermionic isometries will be discussed in the next section). Translations generated by P^\pm and P^I are given by

$$P^\pm : \quad \delta x^\mp = \epsilon^\mp; \quad (2.27)$$

$$P^I : \quad \delta x^- = \mu \sin \mu x^+ \epsilon^I x^I, \quad \delta x^I = \cos \mu x^+ \epsilon^I. \quad (2.28)$$

The rotational currents J^{+I} , J^{ij} and $J^{i'j'}$ (where x^i and $x^{i'}$ parametrise $SO(4)$ and $SO(4)'$ respectively) are associated with the respective transformations

$$J^{+I} : \quad \delta x^- = -\cos \mu x^+ \epsilon^I x^I, \quad \delta x^I = \mu^{-1} \sin \mu x^+ \epsilon^I; \quad (2.29)$$

$$J^{ij} : \quad \delta x^i = -\epsilon^{ij} x^j, \quad \delta \theta = -\frac{1}{4} \epsilon^{ij} \gamma^{ij} \theta; \quad (2.30)$$

$$J^{i'j'} : \quad \delta x^{i'} = -\epsilon^{i'j'} x^{j'}, \quad \delta \theta = -\frac{1}{4} \epsilon^{i'j'} \gamma^{i'j'} \theta, \quad (2.31)$$

where $\epsilon^{ij} = -\epsilon^{ji}$ and $\epsilon^{i'j'} = -\epsilon^{j'i'}$.

Let us consider a finite $P^{r'}$ translation,

$$\begin{aligned} \hat{x}^- &= x^- - \mu \sin(\mu x^+) x^{r'} a^{r'} + \frac{1}{2} \mu \sin(\mu x^+) \cos(\mu x^+) (a^{r'})^2; \\ \hat{x}^{r'} &= x^{r'} - \cos(\mu x^+) a^{r'}, \end{aligned} \quad (2.32)$$

where $a^{r'}$ is a finite global parameter, and the rest of the coordinates are unchanged. One easily checks that this is an isometry: $g_{\hat{x}^\mu \hat{x}^\nu} = g_{x^\mu x^\nu}$.

Now we consider ‘‘standard’’ static boundary conditions in this new coordinate system, namely

$$\partial_\sigma \hat{x}^+| = 0; \quad \hat{x}^{r'}| = 0; \quad \partial_\sigma \hat{x}^-| = 0. \quad (2.33)$$

These boundary conditions correspond to the following conditions in our original coordinate system:

$$\partial_\sigma x^+| = 0; \quad \partial_\sigma x^-| = \mu \sin(\mu x^+) a^{r'} \partial_\sigma x^{r'}|; \quad x^{r'}| = a^{r'} \cos(\mu x^+)|. \quad (2.34)$$

Note that in the modified boundary conditions (2.34) the Dirichlet position $x^{r'}$ is not constant but is explicitly time dependent. This results of course from the explicit time (x^+) dependence of the translation Killing vectors. In the flat space limit ($\mu \rightarrow 0$), the Dirichlet position is purely constant, since the explicit time dependence drops out of the isometry.

One can also act with the other Killing vectors of the background on the standard boundary conditions to generate modified boundary conditions. The action of isometries along the worldvolume is trivial in the sense that this simply maps each point on the worldvolume to another point on the worldvolume, without affecting the transverse positions. Thus the boundary conditions are not affected by the action of these isometries. These isometries generate symmetries of the D-brane, and are discussed in the next section.

The only isometries which will generate distinct embeddings are $P^{r'}$, $J^{+r'}$ and $J^{rr'}$. The action of $J^{+r'}$ will be very similar to that of $P^{r'}$ (effectively exchanging sines and cosines in the expressions given above). Thus a brane which is shifted by a combination of $P^{r'}$ and $J^{+r'}$ transformations will have boundary conditions:

$$x^{r'}| = (a^{r'} \cos(\mu x^+) + b^{r'} \sin(\mu x^+))|. \quad (2.35)$$

Here both $a^{r'}$ and $b^{r'}$ are constant vectors in the Dirichlet directions. When these vectors are orthogonal, say $a^{r'} = a$ and $b^{s'} = b$, the string traces out a trajectory of an ellipse in the $(r's')$ plane, with angular velocity μ with respect to x^+ . Thus these boundary conditions are appropriate to describe a rotating brane. In particular, they describe the branes arising from Penrose limits of giant gravitons.

When either of a or b are zero, the brane oscillates along a line. (2.35) represents the most general symmetry related brane boundary condition, up to the action of the surviving symmetries, and is the condition we will use in what follows. The surviving symmetries can shift, for example, $x^+ \rightarrow x^+ + x_0^+$ which changes the phase of the motion.

The rotational symmetries J^{IJ} in the pp-wave background act exactly as in flat space. Under a finite J^{ij} transformation,

$$\hat{x}^i = x^i \cos a - x^j \sin a, \quad \hat{x}^j = x^j \cos a + x^i \sin a, \quad (2.36)$$

where a is a finite global parameter. If both coordinates are originally Dirichlet they will both remain Dirichlet after the rotation. Furthermore, if the transverse coordinates are zero, the rotation will act trivially. However, if $i = r$ (Neumann) and $j = r'$ (Dirichlet) the boundary conditions will be mixed after the rotation: as we said above, the broken $J^{rr'}$ symmetry does act non-trivially. Since the $J^{rr'}$ transformations are independent of μ the

discussion of such rotated branes mirrors that in flat space (see for example [19]) and we will not consider them further here.

One should contrast the above discussion with the case of branes in flat space. In that case, $\partial_{x^{r'}}$ is Killing vector and one can use this isometry to position a D-brane at any constant transverse position. In the pp-wave background, however, it is $P^{r'}$ rather than $\partial_{x^{r'}}$ that is the Killing vector. Thus, even though the pp-wave geometry is homogeneous, and any two points are related by a symmetry transformation, this does *not* mean that *static* branes at distinct constant values of $x^{r'}$ are related by symmetry. The symmetry transformation relates static boundary conditions to time dependent ones.

2.3 Time-dependent D-branes

We should note here that one can also consider more general time dependent boundary conditions for the open strings. Such boundary conditions are consistent open string boundary conditions in bosonic lightcone gauge, both in flat space and in the pp-wave background. In fact, *any* transverse boundary conditions of the form

$$x^{r'}| = x^{r'}(x^+)| \quad (2.37)$$

are consistent with the field equations (2.20) in light cone gauge. To see this first notice that the choice of bosonic light cone gauge allows arbitrary δx^+ on the boundary. Now consider what variations $\delta x^{r'}$ preserve the boundary conditions (2.37): we need to enforce

$$p^+ \delta x^{r'}| = (\partial_\tau x^{r'}) \delta x^+|, \quad (2.38)$$

on the boundary. For the fermionic boundary conditions let us assume the same condition as for the static D-branes holds, namely (2.15) holds on the boundary, with appropriate Ω that satisfies $\Omega \Pi \Omega \Pi = \pm 1$. Now if we substitute these boundary conditions into the action variation given in (2.6), (2.7), (2.8), (2.9) and (2.10) we are left with the following boundary term

$$\begin{aligned} \delta S &= - \int d\tau (\partial_\sigma x^I \delta x^I + \delta x^+ \partial_\sigma x^-); \\ &= -(p^+)^{-1} \int d\tau \delta x^+ (\partial_\sigma x^I \partial_\tau x^I + p^+ \partial_\sigma x^-) = -(p^+)^{-1} \int d\tau \delta x^+ T_{\tau\sigma}, \end{aligned} \quad (2.39)$$

where in the second line we have used $\partial_\sigma x^r = 0$ for Neumann directions and (2.38) for the transverse ‘‘Dirichlet’’ directions. So the boundary term again vanishes, provided that we impose the Virasoro constraint $T_{\tau\sigma} = 0$ on the boundary. We are then left with the usual field equations (2.20) along with the boundary conditions (2.37) and (2.15). Thus time

dependent transverse boundary conditions are consistent with the usual field equations (even in flat space, since nowhere did we require $\mu \neq 0$) regardless of the explicit x^+ dependence. Furthermore, such boundary conditions also solve the DBI field equations. It was already known in flat space that even arbitrary $x^{r'}(x^+)$ is consistent with preservation of one quarter supersymmetry (half the kinematical supercharges), see for example [20], but later we will show that in fact such branes admit a full compliment of sixteen supercharges.

In this section we have considered a number of generalized Dirichlet boundary conditions for open strings. We should also mention that it would be interesting to consider branes carrying fluxes. It was found in [11] and [16] that certain constant fluxes can displace with D -branes from the origin while preserving the same 16 supercharges as the brane at the origin and certain D_+ 5-branes with non-constant fluxes preserved one quarter of the supersymmetry [11]. We leave for future work the exploration of the related open strings coupling to flux condensates.

3 Canonical Quantization

In this section we present the canonical quantization of open strings with the various boundary conditions just discussed. It is useful to start by recalling the solutions of the field equations (2.20) without imposing any restrictions due to boundary conditions and/or the reality conditions of the fields. Recall that the most general solutions of the equation

$$(\partial_\tau^2 - \partial_\sigma^2 + m^2)\phi = 0, \quad (3.1)$$

are

$$\phi_k(\tau, \sigma) = e^{-i(w_k\tau + k\sigma)}, \quad \tilde{\phi}_k(\tau, \sigma) = e^{-i(w_k\tau - k\sigma)}, \quad (3.2)$$

where $\omega_k = \text{sgn}(k)\sqrt{k^2 + m^2}$, k is an arbitrary complex parameter, and $\text{sgn}(k) = \sqrt{k^2}/k$. Notice that the choices $k = 0$ and $k = \pm im$ are special, because in these cases ϕ_k depends only on one of the two worldsheet coordinates,

$$\phi_0 = e^{-im\tau}, \quad \phi_{im} = e^{m\sigma}, \quad \phi_{-im} = e^{-m\sigma}. \quad (3.3)$$

Given a solution of (3.1) one can construct a solution of the fermionic equations as

$$\theta^1 = id_k\phi_k\Pi\epsilon_k + \tilde{\phi}_l\tilde{\epsilon}_l, \quad \theta^2 = \phi_k\epsilon_k - id_l\tilde{\phi}_l\Pi\tilde{\epsilon}_l, \quad (3.4)$$

where ϵ_k and ϵ_l are (independent) constant spinors, k and l are general complex parameters and

$$d_k = \frac{1}{m}(\omega_k - k). \quad (3.5)$$

3.1 Mode expansions

The mode expansions for the closed and open strings with various boundary conditions follow from the results we just presented by imposing appropriate boundary conditions.

For closed strings the complete set of solutions of (3.1) is given by (3.2) with k integral. The mode expansions of the bosonic and fermionic coordinates were given in [2] and we summarise these results for later convenience in appendix C.

The explicit solution for the bosonic coordinates for a string with $(p-1)$ Neumann boundary conditions and “generalized” Dirichlet boundary conditions on the remaining coordinates is given by

$$x^r(\sigma, \tau) = x_0^r \cos(m\tau) + m^{-1} p_0^r \sin(m\tau) + i \sum_{n \neq 0} \omega_n^{-1} \alpha_n^r e^{-i\omega_n \tau} \cos(n\sigma); \quad (3.6)$$

$$x^{r'}(\sigma, \tau) = x_0^{r'}(\sigma, \tau) + \sum_{n \neq 0} \omega_n^{-1} \alpha_n^{r'} e^{-i\omega_n \tau} \sin(n\sigma), \quad (3.7)$$

where the zero mode part, $x_0^{r'}(\sigma, \tau)$, satisfies (3.1) and its exact form depends on the boundary conditions under consideration. The two types of zero modes we will consider are the following. For a static brane located at $x_0^{r'}$ it is

$$x_0^{r'}(\sigma, \tau) = \frac{x_0^{r'}}{e^{m\pi} + 1} (e^{m\sigma} + e^{m(\pi-\sigma)}), \quad (3.8)$$

For the symmetry related branes, as discussed in section 2.2, the appropriate zero modes can be read off from (2.35). Reality of the bosonic coordinates requires that the zero modes are real and

$$\alpha_n^r = (\alpha_{-n}^r)^\dagger, \quad \alpha_n^{r'} = (\alpha_{-n}^{r'})^\dagger. \quad (3.9)$$

It is convenient to discuss separately D_- and D_+ branes. The Ω appropriate to a single static D-brane is a product of the gamma matrices with indices along transverse directions:

$$\Omega = \prod_{r'=p}^8 \gamma^{r'}. \quad (3.10)$$

For the corresponding anti-brane, $\Omega \rightarrow -\Omega$.

3.1.1 D_- branes

For these there are the following possibilities:

$$\begin{aligned} D7 & : (+, -, 2, 4) \quad (+, -, 4, 2) \\ D5 & : (+, -, 1, 3) \quad (+, -, 3, 1) \\ D3 & : (+, -, 0, 2) \quad (+, -, 2, 0). \end{aligned} \quad (3.11)$$

In these expressions we divide the eight transverse coordinates into two groups of four, reflecting the $SO(4) \times SO(4)'$ symmetry of the background. Following the notation of [11], (m, n) implies that there are m Neumann directions in the first group and n in the second. Solving the fermion equations of motion, the mode expansions are

$$\begin{aligned}\theta^1 &= \theta_0 \cos(m\tau) + \tilde{\theta}_0 \sin(m\tau) + \sum_{n \neq 0} c_n \left(id_n \Pi \theta_n \phi_n + \tilde{\theta}_n \tilde{\phi}_n \right); \\ \theta^2 &= \Pi \tilde{\theta}_0 \cos(m\tau) - \Pi \theta_0 \sin(m\tau) + \sum_{n \neq 0} c_n \left(-id_n \Pi \tilde{\theta}_n \tilde{\phi}_n + \theta_n \phi_n \right),\end{aligned}\tag{3.12}$$

where ϕ_n and $\tilde{\phi}_n$ are given in (3.2), n is an integer and

$$c_n = \frac{1}{\sqrt{1 + d_n^2}}.\tag{3.13}$$

Notice that the zero mode part is also of the form (3.4). In particular, if $\epsilon_0 = \theta_0 + i\tilde{\theta}_0$ then the zero modes of θ_1 and θ_2 are given by $\theta^1 = \text{Re}(\phi_0 \epsilon_0)$ and $\theta^2 = \text{Re}(-id_0 \phi_0 \Pi \epsilon_0)$.

The boundary condition enforces

$$\tilde{\theta}_0 = -\Omega \Pi \theta_0; \quad \tilde{\theta}_n = \Omega \theta_n.\tag{3.14}$$

Combined with the chirality and light cone gauge constraints on θ this reduces the number of independent zero modes to eight. Reality of the spinors requires that

$$\theta_{-n} = (\theta_n)^\dagger.\tag{3.15}$$

For canonical quantisation we introduce canonical momenta

$$P^I = \dot{x}^I, \quad \mathcal{P}^{\mathcal{I}a} = \theta^{\mathcal{I}a},\tag{3.16}$$

where $\mathcal{I} = 1, 2$. Constraints on the fermionic phase space can be incorporated using Dirac brackets. The classical Poisson-Dirac brackets are given by [1]

$$\begin{aligned}[P^I(\sigma), x^J(\sigma')]_{P.B.} &= \pi \delta^{IJ} \delta(\sigma, \sigma'); \\ \{(\theta^1)^\alpha(\sigma), (\theta^1)^\beta(\sigma')\}_{D.B.} &= \frac{i}{4p^+} \pi (\gamma^+)^{\alpha\beta} \delta(\sigma, \sigma'); \\ \{(\theta^2)^\alpha(\sigma), (\theta^2)^\beta(\sigma')\}_{D.B.} &= \frac{i}{4p^+} \pi (\gamma^+)^{\alpha\beta} \delta(\sigma, \sigma'); \\ [x_0^-, (\theta^1)^\alpha]_{D.B.} &= \frac{\pi}{2p^+} (\theta^1)^\alpha, \quad [x_0^-, (\theta^2)^\alpha]_{D.B.} = \frac{\pi}{2p^+} (\theta^2)^\alpha,\end{aligned}\tag{3.17}$$

where x_0^- is the zero mode of x_0^- which satisfies $[p^+, x_0^-] = \pi$. All remaining brackets are zero. We now substitute the mode expansions into these relations to determine the Poisson-Dirac brackets for the oscillators; this involves decomposing the delta functions in terms of

functions which are orthogonal over the domain $[0, \pi]$. We give details of the calculation of oscillator commutation relations in appendix D.

Canonical quantization proceeds by replacing $\{.,.\}$ in the Poisson-Dirac brackets by $-i[.,.]$. The (anti)commutators are found to be

$$\begin{aligned} [a_n^I, a_l^J] &= \text{sgn}(n)\delta_{n+l}\delta^{IJ}, & [\bar{a}_0^r, a_0^s] &= \delta^{rs}, \\ \{\theta_0^\alpha, \theta_0^\beta\} &= \frac{1}{8p^+}(\gamma^+)^{\alpha\beta} & \{\theta_n^\alpha, \theta_m^\beta\} &= \frac{1}{8p^+}(\gamma^+)^{\alpha\beta}\delta_{n+m,0}, \end{aligned} \quad (3.18)$$

with

$$a_0^r = \frac{1}{\sqrt{2m}}(p_0^r + imx_0^r), \quad \bar{a}_0^r = \frac{1}{\sqrt{2m}}(p_0^r - imx_0^r), \quad a_n^I = \sqrt{\frac{1}{|\omega_n|}}\alpha_n^I. \quad (3.19)$$

Notice that the coefficient of the fermionic anti-commutators differs by a factor of two from the closed string result (C.5). This is a factor which is well-known in the flat space limit. For the two types of generalized Dirichlet boundary conditions we consider, (3.8) and (2.35), the zero modes in the Dirichlet bosonic coordinates are found to commute with themselves and all of the other oscillators. Thus $x_0^{r'}$ and the radii for the symmetry related branes, $a^{r'}$ and $b^{r'}$, are effectively c-numbers.

3.1.2 D_+ branes

In this case we have the following possibilities:

$$\begin{aligned} D9 &: (+, -, 4, 4) \\ D7 &: (+, -, 3, 3) \\ D5 &: (+, -, 0, 4) \quad (+, -, 2, 2) \quad (+, -, 4, 0) \\ D3 &: (+, -, 1, 1) \\ D1 &: (+, -, 0, 0). \end{aligned} \quad (3.20)$$

We should add the caveat here that $(+, -, 0, 4)$ and $(+, -, 4, 0)$ constant embeddings do not solve the DBI field equations without the introduction of worldvolume flux [11]. Nevertheless, these branes appear to be on equal footing with the other ones as far as the analysis in this paper is concerned.

For branes in (3.20) the mode expansion of the bosons is exactly as for the D_- branes and for the fermions we find that⁶

$$\theta^1 = \theta_0^+ e^{m\sigma} + \theta_0^- e^{-m\sigma} + \sum_{n \neq 0} c_n \left(id_n \Pi \theta_n \phi_n + \tilde{\theta}_n \tilde{\phi}_n \right); \quad (3.21)$$

⁶Note that possible terms in the mode expansion of the form $\theta_k e^{i\omega_k \tau \pm k\sigma}$, with $k^2 \neq m^2$ are excluded by the boundary conditions.

$$\theta^2 = \Pi\theta_0^+ e^{m\sigma} - \Pi\theta_0^- e^{-m\sigma} + \sum_{n \neq 0} c_n \left(-id_n \Pi \tilde{\theta}_n \tilde{\phi}_n + \theta_n \phi_n \right),$$

Notice that the zero mode part can be rewritten in the same form as the non-zero modes, but now instead of ϕ_0 one should use ϕ_{im} and ϕ_{-im} (using the notation introduced in (3.2)).

The boundary condition enforces

$$\begin{aligned} \theta_0^\pm &= \pm \Omega \Pi \theta_0^\pm; \\ \tilde{\theta}_n &= c_n^2 (\Omega(1 - d_n^2) - 2id_n \Pi) \theta_n. \end{aligned} \tag{3.22}$$

Reality of the fermions is enforced by

$$\theta_{-n} = (\theta_n)^\dagger, \tag{3.23}$$

with θ_0^\pm real.

Canonical quantization proceeds as before by introducing canonical momenta; details of the analysis are given in the appendix. The (anti)commutators are given by

$$\begin{aligned} [a_n^I, a_l^J] &= \text{sgn}(n) \delta_{n+l,0} \delta^{IJ}, \quad [\bar{a}_0^r, a_0^s] = \delta^{rs}; \\ \{\theta_n^\alpha, \theta_m^\beta\} &= \frac{1}{8p^+} (\gamma^+)^{\alpha\beta} \delta_{n+m,0}; \\ \{\mathcal{P}_+ \theta_0^+, \mathcal{P}_+ \theta_0^+\} &= \frac{\pi m}{4p^+(e^{2\pi m} - 1)} \mathcal{P}_+ \gamma^+; \\ \{\mathcal{P}_- \theta_0^-, \mathcal{P}_- \theta_0^-\} &= \frac{\pi m e^{2\pi m}}{4p^+(e^{2\pi m} - 1)} \mathcal{P}_- \gamma^+. \end{aligned} \tag{3.24}$$

We include in these brackets the projections

$$\mathcal{P}_\pm = \frac{1}{2}(1 \pm \Omega \Pi), \tag{3.25}$$

which enforce the constraints on the zero modes. Note that in the $m \rightarrow 0$ limit, the fermion zero mode terms reproduce those in the previous section, as they must. Again the zero modes in the Dirichlet coordinates do not have conjugate momenta and commute with all of the oscillators.

4 Worldsheet symmetries and conserved charges

In this section we discuss the construction of symmetry generating charges from conserved currents in the presence of worldsheet boundaries. In a flat target space, one often obtains these using a proper doubling of the interval so that classical open string solutions satisfy periodic boundary conditions for $[0, 2\pi]$; one then substitutes the mode expansions into the expressions for the closed string supercharges. Here it is not obvious how to apply

such a doubling trick since the mode expansions involve non-periodic functions (see, for instance, (3.21)). Instead, we will consider the symmetries of the worldsheet action with given boundary conditions from first principles.

In sigma models the isometries of the target spacetime result in conserved worldsheet currents. However, these are not the only global symmetries. In our case the light-cone action is quadratic in the fields and this implies that there are an infinite number of global worldsheet symmetries. Particular cases of these symmetries are some of the target space kinematical symmetries (to be discussed below).

If the closed string action is invariant under a given set of symmetry transformations, the open string action will only be invariant under a subset of these transformations which is compatible with the (given) boundary conditions on the worldsheet fields. A novel feature for our open strings is that in some cases it is a combination of target space and worldsheet symmetries that is compatible with the chosen boundary conditions.

4.1 Closed strings

To obtain the symmetry currents we use the standard Noether method based on localisation of the parameters of global transformations. It is useful to first recall the symmetries of the action (2.19) when the string worldsheet is closed. Suppose that ϵ is a parameter of a global transformation that leaves the action invariant (up to total derivative terms which vanish when the worldsheet is closed); then when ϵ becomes local the variation of the action takes the form

$$\delta S = T \int d^2\sigma \mathcal{G}^a \partial_a \epsilon, \quad (4.1)$$

where \mathcal{G}^a is the corresponding symmetry current. As usual this implies that there is a conserved symmetry current, $\partial_a \mathcal{G}^a = 0$, with a corresponding conserved charge given by

$$G = T \int_0^{2\pi} d\sigma \mathcal{G}^\tau. \quad (4.2)$$

We take all generators to be Hermitian. We normalize the currents such that the Poisson bracket of a bosonic generator with a field Φ yields the corresponding symmetry variation, $\delta\Phi$, but the Poisson bracket of a fermionic generator yields, $-i\delta\Phi$. The factor of i is required by the reality conditions; alternatively one can consider anti-Hermitian fermionic generators.

The isometries of the target space give rise to global worldsheet symmetries. The derivation of the transformation rules and of the corresponding Noether currents and charges has been given by Metsaev in [1]. What we would like to point out here is that some of these symmetries admit an infinite extension.

Let us consider the transformation

$$\delta x^I = \epsilon^I \phi(\tau, \sigma), \quad p^+ \delta x^- = -x^I \epsilon^I \partial_\tau \phi(\tau, \sigma). \quad (4.3)$$

The Lagrangian in (2.19) is invariant (up to spatial total derivatives) under this symmetry provided $\phi(\tau, \sigma)$ satisfies

$$(\partial_\tau^2 - \partial_\sigma^2 + m^2)\phi = 0. \quad (4.4)$$

The associated conserved currents are

$$\begin{aligned} P_\phi^{I\tau} &= \partial_\tau x^I \phi - x^I \partial_\tau \phi; \\ P_\phi^{I\sigma} &= x^I \partial_\sigma \phi - \partial_\sigma x^I \phi. \end{aligned} \quad (4.5)$$

For the closed string, only those ϕ which respect the periodicity of the worldsheet give rise to conserved charges. The complete set of solutions of (4.4) subject to periodic boundary conditions are given by ϕ_n and $\tilde{\phi}_n$, introduced in (3.2), with n an integer. It follows that there is a (countably) infinite number of symmetries. For each n there are two (complex) conserved charges, the first being

$$P_n^{1I} = T \int d\sigma (\partial_\tau x^I \tilde{\phi}_{-n} - x^I \partial_\tau \tilde{\phi}_{-n}), \quad (4.6)$$

and the other P_n^{2I} , which is the same as P_n^{1I} but with $\tilde{\phi}_{-n}$ replaced by ϕ_{-n} . Notice that complex conjugation relates $P_n^{\mathcal{I}I}$, $\mathcal{I} = 1, 2$ to $P_{-n}^{\mathcal{I}I}$.

The $n = 0$ case is special in that we recover symmetries associated with the isometries of the target space. Using $\phi = \cos m\tau$, the transformation (4.3) reduces to the translation isometry given in (2.28), and using $\phi = \mu^{-1} \sin m\tau$ we obtain the rotational isometry given in (2.29). In terms of currents

$$P^I = \text{Re} P_0^I, \quad J^{+I} = \text{Im} P_0^I. \quad (4.7)$$

Notice that only in the $n = 0$ case does the function f depend only on worldsheet time, and so by using the lightcone relation, $x^+ = p^+ \tau$, the symmetries can be expressed in terms of target space coordinates. All the other symmetries are stringy extensions.

There is a corresponding story involving the fermions: a shift in the fermions by a parameter that satisfies the fermionic field equation is a symmetry of the action (2.19). There is again a countably infinite number of such transformations,

$$\begin{aligned} \delta\theta^1 &= -c_n (i d_n \Pi \epsilon_n^2 \phi_n + \epsilon_n^1 \tilde{\phi}_n); \\ \delta\theta^2 &= -c_n (-i \Pi d_n \epsilon_n^1 \tilde{\phi}_n + \epsilon_n^2 \phi_n); \\ \delta x^- &= i(\theta^1 \bar{\gamma}^- \delta\theta^1 + \theta^2 \bar{\gamma}^- \delta\theta^2), \end{aligned} \quad (4.8)$$

where $md_n = (\omega_n - n)$, $c_n = (1 + d_n^2)^{-1/2}$, the constant spinors ϵ_n satisfy $\bar{\gamma}^+ \epsilon_n = 0$, $n > 0$ is integral ($n = 0$ is considered below). The current associated with the ϵ_n^1 symmetry is

$$\begin{aligned} Q_{-n}^{1\tau} &= 2p^+ \bar{\gamma}^- (\theta^1 - id_n \Pi \theta^2) c_n \tilde{\phi}_n; \\ Q_{-n}^{1\sigma} &= 2p^+ \bar{\gamma}^- (\theta^1 + id_n \Pi \theta^2) c_n \tilde{\phi}_n, \end{aligned} \quad (4.9)$$

and that associated with ϵ_n^2 symmetry is

$$\begin{aligned} Q_{-n}^{2\tau} &= 2p^+ \bar{\gamma}^- (\theta^2 + id_n \Pi \theta^1) c_n \phi_n; \\ Q_{-n}^{2\sigma} &= -2p^+ \bar{\gamma}^- (\theta^2 - id_n \Pi \theta^1) c_n \phi_n. \end{aligned} \quad (4.10)$$

Both currents are conserved when the fermions are on-shell.

The $n = 0$ case is special. In this case the transformations are given by

$$\begin{aligned} \delta\theta^1 &= \cos m\tau \epsilon^2 - \sin m\tau \Pi \epsilon^1; \\ \delta\theta^2 &= -(\cos m\tau \epsilon^1 + \sin m\tau \Pi \epsilon^2); \\ \delta x^- &= i(\theta^1 \bar{\gamma}^- \delta\theta^1 + \theta^2 \bar{\gamma}^- \delta\theta^2). \end{aligned} \quad (4.11)$$

Since these only depend on τ they can be written entirely in terms of target space coordinates. These transformations are precisely the kinematical supersymmetry transformations obtained in [1]. The corresponding Noether currents are

$$\begin{aligned} Q^{+1\tau} &= 2p^+ \bar{\gamma}^- (\cos \mu x^+ \theta^2 + \sin \mu x^+ \Pi \theta^1); \\ Q^{+1\sigma} &= 2p^+ \bar{\gamma}^- (-\cos \mu x^+ \theta^2 + \sin \mu x^+ \Pi \theta^1); \\ Q^{+2\tau} &= -2p^+ \bar{\gamma}^- (\cos \mu x^+ \theta^1 - \sin \mu x^+ \Pi \theta^2); \\ Q^{+2\sigma} &= -2p^+ \bar{\gamma}^- (\cos \mu x^+ \theta^1 + \sin \mu x^+ \Pi \theta^2); \end{aligned} \quad (4.12)$$

These expressions agree with the ones given in [1].

The action (2.19) is also invariant under the (dynamical) supersymmetry transformations

$$\begin{aligned} \delta x^I &= 2i(\theta^1 \bar{\gamma}^I \epsilon^1 + \theta^2 \bar{\gamma}^I \epsilon^2); \\ p^+ \delta\theta^1 &= \frac{1}{2}(\gamma^{+I} \partial_- x^I \epsilon^1 + m \Pi \gamma^{+I} x^I \epsilon^2); \\ p^+ \delta\theta^2 &= \frac{1}{2}(\gamma^{+I} \partial_+ x^I \epsilon^2 - m \Pi \gamma^{+I} x^I \epsilon^1), \end{aligned} \quad (4.13)$$

which were also given in [10]. Just as in the flat space transformations ($m = 0$), these transformations involve spinors ϵ which do not satisfy the fermionic lightcone gauge condition $\bar{\gamma}^+ \epsilon = 0$. The ϵ transformations effectively have to be combined with a kappa symmetry

transformation in order to preserve $\bar{\gamma}^+\theta = 0$; this is the origin of the γ^+ factors in the transformations and of the simultaneous x^I transformations. The corresponding currents are given by

$$\begin{aligned}
Q^{-1\tau} &= 2(\partial_- x^I \bar{\gamma}^I \theta^1 - m x^I \bar{\gamma}^I \Pi \theta^2); \\
Q^{-1\sigma} &= 2(\partial_- x^I \bar{\gamma}^I \theta^1 + m x^I \bar{\gamma}^I \Pi \theta^2); \\
Q^{-2\tau} &= 2(\partial_+ x^I \bar{\gamma}^I \theta^2 + m x^I \bar{\gamma}^I \Pi \theta^1); \\
Q^{-2\sigma} &= 2(-\partial_+ x^I \bar{\gamma}^I \theta^2 + m x^I \bar{\gamma}^I \Pi \theta^1).
\end{aligned} \tag{4.14}$$

These expressions agree with the ones given in [1].

The closed string (super)charges generate the superalgebra of the plane wave background, which is given in appendix B. The realization of this algebra in terms of closed string modes was worked out in [2]. As we have seen the kinematical charges, P^I, J^{+I} and Q^{+1}, Q^{+2} , are members of an infinite family of symmetries and thus one may extend the superalgebra to include these charges as well. The charges $P_m^{\mathcal{I}I}, Q_n^{\mathcal{I}I}, \mathcal{I} = 1, 2$, when evaluated on-shell using the closed string mode expansions reviewed in appendix C are given by

$$\begin{aligned}
P_n^{1I} &= 2\alpha_n^{1I}, & P_n^{2I} &= 2\alpha_n^{2I}, \\
Q_n^1 &= 2p^+ \bar{\gamma}^- \theta_n^1, & Q_n^2 &= 2p^+ \bar{\gamma}^- \theta_n^2.
\end{aligned} \tag{4.15}$$

Using these one can compute the extension of the superalgebra. This result will be given in [6].

We should comment that although we have identified all conserved charges realised linearly in the oscillators there are also additional conserved charges realised non-linearly in the oscillators. As can be seen from formulas in appendix C, some of the target space symmetry generators are realized in terms of oscillator bilinears. There are, however, additional symmetries whose on-shell charge is bilinear in the oscillators. An example is the symmetry

$$\begin{aligned}
\delta\theta^1 &= \epsilon^{IJKL} \gamma^{IJKL} \partial_\sigma \theta^1; \\
\delta\theta^2 &= \eta \epsilon^{IJKL} \gamma^{IJKL} \partial_\sigma \theta^2,
\end{aligned} \tag{4.16}$$

where ϵ^{IJKL} is the parameter of the transformation antisymmetric in all indices (note that there is no summation over I, J, K, L in (4.16)). η is equal to plus one for $[\Pi, \gamma^{-IJKL}] = 0$ and minus one for $\{\Pi, \gamma^{-IJKL}\} = 0$. The associated tensorial charge is

$$J^{IJKL} \sim \int d\sigma (\partial_\sigma \theta^1 \gamma^{-IJKL} \theta^1 + \eta \partial_\sigma \theta^2 \gamma^{-IJKL} \theta^2), \tag{4.17}$$

and on-shell is equal to

$$J^{IJKL} \sim \sum_n n(\theta_{-n}^1 \gamma^{-IJKL} \theta_n^1 - \eta \theta_{-n}^2 \gamma^{-IJKL} \theta_n^2). \quad (4.18)$$

There are also other closely related tensorial charges. We will discuss all of them as well as the corresponding extension of the closed string superalgebra together with the related open string charges in [6].

4.2 Open strings

Now we consider symmetries of the open string action, with given boundary conditions on the worldsheet fields. Only those symmetries of the closed string action which are compatible with these boundary conditions will leave the open string action invariant. Suppose we consider an infinitesimal global symmetry transformation of the form

$$\delta z^m = H^m(z^n) \epsilon. \quad (4.19)$$

Only when this preserves the boundary conditions on all the fields z^m does this leave the action (including boundary terms) invariant. To take a specific example, suppose $x^{r'}$ satisfies a Dirichlet boundary condition, $\partial_\tau x^{r'}| = 0$. Then a global $P^{r'}$ transformation manifestly does not preserve this condition. In fact, if we evaluate the variation of the action under such a global transformation (using (2.6) and so on), we find that there is a non-vanishing boundary term

$$\delta S = -\frac{1}{\pi} \int d\tau \partial_\sigma x^{r'} \cos(\mu x^+) \epsilon^{r'}, \quad (4.20)$$

explicitly demonstrating that the action is not preserved.

Symmetries of the open string action will thus only be those which preserve the boundary conditions and for which, by construction, the variation of the action under a local transformation is

$$\delta S = \frac{1}{\pi} \int d^2 \sigma \mathcal{G}^a \partial_a \epsilon, \quad (4.21)$$

where the \mathcal{G}^a corresponding to each preserved symmetry of the action is of the same form as for the closed string. Partially integrating this expression, we get

$$\delta S = -\frac{1}{\pi} \int d^2 \sigma \partial_a G^a \epsilon + \frac{1}{\pi} \int d\tau [G^\sigma \epsilon]_{\sigma=0}^{\sigma=\pi}. \quad (4.22)$$

On shell the bulk term vanishes by virtue of the equations of motion. If this is to be a good symmetry given the boundary conditions the boundary term must also vanish for all ϵ . This requires that

$$[G^\sigma]_{\sigma=\pi} = [G^\sigma]_{\sigma=0} = 0, \quad (4.23)$$

which will hold by construction if the variational problem has been set up correctly (so all boundary terms vanish) and if the conserved symmetries have been correctly identified. Thus it is a consistency check once we have identified the preserved symmetries for specific boundary conditions that this term does indeed vanish. The conserved charges are then given by

$$G = \frac{1}{\pi} \int d\sigma G^\tau. \quad (4.24)$$

These are conserved since

$$\pi \partial_\tau G = \int d\sigma \partial_\tau G^\tau = - \int d\sigma \partial_\sigma G^\sigma = -[G^\sigma]_{\sigma=0}^{\sigma=\pi} = 0, \quad (4.25)$$

where the last equality holds by construction. Let us now discuss in detail which are the preserved charges for each of the boundary conditions given in the previous section.

4.2.1 D-branes

In order to respect the boundary conditions chosen for the D-brane the symmetry transformations should be such that $\delta x^{r'}| = 0$ and $\delta\theta^1| = \Omega\delta\theta^2|$ at the boundary; the other variations $\delta x^+|$, $\delta x^-|$ and $\delta x^r|$ are arbitrary. Looking at the symmetry transformations of the closed string action in (2.28), (2.30), (4.11) and (4.13), it is already apparent that some of these variations are not compatible with these conditions. As we have discussed, under a global transformation corresponding to $P^{r'}$, the transformation does not preserve $\delta x^{r'}| = 0$ on the boundary, and hence does not preserve the action (including boundary terms). Thus $P^{r'}$ cannot be a conserved charge of the D-brane. This is of course apparent since the D-brane explicitly breaks the translational symmetry.

The other translational charges P^+ , P^- and P^r are compatible with the boundary conditions and are conserved charges of the D-brane. As mentioned above it is a nice consistency check to explicitly verify that (4.23) holds for these charges. Take for example P^r : then using the explicit forms for the currents given in the appendix

$$\mathcal{P}^{r\sigma} = -(\cos(\mu x^+) \partial_\sigma x^r + \mu \sin(\mu x^+) x^r \partial_\sigma x^+). \quad (4.26)$$

This vanishes on the boundary by virtue of the Neumann conditions on both x^+ and x^r . In what follows we have explicitly checked that each charge which we identify as being conserved satisfies $G^\sigma(0) = G^\sigma(\pi) = 0$.

Similarly to P^I , some of the rotational J^{+I} symmetries are preserved by the boundary conditions, i.e. the J^{+r} ones, and some are broken, i.e. the $J^{+r'}$ ones. In the closed string case we showed that P^I and J^{+I} were the $n = 0$ members of an infinite number of

symmetries. The same holds for open strings as well. The argument for the symmetries is the same as that given in the closed string case, i.e. a shift in the coordinates x^I by a parameter ϵ^I that satisfies the x^I field equations is a symmetry of the worldsheet action provided the boundary terms vanish. Consider first the case of transformations that involve a shift of the Neumann coordinates,

$$\delta x^r = \epsilon^r f_n, \quad p^+ \delta x^- = -x^r \epsilon^r \partial_\tau f_n \quad (4.27)$$

where $\sqrt{|\omega_n|} f_n = e^{-i\omega_n \tau} \cos n\sigma$ (the prefactors are introduced for later convenience). This transformation leaves the action invariant including boundary terms. The corresponding conserved charge is given by (4.6) but with $\tilde{\phi}_{-n}$ replaced by f_n .

Consider now the similar transformation involving the Dirichlet direction, i.e. (4.27) but with x^r replaced by $x^{r'}$ and f_n replaced by $\sqrt{|\omega_n|} \tilde{f}_n = -ie^{-i\omega_n \tau} \sin n\sigma$. This transformation, however, leads to a non-zero boundary term if $x_0^{r'}(\sigma, \tau)$ is non-zero at the string endpoints, where $x_0^{r'}(\sigma, \tau)$ is defined in (3.7). So unless the brane is located at the origin of transverse space, these transformations do not lead to a conserved charge. However, one can modify the transformation rules such that there is conserved charge even when the brane is located at arbitrary constant position. Indeed, consider the transformations

$$\delta x^{r'} = \epsilon^{r'} \tilde{f}_n, \quad p^+ \delta x^- = -\epsilon^{r'} (x^{r'} \partial_\tau \tilde{f}_n + P_{(0)\tilde{f}_n}^{r'\tau}) \quad (4.28)$$

where $P_{(0)\tilde{f}_n}^{r'\tau}$ is given by (4.5) but with $x^{r'}$ replaced by $x_0^{r'}(\sigma, \tau)$. One may verify that these transformations leave the action invariant, including boundary terms, even when the brane is located at a non-zero transverse position. The corresponding conserved charge is then given by (4.6) but with $x^{r'}$ replaced by $(x^{r'} - x_0^{r'})$ and $\tilde{\phi}_{-n}$ by \tilde{f}_n . It follows that when the charge is evaluated in terms of modes it takes the same form as for the conserved charge associated with the brane located at the origin of transverse space.

This is a phenomenon that we will see repeatedly in what follows: certain symmetries appear to be broken when the brane is located away from the origin of transverse space, and the breaking terms solely depend on quantities fully determined by the boundary conditions (in the case above the breaking terms depend only on the boundary value of the Dirichlet coordinates). In these cases, however, one can find modified transformation rules (involving worldsheet symmetries) that lead to a conserved charge. The latter when evaluated in terms of modes is exactly equal to the conserved charge of the brane located at the origin.

The $n = 0$ case is special, as for the closed string case. In this case the real and imaginary part of the transformations in the Neumann direction give the P^r and J^{+r} transformations.

In addition, the Dirichlet directions do not yield any conserved charge when $n = 0$, in agreement with the fact that the D-branes break the $P^{r'}$ and $J^{+r'}$ symmetries.

We now move to the rotational symmetries J^{IJ} . Consider first the J^{rs} case. The fermionic variations are compatible with the boundary conditions since

$$|\delta\theta^1| = -\frac{1}{4}\epsilon^{rs}\gamma^{rs}\theta^1| = -\frac{1}{4}\epsilon^{rs}\gamma^{rs}\Omega\theta^2| = \Omega\delta\theta^2|, \quad (4.29)$$

where in the second equality we use the relation between fermions on the boundary and in the final equality we use $[\Omega, \gamma^{rs}] = 0$. This follows from the explicit form of Ω given in (3.10). Since δx^r is arbitrary on the boundary the bosonic variations are also compatible with the boundary conditions and J^{rs} is a conserved charge. However, the $J^{rr'}$ symmetry is broken since the bosonic variations mix Dirichlet and Neumann coordinates (and furthermore the fermionic variations do not preserve the fermion boundary conditions for reasons akin to above).

The $J^{r's'}$ symmetry is another instance where we will use the worldsheet symmetries we just discussed in order to restore a seemingly broken symmetry. The fermionic variations do preserve the fermion boundary conditions (as in (4.29), Ω commutes with $\gamma^{r's'}$) and there is actually a conserved current associated with these transformations. The bosonic variations, however, will only preserve the boundary condition if the brane is located at the origin. That is, $\delta x^{r'}| = 0 = \epsilon^{r's'}x^{s'}|$ can be satisfied on the boundary only if $x_0^{s'} = 0$. So even though there is a conserved current for any value of transverse position, $x_0^{r'}$, the transformation rules change that value. Thus, the $J^{r's'}$ symmetry is broken by the position of the brane unless the brane is located at the origin of transverse space.

We will now see that even at arbitrary transverse position one can obtain a good symmetry by combining the $J^{r's'}$ symmetry with a certain worldsheet symmetry. Consider the worldsheet symmetry given by $\delta x^{r'} = \epsilon^{r's'}x_0^{s'}(\sigma, \tau)$, where $x_0^{r'}(\sigma, \tau)$ is defined in (3.7). By construction, $x_0^{s'}(\sigma, \tau)$ satisfies (3.1), so this transformation leaves the action invariant⁷. We now combine the $J^{r's'}$ symmetry with this worldsheet symmetry by considering the transformations,

$$\delta x^{r'} = \epsilon^{r's'}(x^{s'} - x_0^{s'}(\sigma, \tau)); \quad \delta\theta = -\frac{1}{4}\epsilon^{r's'}\gamma^{r's'}\theta. \quad (4.30)$$

These transformations leave invariant the action, including boundary terms, and respect both fermionic and bosonic boundary conditions. It follows that this is a good symmetry.

⁷In general, the fact that the parameter satisfies the free field equation only implies that the Lagrangian is invariant up to total derivatives. The cancellation of the total derivative terms depends on the boundary conditions imposed on the fields. The worldsheet symmetry $\delta x^{r'}$ is actually a symmetry including boundary terms. Notice that the $J^{r's'}$ transformations also leave the action invariant including boundary terms.

The associated conserved current is just $J'^{s'}$, with $x^{r'}$ replaced by $(x^{r'} - x_0^{r'}(\sigma, \tau))$ and similarly for $x^{s'}$. We shall show in [6] that the conserved charge when realised in terms of the oscillators is precisely that for the branes at the origin.

In the above we have used worldsheet symmetries to subtract from symmetry currents terms which do not preserve the boundary conditions. However this trick can be applied much more generally: we may also combine good worldsheet symmetries with good target space symmetries (i.e. target space symmetries that are compatible with the boundary conditions). In this way one can obtain Noether charges that are equal to the original charge associated with a superisometry but with certain terms subtracted.

A specific example that will arise in the open string algebras in [6] is the following. Suppose we first isolate the part of the (conserved) Hamiltonian symmetry current which involves only the Dirichlet zero modes and which is conserved by itself; call this ΔH^a . Then under a symmetry transformation

$$\delta x^+ = -\epsilon^+; \quad p^+ \delta x^- = -\Delta H^\tau \epsilon^+, \quad (4.31)$$

the action is preserved and the corresponding Noether charge is

$$\hat{H} = H - \frac{1}{\pi} \int_0^\pi d\sigma \Delta H^\tau \equiv H - \Delta H, \quad (4.32)$$

which when evaluated in terms of modes is precisely the original Hamiltonian with the Dirichlet zero mode (c-number) contribution subtracted.

We next consider the transformations associated with the supercharges. As in previous sections, it will be convenient to discuss separately the D_- and D_+ branes.

4.2.2 D_- branes

First consider the kinematical supercharges Q^+ . The corresponding transformation rules are given in (4.11). Under global transformations corresponding to Q^+ the action is invariant up to the total derivative term

$$\delta S = \frac{1}{\pi} \int d\tau i p^+ (\delta\theta^1 \bar{\gamma}^- \theta^1 - \delta\theta^2 \bar{\gamma}^- \theta^2). \quad (4.33)$$

Thus provided that the variations are compatible with the fermion boundary conditions, that is, $\delta\theta^1 = \Omega \delta\theta^2$ on the boundary, the action remains invariant. We find that the fermion boundary conditions are compatible with (4.11) only when

$$\epsilon^2 = -\Omega \epsilon^1, \quad \Pi \Omega \Pi \Omega = -1. \quad (4.34)$$

Note that the second condition would not have to be imposed in flat space ($\mu = 0$) but is necessary here to enforce $\delta\theta^1| = \Omega\delta\theta^2|$ at all x^+ . Thus the boundary conditions can preserve at most one half of the Q^+ supercharges, the combinations

$$q^+ = \frac{1}{2}(-Q^{+2} + \bar{\Omega}Q^{+1}). \quad (4.35)$$

(The normalization of q^+ is for later convenience). We conclude that the D_- branes preserve one half of the kinematical supersymmetries of the closed string. Let us note at this stage that these symmetries are preserved regardless of the Dirichlet boundary conditions: even D_- branes with general time dependent boundary conditions $x^{r'}(x^+)$ preserve these same charges.

The Q^+ symmetry preserved by the D_- branes is just the $n = 0$ member of a family of infinite number of symmetries. The additional symmetry transformations are given by (4.8) but with parameters satisfying

$$\epsilon_n^1 = \Omega\epsilon_n^2. \quad (4.36)$$

The corresponding Noether charges can be obtained from (4.9) and (4.10) by taking combinations $Q_n = \frac{1}{2}(Q_n^1 + \bar{\Omega}Q_n^2)$.

Let us now consider the dynamical supercharge (Q^-) transformations. Once again the action is only invariant up to a total derivative term and the boundary conditions are not in general consistent with (4.13). Let us consider first the δx^I transformations: if these are to preserve the Dirichlet boundary conditions then we must consider only such transformations satisfying

$$\delta x^{r'}| = 0 = \theta^2|(\Omega^t\bar{\gamma}^{r'}\epsilon^1 + \bar{\gamma}^{r'}\epsilon^2) \quad \rightarrow \quad \epsilon^1 = \Omega\epsilon^2, \quad (4.37)$$

where we have used $\{\Omega, \gamma^{r'}\} = 0$ (which follows from (3.10)). Thus the D-brane can preserve at most one half of the Q^- supersymmetries. Now under transformations (4.13) the action is invariant up to the boundary term

$$- \int d\tau (\partial_+ x^I \theta^1 \bar{\gamma}^I \epsilon^1 - \partial_- x^I \theta^2 \bar{\gamma}^I \epsilon^2 + m x^I (\theta^1 \bar{\Pi} \bar{\gamma}^I \epsilon^2 + \theta^2 \bar{\Pi} \bar{\gamma}^I \epsilon^1)). \quad (4.38)$$

This term does not in general vanish even given the relation (4.37), along with the conditions $\theta^1 = \Omega\theta^2$, $[\gamma^r, \Omega] = 0$, $\partial_\sigma x^r = 0$, and $\{\gamma^{r'}, \Omega\} = 0$.

This boundary term vanishes only when the fermionic variations in (4.13) are consistent with the fermion boundary conditions. So one should substitute $\epsilon^1 = \Omega\epsilon^2$ into the fermionic

variations and demand that they preserve $\theta^1 = \Omega\theta^2$ on the boundary. Now

$$\begin{aligned}
p^+(\delta\theta^1 - \Omega\delta\theta^2)| &= - \left(\Omega \sum_{r=1}^{p-1} \partial_\sigma x^r \gamma^{+r} + \Omega \sum_{r'=p}^8 \partial_\tau x^{r'} \gamma^{+r'} \right) \epsilon^2 \Big| \\
&\quad + \frac{1}{2} m \left(\sum_{r=1}^{p-1} (\Pi + \Omega\Pi\Omega) \gamma^{+r} x^r + \sum_{r'=p}^8 (\Pi - \Omega\Pi\Omega) \gamma^{+r'} x^{r'} \right) \epsilon^2 \Big|.
\end{aligned} \tag{4.39}$$

This must vanish for any of the Q^- supersymmetry to be preserved. To evaluate this expression we have used only $[\Omega, \gamma^r] = \{\Omega, \gamma^{r'}\} = 0$; the first two terms manifestly drop out when x^r is pure Neumann and $x^{r'}$ is pure Dirichlet, but we will use the more general expression later for mixed boundary conditions.

For our case (i.e. $\Pi\Omega\Pi\Omega = -1$), (4.39) reduces to

$$\sum_{r'=p}^8 \left(-\partial_\tau x_0^{r'}(\sigma, \tau) \Omega + m x_0^{r'}(\sigma, \tau) \Pi \right) \gamma^{+r'} \epsilon^2 \Big|. \tag{4.40}$$

Note that the Neumann directions drop out of the condition. (4.40) can only be satisfied if the transverse positions are all zero. Thus we see that when the D_- brane is located at the origin, (4.40) vanishes and half of the dynamical supersymmetries are preserved, namely the combinations

$$q^- = \frac{1}{2}(Q^{-1} + \bar{\Omega}Q^{-2}), \tag{4.41}$$

but when the brane is located at some other position all Q^- supersymmetries are broken. This is exactly what we found in the probe analysis [11]. We shall now show, however, that there are eight fermionic symmetries, which are combinations of worldsheet and Q^- supersymmetries, that are preserved by the brane even when it is located away from the origin. So in effect the brane away from the origin is as supersymmetric as the brane at the origin.

Recall that under the fermionic transformation,

$$\begin{aligned}
p^+ \delta\theta^1 &= id_k \Pi \epsilon_k^2 \phi_k + \epsilon_k^1 \tilde{\phi}_k; \\
p^+ \delta\theta^2 &= -i \Pi d_k \epsilon_k^1 \tilde{\phi}_k + \epsilon_k^2 \phi_k, \\
\delta x^- &= i(\theta^1 \bar{\gamma}^- \delta\theta^1 + \theta^2 \bar{\gamma}^- \delta\theta^2),
\end{aligned} \tag{4.42}$$

where k is a complex parameter, the worldsheet Lagrangian transforms as a total σ -derivative (since fermionic variations satisfy the fermionic field equations). When k is an integer, these are just the closed string worldsheet symmetries discussed in (4.8) and when the relation (4.36) is imposed these become the stringy extensions of Q^+ transformations

discussed above. For general k , however, the boundary terms are non-vanishing, and (4.42) does not lead to a conserved charge.

Now, since $x_0^{r'}(\sigma, \tau)$ satisfies the bosonic field equation, it can always be expanded as

$$x_0^{r'}(\sigma, \tau) = \sum_k (c_k^{r'} \phi_k + \tilde{c}_k^{r'} \tilde{\phi}_k), \quad (4.43)$$

where the coefficients $c_k^{r'}$ and $\tilde{c}_k^{r'}$ are constants. For example, for a brane located at arbitrary transverse position, $x_0^{r'}$, $x_0^{r'}(\sigma, \tau)$ is given in (3.8) and can be rewritten as

$$x_0^{r'}(\sigma, \tau) = \frac{x_0^{r'}}{e^{m\pi} + 1} (\phi_{im} + e^{m\pi} \phi_{-im}). \quad (4.44)$$

Let us now try to find a (4.42) transformation such that combined with the Q^- transformation it yields a transformation which leaves the action invariant, including boundary terms, and so leads to a conserved charge. In other words, we want to choose ϵ_k^1 and ϵ_k^2 such that

$$0 = p^+ (\delta\theta^1 - \Omega d\theta^2) \Big| = \sum_k \left(\phi_k \left(c_k^{r'} (i\omega_k \Omega + m\Pi) \gamma^{+r'} \epsilon^2 + (id_k \Pi - \Omega) \epsilon_k^2 \right) \Big| \right. \\ \left. + \tilde{\phi}_k \left(\tilde{c}_k^{r'} (i\omega_k \Omega + m\Pi) \gamma^{+r'} \epsilon^2 + (1 - i\Omega \Pi d_k) \epsilon_k^1 \right) \Big| \right). \quad (4.45)$$

This equation has a solution provided that $d_k^2 \neq 1$ which implies $k \neq 0$:

$$\epsilon_k^1 = -\frac{1}{d_k^2 - 1} \tilde{c}_k^{r'} (i(2\omega_k - k)\Omega + (m + d_k \omega_k)\Pi) \gamma^{+r'} \epsilon^2 \\ \epsilon_k^2 = \frac{1}{d_k^2 - 1} c_k^{r'} (-ki + (m - d_k \omega_k)\Pi\Omega) \gamma^{+r'} \epsilon^2 \quad (4.46)$$

It follows that in all cases in which $x_0^{r'}(\sigma, \tau)$ has an expansion that does not involve the $k = 0$ modes, the D-brane preserves eight fermionic symmetries, namely the combination of Q^- and the special worldsheet transformations (4.42) with the parameter given by (4.46). $k = 0$ is a special case in that the worldsheet symmetry Q_0 is precisely the spacetime kinematical supersymmetry and the terms in $x_0^{r'}$ involve $\sin(m\tau)$ and $\cos(m\tau)$. This case is thus relevant for D-branes translated away from the origin using the broken translational or rotational symmetries. It is convenient to discuss these in detail separately in the next section.

In the specific case of an D_- brane located at constant position $x_0^{r'}$, $x_0^{r'}(\sigma)$ is given in (4.44) and (4.46) becomes,

$$\epsilon_{im}^2 = -\frac{1}{2} \frac{m}{e^{m\pi} + 1} (1 + \Pi\Omega) \gamma^{+r'} \epsilon^2, \quad \epsilon_{-im}^2 = \frac{1}{2} \frac{m e^{m\pi}}{e^{m\pi} + 1} (1 - \Pi\Omega) \gamma^{+r'} \epsilon^2, \quad (4.47)$$

The Noether charge associated with this symmetry is

$$\hat{Q}^- = \frac{1}{\pi} \int_0^\pi d\sigma \left(\sum_{r=1}^{p-1} (\partial_\tau x^r \bar{\gamma}^r \theta_N - \partial_\sigma x^r \bar{\gamma}^r \theta_D + m x^r \bar{\gamma}^r \Omega \Pi \theta_N) \right) \quad (4.48)$$

$$\sum_{r'=p}^8 (\partial_\tau(x^{r'} - x_0^{r'})\bar{\gamma}^{r'}\theta_D - \partial_\sigma(x^{r'} - x_0^{r'})\bar{\gamma}^{r'}\theta_N - m(x^{r'} - x_0^{r'})\bar{\gamma}^{r'}\Omega\Pi\theta_D)$$

where θ_N and θ_D are defined in appendix D. This charge when realised in terms of modes reproduces precisely q^- .

The result however holds much more generally: branes with arbitrary boundary conditions $x^{r'}(x^+)$ preserve one half of the closed string kinematical supersymmetries as well as eight additional dynamical supersymmetries which descend from a combination of closed string and worldsheet symmetries.

4.2.3 D_+ branes

We have seen in the previous section that the conditions for having unbroken kinematical supersymmetry Q^+ are (4.34). For the D_+ branes the second condition is not satisfied and all of the Q^+ supersymmetries are necessarily broken. However, this symmetry is replaced by another fermionic symmetry. As we have discussed, the Q^+ symmetry is a particular case of a worldsheet symmetry and the latter originate from the fact that the worldsheet action is quadratic in the fields. This is true for the D_+ branes as well and it turns out that these branes possess kinematical supersymmetries too.

The new kinematical supersymmetry rules for $n \neq 0$ are still given by (4.8) but the parameters now satisfy

$$\epsilon_n^1 = c_n^2(\Omega(1 - d_n^2) - 2id_n\Pi)\epsilon_n^2 \quad (4.49)$$

where $md_n = (\omega_n - n)$, $c_n = (1 + d_n^2)^{-1/2}$, $\omega_n = \text{sgn}(n)\sqrt{n^2 + m^2}$ and n is an integer. The corresponding Noether charges can be obtained from (4.9) and (4.10) by combining $(Q_n^1 + c_n^2(\bar{\Omega}(1 - d_n^2) - 2id_n\bar{\Pi})Q_n^2)$.

There are some special cases as usual. Firstly, we cannot find a solution for $n = 0$, which reflects the total breaking of the closed string symmetry Q^+ . However, there is a solution for $k = \pm im$:

$$\begin{aligned} \delta\theta^1 &= \mathcal{P}_+\epsilon^+e^{m\sigma} + \mathcal{P}_-\epsilon^-e^{-m\sigma}; \\ \delta\theta^2 &= \Pi\mathcal{P}_+\epsilon^+e^{m\sigma} - \Pi\mathcal{P}_-\epsilon^-e^{-m\sigma}; \\ p^+\delta x^- &= i(\theta^1\bar{\gamma}^-\delta\theta^1 + \theta^2\bar{\gamma}^-\delta\theta^2), \end{aligned} \quad (4.50)$$

where

$$\mathcal{P}_\pm = \frac{1}{2}(1 \pm \Omega\Pi). \quad (4.51)$$

It is these transformations that effectively replace the closed string kinematical supersymmetries Q^+ . Notice that the transformations are similar to (4.11) in that they only depend

on one worldsheet coordinate but in this case it is the σ coordinate rather the τ coordinate that features in the formulas. Thus, unlike (4.11), the transformation rules (4.50) cannot be expressed entirely in terms of target space coordinates. The corresponding Noether current is

$$\begin{aligned}\hat{Q}^{+\tau} &= \frac{2p^+}{\pi}\bar{\gamma}^- (e^{m\sigma}(\mathcal{P}_+\theta^1 + \mathcal{P}_+\Pi\theta^2) + e^{-m\sigma}(\mathcal{P}_-\theta^1 - \mathcal{P}_-\Pi\theta^2)); \\ \hat{Q}^{+\sigma} &= \frac{2p^+}{\pi}\bar{\gamma}^- (e^{m\sigma}(\mathcal{P}_+\theta^1 - \mathcal{P}_+\Pi\theta^2) + e^{-m\sigma}(\mathcal{P}_-\theta^1 + \mathcal{P}_-\Pi\theta^2)).\end{aligned}\quad (4.52)$$

As we shall show in [6], the supercharge \hat{Q}^+ is proportional to the D_+ brane fermionic zero modes exactly as the supercharge Q^+ is for the D_- branes.

We should comment here that although \hat{Q}^+ is just one of the infinite set of symmetries Q_n it does play a distinguished role. This will become apparent when we discuss the superalgebra and spectrum [6]. The superalgebra of the preserved closed string supercharges plus \hat{Q}^+ closes (the introduction of Q_n leads to an infinite extension). \hat{Q}^+ commutes with the Hamiltonian so physical states form corresponding supermultiplets. In the $\mu \rightarrow 0$ limit, it becomes precisely a target space supercharge. Although both \hat{Q}^+ and Q_n act on the spectrum, the former being realised in terms of zero modes is exceptional.

Let us now consider the dynamical supersymmetries. For D_+ branes (4.39) reduces to

$$\frac{1}{2}m \sum_{r=1}^{p-1} (\Pi + \Omega\Pi\Omega)\gamma^{+r} x^r \epsilon^2 = m \sum_{r=1}^{p-1} \Pi\gamma^{+r} x^r \epsilon^2. \quad (4.53)$$

where we consider standard Dirichlet boundary conditions, i.e. $\partial_\tau x_0^r(\sigma, \tau) = 0$. If there are no Neumann directions amongst the x^I , i.e. for a D1-brane, (4.53) vanishes and half of the Q^- are preserved, the combinations $(Q^{-1} + \bar{\Omega}Q^{-2})$, regardless of transverse location. For all other possibilities the Q^- supersymmetry is broken and the boundary term in (4.38) does not vanish.

One may hope that similar considerations as in the previous cases would allow us to find eight additional fermionic symmetries. In this case, however, the violation of the symmetry in (4.53) involves the restriction of the Neumann coordinates to the boundary which is unknown. In contrast, in all previous cases the violating terms were involving terms fully determined by the boundary conditions. We conclude that all D_+ branes but the D1 brane do not preserve any dynamical supersymmetry.

4.2.4 Symmetry-related D-branes

As discussed in section 2.2, one can act by broken spacetime generators on a static brane to obtain a new brane. For every symmetry of the static brane there is necessarily a cor-

responding symmetry of the translated brane and the algebra of the corresponding charges is the same in both cases.

Recall that for a D-brane translated using the background Killing symmetries $P^{r'}$ and $J^{+r'}$ the corresponding Dirichlet boundary condition is (2.35), which couples the x^+ and $x^{r'}$ variations on the boundary as

$$\delta x^{r'}| = (-a^{r'} \mu \sin(\mu x^+) + b^{r'} \mu \cos(\mu x^+)) \delta x^+|. \quad (4.54)$$

In this section we will use t' to denote an ordinary Dirichlet direction and reserve r' and s' to denote directions in which there are modified Dirichlet boundary conditions (2.35). Throughout the section when discussing conserved charges we assume that all constant Dirichlet positions are zero; if any are constant but non-zero we simply replace the conserved symmetry we identify by the same symmetry combined with the appropriate worldsheet symmetries as discussed in the previous section.

Given such boundary conditions, the conserved bosonic charges of the D-brane certainly include $(P^+, P^r, J^{+r}, J^{rs}, J^{t'u'})$, following the same analysis as before. The other bosonic charges are more subtle, however. The Hamiltonian is not conserved; to preserve the boundary condition, we must consider a combination of P^- , $J^{+r'}$ and $P^{r'}$ transformations. That the Hamiltonian by itself is not a good conserved charge follows from the time dependence of the boundary conditions. The action of the combined transformation

$$\hat{P}^- = P^- + \sum_{r'} (\mu b^{r'} P^{r'} - \mu^2 a^{r'} J^{+r'}), \quad (4.55)$$

is

$$\begin{aligned} \delta x^+ &= \epsilon^+; & \delta x^{r'} &= -a^{r'} \mu \sin(\mu x^+) \epsilon^+ + \mu b^{r'} \cos(\mu x^+) \epsilon^+; \\ \delta x^- &= \mu^2 \sum_{r'} (a^{r'} \cos(\mu x^+) x^{r'} + b^{r'} \sin(\mu x^+) x^{r'}) \epsilon^+, \end{aligned} \quad (4.56)$$

which manifestly preserves (4.54). The other modified charges are

$$\begin{aligned} \hat{J}^{r's'} &= J^{r's'} - a^{r'} P^{s'} - \mu b^{r'} J^{+s'} + a^{s'} P^{r'} + \mu b^{s'} J^{+r'}; \\ \hat{J}^{r't'} &= J^{r't'} - a^{r'} P^{t'} - \mu b^{r'} J^{+t'}. \end{aligned} \quad (4.57)$$

Now let us consider the supercharges. Consider first the D_- branes. The kinematical supercharges q^+ are unaffected by the time dependent boundary conditions; the same is true for the fermionic charges Q_n . The combination of dynamical supersymmetry transformations with $\epsilon^1 = \Omega \epsilon^2$ preserves the time dependent $x^{r'}$ boundary condition but following

(4.39)

$$p^+(\delta\theta^1 - \Omega\delta\theta^2)| = m\gamma^+ \sum_{r'} (\sin(m\tau)(b^{r'}\bar{\Pi}\bar{\gamma}^{r'} - a^{r'}\bar{\gamma}^{r'}\Omega)\epsilon^2 + \cos(m\tau)(b^{r'}\bar{\gamma}^{r'}\Omega + a^{r'}\bar{\Pi}\bar{\gamma}^{r'}))\epsilon^2. \quad (4.58)$$

On its own this does not preserve the fermion boundary condition but it is a good symmetry when combined with Q^+ transformations with parameters $\hat{\epsilon}^1$ and $\hat{\epsilon}^2$ such that

$$(\delta\theta^1 - \Omega\delta\theta^2)| = (\cos(m\tau) + \sin(m\tau)\Omega\Pi)(\hat{\epsilon}^2 + \Omega\hat{\epsilon}^1) \quad (4.59)$$

provided that

$$(\hat{\epsilon}^2 + \Omega\hat{\epsilon}^1) = \mu\gamma^+ \sum_{r'} (b^{r'}\bar{\Omega}\bar{\gamma}^{r'} - a^{r'}\bar{\Pi}\bar{\gamma}^{r'})\epsilon^2. \quad (4.60)$$

The corresponding Noether charge is then

$$\hat{q}^- = \frac{1}{2} \left(Q^{-1} + \bar{\Omega}Q^{-2} + \frac{1}{2}\mu \sum_{r'} (b^{r'}\bar{\gamma}^{r'+} - a^{r'}\bar{\gamma}^{r'}\Omega\Pi\gamma^+)(Q^{+2} + \bar{\Omega}Q^{+1}) \right). \quad (4.61)$$

That the symmetry involves only this specific combination of Q^+ transformations follows from the fact that kinematical transformations with $(\hat{\epsilon}^2 + \Omega\hat{\epsilon}^1) = 0$ automatically preserve the fermion boundary conditions by themselves. The charge (4.61) has eight independent components in total. It is interesting to prove this same result from a rather different perspective using the kappa symmetry projections; we include this in appendix E.

Thus the brane preserves in total one half of the closed string supersymmetries, as indeed it must, being related by symmetry to the half supersymmetric branes in the previous subsection. Under the symmetry transformation, the preserved dynamical charge is shifted by a term proportional to the kinematical charge. This is to be expected since from the spacetime superalgebra the commutator of the translation and Q^- is proportional to Q^+ . We will make this statement more precise in [6] when we discuss the embedding of the symmetry-related brane algebra into the closed string algebra.

There is one special case where these preserved dynamical charges are actually the same ones as for the static brane: this requires

$$\sum_{r'} (b^{r'}\bar{\gamma}^{r'} - a^{r'}\bar{\gamma}^{r'}\Omega\Pi) = 0. \quad (4.62)$$

If one considers a $(+, -, 2, 0)$ D3-brane, and takes $a^3 = r$, $b^4 = r$, so the brane traces out a circle in the (34) plane, this condition is satisfied. Such a brane arises from taking a specific Penrose limit of a giant graviton in $AdS_5 \times S^5$ [21] which are indeed known to be half supersymmetric rotating branes [22, 23]. Penrose limits of giant gravitons were also discussed in [11, 15, 16].

What is particularly interesting, however, is that we now see that all D_- branes still preserve one half of the closed string supersymmetries when they rotate in any transverse plane. This suggests that there are many more half supersymmetric rotating branes in AdS than were previously known. Furthermore, translated or boosted branes (by which we mean those with $b^{r'} = 0$ and $a^{r'} = 0$, respectively) may also originate from as yet unknown time dependent embeddings in AdS which are half supersymmetric. Following the same approach as here, such branes could be obtained from the half supersymmetric static embeddings given in [11] by the action of the symmetries of the AdS background which are broken by the static brane.

Now consider the D_+ branes. As for the static branes closed string kinematical supersymmetries are completely broken by the fermion boundary conditions, but the branes possess different kinematical supersymmetries instead. The preserved worldsheet fermionic symmetries Q_n are also the same as for the static branes. Moving now to the dynamical supersymmetry, we observe that for the symmetry-related D1-branes the preserved Noether charges are the combinations

$$\hat{q}^- = \frac{1}{2}(Q^{-1} + \bar{\Omega}Q^{-2}) + \frac{1}{4}\mu \sum_{r'} \left((b^{r'}\gamma^{r'+} - a^{r'}\bar{\gamma}^{r'}\Omega\Pi\gamma^+)Q^{+2} + (b^{r'}\gamma^{r'+}\bar{\Omega} + a^{r'}\bar{\gamma}^{r'+}\bar{\Pi})Q^{+1} \right). \quad (4.63)$$

Thus a D1-brane carrying angular momentum in a transverse plane still preserves one half of the dynamical supersymmetry.

This result fits in neatly with [11], where it was noted that $(+, 1, 0)$ D1-branes preserve one quarter of the closed string supersymmetries. It was then argued that these should arise from the Penrose limits of rotating D-strings in $AdS_5 \times S^5$. A distinct Penrose limit of these same rotating D-strings gives the $(+, -, 0, 0)$ branes carrying angular momentum in a single transverse plane, which we have just shown also preserve one quarter of the closed string supersymmetries (though let us emphasise that like the static branes they do possess eight additional kinematical supersymmetries as well).

5 Conclusions

We studied in this paper D-branes in the maximally supersymmetric plane wave background of IIB supergravity. We found both time-dependent and static branes. The branes fall into equivalence classes depending on whether they are related by the action of target space isometries. In particular, static branes localized at the origin are in the same equivalence class as certain time-dependent branes. The symmetries of all branes in the same equivalence

class are the same, but the relation of the conserved charges with the conserved charges of the closed string is different for different members of the equivalence class.

Let us summarize the results for the supersymmetric static Dp-branes. The D-branes fall into two categories: the D_- branes satisfying $\Pi\Omega\Pi\Omega = -1$, where Ω is related to the fermion boundary conditions (see (2.15)), and D_+ branes satisfying $\Pi\Omega\Pi\Omega = 1$. The D_- branes have worldvolume coordinates $(+, -, m, m \pm 2)$ and the D_+ are the remaining $(+, -, m, n)$ branes.

D_- branes always preserve eight kinematical Q^+ supersymmetries irrespective of location. They preserve eight dynamical supersymmetries Q^- only if the brane is located at the origin, and away from the origin Q^- is broken but we find instead eight new dynamical supercharges, \hat{Q}^- . The latter are a linear combination of Q^- with a certain fermionic worldsheet symmetry. \hat{Q}^- anticommute to the Hamiltonian much the same way as Q^- do for the branes at the origin. Actually, as we will show in [6], \hat{Q}^- evaluated on-shell is equal to Q^- on-shell (but the latter is relevant for branes at the origin and the former for branes away from the origin). Furthermore, we have seen that a far stronger result also holds: D_- branes with arbitrary time dependent Dirichlet boundary conditions of the form $x^{r'}(x^+)$ will also preserve not just half of the closed string kinematical supersymmetries but in addition eight dynamical supersymmetries, the combination of closed string with worldsheet symmetries.

In D_+ branes all of the Q^+ supersymmetries are broken, but a different set of 8 kinematical supersymmetries \hat{Q}^+ is present and one half of the Q^- are preserved if and only if there are no Neumann directions transverse to the lightcone, i.e. it is a D1-brane⁸. Thus, all D_+ branes but the D1-brane preserve 8 supercharges, and the latter preserves 16 supercharges. The number of supersymmetries that directly originate from the closed string supersymmetries agrees with the probe analysis [11] in all cases.

We have seen that the light-cone action admits an infinite number of worldsheet symmetries. These symmetries should also be present before gauge fixing the κ -symmetry. It would be interesting to understand this in detail, since this issue is closely related to the question of under what conditions a given sigma model is related to an integrable theory. For a related discussion we refer to [24].

A motivation for this work was to gain intuition about string theory on curved backgrounds with RR fluxes. One of the main messages of this paper is that even the question of which are the possible supersymmetric states of string theory on such backgrounds (su-

⁸We do not consider branes with worldvolume flux in this paper. We note that the probe analysis in [11] yielded D_+ $(+, -, 0, 4)$ and $(+, -, 4, 0)$ branes that preserve 1/2 of dynamical supersymmetry.

persymmetric branes in our case) is subtle. We have seen that tree-level open string theory yields more supersymmetric branes than the probe analysis. The additional supersymmetries were based on specific properties of the plane wave background. An important issue is whether the extra symmetries are respected by interactions. Provided that the answer is affirmative, consistency requires that the worldvolume theory of D-branes in the plane wave background exhibits more supersymmetries than the generic kappa-symmetry analysis yields (but if string interactions invalidate the extra symmetries only the quadratic part of the worldvolume theory that captures the spectrum of low energy excitations should exhibit more supersymmetries). In other words, our results indicate that the generic analysis of the supersymmetry of embeddings only yields a lower bound on the number of supersymmetries rather than the exact number.

Given that the worldsheet theory is solvable only for very specific backgrounds but the probe analysis is possible more generally, it is important to understand under which conditions one needs to amend the standard kappa-symmetry analysis. More generally it would be interesting to understand which backgrounds exhibit similar symmetry restoration mechanisms.

In this paper we have studied open strings in the light-cone gauge. It would be interesting to extend the analysis to more covariant gauges which would allow, amongst other things, the study of branes not accessible in light-cone gauge, such as the supersymmetric branes found in [11] where only one of the lightcone directions is a worldvolume coordinate. A description of the closed string in the plane wave background where all target space symmetries are manifest was given in [25] using the pure spinor formalism. Worldsheet superconformal field theories associated with the plane wave (and other RR backgrounds) were presented in [26, 27] using the $U(4)$ hybrid formalism. It would be interesting to analyze D-branes using these formalisms, particularly to understand the relation between the worldsheet symmetries we find here and worldsheet superconformal symmetries. Furthermore, quantum properties of the D-branes may be more transparent in the $U(4)$ formalism.

In [6] we continue the analysis started here by computing the corresponding superalgebras and the spectra of branes in all cases and comparing with the spectrum of small fluctuations around the corresponding D-brane embedding in the DBI. Finally let us comment that finding the corresponding boundary states and checking open/closed duality for the branes found here which are not included in the analysis of [8] is an important issue which we will also address elsewhere.

Acknowledgments

We thank Università di Roma “Tor Vergata” for warm hospitality at the initial stages of this work. KS would also like to thank the Isaac Newton Institute, the Amsterdam Summer Workshop, and the Aspen Center for Physics for hospitality during the completion of this work. This material is based upon work supported by the National Science Foundation under Grant No. PHY-9802484. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

A Conventions

We follow closely the conventions of [1]. The Dirac matrices in ten dimensions Γ^μ are decomposed in terms of 16-dimensional gamma matrices γ^μ such that

$$\Gamma^\mu = \begin{pmatrix} 0 & \gamma^\mu \\ \bar{\gamma}^\mu & 0 \end{pmatrix}, \quad (\text{A.1})$$

where

$$\gamma^\mu \bar{\gamma}^\nu + \gamma^\nu \bar{\gamma}^\mu = 2\eta^{\mu\nu}, \quad \gamma^\mu = (\gamma^\mu)^{\alpha\beta}, \quad \bar{\gamma}^\mu = \gamma_{\alpha\beta}^\mu, \quad (\text{A.2})$$

$$\gamma^\mu = (1, \gamma^I, \gamma^9), \quad \bar{\gamma}^\mu = (-1, \gamma^I, \gamma^9). \quad (\text{A.3})$$

Here (α, β) are $SO(9, 1)$ spinor indices in chiral representation; we use the Majorana representation for Γ such that $C = \Gamma^0$, and so all γ^μ are real and symmetric. We use the convention that $\gamma^{\mu_1 \dots \mu_k}$ are the antisymmetrised product of k gamma matrices with unit weight.

We assume the normalisation $\gamma^0 \bar{\gamma}^1 \dots \gamma^8 \bar{\gamma}^9 = 1$, so that

$$\Gamma_{11} = \Gamma^0 \dots \Gamma^9 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (\text{A.4})$$

We define

$$\Pi^\alpha{}_\beta \equiv (\gamma^1 \bar{\gamma}^2 \gamma^3 \bar{\gamma}^4)^\alpha{}_\beta, \quad (\Pi')^\alpha{}_\beta \equiv (\gamma^5 \bar{\gamma}^6 \gamma^7 \bar{\gamma}^8)^\alpha{}_\beta, \quad \gamma^0 \bar{\gamma}^9 = \gamma^{+-}. \quad (\text{A.5})$$

Other useful relations are

$$\gamma^{+-} \Pi \Pi' = 1, \quad (\gamma^{+-})^2 = \Pi^2 = (\Pi')^2 = 1, \quad (\text{A.6})$$

$$\gamma^{+-} \gamma^\pm = \pm \gamma^\pm, \quad \bar{\gamma}^\pm \gamma^{+-} = \mp \bar{\gamma}^\pm, \quad \gamma^+ \bar{\gamma}^+ = \gamma^- \bar{\gamma}^- = 0. \quad (\text{A.7})$$

Abbreviated notations are used, in which the spinor indices are indicated by the positioning of matrices; frequently used expressions in the text include

$$\Omega Q \equiv \bar{\Omega}_\alpha{}^\beta Q_\beta; \quad Q\Omega \equiv Q_\alpha \Omega^\alpha{}_\beta; \quad \Omega^t Q \equiv (\bar{\Omega}^t)_\alpha{}^\beta Q_\beta; \quad Q\Omega^t \equiv Q_\alpha (\Omega^t)^\alpha{}_\beta; \quad (\text{A.8})$$

$$\Omega^\alpha{}_\beta = (\gamma^{i_1})^{\alpha\gamma} \dots (\bar{\gamma}^{i_q})_{\delta\beta}; \quad (\Omega^t)^\alpha{}_\beta = (\gamma^{i_q})^{\alpha\gamma} \dots (\bar{\gamma}^{i_1})_{\delta\beta}; \quad (\text{A.9})$$

$$\bar{\Omega}_\alpha{}^\beta = (\bar{\gamma}^{i_1})_{\alpha\gamma} \dots (\gamma^{i_q})^{\delta\beta}; \quad (\bar{\Omega}^t)_\alpha{}^\beta = (\bar{\gamma}^{i_q})_{\alpha\gamma} \dots (\gamma^{i_1})^{\delta\beta}. \quad (\text{A.10})$$

B Symmetry currents and closed string superalgebra

Here we give the explicit forms of the bosonic symmetry currents in lightcone gauge; the fermion currents are given in the main text. The momenta are

$$\mathcal{P}^{+\tau} = p^+; \quad P^{+\sigma} = 0; \quad (\text{B.1})$$

$$\mathcal{P}^{I\tau} = (\cos(m\tau)\partial_\tau x^I + m \sin(m\tau)x^I); \quad (\text{B.2})$$

$$\mathcal{P}^{I\sigma} = -\cos(m\tau)\partial_\sigma x^I. \quad (\text{B.3})$$

The rotation currents are

$$\mathcal{J}^{+I\tau} = (\mu^{-1} \sin(m\tau)\partial_\tau x^I - p^+ x^I \cos(m\tau)); \quad (\text{B.4})$$

$$\mathcal{J}^{+I\sigma} = -\mu^{-1} \sin(m\tau)\partial_\sigma x^I; \quad (\text{B.5})$$

$$\mathcal{J}^{ij\tau} = (x^i \partial_\tau x^j - x^j \partial_\tau x^i - \frac{1}{2} i p^+ (\theta^1 \gamma^{-ij} \theta^1 + \theta^2 \gamma^{-ij} \theta^2)); \quad (\text{B.6})$$

$$\mathcal{J}^{ij\sigma} = -(x^i \partial_\sigma x^j - x^j \partial_\sigma x^i + \frac{1}{2} i p^+ (\theta^1 \gamma^{-ij} \theta^1 - \theta^2 \gamma^{-ij} \theta^2)), \quad (\text{B.7})$$

with a corresponding expression for $\mathcal{J}^{i'j'}$. The remaining charge is the lightcone Hamiltonian.

$$\mathcal{P}^{-\tau} = (\partial_\tau x^- + i(\theta^1 \bar{\gamma}^- \partial_+ \theta^1 + \theta^2 \bar{\gamma}^- \partial_- \theta^2) - \mu m (x^I)^2 - 4im\theta^1 \bar{\gamma}^- \Pi \theta^2); \quad (\text{B.8})$$

$$\mathcal{P}^{-\sigma} = -(\partial_\sigma x^- + i(\theta^1 \bar{\gamma}^- \partial_+ \theta^1 - \theta^2 \bar{\gamma}^- \partial_- \theta^2)). \quad (\text{B.9})$$

Using the Virasoro constraint, and the fermion field equations, we find that the onshell Hamiltonian density is

$$\mathcal{H} = -\mathcal{P}^{-\tau} = \frac{1}{2p^+} ((\partial_\tau x^I)^2 + (\partial_\sigma x^I)^2 + m^2 (x^I)^2) + i(\theta^1 \bar{\gamma}^- \partial_\tau \theta^1 + \theta^2 \bar{\gamma}^- \partial_\tau \theta^2). \quad (\text{B.10})$$

Finally, let us remark that onshell $\mathcal{P}^{-\sigma} = -\partial_\sigma x^-$.

The symmetry superalgebra of the pp-wave background is as follows. The commutators of the bosonic generators are⁹

$$[P^-, P^I] = i\mu^2 J^{+I}, \quad [P^I, J^{+J}] = i\delta^{IJ} P^+, \quad [P^-, J^{+I}] = -iP^I, \quad (\text{B.11})$$

⁹Notice our rotational generators differ from the ones in [1] by a factor of i .

$$\begin{aligned}
[P^i, J^{jk}] &= -i(\delta^{ij}P^k - \delta^{ik}P^j), & [P^{i'}, J^{j'k'}] &= -i(\delta^{i'j'}P^{k'} - \delta^{i'k'}P^{j'}), \\
[J^{+i}, J^{jk}] &= -i(\delta^{ij}J^{+k} - \delta^{ik}J^{+j}), & [J^{+i'}, J^{j'k'}] &= -i(\delta^{i'j'}J^{+k'} - \delta^{i'k'}J^{+j'}), \\
[J^{ij}, J^{kl}] &= -i(\delta^{jk}J^{il} + \dots), & [J^{i'j'}, J^{k'l'}] &= -i(\delta^{j'k'}J^{i'l'} + \dots),
\end{aligned}$$

where the ellipses denote permutations, whilst the commutation relations between the even and odd generators are

$$\begin{aligned}
[J^{ij}, Q^\pm] &= -\frac{1}{2}iQ^\pm(\gamma^{ij}), & [J^{i'j'}, Q^\pm] &= -\frac{1}{2}iQ^\pm(\gamma^{i'j'}), \\
[J^{+I}, Q^-] &= -\frac{1}{2}iQ^+(\gamma^{+I}), \\
[P^I, Q^-] &= \frac{1}{2}\mu Q^+(\Pi\gamma^{+I}), & [P^-, Q^+] &= \mu Q^+\Pi,
\end{aligned} \tag{B.12}$$

and the anticommutation relations are

$$\begin{aligned}
\{Q^+, \bar{Q}^+\} &= 2P^+\bar{\gamma}^-, \\
\{Q^+, \bar{Q}^-\} &= (\bar{\gamma}^-\gamma^+\bar{\gamma}^I)P^I - i\mu(\bar{\gamma}^-\gamma^+\bar{\gamma}^I\Pi)J^{+I}, \\
\{Q^-, \bar{Q}^+\} &= (\bar{\gamma}^+\gamma^-\bar{\gamma}^I)P^I - i\mu(\bar{\gamma}^+\gamma^-\bar{\gamma}^I\Pi)J^{+I}, \\
\{Q^-, \bar{Q}^-\} &= 2\bar{\gamma}^+P^- + i\mu(\gamma^{+ij}\Pi)J^{ij} + i\mu(\gamma^{+i'j'}\Pi')J^{i'j'}.
\end{aligned} \tag{B.13}$$

C Mode expansions for closed strings

In our conventions the closed string mode expansions are given by

$$x^I(\sigma, \tau) = \cos(m\tau)x_0^I + m^{-1}\sin(m\tau)p_0^I + i\sum_{n\neq 0}\omega_n^{-1}(\alpha_n^{II}\tilde{\phi}_n + \alpha_n^{2I}\phi_n); \tag{C.1}$$

$$\theta^1(\sigma, \tau) = \theta_0^1\cos(m\tau) + \Pi\theta_0^2\sin(m\tau) + \sum_{n\neq 0}c_n\left(id_n\Pi\theta_n^2\phi_n + \theta_n^1\tilde{\phi}_n\right); \tag{C.2}$$

$$\theta^2(\sigma, \tau) = \theta_0^2\cos(m\tau) - \Pi\theta_0^1\sin(m\tau) + \sum_{n\neq 0}c_n\left(-id_n\Pi\theta_n^1\tilde{\phi}_n + \theta_n^2\phi_n\right). \tag{C.3}$$

After canonical quantization we get the following (anti)commutators

$$[p_0^I, x_0^J] = -i\delta^{IJ}, \quad [\alpha_m^{\mathcal{I}I}, \alpha_n^{\mathcal{J}J}] = \frac{1}{2}\omega_m\delta_{n+m,0}\delta^{\mathcal{I}\mathcal{J}}\delta^{IJ}, \tag{C.4}$$

$$\{\theta_0^{\mathcal{I}}, \theta_0^{\mathcal{J}}\} = \frac{1}{4p^+}(\gamma^+)\delta^{\mathcal{I}\mathcal{J}}, \quad \{\theta_m^{\mathcal{I}}, \theta_n^{\mathcal{J}}\} = \frac{1}{4p^+}(\gamma^+)\delta^{\mathcal{I}\mathcal{J}}\delta_{m+n,0}, \tag{C.5}$$

where $\mathcal{I} = 1, 2$. It is convenient to introduce creation and annihilation operators

$$a_0^I = \frac{1}{\sqrt{2m}}(p_0^I + imx_0^I), \quad \bar{a}_0^I = \frac{1}{\sqrt{2m}}(p_0^I - imx_0^I), \quad [\bar{a}_0^I, a_0^J] = \delta^{IJ}. \tag{C.6}$$

Expressed in terms of these modes the spacetime charges are

$$P^+ = p^+, \quad P^I = p_0^I, \quad J^{+I} = -x_0^I p^+, \tag{C.7}$$

$$Q^{+1} = 2p^+\bar{\gamma}^-\theta_0^2, \quad Q^{+2} = -2p^+\bar{\gamma}^-\theta_0^1. \tag{C.8}$$

Note that the complex Q^+ appearing in the closed string algebra is

$$Q^+ = \frac{1}{\sqrt{2}}(iQ^{+1} - Q^{+2}) = 2p^+ \bar{\gamma}^- \theta_0. \quad (\text{C.9})$$

The rotation charges are

$$\begin{aligned} J^{IJ} &= -i(a_0^I \bar{a}_0^J - a_0^J \bar{a}_0^I + \frac{1}{2} \sum_{\mathcal{I}=1,2} \theta_0^{\mathcal{I}} \gamma^{-IJ} \theta_0^{\mathcal{I}}) \\ &\quad - i \sum_{\mathcal{I}=1,2} \sum_{n>0} (2\omega_n^{-1} (\alpha_{-n}^{\mathcal{I}I} \alpha_n^{\mathcal{I}J} - \alpha_{-n}^{\mathcal{I}J} \alpha_n^{\mathcal{I}I}) + p^+ \theta_{-n}^{\mathcal{I}} \gamma^{-IJ} \theta_n^{\mathcal{I}}). \end{aligned} \quad (\text{C.10})$$

The Hamiltonian is

$$\begin{aligned} H &= \frac{1}{2p^+} (p_0^2 + m^2 x_0^2) + 2im(\theta_0^1 \bar{\gamma}^- \Pi \theta_0^2) \\ &\quad + \sum_{\mathcal{I}=1,2} \sum_{n \neq 0} \left(\frac{1}{p^+} \alpha_{-n}^{\mathcal{I}I} \alpha_n^{\mathcal{I}I} + \omega_n \theta_{-n}^{\mathcal{I}} \bar{\gamma}^- \theta_n^{\mathcal{I}} \right). \end{aligned} \quad (\text{C.11})$$

Finally the dynamical supercharges take the form

$$\begin{aligned} Q^{-1} &= (2p_0^I \bar{\gamma}^I \theta_0^1 - 2mx_0^I \bar{\gamma}^I \Pi \theta_0^2) \\ &\quad + \sum_{n>0} \left(4c_n \alpha_{-n}^{I1} \bar{\gamma}^I \theta_n^1 + \frac{2im}{\omega_n c_n} \alpha_{-n}^{2I} \bar{\gamma}^I \Pi \theta_n^2 + h.c. \right) \\ Q^{-2} &= (2p_0^I \bar{\gamma}^I \theta_0^2 + 2mx_0^I \bar{\gamma}^I \Pi \theta_0^1) \\ &\quad + \sum_{n>0} \left(4c_n \alpha_{-n}^{I2} \bar{\gamma}^I \theta_n^2 - \frac{2im}{\omega_n c_n} \alpha_{-n}^{1I} \bar{\gamma}^I \Pi \theta_n^1 + h.c. \right) \end{aligned} \quad (\text{C.12})$$

The complex Q^- appearing in the algebra is $(Q^{-1} + iQ^{-2})/\sqrt{2}$.

D Poisson-Dirac brackets for oscillators

Each of the classical commutation relations is of the generic form

$$[z^m(\sigma), z^n(\sigma')] = h^{mn} \delta(\sigma, \sigma'), \quad (\text{D.1})$$

where z^m denotes both scalars and fermions. To evaluate the Poisson-Dirac brackets for the oscillators, we should use the following procedure.

Firstly recall that given an orthonormal complete set of functions, $u_n(\sigma)$, over some domain, the delta function can be represented as,

$$\delta(\sigma, \sigma') = \sum u_n(\sigma) u_n(\sigma') \quad (\text{D.2})$$

In the domain $[0, \pi]$ there are several distinct sets of complete orthonormal functions:

$$u_0 = \frac{1}{\sqrt{\pi}}, \quad u_n = \sqrt{\frac{2}{\pi}} \cos n\sigma, \quad n = 1, 2, \dots \quad (\text{D.3})$$

$$v_n = \sqrt{\frac{2}{\pi}} \sin n\sigma, \quad n = 1, 2, \dots \quad (\text{D.4})$$

These give rise to the following representations of the delta function

$$\begin{aligned}\delta(\sigma, \sigma') &= \frac{1}{\pi} + \sum_{n=1}^{\infty} \frac{2}{\pi} \cos(n\sigma) \cos(n\sigma') = \sum_{-\infty < n < \infty} \frac{1}{\pi} \cos(n\sigma) \cos(n\sigma'), \quad (\text{D.5}) \\ &= \sum_{n=1}^{\infty} \frac{2}{\pi} \sin(n\sigma) \sin(n\sigma') = \sum_{-\infty < n < \infty} \frac{1}{\pi} \sin(n\sigma) \sin(n\sigma').\end{aligned}$$

Now to evaluate commutation relations for fields which are already expanded in one of these bases is straightforward. For example, scalars satisfying Neumann or Dirichlet boundary conditions (3.6) and (3.7) are expanded in terms of cosines and sines respectively. Thus we can substitute the delta function Fourier expansions directly into their commutation relations (3.17) to determine the commutation relations for the modes. This gives the first terms in (3.18).

D.1 D_- branes

The fermion mode expansions (3.12) and (3.21) do not however involve mutually orthogonal functions. This reflects the fact that θ^1 and θ^2 do not satisfy pure Neumann or pure Dirichlet boundary conditions at the string endpoints. However, certain combinations of θ^1 and θ^2 will satisfy such conditions.

Let us consider first the case $\Omega\Pi\Omega\Pi = -1$; then by construction $(\theta^1 - \Omega\theta^2)$ satisfies a Dirichlet condition at the endpoints whilst $(\theta^1 + \Omega\theta^2)$ satisfies a Neumann condition at the endpoints. In fact using (3.12) we find that

$$\theta_D(\sigma, \tau) = (\theta^1 - \Omega\theta^2)(\sigma, \tau) = 2 \sum_n c_n e^{-i\omega_n \tau} (d_n \Pi + i\Omega) \sin(n\sigma) \theta_n; \quad (\text{D.6})$$

$$\begin{aligned}\theta_N(\sigma, \tau) &= (\theta^1 + \Omega\theta^2)(\sigma, \tau) \quad (\text{D.7}) \\ &= 2(\theta_0 \cos(m\tau) - \Omega\Pi\theta_0 \sin(m\tau)) + 2 \sum_{n \neq 0} c_n e^{-i\omega_n \tau} (id_n \Pi + \Omega) \cos(n\sigma) \theta_n,\end{aligned}$$

which are manifestly both expansions in the orthogonal functions. Using the commutation relations (3.17) we find that

$$\{\theta_D(\sigma), \theta_D(\sigma')\}_{D.B.} = \frac{i}{2p^+} \pi \gamma^+ \delta(\sigma, \sigma'); \quad \{\theta_N(\sigma), \theta_N(\sigma')\}_{D.B.} = \frac{i}{2p^+} \pi \gamma^+ \delta(\sigma, \sigma'), \quad (\text{D.8})$$

and substituting in appropriate mode expansions for the Dirac delta function we are able to fix all fermionic oscillator anticommutators. This gives

$$\{\theta_n^\alpha, \theta_m^\beta\}_{D.B.} = \frac{1}{8p^+} (\gamma^+)^{\alpha\beta} \delta_{n+m,0}, \quad (\text{D.9})$$

which is in fact the same expression as in flat space.

D.2 D_+ branes

Now let us consider the case $\Omega\Pi\Omega\Pi = 1$ which is rather more subtle. By construction $(\theta^1 - \Omega\theta^2)$ again satisfies a pure Dirichlet boundary condition. Using the mode expansions (3.21) we find the explicit form to be

$$\theta_D(\sigma, \tau) = (\theta^1 - \Omega\theta^2)(\sigma, \tau) = 2 \sum_n c_n e^{-i\omega_n \tau} (d_n \Pi + i\Omega) \sin(n\sigma) \theta_n, \quad (\text{D.10})$$

which is in fact the same as in (D.6); it also satisfies the commutation relation given in (D.8). Thus for the non-zero modes, the Poisson-Dirac brackets are the same as in the $\Pi\Omega\Pi\Omega = -1$ case, namely (D.9). To extract the brackets for the zero modes is rather harder, because they drop out of both this Dirichlet combination and of the natural Neumann combination which is

$$\theta_N(\sigma, \tau) = (\theta^1 + \Omega\theta^2)(\sigma, \tau) - m \int^\sigma d\sigma' (\Pi\theta^2 + \Omega\Pi\theta^1)(\sigma', \tau). \quad (\text{D.11})$$

This is Neumann because using the field equations, and the boundary conditions, its sigma derivative vanishes on the boundary. Explicitly,

$$\partial_\sigma \theta_N(\sigma, \tau) = 2 \sum_n c_n \omega_n e^{-i\omega_n \tau} (i d_n \Pi - \Omega) \sin(n\sigma) \theta_n. \quad (\text{D.12})$$

To derive the zero mode commutation relations we should work directly with the fields θ^1, θ^2 which are not expanded in terms of orthogonal functions.

Explicitly evaluating the anticommutators using the mode expansions (3.21) we find

$$\begin{aligned} \{\theta^1(\sigma, \tau), \theta^1(\sigma', \tau)\}_{D.B.} &= \frac{i}{4p^+} \pi(\gamma^+) \delta(\sigma, \sigma') \\ &= \{\mathcal{P}_+\theta_0^+, \mathcal{P}_+\theta_0^+\} e^{m(\sigma+\sigma')} + \{\mathcal{P}_+\theta_0^+, \mathcal{P}_-\theta_0^-\} e^{m(\sigma-\sigma')} \\ &\quad + \{\mathcal{P}_-\theta_0^-, \mathcal{P}_+\theta_0^+\} e^{-m(\sigma-\sigma')} + \{\mathcal{P}_-\theta_0^-, \mathcal{P}_-\theta_0^-\} e^{-m(\sigma+\sigma')} \\ &\quad + \frac{1}{p^+} \left(\sum_{n>0} \frac{1}{4} i \gamma^+ \cos n(\sigma - \sigma') - i \sum_{n>0} d_n^2 c_n^4 \gamma^+ \cos n(\sigma + \sigma') \right. \\ &\quad \left. + \frac{1}{2} i \sum_{n>0} d_n c_n^4 (1 - d_n^2) \gamma^+ \Omega \Pi \sin n(\sigma + \sigma') \right) \end{aligned} \quad (\text{D.13})$$

In this expression we have substituted the brackets for the non-zero modes. We have also set the anticommutators of the zero modes with non-zero modes to zero. Had these anticommutators being non-zero the right-hand side of the above expression would be time-dependent. The other anticommutators are

$$\begin{aligned} \{\theta^1(\sigma, \tau), \Pi\theta^2(\sigma', \tau)\}_{D.B.} &= 0 \\ &= \{\mathcal{P}_+\theta_0^+, \mathcal{P}_+\theta_0^+\} e^{m(\sigma+\sigma')} - \{\mathcal{P}_+\theta_0^+, \mathcal{P}_-\theta_0^-\} e^{m(\sigma-\sigma')} \end{aligned} \quad (\text{D.14})$$

$$\begin{aligned}
& +\{\mathcal{P}_-\theta_0^-, \mathcal{P}_+\theta_0^+\}e^{-m(\sigma-\sigma')} - \{\mathcal{P}_-\theta_0^-, \mathcal{P}_-\theta_0^-\}e^{-m(\sigma+\sigma')} \\
& +\frac{1}{p^+} \left(i \sum_{n>0} c_n^4 (1-d_n^2)^2 \Omega \Pi \gamma^+ \cos n(\sigma+\sigma') \right. \\
& \left. +\frac{1}{2}i \sum_{n>0} d_n c_n^4 (1-d_n^2) \gamma^+ \sin n(\sigma+\sigma') \right),
\end{aligned}$$

and

$$\begin{aligned}
\{\Pi\theta^2(\sigma, \tau), \Pi\theta^2(\sigma', \tau)\}_{D.B.} &= \frac{i}{4p^+} \pi(\gamma^+) \delta(\sigma, \sigma') \tag{D.15} \\
&= \{\mathcal{P}_+\theta_0^+, \mathcal{P}_+\theta_0^+\}e^{m(\sigma+\sigma')} - \{\mathcal{P}_+\theta_0^+, \mathcal{P}_-\theta_0^-\}e^{m(\sigma-\sigma')} \\
&\quad - \{\mathcal{P}_-\theta_0^-, \mathcal{P}_+\theta_0^+\}e^{-m(\sigma-\sigma')} + \{\mathcal{P}_-\theta_0^-, \mathcal{P}_-\theta_0^-\}e^{-m(\sigma+\sigma')} \\
&+ \frac{1}{p^+} \left(\sum_{n>0} \frac{1}{4} i \gamma^+ \cos n(\sigma-\sigma') - i \sum_{n>0} d_n^2 c_n^4 \gamma^+ \cos n(\sigma+\sigma') \right. \\
&\left. +\frac{1}{2}i \sum_{n>0} d_n c_n^4 (1-d_n^2) \gamma^+ \Omega \Pi \sin n(\sigma+\sigma') \right),
\end{aligned}$$

where again we have used the relations for non-zero modes. It is also convenient to insert factors of Π into the brackets as we have done. Note that (D.13) and (D.15) can be simplified using the identity

$$\frac{1}{4}i\pi\delta(\sigma, \sigma') = \frac{1}{8}i + \frac{1}{4}i \sum_{n>0} \cos(n(\sigma-\sigma')). \tag{D.16}$$

Now the Poisson-Dirac brackets for the fermionic zero modes must satisfy all three expressions. Comparing (D.13) and (D.15) we find that

$$\{\mathcal{P}_+\theta_0^+, \mathcal{P}_-\theta_0^-\}_{D.B.} = 0. \tag{D.17}$$

This means that (D.13) can be simplified to give

$$\frac{1}{8} = a\mathcal{P}_+e^{m(\sigma+\sigma')} + b\mathcal{P}_-e^{-m(\sigma+\sigma')} - \sum_{n>0} \frac{m}{4(m^2+n^2)} (m \cos n(\sigma+\sigma') - n\Omega\Pi \sin n(\sigma+\sigma')). \tag{D.18}$$

Similarly (D.14) yields,

$$0 = a\mathcal{P}_+e^{m(\sigma+\sigma')} - b\mathcal{P}_-e^{-m(\sigma+\sigma')} + \sum_{n>0} \frac{n}{4(m^2+n^2)} (m \sin n(\sigma+\sigma') + n\Omega\Pi \cos n(\sigma+\sigma')). \tag{D.19}$$

where we anticipate that

$$\begin{aligned}
\{\mathcal{P}_+\theta_0^+, \mathcal{P}_+\theta_0^+\}_{D.B.} &= \frac{i}{p^+} a \mathcal{P}_+ \gamma^+. \tag{D.20} \\
\{\mathcal{P}_-\theta_0^-, \mathcal{P}_-\theta_0^-\}_{D.B.} &= \frac{i}{p^+} b \mathcal{P}_- \gamma^+.
\end{aligned}$$

and we used the relations

$$d_n^2 c_n^4 = \frac{m^2}{4(n^2 + m^2)}; \quad (\text{D.21})$$

$$d_n(1 - d_n^2)c_n^4 = \frac{mn}{2(n^2 + m^2)},$$

$$c_n^4(1 - d_n^2)^2 = \frac{n^2}{m^2 + n^2} \quad (\text{D.22})$$

Notice that (D.19) is equal to the σ derivative of (D.18), so we only need to solve the latter. Since a and b are each multiplied by orthogonal projection operators, we can obtain two decoupled equations by multiplying (D.18) by \mathcal{P}_+ and \mathcal{P}_- , respectively. The result is

$$\frac{1}{8} = ae^{m(\sigma+\sigma')} - \sum_{n>0} \frac{m}{4(n^2 + m^2)} (m \cos n(\sigma + \sigma') - n \sin n(\sigma + \sigma')), \quad (\text{D.23})$$

$$\frac{1}{8} = be^{-m(\sigma+\sigma')} - \sum_{n>0} \frac{m}{4(n^2 + m^2)} (m \cos n(\sigma + \sigma') + n \sin n(\sigma + \sigma')),$$

Notice that the equation for b is identical to the equation for a but with $m \rightarrow -m$. Thus, it is sufficient to solve (D.23).

To solve (D.23) we note that the variable $\sigma + \sigma'$ takes values in $[0, 2\pi]$. The Fourier transform of the exponential is given by

$$e^{m(\sigma+\sigma')} = \frac{1}{\pi} (e^{2\pi m} - 1) \left(\frac{1}{2m} + \sum_{n>0} \frac{m}{n^2 + m^2} (m \cos n(\sigma + \sigma') - n \sin n(\sigma + \sigma')) \right) \quad (\text{D.24})$$

Inserting in (D.23) we determine a ,

$$a = \frac{\pi m}{4(e^{2\pi m} - 1)}; \quad (\text{D.25})$$

$$b = \frac{\pi m e^{2\pi m}}{4(e^{2\pi m} - 1)}.$$

where we used $b = a(-m)$. Note that these do reduce to the usual flat space anticommutators in the flat space ($m \rightarrow 0$) limit.

Notice that the $(+, -, 0, 4)$ and $(+, -, 4, 0)$ branes are special in that $\Omega\Pi = 1$, which implies $\mathcal{P}_- = 0$ and we only have the zero modes θ_0^+ . The derivation we presented still holds, but we need to set $\mathcal{P}_- = 0$ and $\mathcal{P}_+ = 1$ in all formulas. In particular,

$$\{\theta_0^+, \theta_0^+\}_{D.B.} = \frac{i}{p^+} a \gamma^+. \quad (\text{D.26})$$

with the same a as in (D.25). For the corresponding anti-branes with $\Omega\Pi = -1$, it is θ_0^- and b which are non-zero.

E Kappa symmetry projections for time dependent branes

In this appendix we use the kappa symmetry projection to show the number of (closed string) supersymmetries preserved by rotating and time-dependent D3-branes.

In the plane wave background the curved space gamma matrices, denoted here by $\Gamma_m^{(c)}$, are related to the 32 dimensional flat space matrices given in appendix A as

$$\Gamma_+^{(c)} = -\mu\Gamma_+ + \frac{1}{2}\mu \sum_{I=1}^8 (x^I)^2 \Gamma_-; \quad \Gamma_-^{(c)} = -\mu^{-1}\Gamma_-; \quad \Gamma_I^{(c)} = \Gamma_I. \quad (\text{E.1})$$

The Killing spinors of the background may conveniently be written as

$$\begin{aligned} \epsilon &= (1 + iP)(\Psi_R + i\Psi_I); \\ P &= -\frac{1}{2}\Gamma_- \left(\sum_{i=1}^4 x^i \Gamma_i \Pi + \sum_{i'=5}^8 x^{i'} \Gamma_{i'} \Pi' \right); \\ \Psi_R &= (\lambda_+ + \cos(\mu x^+) \lambda_- + \Pi \sin(\mu x^+) \eta_-); \\ \Psi_I &= (\eta_+ + \cos(\mu x^+) \eta_- - \Pi \sin(\mu x^+) \lambda_-), \end{aligned} \quad (\text{E.2})$$

where the 32 dimensional constant negative chirality spinors η_{\pm} and λ_{\pm} satisfy $\Gamma_{\pm} \eta_{\pm} = \Gamma_{\pm} \lambda_{\pm} = 0$.

The kappa symmetry projection associated with any brane embedding is defined by [28]

$$d^{p+1} \xi \Gamma = -e^{-\Phi} \mathcal{L}_{DBI}^{-1} e^{\mathcal{F}} \wedge X|_{vol}, \quad (\text{E.3})$$

where $X = \bigoplus_n \Gamma_{(2n)} K^n I$, $|_{vol}$ indicates that one should pick the terms proportional to the volume form, and the operations I and K act on spinors as $I\psi = -i\psi$ and $K\psi = \psi^*$. \mathcal{L}_{DBI}^{-1} is the value of the DBI Lagrangian evaluated on the background. Here we have used the notation

$$\Gamma_{(n)} = \frac{1}{n!} d\xi^{i_1} \wedge \dots \wedge d\xi^{i_n} \Gamma_{i_1 \dots i_n}, \quad (\text{E.4})$$

where $\Gamma_{i_1 \dots i_n}$ is the pullback for the target space gamma matrices

$$\Gamma_{i_1 \dots i_n} = \partial_{i_1} X^{m_1} \dots \partial_{i_n} X^{m_n} \Gamma_{m_1 \dots m_n}^{(c)}. \quad (\text{E.5})$$

For the case in hand, a D3-brane whose worldvolume directions are (x^+, x^-, x^1, x^2) and whose transverse positions are $x^{r'}$ (x^+), the kappa symmetry projection is $\Gamma \epsilon = \epsilon$ where

$$\Gamma = -i(Q + R); \quad Q = \Gamma_{+-12}; \quad R = \sum_{r'} \mu^{-1} \partial_+ x^{r'} \Gamma_{-12r'}. \quad (\text{E.6})$$

and ϵ is the killing spinor in (E.2) evaluate on the worldvolume of the D3-brane. From the real and imaginary parts of the projection we get two independent equations which are

$$\begin{aligned} Q\Psi_R + R\lambda_+ - QP\eta_+ &= -\Psi_I - P\eta_+; \\ Q\Psi_I + R\eta_+ + QP\lambda_+ &= \Psi_R - P\eta_+, \end{aligned} \quad (\text{E.7})$$

where we have used $R\lambda_- = R\psi_- = P\lambda_- = P\psi_- = 0$. Inserting Ψ_R and Ψ_I from (E.2) we get

$$\begin{aligned} (Q\lambda_+ + \eta_+) + (Q\lambda_- + \eta_-) \cos(\mu x^+) + \Pi \sin(\mu x^+) (Q\psi_- - \lambda_-) + (R + P)\lambda_+ &= QP\psi_+; \\ (Q\eta_+ - \lambda_+) + (Q\eta_- - \lambda_-) \cos(\mu x^+) - \Pi \sin(\mu x^+) (Q\lambda_- + \psi_-) + (R + P)\eta_+ &= -QP\lambda_+ \end{aligned} \quad (\text{E.8})$$

These equations should hold for any values of the worldvolume coordinates. This implies that the part that is independent of the worldvolume coordinates should vanish separately,

$$Q\lambda_+ = -\psi_+. \quad (\text{E.9})$$

This relates the real and imaginary parts of the parts of the spinor satisfying $\Gamma_+\epsilon = 0$. The remaining equations can be satisfied by imposing separately:

$$Q\lambda_- = -\psi_-; \quad RQ\psi_+ = [Q, P]\psi_+, \quad \{Q, R\} = 0. \quad (\text{E.10})$$

The last equation holds automatically for the R and Q given in (E.6). The first one can always be satisfied for non-zero λ_- independently of $x^{r'}(x^+)$; thus any time dependent branes always preserve at least one quarter of the closed string supersymmetries, that is, half the Q^+ . The second equation in (E.10) imposes further constraints on η_+ and determines which closed string dynamical supercharges are preserved. As noted in [11], this equation is satisfied automatically if the constant transverse positions are at the origin whilst if any are not we project out the η_+ and λ_+ and break all the Q^- .

The solution in (E.10) is not the most general solution of (E.8), however. R and P may depend on x^+ in such a way that these terms may be combined with the remaining terms in (E.8) rather than treated separately as is done in (E.10). This is exactly what happens for the symmetry-related branes. Recall that for these branes the relevant boundary conditions are (2.35). For these boundary conditions, we still have eight independent solutions to (E.8) satisfying

$$Q\lambda_- = -\eta_-; \quad \eta_+ = 0. \quad (\text{E.11})$$

These are the eight kinematical supercharges which are preserved irrespective of the time dependent boundary conditions. Furthermore, with the boundary conditions (2.35) there are eight additional solutions to (E.8) (provided that the other transverse scalars are zero) for which

$$\eta_- + Q\lambda_- = \sum_{r'} (b^{r'} \Gamma_{-r'} - a^{r'} \Pi \Gamma_{-12r'}) \eta_+. \quad (\text{E.12})$$

with (E.9) also satisfied. The total number of preserved supersymmetries is thus sixteen as expected. Furthermore the specific combination of supersymmetries preserved in (E.12) agrees with that found in the open string analysis.

References

- [1] R. R. Metsaev, “Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background,” Nucl. Phys. B **625**, 70 (2002) [arXiv:hep-th/0112044].
- [2] R. R. Metsaev and A. A. Tseytlin, “Exactly solvable model of superstring in Ramond-Ramond background,” Phys. Rev. D **65** (2002) 126004 [arXiv:hep-th/0202109].
- [3] M. Blau, J. Figueroa-O’Farrill, C. Hull and G. Papadopoulos, “Penrose limits and maximal supersymmetry”, Class. Quant. Grav. **19** L87 (2002) [arXiv: hep-th/0201081]; M. Blau, J. Figueroa-O’Farrill, C. Hull and G. Papadopoulos, “A new maximally supersymmetric background of type IIB superstring theory”, JHEP **0201** 047 (2001) [arXiv: hep-th/0110242].
- [4] M. Blau, J. Figueroa-O’Farrill and G. Papadopoulos, “Penrose limits, supergravity and brane dynamics,” Class. Quant. Grav. **19** 4753 (2002) [arXiv:hep-th/0202111].
- [5] D. Berenstein, J. Maldacena, and H. Nastase, “Strings in flat space and pp waves from $\mathcal{N} = 4$ Super Yang Mills,” JHEP **0204** (2002) 013 [arXiv:hep-th/0202021].
- [6] K. Skenderis and M. Taylor, “ Open strings in the plane wave background II: superalgebras and spectrum”, in preparation.
- [7] M. Billó and I. Pesando, “Boundary states for GS superstring in an Hpp wave background,” Phys. Lett. **B532**, 206 (2002) [arXiv:hep-th/0203028].
- [8] O. Bergman, M. Gaberdiel and M. Green, “D-brane interactions in type IIB plane-wave background”, arXiv:hep-th/0205183.
- [9] A. Dabholkar and S. Parvizi, “Dp branes in pp-wave background,” Nucl. Phys. B **641** (2002) 223 [arXiv:hep-th/0203231].
- [10] P. Bain, K. Peeters and M. Zamaklar, “D-branes in a plane wave from covariant open strings”, arXiv:hep-th/0208038.
- [11] K. Skenderis and M. Taylor, “Branes in AdS and pp-wave spacetimes,” JHEP **06** (2002) 025 [arXiv:hep-th/0204054].
- [12] P. Bain, P. Meessen and M. Zamaklar, “Supergravity solutions for D-branes in Hpp-wave backgrounds,” arXiv:hep-th/0205106.

- [13] P. Lee and J. Park, “Open Strings in PP-Wave Background from Defect Conformal Field Theory,” arXiv:hep-th/0203257.
- [14] D. Berenstein, E. Gava, J. Maldacena, K. Narain, and H. Nastase, “Open strings on plane waves and their Yang-Mills duals,” arXiv:hep-th/0203249.
- [15] V. Balasubramanian, M. Huang, T. Levi and A. Naqvi, “Open strings from N=4 Super Yang Mills”, JHEP**0208** (2002) 037 [arXiv:hep-th/0204196].
- [16] H. Takayanagi and T. Takayanagi, “Notes on giant gravitons on pp-waves”, arXiv:hep-th/0209160.
- [17] C. S. Chu and P. M. Ho, “Noncommutative D-brane and open string in pp-wave background with B-field,” Nucl. Phys. B **636** (2002) 141 [arXiv:hep-th/0203186]; A. Kumar, R. Nayak and Sanjay, “D-brane solutions in pp-wave background”, Phys. Lett. B **541** (2002) 183 [arXiv:hep-th/0204025]; M. Alishahiha and A. Kumar, “D-brane solutions from new isometries of pp-waves”, Phys. Lett. B **542** (2002) 130 [arXiv:hep-th/0205134]; S. Seki, “D5-brane in Anti-de Sitter space and Penrose limit”, Phys. Lett. B **542** (2002) 162 [arXiv:hep-th/0205266]; D. Mateos and S. Ng, “Penrose limits of the baryonic D5-brane”, JHEP **0208** (2002) 005 [arXiv:hep-th/0205291]; Y. Moschita, “D-branes in NSNS and RR pp-wave backgrounds and S-duality”, arXiv:hep-th/0206131; K. Sugiyama and K. Yoshida, “Type IIA String and Matrix String on PP-wave”, Nucl. Phys. **B644** (2002) 128 [arXiv:hep-th/0208029]; A. Biswas, A. Kumar and K. Panigrahi, “p-p’ branes in pp-wave background”, arXiv:hep-th/0208042; O. Ganor and U. Varadarajan, “Nonlocal effects on D-branes in plane wave backgrounds”, arXiv:hep-th/0210035; S. Hyun and H. Shin, “Solvable $N = (4,4)$ type IIA string theory in plane wave background and D-branes”, arXiv:hep-th/0210158; Y. Hikida and S. Yamaguchi, “D-branes in PP-Waves and Massive Theories on Worldsheet with Boundary”, arXiv:hep-th/0210262.
- [18] N. D. Lambert and P. C. West, “D-branes in the Green-Schwarz formalism,” Phys. Lett. B **459**, 515 (1999) [arXiv:hep-th/9905031].
- [19] M. Berkooz, M. R. Douglas and R. G. Leigh, “Branes intersecting at angles,” Nucl. Phys. B **480**, 265 (1996) [arXiv:hep-th/9606139].
- [20] B. Acharya, J. Figueroa-O’Farrill, B. Spence and S. Stanciu, “Planes, branes and automorphisms II: branes in motion,” JHEP **9807** (2002) 005 [arXiv:hep-th/9805176].

- [21] J. McGreevy, L. Susskind and N. Toumbas, “Invasion of the giant gravitons from anti-de Sitter space,” *JHEP* **0006**, 008 (2000) [arXiv:hep-th/0003075].
- [22] A. Hashimoto, S. Hirano and N. Itzhaki, “Large branes in AdS and their field theory dual,” *JHEP* **0008**, 051 (2000) [arXiv:hep-th/0008016].
- [23] M. T. Grisaru, R. C. Myers and O. Tafjord, “SUSY and Goliath,” *JHEP* **0008**, 040 (2000) [arXiv:hep-th/0008015].
- [24] J. Maldacena and L. Maoz, “Strings on pp-waves and massive two dimensional field theories”, arXiv:hep-th/0207284.
- [25] N. Berkovits, “Conformal field theory for the superstring in a Ramond-Ramond plane wave background,” *JHEP* **0204** (2002) 037 [arXiv:hep-th/0203248].
- [26] N. Berkovits and J. Maldacena, “ $N = 2$ superconformal description of superstring in Ramond-Ramond plane wave backgrounds,” *JHEP* **0210**, 059 (2002) [arXiv:hep-th/0208092].
- [27] N. Berkovits, “ $N = 2$ sigma models for Ramond-Ramond backgrounds,” *JHEP* **0210**, 071 (2002) [arXiv:hep-th/0210078].
- [28] M. Cederwall, A. von Gussich, B. Nilsson and A. Westerberg, “The Dirichlet super-three-brane in ten-dimensional type IIB supergravity,” *Nucl. Phys. B* **490**, 163 (1997) [arXiv:hep-th/9610148]; M. Aganagic, C. Popescu and J. Schwarz, “D-brane actions with local kappa symmetry,” *Phys. Lett. B* **393**, 311 (1997) [arXiv:hep-th/9610249]; M. Cederwall, A. von Gussich, B. Nilsson, P. Sundell and A. Westerberg, “The Dirichlet super-p-branes in ten-dimensional type IIA and IIB supergravity,” *Nucl. Phys. B* **490**, 179 (1997) [arXiv:hep-th/9611159]; E. Bergshoeff and P. Townsend, “Super D-branes,” *Nucl. Phys. B* **490**, 145 (1997) [arXiv:hep-th/9611173]; M. Aganagic, C. Popescu and J. Schwarz, “Gauge-invariant and gauge-fixed D-brane actions,” *Nucl. Phys. B* **495**, 99 (1997) [arXiv:hep-th/9612080]; E. Bergshoeff, R. Kallosh, T. Ortin and G. Papadopoulos, “Kappa-symmetry, supersymmetry and intersecting branes,” *Nucl. Phys. B* **502**, 149 (1997) [arXiv:hep-th/9705040].