

# Wave function of string and membrane and spacetime geometry

Hongwei Xiong, Shujuan Liu

*Department of Applied Physics, Zhejiang University of Technology, Hangzhou, 310032, China*

(January 2, 2019)

## Abstract

A first-quantized string (and membrane) theory is developed here by using a general wave function of the string (and membrane), analogously to the first-quantized quantum theory of a point particle. From the general wave function of the string (and membrane), the properties of the string (and membrane) such as its relation to Bosons, Fermions and spacetime are investigated. The string and membrane wave functions are found to be very useful and we can deduce Klein-Gordon equation, Dirac equation and the fundamental property of the spacetime from this new starting point.

PACS number(s): 03.65.-w, 11.25.-w, 03.65.Pm

*Keywords:* Wave function; String; Membrane; Klein-Gordon equation; Dirac equation; Spacetime

arXiv:hep-th/0211015v2 18 Nov 2002

During the last twenty years, we witnessed the rapid development [1,2] of superstring theory which is trying to become a theory of everything. Due to the first and second revolutions of the superstring, although a lot of miraculous characteristics of the superstring such as string duality [3,4,5] are found, we still do not know the first law of the superstring. To describe the string based on the general quantum mechanical postulates, there are two approaches to investigate the string theory. One is the first-quantized string theory where the string is firstly regarded as a classical object, and the quantum property of the string is obtained by using the quantization method. The other is the second-quantized string field theory where the dynamical variables are regarded as the functional of the string coordinates. In the present work, we try to develop a first-quantized string (and membrane) theory by proposing a general wave function of the string (and membrane), analogously to the wave function of the point particle. The string (and membrane) is regarded as an inner structure of the membrane (and string) wave function, and is found to be relevant closely to the properties of bosons, fermions and spacetime. Through the investigation of the differential equation of the wave functions of membranes and strings, the Klein-Gordon equation for bosons and Dirac equation for fermions are deduced in a nature way.

It is well known that the wave function of a point particle is  $\Psi(X^\mu)$  ( $\mu = 1, \dots, N$  for a  $N$ -dimensional spacetime.). This sort of wave function is generalized to the case of the multi-component wave function  $\Psi_i(X^\mu)$  ( $i = 1, \dots, M$ ) which describes the particles with nonzero spin. In this paper, the wave function description of the point particle is generalized to develop a general membrane (and string) theory. The wave function of the membrane is supposed here to be  $\Psi_i(X^\mu(\tau, \sigma_1, \dots, \sigma_l))$ , and this would be the starting point to describe both bosons and fermions. Here  $\tau$  and  $\sigma_1, \dots, \sigma_l$  are real parameters to show the inner freedom of the membrane. As a special case, the string can be investigated when two parameters  $\tau$  and  $\sigma$  are used. Note that, when this sort of wave function of the membrane is used, the membrane is no more regarded as an entity of the ordinary membrane theory. In the ordinary first-quantized membrane theory, the membrane is firstly regarded as a classical oscillating membrane, and the method of quantization is used to reveal the

quantum character of the membrane. In this paper, we will try to reveal the character of the membrane and its relation to the spacetime directly from the wave function of the membrane. This can be regarded as a postulate in the present theory.

Firstly, we will investigate the quantum properties of strings from the string wave function. Suppose  $\Psi_M^T = (\Psi_1(X^\mu(\tau, \sigma)), \dots, \Psi_M(X^\mu(\tau, \sigma)))$ . Here  $T$  is a transpose of matrixes. For this sort of string wave function, we consider here the following general linear second-order partial differential equation:

$$\Xi_1 \frac{\partial^2 \Psi_M(X^\mu(\tau, \sigma))}{\partial \tau^2} + \Xi_2 \frac{\partial^2 \Psi_M(X^\mu(\tau, \sigma))}{\partial \sigma^2} = m^2 \Psi_M(X^\mu(\tau, \sigma)), \quad (1)$$

where  $\Xi_1$  and  $\Xi_2$  are  $M \times M$  matrixes. After a simple calculation, the above equation takes the form

$$\Xi_1 \frac{\partial^2 \Psi_M}{\partial X^\mu \partial X^\nu} \frac{\partial X^\mu}{\partial \tau} \frac{\partial X^\nu}{\partial \tau} + \Xi_2 \frac{\partial^2 \Psi_M}{\partial X^\mu \partial X^\nu} \frac{\partial X^\mu}{\partial \sigma} \frac{\partial X^\nu}{\partial \sigma} + \Xi_1 \frac{\partial \Psi_M}{\partial X^\mu} \frac{\partial^2 X^\mu}{\partial \tau^2} + \Xi_2 \frac{\partial \Psi_M}{\partial X^\mu} \frac{\partial^2 X^\mu}{\partial \sigma^2} = m^2 \Psi_M. \quad (2)$$

From the above equation, we have

$$\Xi_1 \frac{\partial \Psi_M}{\partial X^\mu} \frac{\partial^2 X^\mu}{\partial \tau^2} + \Xi_2 \frac{\partial \Psi_M}{\partial X^\mu} \frac{\partial^2 X^\mu}{\partial \sigma^2} = 0, \quad (3)$$

and

$$\Xi_1 \frac{\partial^2 \Psi_M}{\partial X^\mu \partial X^\nu} \frac{\partial X^\mu}{\partial \tau} \frac{\partial X^\nu}{\partial \tau} + \Xi_2 \frac{\partial^2 \Psi_M}{\partial X^\mu \partial X^\nu} \frac{\partial X^\mu}{\partial \sigma} \frac{\partial X^\nu}{\partial \sigma} = m^2 \Psi_M. \quad (4)$$

To have a nontrivial solution of  $\Psi_M$  and  $X^\mu(\tau, \sigma)$ , from Eq. (3), we have  $\Xi_1 = -\Xi_2$  and get the following equation about strings:

$$\partial^2 X^\mu / \partial \tau^2 - \partial^2 X^\mu / \partial \sigma^2 = 0. \quad (5)$$

It is not difficult to confirm that there is not a nontrivial solution of strings if  $\Xi_1 = \Xi_2$ . We see that the equation of the string is a nature result when the string wave function is used to describe the microscopic world. In this situation, the parameters  $\tau$  and  $\sigma$  of the string wave function are symmetry breaking. This implies the symmetry breaking of the

spacetime, i.e., for a  $N$ -dimensional spacetime, there should be a one-dimensional time and  $(N - 1)$ -dimensional space.

For Eq. (5), at first, we consider the open string with Neumann boundary condition  $X'^{\mu}(\tau, \sigma)|_{\sigma=0, \pi} = 0$ . Under this boundary condition, the solution becomes:

$$X^{\mu}(\tau, \sigma) = \xi_0^{\mu} + \xi_1^{\mu} \tau + \sum_{n=1}^{\infty} \left( \eta_{n_1}^{\mu} \cos n\tau + \eta_{n_2}^{\mu} \sin n\tau \right) \cos n\sigma. \quad (6)$$

Note that for a classical string, the parameters  $\xi_0^{\mu}$ ,  $\xi_1^{\mu}$ ,  $\eta_{n_1}^{\mu}$ ,  $\eta_{n_2}^{\mu}$  are determined by the initial conditions  $(X^{\mu}(\tau = 0, \sigma)$  and  $X'^{\mu}(\tau = 0, \sigma)|_{\sigma}$ ) of the oscillating string. In the case of quantum string introduced here, however, there would be a much strong confinement condition on these parameters through the investigation of Eq. (2), and this confinement would be relevant closely to the properties of the spacetime.

For the oscillating mode  $n$  of the open string,  $X_n^{\mu}(\tau, \sigma) = \left( \eta_{n_1}^{\mu} \cos n\tau + \eta_{n_2}^{\mu} \sin n\tau \right) \cos n\sigma$ . Substituting it into Eq. (4), we have

$$\Xi_1 \frac{\partial^2 \Psi_M}{\partial X^{\mu} \partial X^{\nu}} \chi_n^{\mu\nu} = m^2 \Psi_M, \quad (7)$$

where

$$\begin{aligned} \chi_n^{\mu\nu} = & \left( \eta_{n_1}^{\mu} \sin n\tau - \eta_{n_2}^{\mu} \cos n\tau \right) \left( \eta_{n_1}^{\nu} \sin n\tau - \eta_{n_2}^{\nu} \cos n\tau \right) n^2 \cos^2 n\sigma - \\ & \left( \eta_{n_1}^{\mu} \cos n\tau + \eta_{n_2}^{\mu} \sin n\tau \right) \left( \eta_{n_1}^{\nu} \cos n\tau + \eta_{n_2}^{\nu} \sin n\tau \right) n^2 \sin^2 n\sigma. \end{aligned} \quad (8)$$

We see that for  $\Psi_M$  having a nontrivial solution,  $m$  must be zero and  $\eta_{n_1}^{\mu} = \eta_{n_2}^{\mu} (= \eta_n^{\mu})$  (or  $\eta_{n_1}^{\mu}, \eta_{n_2}^{\mu} = 0$ ). In this case, for the oscillating mode  $n$ , every component of  $\Psi_M$  should satisfy the following equation:

$$\frac{\partial^2 \Psi_i}{\partial X^{\mu} \partial X^{\nu}} \Gamma_n^{\mu\nu} = 0, \quad (9)$$

where  $\Gamma_n^{\mu\nu} = \eta_n^{\mu} \eta_n^{\nu}$ .

Eq. (9) is the differential equation of  $\Psi_i$  about the spacetime coordinates  $X^{\mu}$ . In a sense, in this equation, the inner freedom of the string is omitted for the time being. However, the

inner motion of the string will give a strong confinement on the above equation and even the properties of the spacetime. There are several results we can deduce from the above equation: (i)  $\Gamma_n^{\mu\nu}$  is a second-rank symmetric tensor. For this sort of symmetric tensor, after a matrix transformation, it becomes a diagonal matrix which can be regarded as exactly the metric tensor  $g^{\mu\nu}$  ( $g^{\mu\nu} = U^T \Gamma_n^{\mu\nu} U$  with  $U$  a transformation matrix) of the spacetime. We see that the geometry of the spacetime is determined by the oscillating properties of the string due to the fact that the magnitude of the oscillating string determines the metric tensor of the spacetime. The invariance of Eq. (9) under Lorentz transformation is very obvious based on the analysis here. (ii) The mass of this sort of string is zero, even for any oscillating mode of the string. Thus, Eq. (9) is exactly the well-known Klein-Gordon equation describing massless bosons. (iii) From Eqs. (4) and (6), when various oscillating modes coexist, it is not difficult to confirm that the magnitude of various oscillating modes must be identical and therefore correspond to identical spacetime geometry. If various oscillating modes of the string are regarded as various type of particles, the researches here show that various type of particles must experience identical spacetime geometry. A character obtained directly is the quantization of the magnitude of various oscillating modes. Another implication would be the pith of the general relativity, i.e., for any type of particles, its motion can be described by the pure geometry of the spacetime.

In the case of a closed string with the periodicity condition  $X^\mu(\tau, \sigma) = X^\mu(\tau, \sigma = \pi)$ , it is straightforward to confirm that the discussions and results of the open string hold also for the closed string. For both open and closed strings, when the second-order partial differential equation of the string wave function is used, the spin would be an integer number by calculating the angular momentum of the string, i.e.,  $\Psi_M$  given by Eq. (1) describes massless bosons. In addition, these discussions and results hold too when the general membrane wave function is investigated.

Now we turn to investigate the first-order partial differential equation of  $\Psi_M$ . In this situation, we have

$$\Lambda_1 \frac{\partial \Psi_M}{\partial X^\mu} \frac{\partial X^\mu}{\partial \tau} + \Lambda_2 \frac{\partial \Psi_M}{\partial X^\mu} \frac{\partial X^\mu}{\partial \sigma} = m \Psi_M. \quad (10)$$

To have a solution of  $\Psi_M$ ,  $X^\mu$  should satisfy  $\partial X^\mu / \partial \tau = \text{const}$  and  $\partial X^\mu / \partial \sigma = \text{const}$ . In this situation, if the boundary condition of the open or closed strings is used, we have

$$X^\mu(\tau, \sigma) = \xi_0^\mu + \xi_1^\mu \tau. \quad (11)$$

From the above equation, Eq. (10) takes the form

$$\Lambda^\mu \frac{\partial \Psi_M}{\partial X^\mu} = m \Psi_M, \quad (12)$$

where  $\Lambda^\mu = \Lambda_1 \xi_1^\mu$ .

From the form of  $\Lambda^\mu$ , we can see clearly the covariant property of the matrixes  $\Lambda^\mu$ . We know Eq. (12) is the well-known Dirac equation which is used to describe the fermions with nonzero mass. Note that in the investigation here about fermions, there is in fact a wrong discussion. The matrix in Eq. (12) is  $\Lambda^\mu = \Lambda_1 \xi_1^\mu$ . It is obvious that  $\Lambda^\mu$  has only one linear independent matrix, while in the Dirac equation, four linear independent matrixes are needed to show the spin freedom of fermions. To generate four linear independent matrixes, the membrane wave function should be  $\Psi_M^T = (\Psi_1(X^\mu(\tau, \sigma_1, \sigma_2, \sigma_3)), \dots, \Psi_M(X^\mu(\tau, \sigma_1, \sigma_2, \sigma_3)))$ . In this situation,  $X^\mu(\tau, \sigma_1, \sigma_2, \sigma_3) = \xi_0^\mu + \xi_1^\mu \tau + \xi_2^\mu \sigma_1 + \xi_3^\mu \sigma_2 + \xi_4^\mu \sigma_3$  and  $\Lambda^\mu = \Lambda_1 \xi_1^\mu + \Lambda_2 \xi_2^\mu + \Lambda_3 \xi_3^\mu + \Lambda_4 \xi_4^\mu$ . For this sort of three-dimensional membrane to exist in the world we live, the dimensionality of the spacetime must be larger than or equal to four (three-dimensional space and one-dimensional time). Note that when the case of fermions is discussed, the non-trivial solution is a nonoscillating string. From the result given by  $X^\mu(\tau, \sigma_1, \sigma_2, \sigma_3)$ , there is not a boundary condition for the membrane. In fact, the size of the membrane can be finite or even infinite. This will not lead to conceptual difficulty when the membrane is regarded as an inner structure of the membrane wave function. We know for the wave function of a point particle, the wave function can be nonzero in the whole world and this does not lead to any problem. The membrane of the fermions discussed here is a nonoscillating one. Maybe this would make superstring theorists a little distressed. Nevertheless, the membrane

can still oscillate in the extra warped space, but not in the macroscopic four-dimensional spacetime. We see that, on the one hand the finite size of the membrane is necessary to generate the Dirac equation, and on the other hand this unique characteristic can be used exactly to overcome the well-known ultraviolet divergences .

In conclusion, the membrane wave function is proposed and used to give an unified description of bosons and fermions. The general wave function of the membrane (and string) proposed here is quite different from the wave functional description of the membrane in the ordinary string field theory, and can be regarded as a first-quantized string theory. Starting from the membrane wave function, the fundamental property of the spacetime can be investigated in a concise way. These results show that membrane (and string) is necessary to describe the world we live, even when the problem of the quantum gravity is not considered. Of course we do not propose a theory of everything, what we try to argue here is that it is very useful to regard the membrane as an inner structure of the general membrane wave function, and expect that this new starting point would be helpful to answer the question what the membrane is. Obviously, the strength of the superstring and beautiful mathematics of the warped high-dimensional spacetime developed in the last twenty years would still play an important role in the theory proposed here. In addition, by investigating the general membrane wave function, it would be an interesting future work to reconsider the quantum-mechanical postulates.

#### **ACKNOWLEDGMENTS**

This work was supported by Natural Science Foundation of China under grant Number 10205011.

## REFERENCES

- [1] M.B. Green, J.H. Schwarz, E. Witten, Superstring Theory (Cambridge: Cambridge University Press, 1987).
- [2] J. Polchinski, String Theory (Cambridge: Cambridge University Press, 1998).
- [3] C.M. Hull, P.K. Townsend, Nucl. Phys. B 438 (1995) 109.
- [4] P.K. Townsend, Phys. Lett. B 350 (1995) 184.
- [5] E. Witten, Nucl. Phys. B 443 (1995) 85.