

# Fate of the Born-Infeld solution in string theory

G. KARATHEODORIS, A. PINZUL AND A. STERN

Department of Physics, University of Alabama,  
Tuscaloosa, Alabama 35487, USA

## ABSTRACT

We argue that the Born-Infeld solution on the D9-brane is unstable under inclusion of derivative corrections to Born-Infeld theory coming from string theory. More specifically, we find no electrostatic solutions to the first order corrected Born-Infeld theory on the D9-brane which give a finite value for the Lagrangian.

The Born-Infeld (BI) nonlinear description of electrodynamics is of current interest due to its role as an effective action for Dp-branes.[1],[2] One of its main features is the existence of a maximum allowed value  $|E|_{max}$  for electrostatic fields.\* In the 30's, Born and Infeld promoted this theory because it has a spherically symmetric solution with finite classical self-energy.[3] It agrees with the Coulomb solution at large distances, but has a finite limit for the radial component of the electric field at the origin. This limit is just  $|E|_{max}$ . As a result, the vector field is singular at the origin. Equivalently, the solution is thus defined on a Euclidean manifold with one point (corresponding to the 'source') removed, despite the fact that the energy density is well behaved at the point. The theory contains one dimensionful parameter (namely,  $|E|_{max}$ ), which was determined by Born and Infeld after fixing the classical self-energy with the electron mass. For the Dp-brane the dimensionful parameter is the string tension, and so any BI-type solution (or BIon[4]) appearing there should correspond to a charged object with characteristic mass at the string tension. The fact that energetics admits vector fields that are not everywhere defined indicates that BI theory cannot be a complete description.[5] Moreover, in string theory, the BI effective action is only valid for slowly varying fields, i.e. it is the lowest order term in the derivative expansion for the full effective Dp-brane action. Since derivatives of the fields are not 'small' in the interior of the BI solution, the validity of such a solution in string theory can be questioned. It is therefore of interest to know whether or not analogues of the BI solutions survive for the full effective Dp-brane action. A derivative expansion has been recently carried out to obtain lowest order string corrections to the BI action for the space-filling D9-brane.[6],[7] With the restriction to electrostatic configurations, we find that the BI solution on the D9-brane is unstable with the inclusion of such corrections, indicating that such singular field configurations may not follow from the full effective action. The reason is basically due to the result that derivative corrections make it difficult for the electrostatic field to attain its maximum value  $|E|_{max}$ . Here we only allow for field configurations that lead to a finite value for the Lagrangian.† This is a reasonable requirement from the point of view of the path integral, where one expects to recover the classical solutions in the WKB approximation. Moreover, we find no nontrivial electrostatic solutions on the D9-brane associated with a finite Lagrangian.

The situation here is in contrast to skyrmion physics, where there are no nontrivial solutions to the zeroth order effective action for QCD. Higher order derivative corrections, like the Skyrme term, are necessary to stabilize the skyrmion. On the other hand, BIONS appear at lowest order, but become unstable upon including the next order electrostatic corrections. There remains the possibility, however, of stabilizing the BION with the inclusion of other degrees of freedom. For example, if we allow for magnetic effects one might find that the higher order corrections give a magnetic dipole moment to the BION. Another possibility which is of current interest, concerns BI-type solutions that appear after dimensional reduction. In this

---

\*Here we don't consider the excitation of transverse modes of the brane.

†We are unable to make a similar requirement for the energy, due to the fact that the corrections to the action contain terms with multiple time derivatives, making it problematic to find the corresponding Hamiltonian.

case, the BI action is replaced by the Dirac-Born-Infeld action (DBI), containing degrees of freedom associated with the transverse modes of the brane. Classical solutions to the DBI action were found in [4],[8], [9], and they represent fundamental strings attached to branes.<sup>‡</sup> It is of interest to examine how such solutions are affected by derivative corrections[6]. (One set of solutions (the BPS solutions) were found to be unaffected to all orders[10].) We hope to address these issues in future works.

We begin with a review of the Born-Infeld electrostatics. The BI action is expressed in terms of the determinant of the matrix with elements

$$h_{\mu\nu} = \eta_{\mu\nu} + (2\pi\alpha')F_{\mu\nu} , \quad (1)$$

where  $F_{\mu\nu} = \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu$  is the field strength and one assumes the flat metric  $\eta_{\mu\nu}$ . On the D9-brane it is given by

$$\mathcal{S}_{BI}^{(0)} = \frac{1}{(4\pi^2\alpha')^5 g_s} \int d^{d+1}x \mathcal{L}_{BI}^{(0)} , \quad \mathcal{L}_{BI}^{(0)} = 1 - \sqrt{-\det[h_{\mu\nu}]} , \quad (2)$$

where  $d$  is 9. In the absence of magnetic fields  $\mathcal{L}_{BI}^{(0)}$  simplifies to

$$\mathcal{L}_{BI}^{(0)} = 1 - \sqrt{1 - \vec{f}^2} , \quad (3)$$

where  $\vec{f} = 2\pi\alpha'\vec{E}$ , and  $\vec{E}$  is the electric field strength. It is only defined for  $|\vec{E}|$  below  $|E|_{max} = (2\pi\alpha')^{-1}$ . Field strengths above this critical value are said to be associated with string instabilities. For nine dimensional spherically symmetric electrostatic configurations, we define the angular variable  $\theta$ , taking values between  $\pm\pi/2$ . It is a function of the radial coordinate, with  $\vec{f} = \hat{r} \sin\theta(r)$ . Then the Lagrangian can be written

$$L_{BI}^{(0)} = \int d^d x \mathcal{L}_{BI}^{(0)} = \Omega^{d-1} \int dr r^{d-1} \left( 1 - \cos\theta(r) \right) , \quad (4)$$

where  $\Omega^{d-1}$  is the volume of a unit  $d-1$ -sphere. The relevant degree of freedom is the potential  $\mathcal{A}_0(r)$ , where  $\sin\theta(r) = \mathcal{A}'_0(r)$ , the prime denoting a derivative in  $r$ . The action is extremized with respect to  $\mathcal{A}_0$  by

$$\theta(r) = \tan^{-1} \frac{Q}{r^{d-1}} . \quad (5)$$

The integration constant  $Q$  is the charge. (5) approaches the Coulomb solution when  $r \rightarrow \infty$ . When  $r \rightarrow 0$ ,  $\theta \rightarrow \frac{\pi}{2} \text{sign}(Q)$ , and so unlike in Maxwell theory, the Lagrangian density has a finite value at the location of the source.

Concerning the energy, one can apply the canonical formalism starting from the action (2). The Hamiltonian density is

$$\mathcal{H}_{BI}^{(0)} = \Pi^i \partial_0 \mathcal{A}_i - \mathcal{L}_{BI}^{(0)}$$

---

<sup>‡</sup>As no such interpretation is possible for the BIon on the D9-brane, it is convenient that we find it to be unstable.

$$= \Pi^i F_{0i} + \sqrt{-\det[h_{\mu\nu}]} - 1 + \Pi^i \partial_i \mathcal{A}_0, \quad (6)$$

where  $i = 1, 2, \dots, d$ ,

$$\Pi^i = -\pi\alpha' \sqrt{-\det[h_{\mu\nu}]} (h^{i0} - h^{0i}), \quad (7)$$

are the momenta conjugate to  $\mathcal{A}_i$  and  $h^{\mu\nu} h_{\nu\rho} = \delta_\rho^\mu$ . As usual, the momentum conjugate to  $\mathcal{A}_0$  is constrained to be zero. In the absence of magnetic fields, (7) reduces to

$$\vec{\pi} = \frac{\vec{\Pi}}{2\pi\alpha'} = \frac{\vec{f}}{\sqrt{1 - \vec{f}^2}}, \quad (8)$$

and so substituting into (6) gives

$$\begin{aligned} \mathcal{H}_{BI}^{(0)} &= \sqrt{1 + \vec{\pi}^2} - 1 - (2\pi\alpha') \partial_i \pi^i \mathcal{A}_0 \\ &= \frac{1}{\sqrt{1 - \vec{f}^2}} - 1 - (2\pi\alpha') \partial_i \pi^i \mathcal{A}_0, \end{aligned} \quad (9)$$

where we integrated by parts. The coefficient of  $\mathcal{A}_0$  gives the Gauss law constraint. The remaining terms can be used to identify the self-energy of the BI solution

$$E_{BI}^{(0)} = \frac{\Omega^{d-1}}{(4\pi^2\alpha')^5 g_s} \int dr r^{d-1} \left( \sec \theta(r) - 1 \right), \quad (10)$$

which is finite, despite the fact that the vector field  $\vec{f}$  is ill-defined at the origin.

In addition to the nine dimensional spherically symmetric solution, there are also ‘axially symmetric’ BI solutions. They correspond to (5) with  $d < 9$ . Then instead of a point singularity on the  $D9$ -brane, there would be  $(9 - d)$ -dimensional Euclidean surface where  $\vec{f}$  is singular. In that case, (10) is the energy per unit volume along the surface, and  $r$  is the distance from the singular surface.

A derivative expansion has been carried out in [6],[7] to obtain corrections to the BI action for the  $D9$ -brane. At first order, one obtains terms involving first and second derivatives of  $F_{\mu\nu}$ . They are contained in the rank-4 tensor

$$S_{\beta\gamma\nu\rho} = 2\pi\alpha' \partial_\beta \partial_\gamma F_{\nu\rho} + (2\pi\alpha')^2 h^{\alpha\delta} (\partial_\beta F_{\nu\alpha} \partial_\gamma F_{\rho\delta} - \partial_\beta F_{\rho\alpha} \partial_\gamma F_{\nu\delta}), \quad (11)$$

which is antisymmetric in the last two indices. Up to first order the action is

$$\begin{aligned} \mathcal{S}_{BI}^{(0)} + \mathcal{S}_{BI}^{(1)} &= \frac{1}{(4\pi^2\alpha')^5 g_s} \int d^{10}x \left\{ 1 - \sqrt{-\det[h_{\mu\nu}]} \left( 1 + \frac{\kappa}{4} \Delta \right) \right\}, \\ \Delta &= h^{\mu\nu} h^{\rho\sigma} h^{\alpha\beta} h^{\gamma\delta} (S_{\nu\rho\alpha\beta} S_{\sigma\mu\gamma\delta} - 2S_{\beta\gamma\nu\rho} S_{\delta\alpha\sigma\mu}), \end{aligned} \quad (12)$$

where  $\kappa = \frac{(2\pi\alpha')^2}{48}$ . The derivatives  $\partial_\mu$  appearing in (11) must be covariant derivatives with respect to diffeomorphism in the ten dimensions space-time for the action to be invariant

under such diffeos.<sup>§</sup> We avoid this complication and work in Cartesian coordinates  $x_i$ . For electrostatic fields  $\vec{E}(\vec{x}) = \vec{f}(\vec{x})/(2\pi\alpha')$ ,

$$\begin{aligned} S_{ijk0} = -S_{ij0k} &= \partial_i \partial_j f_k + \frac{f_\ell}{1 - \vec{f}^2} (\partial_i f_k \partial_j f_\ell + \partial_i f_\ell \partial_j f_k) \\ S_{ijk\ell} &= -\frac{1}{1 - \vec{f}^2} (\partial_i f_k \partial_j f_\ell - \partial_i f_\ell \partial_j f_k) \end{aligned} \quad (13)$$

Then exploiting the symmetry of the indices we can write

$$-\frac{1}{4}\Delta = \frac{1}{2}S_{ijk\ell}S^{ijk\ell} + \frac{1}{1 - \vec{f}^2}S_{ijk0}S^{ijk}_0, \quad (14)$$

where the spatial indices are raised using  $h^{ij} = \delta_{ij} + f_i f_j / (1 - \vec{f}^2)$ .

Next we again assume spherical symmetry  $\vec{f} = \hat{r} \sin \theta(r)$ ,  $-\pi/2 \leq \theta \leq \pi/2$ . After substituting into (13)

$$\begin{aligned} S_{ijk\ell}S^{ijk\ell} &= \frac{2(d-1)}{r^4} \frac{\tan \theta}{\cos^3 \theta} [2(rH)^2 + (d-2)\sin^2 \theta] \\ S_{ijk0}S^{ijk}_0 &= \frac{1}{\cos^4 \theta} \left[ H'^2 + \frac{d-1}{r^4} \left( 2(rH - \sin \theta)^2 + (rH \cos^2 \theta - \sin \theta)^2 \right) \right], \end{aligned}$$

where  $H = \theta' / \cos \theta$  and again  $d = 9$ . Then the first order (in  $\kappa$ ) corrected Lagrangian is given by

$$\begin{aligned} L_{BI}^{(0)} + L_{BI}^{(1)} &= \int d^{d-1}x (\mathcal{L}_{BI}^{(0)} + \mathcal{L}_{BI}^{(1)}) \\ &= \Omega^{d-1} \int dr r^{d-1} \left\{ 1 - \cos \theta + \frac{\kappa}{\cos^3 \theta} \left( H'^2 + \frac{d-1}{r^4} \Sigma \right) \right\}, \end{aligned} \quad (15)$$

where

$$\Sigma = 3(rH - \sin \theta)^2 + (rH \sin^2 \theta)^2 + 2rH \sin^3 \theta + (d-2)\sin^4 \theta \quad (16)$$

The correction to the BI Lagrangian is positive (assuming  $-\pi/2 \leq \theta \leq \pi/2$ ), and so the appearance of the  $\cos^3 \theta$  in the denominator makes it challenging to find solutions having a limiting value of  $\pm\pi/2$  for  $\theta$  and a finite value for  $L_{BI}^{(0)} + L_{BI}^{(1)}$ . The action is extremized with respect to the potential  $\mathcal{A}_0(r)$ , where again  $\sin \theta(r) = \mathcal{A}'_0(r)$ , for

$$\begin{aligned} \frac{1}{\kappa}(Q - r^{d-1} \tan \theta) \cos^2 \theta &= 2 \left( \frac{r^{d-1} H'}{\cos^3 \theta} \right)'' + 3 \frac{r^{d-1} \sin \theta H'^2}{\cos^3 \theta} \\ &\quad - \frac{d-1}{4 \cos^3 \theta} \left\{ [27 - 4 \cos 2\theta + \cos 4\theta](r^{d-3} H)' \right. \\ &\quad \left. + \frac{1}{2} r^{d-5} \sin \theta \left[ 126 - 53d + 85r^2 H^2 + 4 \cos 2\theta(5 + d - 3r^2 H^2) + \cos 4\theta(d - 2 - r^2 H^2) \right] \right\} \end{aligned} \quad (17)$$

We note that by rescaling  $r$  and  $Q$  we can set  $\kappa$  equal to unity. Also (17) is invariant under  $Q \rightarrow -Q$  and  $\theta(r) \rightarrow -\theta(r)$ , and this gives a prescription for mapping any possible charged

<sup>§</sup>This fact was overlooked in previous versions of the paper.

solution to the anti-solution. The left hand side of (17) vanishes for the original BI solution (5), while the right hand side represents derivative corrections. Substituting (5) into the left hand side gives a vanishingly small correction as  $r \rightarrow \infty$ , but it is singular for  $r \rightarrow 0$ . The BI solution therefore cannot be trusted near the origin.

Due to the presence of high order time derivatives in the correction terms in the action (12), the computation of the Hamiltonian for the system, and consequently the correction to the electrostatic energy (10), is problematic. For this reason we only require solutions to (17) to be associated with a finite value for the Lagrangian (15) instead of the energy. Since the BI (actually, Coulumb) solution is valid at large  $r$ ,

$$\theta(r) \rightarrow \frac{Q}{r^{d-1}}, \quad \text{as } r \rightarrow \infty, \quad (18)$$

we can use this as an initial condition at some large value of  $r$  and numerically integrate to small  $r$ . We find that for  $d = 9$  and  $Q > 0$ ,  $\theta(r)$  tends to the limit of  $\pi/2$  as  $r \rightarrow 0$ , as with the BI solution. However, the corresponding radial Lagrangian density diverges as  $r \rightarrow 0$  (faster than  $1/r$ ). We plot the results below for  $Q = 1$  (and  $\kappa = 1$ ):




fig.1  $\theta(r)$  vs.  $r$




fig.2  $r^9(\mathcal{L}_{BI}^{(0)} + \mathcal{L}_{BI}^{(0)})$  vs.  $r$

In Fig.2 we plot the radial Lagrangian density times  $r$ . It blows up at the origin, and so it appears that this solution does not lead to a finite value of the Lagrangian. On the other hand, the numerical integration procedure breaks down near the origin. A more careful analysis requires that we look for a solution near the origin, consistent with the requirement of a finite Lagrangian, and try to match it with the above solution for some range of  $r$ . The requirement of a finite Lagrangian means that the limit of  $\theta(r) \rightarrow \pi/2$  as  $r \rightarrow 0$  has to be approached slow enough so that the integral over the correction term in (15) is finite. Upon writing  $\theta(r) = \pi/2 - \epsilon(r)$ , and keeping only the lowest order terms in  $\epsilon$ , (17) reduces to

$$(\epsilon Q - r^{d-1}) \epsilon^4 = 2\epsilon^3 \left( \frac{r^{d-1} H'}{\epsilon^3} \right)'' + 3r^{d-1} H'^2 - (d-1) \left\{ 8(r^{d-3} H)' + r^{d-5} [13 - 7d + 12r^2 H^2] \right\}, \quad (19)$$

where  $H' \rightarrow -(\log \epsilon)''$  and we again set  $\kappa$  equal to one. Taking  $d = 9$  and  $Q > 0$ , one has the following solution near the origin

$$\epsilon(r) \rightarrow \left( \frac{76432 r^4}{125 Q} \right)^{\frac{1}{5}}, \quad \text{as } r \rightarrow 0 \quad (20)$$

It is easy to check that it gives a finite contribution to the Lagrangian as  $r \rightarrow 0$ . (For this one only needs that  $\epsilon(r)$  goes to zero slower than  $r^{5/3}$ .) On the other hand, after numerically integrating this solution starting from some initial value  $r_0$  to increasing values of  $r$  we find that for any value of  $Q$ ,  $\epsilon$  goes quickly to  $\pi/2$  or  $-\pi/2$ , and so the Lagrangian density is poorly behaved for large values of  $r$ . Below we plot the results for  $Q = 1$ :




fig.3  $\theta(r)$  vs.  $r$




fig.4  $r^8(\mathcal{L}_{BI}^{(0)} + \mathcal{L}_{BI}^{(0)})$  vs.  $r$

From fig. 3,  $\theta' > 0$  to the right of the minimum, and so no match with fig. 1 is possible there. To the left of the minimum,  $\theta'$  tends to  $-\infty$  as  $r \rightarrow 0$  according to (20), while from fig. 1 it appears to vanish in the limit. To make matters worse, the graphs we obtain for  $\theta$  near the origin are highly sensitive to the initial value  $r_0$  of the integration, and this persists for all values of  $Q$ . We have checked that this is not due to the neglect of higher order terms in (20). So not only is there no agreement between the two numerical integration procedures, the validity of the solution near the origin is questionable. We are thus unable to find any spherically symmetric charged solutions to (17) consistent with the requirement of a finite Lagrangian.

We also find no axially symmetric charged solutions with finite Lagrangian density everywhere. Again, they correspond to  $d < 9$ , and at zeroth order were characterized by a  $(9 - d)$ -dimensional Euclidean surface where  $\vec{f}$  is singular. Using (18) we can once again numerically integrate to small values of  $r$ . The conclusions are the same as for  $d = 9$ . For  $Q > 0$ ,  $\theta(r)$  tends to the limit of  $\pi/2$  as  $r \rightarrow 0$ , and the corresponding radial Lagrangian density diverges as  $r \rightarrow 0$  (faster than  $1/r$ ). For  $Q > 0$  and  $d \geq 6$ , the solution to (19) near the origin has the form

$$\epsilon(r) \rightarrow \left( \frac{Cr^{d-5}}{125Q} \right)^{\frac{1}{5}}, \quad \text{as } r \rightarrow 0, \quad (21)$$

where  $C = 20008, 33324, 51836$  for  $d = 6, 7, 8$ , respectively. As with the case  $d = 9$ , it is consistent with the requirement of a finite Lagrangian for small  $r$ , but the Lagrangian density diverges after numerically integrating to increasing large  $r$ , and the problem of the sensitivity to the initial value  $r_0$  persists. So again we find no agreement between the two numerical integration procedures. For  $d < 6$ , we find no solutions to (19) near the origin satisfying the requirement of a finite Lagrangian.

Above we found negative results for all configurations associated with some nonzero value for the charge. Although there are no  $Q = 0$  soliton solutions of the zeroth order Born-Infeld system, this need not be a priori true at higher orders. In this case,  $\theta(r)$  for any spherically symmetric solution would not fall off as a power as  $r \rightarrow \infty$ , but rather

$$\theta(r) \rightarrow \left\{ A \cos \alpha r + B \sin \alpha r \right\} \frac{e^{-\alpha r}}{r^{(d-1)/4}}, \quad \text{as } r \rightarrow \infty, \quad (22)$$

where  $\alpha = 8^{-1/4}$ ,  $A$  and  $B$  are integration constants and we again set  $\kappa = 1$ . Here we look for nonsingular soliton solutions, and so  $\theta(r)$  should go to zero as  $r \rightarrow 0$ . Such configurations then give a well defined vector field at the origin. Two types of behavior are allowed near the origin:

$$\theta(r) \rightarrow Cr \quad \text{or} \quad \theta(r) \rightarrow Dr^3, \quad \text{as } r \rightarrow 0, \quad (23)$$

where  $C$  and  $D$  are integration constants. Next one can numerically integrate (17) starting from both (22) and (23), and attempt to match  $\theta(r)$  and its derivatives for some range of  $r$ . We found no values for the constants  $A$ ,  $B$  and  $C$  or  $D$  where this is possible.

### Acknowledgement

We are grateful for correspondence with Koji Hashimoto. This work was supported by the joint NSF-CONACyT grant E120.0462/2000 and the U.S. Department of Energy under contract number DE-FG05-84ER40141.

### REFERENCES

- [1] E. S. Fradkin and A. A. Tseytlin, Phys. Lett. **B163** (1985) 123; A. Abouelsaood, C. G. Callan, C. R. Nappi, and S. A. Yost, Nucl. Phys. **B280** (1987) 599; E. Bergshoeff, E. Sezgin, C. N. Pope, and P. K. Townsend, Phys. Lett. **188B** (1987) 70; R. G. Leigh, Mod. Phys. Lett. **A4** (1989) 2767.
- [2] For reviews, see C. P. Bachas, arXiv:hep-th/9806199; A. A. Tseytlin, arXiv:hep-th/9908105; C. V. Johnson, arXiv:hep-th/0007170; I. V. Vanea, arXiv:hep-th/0109029; R. J. Szabo, arXiv:hep-th/0207142.
- [3] M. Born and L. Infeld, Proc. Roy. Soc. London **A144** 425 (1934).
- [4] G. W. Gibbons, Nucl. Phys. B **514**, 603 (1998).
- [5] E. Feenberg, Phys. Rev. **47**, 148 (1935); H. L. M. Pryce, Proc. Roy. Soc. **A155**, 597 (1936); A. A. Chernitsky, Helv. Phys. Acta **71**, 274 (1998); D. Chruscinski, Phys. Lett. A **240**, 8 (1998).
- [6] N. Wyllard, Nucl. Phys. B **598**, 247 (2001); JHEP **0108**, 027 (2001).
- [7] S. R. Das, S. Mukhi and N. V. Suryanarayana, JHEP **0108**, 039 (2001).

- [8] C. G. Callan and J. M. Maldacena, Nucl. Phys. B **513**, 198 (1998).
- [9] P. S. Howe, N. D. Lambert and P. C. West, Nucl. Phys. B **515**, 203 (1998).
- [10] L. Thorlacius, Phys. Rev. Lett. **80**, 1588 (1998).