

# Abelianization of Constraints in SU(N) Yang-Mills Theory

F. Loran\*

*Department of Physics, Isfahan University of Technology (IUT)  
Isfahan, Iran*

## Abstract

The abelian form of the first class constraints of SU(N) Yang-Mills theory in  $D = 3 + 1$  is obtained by perturbation. Considering the abelian constraints, we show that Coulomb gauge does not lead to Gribov ambiguity.

It is well known that SU(N) Yang-Mills theory is a constraint system possessing non-abelian first class constraints  $\phi_a$ ,

$$\phi_a = \nabla \cdot \vec{\Pi}_a - gf_{abc} \vec{A}_b \cdot \vec{\Pi}_c \approx 0, \quad a = 1, \dots, N^2 - 1, \quad (1)$$

in which,  $f_{abc}$  are the structure coefficients of SU(N) algebra,  $\vec{\Pi}_a$ 's are momenta conjugate to gauge fields  $\vec{A}_a$ 's,

$$\{A_a^i(x), \Pi_b^j(y)\} = \delta_{ab} \delta^{ij} \delta(x - y), \quad i, j = 1, 2, 3, \quad (2)$$

and  $g$  is the coupling constant of the gauge field self interaction [1, 2]. The constraints  $\phi_a$ 's (1) form a representation of SU(N) algebra, i.e.

$$\{\phi_a^{g_1}, \phi_b^{g_2}\} = gf_{abc} \phi_c^{g_1 g_2}, \quad (3)$$

where  $g_1$  and  $g_2$  are smooth real functions and  $\phi_a^g = \int_x g(x) \phi_a(x)$ . In 1978, Gribov showed that Coulomb gauge  $\nabla \cdot \vec{A}_a = 0$  is insufficient to fix the gauge freedom of the action generated by non-abelian constraints  $\phi_a$ 's (1). He observed that for SU(2) Yang-Mills theory there exist at least two points on the gauge orbit that satisfy Coulomb gauge [3]. This effect is in general called Gribov ambiguity and became a serious drawback for the quantization of Yang-Mills theory. Gribov ambiguities can be remedied in the following three ways:

1. Considering proper gauge fixing condition like the axial gauge  $A_a^3 = 0$ , that does not lead to Gribov ambiguity [4],
2. Applying quantization approaches in which there is no need to do gauge fixing [5],
3. Using the celebrated BRST-BFV approach [6, 7] where one considers BRS transformation instead of gauge transformation. For a review see reference [2].

---

\*e-mail: loran@cc.iut.ac.ir

One may add another approach to the above list which we discuss here.

It is proved that non-abelian constraints become abelian if one maps each constraint to the surface of the other ones [8]. To be explicit, consider two independent constraints  $\phi$  and  $\psi$  which satisfy the following algebra:

$$\{\phi, \psi\} = C\phi + D\psi, \quad (4)$$

where  $C$  and  $D$  are some functions of phase space coordinates. One can show that the constraints  $\phi' = \phi|_{(\psi=0)}$  and  $\psi' = \psi|_{(\phi=0)}$  are equivalent to  $\phi$  and  $\psi$  and commute with each other, i.e.

$$\{\phi', \psi'\} = 0. \quad (5)$$

The non-abelian constraints  $\phi_a$ 's (1) can be made abelian in a similar way. Using the Helmholtz theorem in vector analysis [9], one can write the vector  $\vec{\Pi}_a$  as,

$$\vec{\Pi}_a(x) = -\nabla \int_y \frac{\nabla \cdot \vec{\Pi}_a(y)}{4\pi |\vec{x} - \vec{y}|} + \nabla \times \int_y \frac{\nabla \times \vec{\Pi}_a(y)}{4\pi |\vec{x} - \vec{y}|}, \quad (6)$$

up to some surface terms. Inserting  $\vec{\Pi}_a$  from Eq.(6) into Eq.(1), one can obtain a set of new constraints, say  $\phi_a^1$ 's, equivalent to  $\phi_a$ 's (1), defined as follows,

$$\phi_a^1(x) = \nabla \cdot \vec{\Pi}_a - g f_{abc} \vec{A}_b \cdot \nabla \times \int_y \frac{\nabla \times \vec{\Pi}_c(y)}{4\pi |\vec{x} - \vec{y}|} + g^2 R_a^{(2)}(x), \quad (7)$$

where

$$\begin{aligned} R_a^{(2)}(x) &= \frac{1}{g^2} \left( g f_{abc} \vec{A}_b \cdot \nabla \int_y \frac{\nabla \cdot \vec{\Pi}_c(y)}{4\pi |\vec{x} - \vec{y}|} \right) \\ &= f_{abc} f_{cde} \vec{A}_b \cdot \nabla \int_y \frac{\vec{A}_d \cdot \vec{\Pi}_e}{4\pi |\vec{x} - \vec{y}|}. \end{aligned} \quad (8)$$

One should note that to obtain the second equality we have considered  $\phi_a = 0$  (1). It can be easily verified that  $\{\phi_a^1, \phi_b^1\} = \mathcal{O}(g^2)$ . Inserting  $\vec{\Pi}_a$  from Eq.(6) into Eq.(7) and using Eq.(1) again, one obtains  $\phi_a^2$ 's, a new set of constraints equivalent to  $\phi_a$ 's, which satisfy the following algebra

$$\{\phi_a^2, \phi_b^2\} = \mathcal{O}(g^3). \quad (9)$$

At  $N$ th step, one finds  $\phi_a^N$ 's,

$$\phi_a^N(x_1) = \nabla \cdot \vec{\Pi}_a(x_1) + \sum_{n=1}^N \left( \frac{-g}{4\pi} \right)^n P_a^{(n)}(x_1) + (-g)^{(N+1)} R_a^{(N+1)}(x_1), \quad (10)$$

where

$$P_a^{(n)}(x_1) = Q_{a,a_{n+1}}^{(n)}(x_1) \times \int_{x_{n+1}} \frac{\nabla_{x_{n+1}} \times \vec{\Pi}_{a_{n+1}}(x_{n+1})}{|\vec{x}_n - \vec{x}_{n+1}|}, \quad (11)$$

and

$$R_a^{(N+1)}(x_1) = f_{a_{N+1} b_{N+1} a_{N+2}} Q_{a,a_{N+1}}^{(N)}(x_1) \int_{x_{N+1}} \frac{\vec{A}_{b_{N+1}}(x_{N+1}) \cdot \vec{\Pi}_{a_{N+2}}(x_{N+1})}{|\vec{x}_N - \vec{x}_{N+1}|}, \quad (12)$$

in which

$$\begin{aligned} Q_{a,a_{n+1}}^{(n)}(x_1) &= f_{a_n,b_n,a_{n+1}} Q_{a,a_n}^{(n-1)}(x_1) \int_{x_n} \frac{1}{|\vec{x}_{n-1} - \vec{x}_n|} \vec{A}_{b_n}(x_n) \cdot \nabla_{x_n}, & n > 1, \\ Q_{a,a_2}^{(1)}(x_1) &= f_{ab_1a_2} \vec{A}_{b_1}(x_1) \cdot \nabla_{x_1}. \end{aligned} \quad (13)$$

Since

$$\phi_a^N = \phi_a^N|_{(\phi_b^N=0)} + \mathcal{O}(g^{N+1}), \quad b \neq a, \quad (14)$$

one verifies that  $\{\phi_a^N, \phi_b^N\} = \mathcal{O}(g^{N+1})$ , (see Eqs.(4,5)). In this way, non-abelian constraints  $\phi_a$ 's (1), can be made abelian by perturbation. The generator of gauge transformation is

$$G_\epsilon^N = \int_x \epsilon_a(x) \phi_a^N(x) + \mathcal{O}(g^{N+1}), \quad (15)$$

where  $\epsilon_a(x)$  is some infinitesimal real smooth function. Since  $\{\nabla \cdot \vec{A}_a(x), \nabla \times \vec{\Pi}_b(y)\} = 0$ , one verifies that

$$\begin{aligned} \delta_\epsilon (\nabla \cdot \vec{A}_a(x)) &= \{\nabla \cdot \vec{A}_a(x), G_\epsilon^N\} \\ &= \{\nabla \cdot \vec{A}_a(x), \int_y \epsilon_a(y) \nabla \cdot \vec{\Pi}_a(y)\} \\ &= -\nabla^2 \epsilon_a(x) + \mathcal{O}(g^{N+1}), \end{aligned} \quad (16)$$

for arbitrary  $N$ . We conclude that Coulomb gauge intersects the gauge orbit only once as is the case in QED. Therefore, Coulomb gauge does not lead to Gribov ambiguity. Of course some non-perturbative effects may emerge that should be added to the other non-perturbative effects in quantization of the theory.

It is interesting to calculate Ward identities corresponding to the gauge transformation generated by  $G_\epsilon^N$  (15). In this way one can compare Dirac quantization [10] by BRST quantization and verify their equivalence explicitly.

## Acknowledgement

The author would like to express his thanks to M. Haghghat for useful discussions.

## References

- [1] K. Sundermeyer, "Constrained Dynamics, Lecture Notes in Physics" Springer-Verlag, Berlin-Heidelberg, 1982;  
M. Henneaux and C. Teitelboim "Quantization of Gauge System" Princeton University Press, Princeton, New Jersey, 1992.
- [2] M. Henneaux, Phys. Rep. **126**, (1985) 1.
- [3] V. N. Gribov, Nucl. Phys. **B139**, (1978) 1;  
I. M. Singer, Comm. Math. Phys. **60**, (1978) 7.

- [4] R. Arnowitt and S. I. Fickler, Phys. Rev. **127**, (1962) 1821;  
R. N. Mohapatra, Phys. Rev. **D4**, (1971) 2215;  
I. Bars and F. Green, Nucl. Phys. **B142**, (1978) 157.
- [5] J. R. Klauder, Lect. Notes Phys. **572**, (2001) 143, hep-th/0003297;  
J. R. Klauder and S. V. Shabanov, Phys. Lett. **B398**, (1997) 116, hep-th/9608027;  
J. Govaerts and J. R. Klauder, Annals Phys. **274**, (1999) 251, hep-th/9809119;  
V. M. Villanueva, J. Govaerts, J-L. Lucio-Martinez, J.Phys. **A33**, (2000) 4183, hep-th/9909033.
- [6] C. Becchi, A. Rouet and R. Stora, Comm. Math. Phys. **42**, (1975) 127.
- [7] E. S. Fradkin and G. A. Vilkovisky, Phys. Lett. **B55**, (1975) 224;  
I. A. Batalin and G. A. Vilkovisky, Phys. Lett. **B69**, (1977) 309;  
E. S. Fradkin and E. S. Fradkina, Phys. Lett. **B72**, (1978) 343;  
I. A. Batalin and E. S. Fradkin, Phys. Lett. **B122**, (1983) 157.
- [8] F. Loran, Phys. Lett. **B547**, (2002) 63, hep-th/0209180;  
F. Loran, Phys. Lett. **B554**, (2003) 207, hep-th/0212341.
- [9] G. Arfken, *Mathematical Methods for Physicists*, second edition, Academic Press, Inc., New York, 1970.
- [10] P. A. M. Dirac, Can. J. Math. **2**, (1950) 129;  
Proc. R. Soc. London Ser. A **246**, (1958) 326;  
*"Lectures on Quantum Mechanics"*, Yeshiva University Press, New York, 1964.