

The scientific work of Sonia Stanciu

Constantin Bachas

Laboratoire de Physique Théorique de l' Ecole Normale Supérieure
24 rue Lhomond, 75231 Paris cedex, France

Abstract. The Leuven workshop on the 'Quantum Structure of Space-time and the Geometrical Nature of the Fundamental Interactions' had a special session dedicated to the memory of Sonia Stanciu. This is the summary of a talk delivered by the author on this occasion.

I was asked by the organizers to say a few words about the scientific life of Sonia Stanciu, a young string theorist and active member of this European Network, who passed away a few months ago after a short but devastating illness.

Sonia started her scientific career in Bonn, where she arrived from her native Romania to work on a doctoral degree under the supervision of Vladimir Rittenberg. Her thesis [1] was on hierarchies of integrable equations, in relation with what are nowadays referred to as 'old matrix models'. These describe certain types of matter fields living on random triangulated surfaces, *i.e.* interacting with regularized 2d quantum gravity. The models simplify considerably in the continuum or 'double scaling' limit [2], in which their partition and correlation functions can be obtained as solutions of differential equations. Matrix models thus raised the hope of a non-perturbative approach to string theory, but this was quickly frustrated by two important and still open problems: the approach is restricted to bosonic strings, and to 'spacetimes' with at most two dimensions.

Sonia studied supersymmetric extensions of the KdV hierarchy as a possible road towards lifting the above restrictions. Her thesis begins with Scott-Russel's remarkable description of his first encounter with a solitary wave in the Union canal [3] and continues with a nice introduction to the history of classical integrability. She then goes on to describe her own research, initiated in two articles with Jose Figueroa-O'Farrill [4]. Jose was in fact the real scientific mentor of Sonia's thesis. He would also become her life-long collaborator, and her husband.

Sonia spent the last year of her graduate studies visiting Queen Mary College in London, and then moved on to a postdoctoral position at the ICTP in Trieste. During this time she continued to collaborate with Jose, first on possible obstructions to gaugings of Wess-Zumino-Witten models [5], and then on σ -models based on non-semi-simple Lie algebras [6] which generalized the Nappi-Witten plane-wave background [7]. As with most of their other joint work, these articles were characterized by mathematical precision and by a special affinity for algebraic structures. They prepared the ground for the next class of problems that Sonia attacked, now as a mature physicist: the

embedding of D-branes in exact curved-spacetime backgrounds. This is the subject on which she had the highest impact, and the one on which she worked until the end.

String theory predicts two kinds of quantum corrections to classical geometry: string-loop corrections and α' corrections. Though dualities may in some cases exchange the two, the α' effects are more accessible thanks to the powerful algebraic techniques of conformal field theory. This is illustrated by the success of boundary CFT in describing the D-branes of certain exact (super)string backgrounds. In the semiclassical regime, D-branes are submanifolds which may carry a non-trivial gauge connection, and are solutions of the Dirac-Born-Infeld equations [8]. These equations, together with an infinite series of α' corrections are, on the other hand, equivalent to the condition of conformal invariance on the boundary of a string worldsheet,

$$T_{zz} = T_{\bar{z}\bar{z}} \quad \text{at} \quad \text{Re}z = 0 . \quad (1)$$

Here T_{ab} is the energy-momentum tensor, and (1) says that the net flow of energy to the boundary of the half-plane must be zero. The point is that in many exact CFTs, such as minimal models or WZW models, one can bypass the geometric intuition and solve the above equation algebraically. In this approach one must supplement the condition (1) with certain consistency or ‘sewing’ constraints [9, 10, 11] that translate the locality of the underlying field theory.

Sonia was among the first to realize that this was a very fertile and worth-pursuing direction. I remember listening to her talk at a meeting of Dutch particle theorists, at NIKHEF in the Fall of 1999. She was moving from Imperial College to the Spinoza Institute in Utrecht at that time, and was invited to present her results. I came out thinking that these were interesting, but that I would not want to work on the subject myself. Surely enough, this is what I was doing for the next couple of years. It gave me the opportunity to discuss with Sonia on many occasions, at conferences, via email, and during the Fall 2000 semester at the Institut Henri Poincaré.

I will not attempt to summarize the applications of BCFT, and the interplay of geometric and algebraic techniques, in the study of curved-spacetime D-branes in string theory. There exist already many complimentary reviews on this subject [12–16]. Here I will limit myself to some comments on Sonia’s contributions in the field.

The simplest but rich example of quantum geometry, which has served as prototype for many studies, is the WZW model based on the manifold of $SU(2)$. The algebraic construction of D-branes in this model has been known essentially since the pioneering work of Cardy [9], while their geometric interpretation was only clarified in recent years [17, 18, 19]. The algebraic construction proceeds by first imposing the symmetry-preserving boundary conditions on the affine currents :

$$J_z^a = -\Omega^a_b J_{\bar{z}}^b \quad \text{at} \quad \text{Re}z = 0 , \quad (2)$$

where Ω is an automorphism of the $SU(2)$ algebra. Because of the Sugawara form of the energy-momentum tensor, eqs. (2) imply automatically eq. (1). The sewing constraints can then be solved by Cardy’s prescription, valid for any rational CFT. In

the semiclassical regime these Cardy states correspond to D-branes which wrap spherical conjugacy classes, and are stabilized against shrinking by a non-trivial magnetic flux.

In a series of articles Sonia tried to extend these considerations in several directions : to general compact semi-simple Lie groups [20], to N=2 coset models [21], to the Nappi-Witten plane-wave background [22], and to the WZW model based on the non-compact group $SL(2,R)$ [23]. Each of them presents technical and conceptual challenges, which are still the subject of active research today (see for instance the talks of Angelos Fotopoulos, Lennaert Huiszoon and Marika Taylor in these proceedings). The $SL(2,R)$ WZW model is especially interesting because it is part of the near-horizon geometry of the stringy NS5/F1 black hole [24]. Sonia was the first to study the supersymmetric D-branes of this background from the CFT point of view. We continued her work with Marios Petropoulos, adding a new type of D-brane with anti-de-Sitter geometry to her list [25]. These new branes span twisted conjugacy classes of $SL(2,R)$, and are obtained when Ω is a non-trivial outer automorphism. The extension of the argument in [17] to the case of an outer automorphism was in fact also one of Sonia's nice remarks [20, 18]. The dual holographic description of these D-branes, and the construction of the corresponding boundary states, are challenging problems that are still only partially resolved (see for instance [26, 27] and references therein).

One of the last problems to interest Sonia was the definition of D-brane charge. In the $SU(2)$ WZW model this takes values in the finite group Z_{k+2} , where k is the level of the current algebra. Roughly speaking, the charge is the magnetic flux through the D-brane, defined modulo large gauge transformations of the Neveu-Schwarz B -field (see [28, 29] for various related 'physicist' arguments). A mathematically more precise formulation, in terms of relative cohomology, was given in the last paper of Sonia with Jose [30] (see also [31, 32, 33] for earlier work). In this approach one considers topological obstructions to defining the σ -model on open worldsheets. With hindsight, the appropriate mathematical framework in which to define the invariant D-brane charge is most likely some twisted version of K-theory [34]. It is, however, not yet clear how this ties together with the boundary-state CFT approach, even for the next example in the list, *i.e.* the WZW model on the group manifold of $SU(3)$ [35, 29, 36]. Sonia might have talked about this today, had illness not cut her life so short.

Most of us will remember Sonia as a quiet colleague, sharing a real passion for the field to which she has contributed in her own discreet and elegant way. She will be greatly missed in this and in all future network meetings.

References

- [1] S. Stanciu, "Supersymmetric integrable hierarchies and string theory," arXiv:hep-th/9407189.
- [2] E. Brezin and V. A. Kazakov, "Exactly Solvable Field Theories Of Closed Strings," Phys. Lett. B **236** (1990) 144 ; D. J. Gross and A. A. Migdal, "Nonperturbative Two-Dimensional Quantum Gravity," Phys. Rev. Lett. **64**, 127 (1990) ; M. R. Douglas and S. H. Shenker, "Strings In Less Than One-Dimension," Nucl. Phys. B **335**, 635 (1990).

- [3] J. Scott-Russel, “Report on waves,” Report of the 14th Meeting of the British Association for the Advancement of Science, John Murray, London, 1844, p.p. 311-390.
- [4] J. M. Figueroa-O’Farrill and S. Stanciu, “On a new supersymmetric KdV hierarchy in 2-d quantum supergravity,” *Phys. Lett. B* **316**, 282 (1993) [arXiv:hep-th/9302057] ; “New supersymmetrizations of the generalized KdV hierarchies,” *Mod. Phys. Lett. A* **8**, 2125 (1993) [arXiv:hep-th/9303168].
- [5] J. M. Figueroa-O’Farrill and S. Stanciu, “Equivariant Cohomology And Gauged Bosonic Sigma Models,” arXiv:hep-th/9407149 ; ‘ ‘Gauged Wess-Zumino terms and equivariant cohomology,” *Phys. Lett. B* **341**, 153 (1994) [arXiv:hep-th/9407196].
- [6] J. M. Figueroa-O’Farrill and S. Stanciu, “Nonreductive WZW models and their CFTs,” *Nucl. Phys. B* **458**, 137 (1996) [arXiv:hep-th/9506151] ; “Nonreductive WZW models and their CFTs, II: N=1 and N=2 cosets,” *Nucl. Phys. B* **484**, 583 (1997) [arXiv:hep-th/9605111].
- [7] C. R. Nappi and E. Witten, “A WZW model based on a nonsemisimple group,” *Phys. Rev. Lett.* **71**, 3751 (1993) [arXiv:hep-th/9310112].
- [8] J. Dai, R. G. Leigh and J. Polchinski, “New Connections Between String Theories,” *Mod. Phys. Lett. A* **4**, 2073 (1989) ; R. G. Leigh, “Dirac-Born-Infeld Action From Dirichlet Sigma Model,” *Mod. Phys. Lett. A* **4**, 2767 (1989).
- [9] J. L. Cardy, “Boundary Conditions, Fusion Rules And The Verlinde Formula,” *Nucl. Phys. B* **324**, 581 (1989).
- [10] J. L. Cardy and D. C. Lewellen, “Bulk And Boundary Operators In Conformal Field Theory,” *Phys. Lett. B* **259**, 274 (1991) ; D. C. Lewellen, “Sewing constraints for conformal field theories on surfaces with boundaries,” *Nucl. Phys. B* **372**, 654 (1992).
- [11] G. Pradisi, A. Sagnotti and Y. S. Stanev, “Planar duality in SU(2) WZW models,” *Phys. Lett. B* **354**, 279 (1995) [arXiv:hep-th/9503207]. “Completeness Conditions for Boundary Operators in 2D Conformal Field Theory,” *Phys. Lett. B* **381**, 97 (1996) [arXiv:hep-th/9603097].
- [12] M. R. Douglas, “Lectures on D-branes on Calabi-Yau manifolds,” *Prepared for ICTP Spring School on Superstrings and Related Matters, Trieste, Italy, 2-10 Apr 2001* ; see also “D-branes on Calabi-Yau manifolds,” arXiv:math.ag/0009209.
- [13] C. Schweigert, J. Fuchs and J. Walcher, “Conformal field theory, boundary conditions and applications to string theory,” arXiv:hep-th/0011109.
- [14] A. Recknagel, “Boundary conformal field theory and the world-sheet approach to D-branes,” Habilitation thesis, unpublished.
- [15] C. Angelantonj and A. Sagnotti, “Open strings,” *Phys. Rept.* **371**, 1 (2002) [arXiv:hep-th/0204089].
- [16] V. Schomerus, “Lectures on branes in curved backgrounds,” *Class. Quant. Grav.* **19**, 5781 (2002) [arXiv:hep-th/0209241].
- [17] A. Y. Alekseev and V. Schomerus, “D-branes in the WZW model,” *Phys. Rev. D* **60**, 061901 (1999) [arXiv:hep-th/9812193].
- [18] G. Felder, J. Frohlich, J. Fuchs and C. Schweigert, “The geometry of WZW branes,” *J. Geom. Phys.* **34**, 162 (2000) [arXiv:hep-th/9909030].
- [19] C. Bachas, M. R. Douglas and C. Schweigert, “Flux stabilization of D-branes,” *JHEP* **0005**, 048 (2000) [arXiv:hep-th/0003037].
- [20] S. Stanciu, “D-branes in group manifolds,” *JHEP* **0001**, 025 (2000) [arXiv:hep-th/9909163].
- [21] S. Stanciu, “D-branes in Kazama-Suzuki models,” *Nucl. Phys. B* **526**, 295 (1998) [arXiv:hep-th/9708166] ;
- [22] S. Stanciu and A. A. Tseytlin, “D-branes in curved spacetime: Nappi-Witten background,” *JHEP* **9806**, 010 (1998) [arXiv:hep-th/9805006] ; J. M. Figueroa-O’Farrill and S. Stanciu, “More D-branes in the Nappi-Witten background,” *JHEP* **0001**, 024 (2000) [arXiv:hep-th/9909164].
- [23] S. Stanciu, “D-branes in an AdS(3) background,” *JHEP* **9909**, 028 (1999) [arXiv:hep-th/9901122] ; J. M. Figueroa-O’Farrill and S. Stanciu, “D-branes in AdS(3) x S(3) x S(3) x S(1),” *JHEP* **0004**, 005 (2000) [arXiv:hep-th/0001199].

- [24] J. Maldacena and A. Strominger, “AdS(3) black holes and a stringy exclusion principle,” JHEP **9812**, 005 (1998) [arXiv:hep-th/9804085].
- [25] C. Bachas and M. Petropoulos, “Anti-de-Sitter D-branes,” JHEP **0102**, 025 (2001) [arXiv:hep-th/0012234].
- [26] C. Bachas, J. de Boer, R. Dijkgraaf and H. Ooguri, “Permeable conformal walls and holography,” JHEP **0206**, 027 (2002) [arXiv:hep-th/0111210].
- [27] B. Ponsot, V. Schomerus and J. Teschner, “Branes in the Euclidean AdS(3),” JHEP **0202**, 016 (2002) [arXiv:hep-th/0112198].
- [28] W. Taylor, “D2-branes in B fields,” JHEP **0007**, 039 (2000) [arXiv:hep-th/0004141] ; A. Alekseev, A. Mironov and A. Morozov, “On B-independence of RR charges,” Phys. Lett. B **532**, 350 (2002) [arXiv:hep-th/0005244] ; D. Marolf, “Chern-Simons terms and the three notions of charge,” arXiv:hep-th/0006117.
- [29] J. M. Maldacena, G. W. Moore and N. Seiberg, “D-brane instantons and K-theory charges,” JHEP **0111**, 062 (2001) [arXiv:hep-th/0108100].
- [30] J. M. Figueroa-O’Farrill and S. Stanciu, “D-brane charge, flux quantization and relative (co)homology,” JHEP **0101**, 006 (2001) [arXiv:hep-th/0008038] ; S. Stanciu, “A note on D-branes in group manifolds: Flux quantization and D0-charge,” JHEP **0010**, 015 (2000) [arXiv:hep-th/0006145].
- [31] C. Klimcik and P. Severa, “Open strings and D-branes in WZNW models,” Nucl. Phys. B **488**, 653 (1997) [arXiv:hep-th/9609112].
- [32] K. Gawedzki, “Conformal field theory: A case study,” arXiv:hep-th/9904145.
- [33] D. S. Freed and E. Witten, “Anomalies in string theory with D-branes,” arXiv:hep-th/9907189.
- [34] R. Minasian and G. W. Moore, “K-theory and Ramond-Ramond charge,” JHEP **9711**, 002 (1997) [arXiv:hep-th/9710230] ; E. Witten, “D-branes and K-theory,” JHEP **9812**, 019 (1998) [arXiv:hep-th/9810188]. P. Bouwknegt and V. Mathai, “D-branes, B-fields and twisted K-theory,” JHEP **0003**, 007 (2000) [arXiv:hep-th/0002023].
- [35] S. Fredenhagen and V. Schomerus, “Branes on group manifolds, gluon condensates, and twisted K-theory,” JHEP **0104**, 007 (2001) [arXiv:hep-th/0012164].
- [36] S. Stanciu, “An illustrated guide to D-branes in SU(3),” arXiv:hep-th/0111221.