

The Effective Average Action Beyond First Order

H. Ballhausen

Institute for Theoretical Physics, Heidelberg University
Philosophenweg 16, 69120 Heidelberg, Germany

A derivative expansion of the effective average action beyond first order yields renormalization group functional flow equations which are used for the computation of critical exponents of the Ising universality class. The critical exponent in $D = 3$ is consistent with high-precision methods.

The effective average action

Derivative expansion beyond first order

The effective average action $\Gamma_k[\phi]$ (see [1] and [2, 3, 4, 5]) is a functional of the fields ϕ . It interpolates between the classical action S at some microscopic ultraviolet scale and the effective action $\Gamma = S, \lim_{k \rightarrow 0} \Gamma_k = \Gamma$. Its dependence on the momentum scale k is described by an exact renormalization group functional flow equation:

$$\partial_k \Gamma_k = \frac{1}{2} \text{Tr} f \left(\Gamma_k^{(2)} + R_k \right)^{-1} \partial_k R_k g$$

where $\Gamma_k^{(2)}$ is the second functional derivative of Γ_k with respect to the fields and R_k denotes a momentum cutoff function with the properties $\lim_{k \rightarrow 0} R_k = 0$, $\lim_{k \rightarrow \infty} R_k = 1$ and $\lim_{q^2 \rightarrow 0} R_k(q^2) > 0$ ensuring the above limit properties of Γ_k . Universal properties of the renormalization group flow, however, are independent of the actual choice of R_k .

The flow of $\Gamma_k^{(2)}$ and higher n -point-functions can be derived from the flow of Γ_k by functional derivation, e.g.

$$\partial_k \Gamma_k^{(2)} = \text{Tr} f \left(\Gamma_k^{(3)2} \left(\Gamma_k^{(2)} + R_k \right)^{-3} \partial_k R_k g \right. \\ \left. + \frac{1}{2} \text{Tr} f \left(\Gamma_k^{(4)} \left(\Gamma_k^{(2)} + R_k \right)^{-2} \partial_k R_k g \right) \right)$$

where the flow of $\Gamma_k^{(3)}$ would involve $\Gamma_k^{(4)}$ and $\Gamma_k^{(5)}$ and so on. In order to end up with a closed system of functional flow equations one needs to do a finite truncation of Γ_k .

In the following we concentrate on a one-component real scalar field $\phi : \mathbb{R}^D \rightarrow \mathbb{R}$. There are the invariants $\phi^2 = \frac{1}{2} \phi' \phi'$ and $\phi^4 = \frac{1}{2} \phi' \phi' \phi' \phi'$ featuring the $O(1)$ symmetry of the Ising universality class. Here the lowest order truncation of the effective average action reads

$$\Gamma_k = \int d^D x f U_k(\phi) + \frac{1}{2} \int d^D x \phi' \phi' Z_k + O(\phi^2) g$$

in terms of the effective average potential $U_k(\phi)$ and a field independent wave function renormalization Z_k . The effective average propagator

$$\Gamma_k^{(2)}(q^2) + R_k(q^2) = U_k^0(\phi) + 2 U_k^{00}(\phi) + q^2 Z_k + R_k(q^2)$$

gives the flow of the effective average potential

$$\partial_k U_k(\phi) = \frac{1}{2} \int d^D q \frac{R_k(q)}{(2^D)^D U_k^0(\phi) + 2 U_k^{00}(\phi) + q^2 Z_k + R_k(q^2)}$$

and $\partial_k Z_k$ can be determined from $\partial_{q^2} \Gamma_k^{(2)}(q^2) \Big|_{q^2=0}$.

Such lowest order expansions and more elaborate first order expansions have been successful in the quantitative description of critical phenomena. As the flow equation for the effective average action is exact one may hope that results from expansions to higher order will converge to the correct properties of the theory. One such expansion is examined in the following.

In a second order derivative expansion of the effective average action there are contributions from up to four nonvanishing momenta:

$$\Gamma_k = \int d^D x f U_k(\phi) \\ + Z_{A;k}(\phi) \frac{1}{2!} \phi' \phi' \\ + Z_{B;k}(\phi) \frac{1}{2!} \phi' \phi' \phi' \phi' \\ + Z_{C;k}(\phi) \frac{1}{4!} \phi' \phi' \phi' \phi' \phi' \phi' \\ + Z_{D;k}(\phi) \frac{1}{4!} \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \\ + Z_{E;k}(\phi) \frac{1}{4!} \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \\ + Z_{F;k}(\phi) \frac{1}{4!} \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \\ + Z_{G;k}(\phi) \frac{1}{4!} \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' \phi' + O(\phi^6) g$$

Assuming vanishing fields and momenta at the border of space one may absorb $Z_{B;k}$, $Z_{D;k}$, $Z_{F;k}$ and $Z_{G;k}$ by integration by parts into $Z_{A;k}$, $Z_{C;k}$ and $Z_{E;k}$.

While $Z_{A;k}$ and $Z_{E;k}$ directly contribute to $\Gamma_k^{(2)}$ and thus enter all flow equations as q^2 and q^4 terms in the denominators of the integrands, $Z_{C;k}$ only gives an additive contribution to the four-point-vertex.

Also, its flow requires $\partial_k \Gamma_k^{(4)}$. This is why we neglect $Z_{C;k}$ in this paper which corresponds to a truncation beyond first and below second order only taking contributions to the four-point-vertex with at most two nonvanishing momenta into account.

Substituting $U := U_k$, $Z_0 := Z_{A;k}$ and $Z_1 := \frac{2!}{4!} \phi' \phi' Z_{E;k}$ the truncation thus reads:

$$\Gamma_k = \int d^D x f U(\phi) \\ + Z_0(\phi) \frac{1}{2!} \phi' \phi' \\ + Z_1(\phi) \frac{1}{2!} \phi' \phi' \phi' \phi' + O(\phi^4) g$$

and the next step is to derive $\partial_k U$, $\partial_k Z_0$ and $\partial_k Z_1$.

Renormalization group flow equations

Expanding the fields $\phi(x) = \phi_0 + \phi_1(x)$ in small fluctuations $\phi_1(x) = \int \frac{d^D q}{(2\pi)^D} e^{iqx} \phi(q)$ in momentum space around a constant background ϕ_0 yields the n-point-functions $\langle \phi_1 \dots \phi_n \rangle = \frac{\delta^n \Gamma[\phi_0]}{\delta \phi_0^n} = \frac{\delta^n \Gamma[\phi_0]}{\delta \phi_0^n} \quad j=0:$

$$\begin{aligned} (2) \quad \langle \phi; \phi \rangle &= U^0 + 2 U^{\text{00}} + q^2 Z_0 + q^4 Z_1 \\ (2) \quad \langle \phi; p \rangle &= U^0 + 2 U^{\text{00}} + p^2 Z_0 + p^4 Z_1 \\ (3) \quad \langle \phi; \phi; p \rangle &= p \frac{\delta \Gamma}{\delta p} = 3U^{\text{00}} + 2 U^{\text{000}} \\ &+ (p^2 + (pq) + q^2) Z_0 \\ &+ (p^4 + 2p^2(pq) + 3p^2 q^2 + 2(pq)q^2 + q^4) Z_1 \\ (4) \quad \langle \phi; \phi; \phi; p \rangle &= 3U^{\text{00}} + 12 U^{\text{000}} + 4 U^{\text{0000}} \\ &+ (p^2 + q^2) (Z_0 + 2 Z_1^{\text{00}}) \\ &+ (p^4 + 4p^2 q^2 + q^4) (Z_1 + 2 Z_1^{\text{00}}) \end{aligned}$$

These are inserted into

$$\begin{aligned} \mathcal{G}_k &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} \frac{\mathcal{G}_k R(p^2)}{(2) \langle \phi; p \rangle + R(p^2)} \\ \mathcal{G}_k^{(2)}(\phi; \phi) &= R \frac{d^D p}{(2\pi)^D} \frac{(3) \langle \phi; \phi; p \rangle (3) \langle \phi + \phi; \phi; p \rangle \mathcal{G}_k R(p^2)}{(2) \langle \phi^2 \rangle + R(p^2)^2 (2) \langle (\phi + \phi)^2 \rangle + R((\phi + \phi)^2)} \\ &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} \frac{(4) \langle \phi; \phi; \phi; p \rangle \mathcal{G}_k R(p^2)}{(2) \langle \phi^2 \rangle + R(p^2)^2} \end{aligned}$$

and yield the flow equations

$$\begin{aligned} \mathcal{G}_k U &= \mathcal{G}_k \\ \mathcal{G}_k Z_0 &= \frac{\mathcal{G}_k}{\mathcal{G}_k^2} \mathcal{G}_k^{(2)} \quad \dot{Z}_0 = 0 \\ \mathcal{G}_k Z_1 &= \frac{1}{2} \frac{\mathcal{G}_k}{\mathcal{G}_k^2} \frac{\mathcal{G}_k}{\mathcal{G}_k^2} \mathcal{G}_k^{(2)} \quad \dot{Z}_1 = 0 \end{aligned}$$

In order to be able to perform the q^2 -derivatives one has to expand the denominator containing $(p+q)^2$. Let $N = (2) \langle \phi^2 \rangle + R(p^2)$ and $M = (2) \langle (\phi + \phi)^2 \rangle + R((\phi + \phi)^2)$:

$$\frac{1}{M} = \frac{1}{N} \left[\frac{(2pq + q^2)N}{N^2} + \frac{(2pq + q^2)^2 N^2}{N^3} + \frac{1}{2} \frac{(2pq + q^2)^2 N^2}{N^2} + \dots \right]$$

Then the mixed scalar products (pq) can also be eliminated:

$$\begin{aligned} \frac{1}{2} R \frac{d^D p}{(2\pi)^D} f(p^2; q^2) \quad (pq) &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} f(p^2; q^2) \frac{p^2 q^2}{D} \\ \frac{1}{2} R \frac{d^D p}{(2\pi)^D} f(p^2; q^2) \quad (pq^4) &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} f(p^2; q^2) \frac{p^4 q^4}{D(D+2)} \\ \frac{1}{2} R \frac{d^D p}{(2\pi)^D} f(p^2; q^2) \quad (pq^k) &= 0 \text{ for all odd } k \end{aligned}$$

This finally gives the explicit flow equations:

$$\begin{aligned} \mathcal{G}_k U &= 1 - \frac{h}{N} \\ \mathcal{G}_k Z_0 &= 4 C_0^2 \frac{h}{N^4} + \frac{4}{D} \frac{p^2 N^2}{N^5} - \frac{2}{D} \frac{p^2 N^2}{N^4} + \dots \\ \mathcal{G}_k Z_1 &= 2 C_0^2 \frac{h}{N^5} - \frac{N}{N^4} + \dots \\ \dot{U} f(p^2) &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} f(p^2) \mathcal{G}_k R(p^2) \\ C_0 &= 3U^{\text{00}} + 2 U^{\text{000}} \end{aligned}$$

Scale invariant flow equations

As we are interested in critical phenomena of second order phase transitions corresponding to x points in the renormalization group flow, we write the flow equations in explicitly scale invariant form.

To this end one substitutes $z_0(\lambda) = Z_0(\lambda) = Z_0(k)$ (where k is the running minimum of $U(\lambda)$), $\mathcal{G}_k = Z_0(k)k^{2-D}$, $u = k^D U$, $t = \ln k$, $\mathcal{G}_k = \mathcal{G}_t \ln Z_0(k)$ etc. such that

$$\begin{aligned} \mathcal{G}_t u^0 &= (2 + \gamma) u^0 + (D - 2 + \gamma) u^{\text{00}} \\ C_0 &= \frac{u}{2} \\ C_1 &= \frac{u}{2} \end{aligned}$$

$$\begin{aligned} \mathcal{G}_t Z_0 &= z + (D - 2 + \gamma) \frac{z}{2} \\ &+ 4 C_0 C_0 \left(\frac{0+D}{D} J_4^0 + \frac{2}{D} K_4^1 \right) \\ &+ 8 C_0 C_1 \left(\frac{0+1D}{D} I_3^0 - \frac{2+D}{D} J_4^1 + \frac{2}{D} K_4^2 \right) \\ &+ 4 C_1 C_1 \left(\frac{1+2D}{D} I_3^1 - \frac{4+D}{D} J_4^2 + \frac{2}{D} K_4^3 \right) \\ &+ 8 C_0 C_2 \left(\frac{0+3D}{D} I_3^1 - \frac{4+D}{D} J_4^2 + \frac{2}{D} K_4^3 \right) \\ &+ 4 C_1 C_2 \left(\frac{4+8D}{D} I_3^2 - \frac{12+2D}{D} J_4^3 + \frac{4}{D} K_4^4 \right) \\ &+ 4 C_2 C_2 \left(\frac{4+6D}{D} I_3^3 - \frac{8+D}{D} J_4^4 + \frac{2}{D} K_4^5 \right) \\ C_4 &= \frac{u}{2} \\ C_5 &= \frac{u}{2} \end{aligned}$$

$$\begin{aligned} \mathcal{G}_t Z_1 &= (2 + \gamma) Z_1 + (D - 2 + \gamma) \frac{Z_1}{2} \\ &+ 2 C_0 C_0 \left(\frac{0+2D}{D} J_4^0 + \frac{8+D}{D} K_4^0 \right) \\ &+ 4 C_0 C_1 \left(\frac{0+2D}{D} J_4^0 + \frac{10+4D}{D} J_4^1 + \frac{20+D}{D} K_4^2 \right) \\ &+ 2 C_1 C_1 \left(\frac{0+2D}{D} I_3^0 - \frac{10+4D}{D} J_4^1 + \frac{20+D}{D} K_4^2 \right) \\ &+ 4 C_0 C_2 \left(\frac{0+2D}{D} I_3^0 - \frac{8+6D}{D} J_4^1 + \frac{36+18D+D^2}{D(D+2)} K_4^2 \right) \\ &+ 4 C_1 C_2 \left(\frac{4+8D}{D} I_3^1 - \frac{32+8D}{D} J_4^2 + \frac{64+30D+D^2}{D(D+2)} K_4^3 \right) \\ &+ 2 C_2 C_2 \left(\frac{16+22D}{D} I_3^2 - \frac{72+12D}{D} J_4^3 + \frac{96+42D+D^2}{D(D+2)} K_4^4 \right) \\ C_5 &= \frac{u}{2} \end{aligned}$$

where for abbreviation we have used the constants

$$\begin{aligned} w &= u^0 + 2 u^{\text{00}} \\ C_0 &= w^0 \\ C_1 &= Z_0^0 \\ C_2 &= Z_1^0 \\ C_3 &= w^0 + 2 w^{\text{00}} \\ C_4 &= Z_0^0 + 2 Z_0^{\text{00}} \\ C_5 &= Z_1^0 + 2 Z_1^{\text{00}} \end{aligned}$$

and the momentum integrals

$$\begin{aligned} I_n^m(w; z_0; z_1;) [r] &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} \frac{(p^2)^m \mathcal{G}_t r}{(w + p^2 z_0 + p^4 z_1 + r)^n} \\ J_n^m(w; z_0; z_1;) [r] &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} \frac{(p^2)^m (z_0 + 2p^2 z_1 + r) \mathcal{G}_t r}{(w + p^2 z_0 + p^4 z_1 + r)^n} \\ K_n^m(w; z_0; z_1;) [r] &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} \frac{(p^2)^m 2(z_0 + 2p^2 z_1 + r)^2 \mathcal{G}_t r}{(w + p^2 z_0 + p^4 z_1 + r)^{n+1}} \\ &= \frac{1}{2} R \frac{d^D p}{(2\pi)^D} \frac{(p^2)^m (2z_1 + r) \mathcal{G}_t r}{(w + p^2 z_0 + p^4 z_1 + r)^n} \end{aligned}$$

which will be discussed next:

The m o m e n t u m i n t e g r a l s I , J and K c a n b e c o m p u t e d e f f i c i e n t l y i n c a s e o f a l i n e a r c u t o f f . F o r t h e c h o i c e [6]
 $R_k = Z_k (k^2 - q^2) (k^2 + q^2)$ o n e h a s

$$\begin{aligned} r &= (1 - p^2) - (1 - p^2) \\ \underline{r} &= (1 - p^2) \\ r &= (1 - p^2) \\ \partial_r r &= (2 + p^2) - (1 - p^2) \end{aligned}$$

and I , J and K a r e r e d u c e d t o a s i n g l e i n t e g r a l H :

$$\begin{aligned} I_n^m &= (2 - p^2) H_n^m + H_n^{m+1} \\ J_n^m &= (z_0 - 1) I_n^m + 2z_1 I_n^{m+1} \\ K_n^m &= 2(z_0 - 1)^2 I_{n+1}^m + 8(z_0 - 1)z_1 I_{n+1}^{m+1} + 8z_1^2 I_{n+1}^{m+2} \\ &\quad - 2z_1 I_n^m - v_D (w + z_0 + z_1)^n \end{aligned}$$

where H i s a s p e c i a l c a s e o f a h y p e r g e o m e t r i c f u n c t i o n

$$\begin{aligned} H_n^m(w; z_0; z_1) &= \frac{1}{2} \int_0^1 \frac{d^D p}{(2 - p^2)^D} \frac{(p^2)^m (1 - p^2)}{(w + p^2 z_0 + p^4 z_1 + r)^n} \\ &= v_D \int_0^1 \frac{dx}{((w+1) + (z_0-1)x + z_1 x^2)^n} \\ &= \frac{v_D}{(w+1)^n} \int_0^1 dx \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (1 - bx - cx^2)^{-n} \\ &= \frac{v_D}{(w+1)^n} \int_0^1 dx \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \sum_{k=0}^{\infty} \frac{x^k}{k!} \sum_{l=0}^{\infty} \frac{(n+k-1-l)! k!}{(n-1)! l! (k-2l)!} \frac{c^l b^k z_1^{2l}}{(k-2l)!} \\ &= \frac{v_D}{(w+1)^n} \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{(n+k-1-l)!}{(n-1)!} \frac{c^l b^k z_1^{2l}}{l! (k-2l)!} \int_0^1 dx x^{a+k} \\ &= \frac{v_D}{(w+1)^n} \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{(n+k-1-l)!}{(n-1)!} \frac{c^l b^k z_1^{2l}}{l! (k-2l)!} \frac{1}{a+k+1} \end{aligned}$$

w i t h $v_D = (2^{D+1} - D - 2) (D=2) \int_0^1 dx x^{a+k} \in 0$ for all $x \in [0;1]$, i.e. if the integrand of H has no singularities. This is ensured as the effective average propagator stays finite at all times.

Starting with a quartic potential $u = \frac{1}{2} (\phi^2)^2$ the flow equations are discretized on a grid and numerically solved for different values of ϕ . During the evolution towards $k \rightarrow 0$, the running minimum of the potential ($u^0(\phi) = 0$) may either end up in the massless spontaneously broken phase ($m = 0, \phi \neq 0$) or in the massive symmetric phase ($m^2 = k^2 u^0(0), \phi = 0$). The critical ϕ_c leads to a second order phase transition characterized by ϕ points of all couplings and vanishing masses. Fine tuning around ϕ_c then yields the critical exponent through the scaling law $m \sim |\phi - \phi_c|$ and the established proportionality $\chi \sim |\phi - \phi_c|^{-\gamma}$. The critical exponent γ may be identified with the χ point of the anomalous dimension. It can therefore be directly obtained from the critical trajectory.

The critical exponents for the universality class of the three dimensional Ising model obtained in this way are compared in Table 1 to values obtained in first order ($z_1 = 0$) and lowest order ($z_0^0 = z_0^1 = \dots = 0$). For comparison we also give results from literature. An extensive review of different techniques and recent results can be found in [7]. We adopt their overall estimates of exponents and errors. In addition we quote some of the newest and preciseest results from high temperature expansions, perturbation series at fixed dimension, ϵ -expansions and Monte Carlo simulations plus some additional references.

C o n c l u s i o n a n d o u t l o o k

We have presented the first calculation of a derivative expansion beyond first order of the effective average action. Considering former results from lowest and first order expansions we notice that the critical exponents calculated in this way seem to converge to the values suggested by literature. This observation is very encouraging and the next step will be the inclusion of the third wave function renormalization which will correspond to a calculation in full second order.

effective average action to lowest order	0.6271	0.1120
effective average action to first order	0.6223	0.0806
effective average action beyond first order (this paper)	0.6303	0.0793
literature values from the review [7]	0.6301 (4)	0.0364 (5)
25th-order high-temperature expansion [8], see also [9]	0.63012 (16)	0.03639 (15)
seven-loop perturbation series at fixed $D = 3$ [10], see also [11]	0.6303 (8)	0.0335 (6)
seven-loop order expansion with boundary conditions [11]	0.6305 (25)	0.0365 (50)
Monte Carlo simulation [12], see also [13, 14]	0.6297 (5)	0.0362 (8)

T A B L E 1 : C r i t i c a l e x p o n e n t s o f t h e t h r e e d i m e n s i o n a l I s i n g u n i v e r s a l i t y c l a s s

Electronic address: physics@ballhausen.com

- [1] J. Berges, N. Tetradis and C. Wetterich, Non-perturbative renormalization flow in quantum field theory and statistical physics, Phys.Rept. 363 (2002) 223-386 [hep-ph/0005122].
- [2] C. Wetterich, Average action and the renormalization group equations, Nucl.Phys.B 352 (1991) 529.
- [3] C. Wetterich, The average action for scalar fields near phase transitions, Z.Phys.C 57 (1993) 451.
- [4] C. Wetterich, Exact evolution equation for the effective potential, Phys.Lett.B 301 (1993) 90.
- [5] N. Tetradis, C. Wetterich, Critical exponents from the effective average action, Nucl.Phys.B 422 (1994) 541-592 [hep-ph/9308214].
- [6] D. F. Litin, Optimised renormalisation group flows, Phys.Rev.D 64 (2001) 105007 [hep-th/0103195].
- [7] A. Pelissetto, E. Vicari, Critical Phenomena and Renormalization-Group Theory, Phys.Rept. 368 (2002) 549-727 [cond-mat/0012164].
- [8] M. Campostrini, A. Pelissetto, P. Rossi, E. Vicari, 25th-order high-temperature expansion results for three-dimensional Ising-like systems on the simple cubic lattice, Phys.Rev.E 65 (2002) 066127 [cond-mat/0201180].
- [9] P. Butera, M. Comi, Critical universality and hyperscaling revisited for Ising models of general spin using ext. high-temperature series, Phys.Rev.B 65 (2002) 144431 [hep-lat/0112049].
- [10] F. Jasch, H. Kleinert, Fast-Convergent Resummation Algorithm and Critical Exponents of ϕ^4 -Theory in Three Dimensions, J.Math.Phys. 42 (2001) 52 [cond-mat/9906246].
- [11] R. Guida, J. Zinn-Justin, Critical Exponents of the N-vector model, J.Phys.A 31 (1998) 8103 [cond-mat/9803240].
- [12] M. Hasenbusch, Monte Carlo Studies of the Three-Dimensional Ising Model in Equilibrium, Int.J.Mod.Phys.C 12 (2001) 991.
- [13] H.W. J. Blöte, L.N. Shchur, A.L. Talapov, The Cluster Processor: New Results, Int.J.Mod.Phys.C 10 (1999) 1137 [cond-mat/9912005].
- [14] H.G. Ballesteros, L.A. Fernandez, V. Martı́n-Mayor, A. Muñoz Sudupe, G. Parisi, J.J. Ruiz-Lorenzo, Scaling corrections: site-percolation and Ising model in three dimensions, J.Phys.A 32 (1999) 1 [cond-mat/9805125].