

A Note on Exotic Fractions of Supersymmetries

André Miemiec^{*1} and Igor Schnakenburg^{*2}

^{*1} Department of Theoretical Physics
University of Turin
10125 Torino
Italia
miemiec@to.infn.it

^{*2} Racah Institute of Physics
The Hebrew University
Jerusalem 91904
Israel
igorsc@phys.huji.ac.il

Abstract

In this note we study supersymmetric solutions of 11 dimensional SUGRA. Amongst others, we find a new solution which preserves $3/8$ of the original supersymmetry.

1 Introduction

Solutions of supergravity theories which preserve 1/2 of the original supersymmetry typically occur in the description of brane solutions. Using the techniques of intersecting brane solutions one can reduce this number quite generically to any fraction $1/2^n$, with n an integer. Other fractions are less simple to understand and are called exotic. In [1] the authors discovered a new solution of 11 dimensional supergravity, which preserved a fraction of $\nu = 5/8$ supersymmetries. In a following paper [2] it was shown that this solution fits into a whole class of solutions and the question was raised if further solutions with exotic fractions of preserved supersymmetry can be found within this class. This note confirms in a case by case study that this actually is the case.

The class of solutions proposed in [2] are

$$\begin{aligned} ds_{11}^2 &= -(dt + \omega)^2 + ds^2(\mathbb{R}^{10}) \\ F &= -d\omega \wedge \Omega \end{aligned} \quad (1.1)$$

with Ω defined in terms of complex coordinates $z^a = x^{2a-1} + i x^{2a}$ of $\mathbb{C}^5 = \mathbb{R}^{10}$,

$$\Omega = \frac{i}{2} \sum_{a=1}^5 dz^a \wedge dz^{\bar{a}}, \quad (1.2)$$

and $d\omega = \alpha + \bar{\alpha} \in \Lambda^{2,0}(\mathbb{C}^5) \oplus \Lambda^{0,2}(\mathbb{C}^5)$.

The Einstein equation and the equation of motion for the 4-form field strength read

$$R_{ab} - \frac{1}{12} \left(F_{ac_1c_2c_3} F_b{}^{c_1c_2c_3} - \frac{1}{12} \eta_{ab} F_{c_1c_2c_3c_4} F^{c_1c_2c_3c_4} \right) = 0, \quad (1.3)$$

$$d(*F) + \frac{1}{2} F \wedge F = 0, \quad (1.4)$$

while the Killing spinor equation is given by

$$\partial_\mu \varepsilon - \frac{1}{4} \omega_{\mu ab} \Gamma^{ab} \varepsilon + \frac{1}{288} \left(\Gamma_\mu \Gamma^{\nu_1 \nu_2 \nu_3 \nu_4} - 12 \delta_\mu^{\nu_1} \Gamma^{\nu_2 \nu_3 \nu_4} \right) F_{\nu_1 \nu_2 \nu_3 \nu_4} \varepsilon = 0. \quad (1.5)$$

2 Examples

2.1 First Example

The holomorphic two form α we have chosen is

$$\alpha = \frac{\gamma}{2} (dz^1 \wedge dz^2 + dz^3 \wedge dz^4) \quad (2.1)$$

which leads to

$$\omega = \frac{\gamma}{2} (-x^3 dx^1 + x^1 dx^3 - x^2 dx^4 + x^4 dx^2 - y^3 dy^1 + y^1 dy^3 - y^2 dy^4 + y^4 dy^2) \quad (2.2)$$

The equations of motion (1.3) and (1.4) are both satisfied. The Ricci tensor in a tangent frame is diagonal and given by $R_{ab} = \gamma^2 \cdot \text{diag}(2, 1/2, \dots, 1/2, 0, 0)$ while the contractions of the field strength are:

$$F_{c_1 c_2 c_3 c_4} F^{c_1 c_2 c_3 c_4} = 288 \gamma^2 \quad (2.3)$$

$$F_{ac_1 c_2 c_3} F_b{}^{c_1 c_2 c_3} = \begin{cases} 0 & (a, b) = (0, 0) \\ 30 \gamma^2 & (a, b) = (1, 1), \dots, (8, 8) \\ 24 \gamma^2 & (a, b) = (9, 9), (10, 10) \\ 0 & a \neq b \end{cases} \quad (2.4)$$

Obviously eq. (1.3) is satisfied. The second term in the equation of motion for the 4-form field strength (1.4) reduces to

$$\frac{1}{2} F \wedge F = 2 \gamma^2 (2 dx^{12345678} + dx^{1234569\#} + dx^{1234789\#} + dx^{1256789\#} + dx^{3456789\#})$$

and comparing this with $d(*F)$ one finds that both add up to zero.

The Killing spinor equation (1.5) can be written more symbolically as

$$\partial_\mu \varepsilon - \mathbb{M}_\mu \varepsilon = 0. \quad (2.5)$$

By just solving this equation for the present ansatz one obtains 12 constant Killing spinors. These 12 solutions exhaust the whole set of solutions. Now we give the argument from which we draw this conclusion.

As a preparation we would like to shed some light onto the terms which we gathered in

$$e^\mu{}_c \mathbb{M}_\mu = \frac{1}{4} \underbrace{\omega_{cab} \Gamma^{ab}}_{(3)} - \frac{1}{288} \left(\Gamma_c \underbrace{(\Gamma^{\nu_1 \nu_2 \nu_3 \nu_4} F_{\nu_1 \nu_2 \nu_3 \nu_4})}_{(1)} - 12 e^\mu{}_c \underbrace{\Gamma^{\nu_2 \nu_3 \nu_4} F_{\mu \nu_2 \nu_3 \nu_4}}_{(2)} \right) \quad (2.6)$$

The components of the field strength $F_{\nu_1 \nu_2 \nu_3 \nu_4}$ are constant and those involving the time direction vanish. Also, Gamma matrices with upper spatial indices are identical to the tangent frame Gamma matrices. We conclude that the expressions (1) and (2) are constant. The vielbein in (3) turns out to be constant too. So the only source of coordinate dependence comes from contraction with the vielbein in the last term. Since the field strength components which include a time direction vanish, the only remaining source for a coordinate dependence is set to zero.

This example gives a simple necessary condition, which each solution must satisfy

$$0 = [\partial_\mu, \partial_\nu] \varepsilon = e_\mu{}^a e_\nu{}^b (2 \omega_{[ab]}{}^c \mathbb{M}_c + [\mathbb{M}_b, \mathbb{M}_a]) \varepsilon, \quad (2.7)$$

where $\mathbb{M}_a = e^\mu{}_a \mathbb{M}_\mu$ and $\omega_{[\mu\nu]}{}^a = \partial_{[\mu} e_{\nu]}{}^a$. In deriving eq. (2.7) we used the fact that in the present case all \mathbb{M}_a are constant. Since the only nonvanishing components of $\omega_{[\mu\nu]}{}^a$ are

$$\omega_{[13]}{}^0 = \gamma, \quad \omega_{[24]}{}^0 = -\gamma, \quad \omega_{[57]}{}^0 = \gamma, \quad \omega_{[68]}{}^0 = -\gamma \quad (2.8)$$

equation (2.7) evaluated for $(a, b) = (0, 1)$ just reduces to

$$0 = [\mathbb{M}_1, \mathbb{M}_0] \varepsilon, \quad (2.9)$$

the kernel of which is just the span of the 12 Killing spinors mentioned above. So we already found all solutions.

2.2 Second Example

We now take the following ansatz for

$$\alpha = \frac{\gamma}{2} (dz^1 \wedge dz^2 + dz^1 \wedge dz^3 + dz^2 \wedge dz^3) . \quad (2.10)$$

In this case the Ricci tensor in the tangent frame is no longer diagonal. Nevertheless the equations of motion are satisfied both for the metric and the four form field strength. One finds 16 constant Killing spinors ε_i generating the 16 dimensional kernels of \mathbb{M}_1 to \mathbb{M}_6 . They are contained in the 20 dimensional kernels of \mathbb{M}_0 and \mathbb{M}_7 to \mathbb{M}_{10} . The form of the Killing spinors can be calculated explicitly and is given for convenience in appendix B. The kernels of \mathbb{M}_0 and \mathbb{M}_7 to \mathbb{M}_{10} are all equal and the additional four dimensions are spanned by ϑ , which satisfies

$$0 \neq \mathbb{M}_i(\vartheta) \subset \text{span}(\varepsilon_1, \dots, \varepsilon_{16}), \quad i = 1, \dots, 6$$

and so we can apply the same trick used in [1] to construct a further Killing spinor by setting

$$\varepsilon = \vartheta + x^i \mathbb{M}_i \vartheta .$$

Finally we obtain 20 Killing spinors, i.e. $\nu = 5/8$ of the original supersymmetries are preserved.

3 Conclusion

In the above we have presented two solutions, which preserve two different fractions of supersymmetry. The first example preserves 3/8, the later 5/8. The first example preserves a different exotic fraction of supersymmetry than the example already discovered in [2]. We have also tried different ansätze for α . Their precise form and the amount of supersymmetry they preserve can be found in the table below.

α	ν
$dz^1 \wedge dz^2$	5/8
$dz^1 \wedge dz^2 + dz^1 \wedge dz^3$	5/8
$dz^1 \wedge dz^2 + dz^1 \wedge dz^3 + dz^2 \wedge dz^3$	5/8
$dz^1 \wedge dz^2 + dz^3 \wedge dz^4$	3/8
$dz^1 \wedge dz^2 + dz^3 \wedge dz^4 + dz^1 \wedge dz^5$	3/8
$dz^1 \wedge dz^2 + dz^2 \wedge dz^3 + \dots + dz^5 \wedge dz^1$	3/8

In particular, we have only found solutions which preserve either 3/8 or 5/8 of the original supersymmetry. It would be interesting to see whether this ansatz could also produce solutions with other numbers of preserved supersymmetries.

A Clifford Algebra

The 11 dimensional Clifford Algebra with signature $\eta^{ab} = \text{diag}(-1, 1, \dots, 1)$ satisfies

$$\{\Gamma^a, \Gamma^b\} = 2\eta^{ab} \mathbb{1}_{32} . \quad (\text{A.1})$$

A representation by real matrices Γ^a can be obtained by taking appropriate tensor products of the Pauli matrices

$$\tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \tau_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} .$$

A particular choice is ($\epsilon = i\tau_2$):

$$\begin{aligned} \Gamma^0 &= -\epsilon \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \tau_3 & \Gamma^1 &= \epsilon \otimes \epsilon \otimes \epsilon \otimes \epsilon \otimes \tau_1 \\ \Gamma^2 &= \epsilon \otimes \mathbb{1} \otimes \tau_1 \otimes \epsilon \otimes \tau_1 & \Gamma^3 &= \epsilon \otimes \mathbb{1} \otimes \tau_3 \otimes \epsilon \otimes \tau_1 \\ \Gamma^4 &= \epsilon \otimes \tau_1 \otimes \epsilon \otimes \mathbb{1} \otimes \tau_1 & \Gamma^5 &= \epsilon \otimes \tau_3 \otimes \epsilon \otimes \mathbb{1} \otimes \tau_1 \\ \Gamma^6 &= \epsilon \otimes \epsilon \otimes \mathbb{1} \otimes \tau_1 \otimes \tau_1 & \Gamma^7 &= \epsilon \otimes \epsilon \otimes \mathbb{1} \otimes \tau_3 \otimes \tau_1 \\ \Gamma^8 &= \epsilon \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \epsilon & \Gamma^9 &= \tau_1 \otimes \mathbb{1}_{16} \\ \Gamma^{10} &= \tau_3 \otimes \mathbb{1}_{16} \end{aligned}$$

B Explicite Killing spinors

$$\begin{aligned} \varepsilon &= (a_1 + a_2 + a_3, a_4, a_5, a_6, a_7, a_8, -a_2, -a_4, a_9, a_6, a_2 + a_3 + a_{10}, a_4, a_{11}, a_{12}, a_5 + a_{13} + a_{14}, \\ &\quad a_6, a_5 - a_9 + a_{13}, -a_6, a_{10}, a_4, a_{15}, -a_{12}, a_{14}, a_6, a_1, a_4, a_{13}, -a_6, a_{16}, a_8, -a_3, a_4)^T \\ \vartheta &= (0, a_{17}, 0, a_{18}, 0, a_{19}, 0, -a_{17}, 0, a_{18}, 0, a_{17}, 0, a_{20}, 0, a_{18}, 0, a_{18}, \\ &\quad 0, -a_{17}, 0, a_{20}, 0, -a_{18}, 0, -a_{17}, 0, a_{18}, 0, -a_{19}, 0, -a_{17})^T \end{aligned}$$

References

- [1] J. P. Gauntlett, J. B. Gutowski, C. M. Hull, S. Pakis, and H. S. Reall, “All supersymmetric solutions of minimal supergravity in five dimensions,” [hep-th/0209114](#).
- [2] J. P. Gauntlett and S. Pakis, “The Geometry of D = 11 Killing Spinors,” [hep-th/0212008](#).