

# On type II superstrings in bosonic backgrounds and their T-duality relation

Luca Martucci<sup>1</sup> and Pedro J. Silva<sup>2</sup>

Dipartimento di Fisica dell'Università di Milano,  
Via Celoria 16, I-20133 Milano, Italy

INFN, Sezione di Milano,  
Via Celoria 16, I-20133 Milano, Italy

## ABSTRACT

We derive the actions for type II Green-Schwarz strings up to second order in the fermions, for general bosonic backgrounds. We base our work on the so-called normal coordinate expansion. The resulting actions are  $\kappa$ -symmetric and, for the case of surviving background supersymmetries, supersymmetric. We first obtain the type IIA superstring action from the 11D supermembrane by double dimensional reduction. Then, by means of a generalization of T-duality we derive the type IIB superstring action. The resulting actions are surprisingly simple and elegant.

---

<sup>1</sup>luca.martucci@mi.infn.it

<sup>2</sup>pedro.silva@mi.infn.it

# 1 Introduction

In this short letter we will present the type II superstring actions expanded in fermion coordinates up to second order in general supergravity bosonic backgrounds. In contrast to the usual situation where this expansion is written in a very unfriendly outcast, our result is elegant and economical. The action actually comes in terms of the “natural” operators appearing in the supersymmetric transformation of the supergravity background fields. In the same footing, the well-known symmetries of these actions like  $\kappa$ -symmetry and supersymmetry are almost obvious from the expressions that here we write. This miracle, sort to speak, is the result of using the so-called “normal coordinate expansion” [4] in the superfield formulation. Also, due to the close relation between supersymmetry and T-duality for world-volume theories, our action is also “covariant” under T-duality transformation. In fact, we will re-derive the T-duality rules obtaining a generalization of the Buscher rules that again is elegant and simple. Nevertheless, here we point out that T-duality among string theory actions correspond to a much more complete test for string theory, since not only the zero modes are considered, but also all the excitations.

Due to the simplicity of the results, here we show the actions and leave for latter the actual derivation.

$$S_{(F)} = - \int d^2\xi \sqrt{-\det(g)} + \frac{1}{2} \int d^2\xi \epsilon^{ij} b_{ij} + \frac{i}{2} \int d^2\xi \sqrt{-\det(g)} \bar{y} \tilde{P}_{(-)} \Gamma^i \tilde{D}_i y, \quad (1)$$

where for the type IIa case,

$$\begin{aligned} y &= (y_+ + y_-) \quad \text{with } \Gamma^{11} y_{\pm} = \pm y_{\pm} \quad \text{and } P_{(-)} = \frac{1}{2} \left( 1 - \frac{1}{2\sqrt{-g}} \epsilon^{ij} \Gamma_{ij} \Gamma^{11} \right), \\ \tilde{D}_m &= D_m^{(0)} + W_m, \\ D_m^{(0)} &= \partial_m + \frac{1}{4} \omega_{mab} \Gamma^{ab} + \frac{1}{4 \cdot 2!} H_{mab} \Gamma^{ab} \Gamma^{11}, \\ W_m &= \frac{1}{8} e^{\phi} \left( \frac{1}{2!} \mathbf{F}_{ab}^{(2)} \Gamma^{ab} \Gamma_m \Gamma^{11} + \frac{1}{4!} \mathbf{F}_{abcd}^{(4)} \Gamma^{abcd} \Gamma_m \right). \end{aligned} \quad (2)$$

and for the type IIb case,

$$\begin{aligned} y &= \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \quad \text{with } \Gamma^{11} y_{(1,2)} = y_{(1,2)} \quad \text{and } P_{(-)} = \frac{1}{2} \left( 1 - \frac{1}{2\sqrt{-g}} \sigma_3 \otimes \epsilon^{ij} \Gamma_{ij} \Gamma^{11} \right), \\ \tilde{D}_m &= \begin{pmatrix} \hat{D}_{(1)m}^{(0)} & 0 \\ 0 & \hat{D}_{(2)m}^{(0)} \end{pmatrix} + \begin{pmatrix} 0 & \hat{W}_{(1)m} \\ \hat{W}_{(2)m} & 0 \end{pmatrix}, \\ \hat{D}_{(1,2)m}^{(0)} &= \partial_m + \frac{1}{4} \omega_{mab} \Gamma^{ab} \pm \frac{1}{4 \cdot 2!} H_{mab} \Gamma^{ab}, \\ \hat{W}_{(1,2)m} &= \pm \frac{1}{8} e^{\phi} \left( \mathbf{F}^{(1)}{}_a \Gamma^a \pm \frac{1}{3!} \mathbf{F}_{abc}^{(3)} \Gamma^{abc} + \frac{1}{4!} \mathbf{F}_{abcd} \Gamma^{abcd} \right) \Gamma_m, \end{aligned} \quad (3)$$

where in the expressions for  $\hat{D}^{(0)}$  and  $\hat{W}$ ,  $\pm$  correspond to the label (1,2). Also in all the above, is  $g$  the 10D metric,  $H = db$ , and  $\mathbf{F}^{(n)}$ , are the RR field strength (see [1] and [2] for the supergravity conventions).

In the following we will first derive the type IIa action as a double dimensional reduction from the membrane of 11D. Second, we fully discuss the  $\kappa$ -symmetry transformation and its geometrical meaning from the point of view of the embedding of the 2d-theory into 10d space-time. Third, for the case of supergravity bosonic backgrounds that preserve some supersymmetry, we will give the supersymmetric transformation of the world-sheet variables that correspond to symmetries of the action. Forth, we will derive the type IIb superstring action and give the explicit T-duality rules that interrelates the type IIa and type IIb theories. At last, we will give the  $\kappa$ -symmetry and supersymmetry transformations for the type IIb case. An appendix is enclosed with the T-duality rules of Hassan.

To begin the actual derivation of the above results, our starting point is the supersymmetric membrane action in 11D, i.e.

$$S = - \int d^3\xi \sqrt{\det(-\mathbf{G})} - \frac{1}{6} \int d^3\xi \varepsilon^{ijk} \mathbf{B}_{kji}, \quad (4)$$

where  $i = (0, 1, 2)$  and  $(\mathbf{G}, \mathbf{B})$  are the pull-back to the membrane of the metric and 3-form super-fields of N=1 11D supergravity. Then, we borrow from [3] the expression corresponding to the expansion in normal coordinates, in a general bosonic background, up to second order in the fermions,

$$S = S^{(0)} + S^{(2)}, \quad (5)$$

$$S^{(0)} = - \int d^3\xi \sqrt{-\det(G)} - \frac{1}{6} \int d^3\xi \varepsilon^{ijk} b_{kji}, \quad (6)$$

$$S^{(2)} = \frac{i}{2} \int d^3\xi \left\{ \sqrt{-\det(G)} \left[ \bar{y} \Gamma^i \nabla_i y + \bar{y} T_i \Gamma^i y \right] + \right. \\ \left. - \frac{1}{2} \varepsilon^{ijk} \left[ \bar{y} \Gamma_{ij} \nabla_k y + \bar{y} T_i \Gamma_{jk} y \right] \right\}, \quad (7)$$

where  $(G, b)$  are the bosonic pull-back of  $(\mathbf{G}, \mathbf{B})$ ,  $y$  is a real Majorana 32 component spinor,  $\Gamma$  are real gamma matrices,  $\nabla_i$  is the usual pull-back of the spinor covariant derivative and  $T$  stand for the appropriated components of the supertorsion in 11D (see [3] for more details). To obtain the type IIa string action we perform a double dimensional reduction of the above expression. Using the typical KK-ansatz for the 11D metric

$$ds_{(11)}^2 = e^{-\frac{2}{3}\phi} ds_{(10)}^2 + e^{\frac{3}{4}\phi} (dz - C_m dx^m)^2, \quad (8)$$

after some straight forward but tedious algebra, we get the type IIa string action (1,2) in general bosonic backgrounds expanded up to second order in the fermions.

At this point, it is important to remark that this action is by construction  $\kappa$ -symmetric and supersymmetric (for the case of bosonic backgrounds with surviving supersymmetries). The above statement requires some attention and so, before deriving the type IIb string action, we pause and discuss these symmetries.

The  $\kappa$ -symmetry of the above actions can be understood as inherit from the superfield  $\kappa$ -symmetry of the membrane in 11D [7]. Recall that only by fixing  $\kappa$ -symmetry we get the world-sheet supersymmetric theory transforming the 10D spinors into two dimensional ones. In our case, the resulting expanded action in normal coordinates, after double dimensional reduction, is written in terms of the pull-back of a covariant expansion of the 10d supergravity fields. This expansion comes in terms covariant derivatives, curvature tensor, and the torsion. In principle the above type of Lagrangian contains too much information, related to the generalizations

of concepts like extrinsic curvature, that are by no means related to a pure supersymmetric two-dimensional world-sheet theory. Therefore, the role of  $\kappa$ -symmetry is to eliminate all this redundant information obtaining a world-sheet theory. In other words  $\kappa$ -symmetry is the responsible to eliminate from the action all the geometrical information of the embedding that is not strictly speaking two-dimensional. In practice the expression for the infinitesimal transformations are derivable using the normal coordinate formalism (see for example [4]),

$$\delta_\kappa y^\alpha = (1 + \Gamma_F)\kappa \quad , \quad \delta_\kappa x^m = \frac{i}{2}\bar{y}\Gamma^m(1 + \Gamma_F)y \quad , \quad \delta_\kappa A = \delta_\kappa x^m \partial_m A \quad , \quad (9)$$

where  $\Gamma_F = \frac{1}{2\sqrt{-g}}\epsilon^{ij}\Gamma_{ij}\Gamma^{11}$  and  $A$  is a general field of the supergravity background.

The supersymmetry transformations are also derived within the same approach and are given by,

$$\delta_\epsilon y = \epsilon \quad , \quad \delta_\epsilon x^m = -\frac{i}{2}\bar{y}\Gamma^m y \quad , \quad \delta_\epsilon A = \delta_\epsilon x^m \partial_m A \quad , \quad (10)$$

where again  $A$  is a general field of the supergravity background and we are assuming that the fermion  $\epsilon$  is actually a killing spinor of the bosonic background. Therefore a translation along its orbit corresponds to a supersymmetry transformation of the Lagrangian.

Now that we understand the two more relevant symmetries of the above action, let us concentrate in the type IIb case. In order to obtain the corresponding action for the type IIb string, we perform a T-duality transformation, assuming the existence of a killing direction  $x^9$ . First of all, we rewrite the action (1,2) in the following form,

$$S_{(IIa)} = - \int d^2\xi \sqrt{-\det(G)} + \frac{1}{2} \int d^2\xi \epsilon^{ij} B_{ij} \quad (11)$$

where

$$\begin{aligned} G_{mn} &= g_{mn} - i\bar{y}\Gamma_{(m}D_{n)}y \quad , \\ B_{mn} &= b_{mn} - i\bar{y}\Gamma^{11}\Gamma_{[m}D_{n]}y \quad , \end{aligned} \quad (12)$$

and we have used the fact that this action comes from the pull-back of the normal coordinate expansion of the vielbeins and other supergravity fields obtaining, among other things, the pull-back of the covariant derivative of the fermionic variables  $y$ . Then, as usual in this the context, T-duality is implemented with the help of a Lagrange multiplier  $Z$  and a world-volume vector  $V_i$  (taking the place of  $\partial_i x^9$ ), by introducing a term of the form  $\epsilon^{ij}V_j\partial_i Z$  into the action. Since the action formally has the same form as the bosonic string action, we only get the usual Buscher rules<sup>3</sup> once we solve for  $V_i$ .

$$\begin{aligned} \hat{G}_{99} &= \frac{1}{G_{99}} \\ \hat{G}_{\tilde{m}\tilde{n}} &= G_{\tilde{m}\tilde{n}} - \frac{G_{\tilde{m}9}G_{\tilde{n}9} - B_{\tilde{m}9}B_{\tilde{n}9}}{G_{99}} & \hat{G}_{\tilde{m}9} &= \frac{B_{\tilde{m}9}}{G_{99}} \\ \hat{B}_{\tilde{m}\tilde{n}} &= B_{\tilde{m}\tilde{n}} - \frac{B_{\tilde{m}9}G_{\tilde{n}9} - G_{\tilde{m}9}B_{\tilde{n}9}}{G_{99}} & \hat{B}_{\tilde{m}9} &= \frac{G_{\tilde{m}9}}{G_{99}} \end{aligned} \quad (13)$$

---

<sup>3</sup>See [5] and references therein.

where  $\tilde{m} = 0, \dots, 8$ . The important ingredient comes when we rewrite the fermionic part of the fields,  $(G, B)$  in terms of type IIB variables. At this point, because of the nature of normal coordinate expansion, we can use the transformation rules of Hassan (see the appendix) to obtain the desired result. The corresponding T-duality rules can be found in the appendix, here we write the final form of the transformed fields  $(\hat{G}, \hat{B})^4$ :

$$\begin{aligned}\hat{G}_{mn} &= \hat{g}_{mn} - i\bar{y}\Gamma_{(m}\hat{D}_{n)}y, \\ \hat{B}_{mn} &= \hat{b}_{mn} - i\bar{y}\hat{\Gamma}\Gamma_{[m}\hat{D}_{n]}y,\end{aligned}\tag{14}$$

where  $\hat{\Gamma} = \sigma_3 \otimes \Gamma^{11}$ ,  $(y, \hat{D})$  are define as in (3) and  $(\hat{g}_{mn}, \hat{b}_{mn})$  are transformed metric and NS field in the type IIB theory. Therefore, after some trivial manipulation we find that the expanded up to second order in the fermions type IIB fundamental string action in general bosonic backgrounds is given by equation (1,3).

At last, only remains to obtain the explicit form for the supersymmetry and  $\kappa$ -symmetry transformations for the type IIB action. To derive the rules for these symmetries we adapt a argument presented in [8]. let us begin with the derivation of the  $\kappa$ -symmetry rules for the type IIB superstring. We use that originally the vector field  $V_i$  obeys the constraint  $\epsilon^{ij}\partial_i V_j = 0$ . At this point we know that the action is invariant under the transformation (9). Therefore, since this constraint is invariant under  $\kappa$ -symmetry, the variation of  $S$ , in which we take  $V_i$  as independent variable, should be proportional to  $\epsilon^{ij}\partial_i V_j$ . Such a term can be easily isolated to be

$$\delta_\kappa S \sim \int \epsilon^{ij}\partial_i V_j [\delta_\kappa x^m b_{9m} - \frac{i}{2}\bar{y}\Gamma^{11}(1 + \Gamma_F)\kappa].\tag{15}$$

Hence, to have a  $\kappa$ -invariant action, we must take

$$\delta_\kappa Z = \frac{i}{2}b_{9m}\bar{y}\Gamma^m(1 + \Gamma_F)\kappa - \frac{i}{2}\bar{y}\gamma^{11}(1 + \Gamma_F)\kappa,\tag{16}$$

that, written in IIB T-dual variables<sup>5</sup>, takes the form

$$\delta_\kappa Z = \frac{i}{2}\bar{y}(1 + \hat{\Gamma}_F)\kappa,\tag{17}$$

where  $\hat{\Gamma}_F = \frac{1}{2\sqrt{-g}}\sigma_3 \otimes \epsilon^{ij}\Gamma_{ij}\Gamma^{11}$  and the fermions  $y$  are in the same convention of (3). The other variables T-dualize trivially and it is now easy to complete the derivation of the transformation rules. Hence we get,

$$\delta_\kappa y^\alpha = (1 + \hat{\Gamma}_F)\kappa, \quad \delta_\kappa x^m = \frac{i}{2}\bar{y}\Gamma^m(1 + \hat{\Gamma}_F)y, \quad \delta_\kappa A = \delta_\kappa x^m \partial_m A,\tag{18}$$

where  $A$  is a general field of the supergravity background.

The supersymmetry transformations are derived in the same footing, and are given by

$$\delta_\epsilon y = \epsilon, \quad \delta_\epsilon x^m = -\frac{i}{2}\bar{y}\Gamma^m y, \quad \delta_\epsilon A = \delta_\epsilon x^m \partial_m A,\tag{19}$$

---

<sup>4</sup>We should point out that the transformation rule of the dilaton is obtained with no need of including quantum corrections. Basically it is the curved space-time structure what provides a classical derivation for the dilaton T-duality.

<sup>5</sup> $\Gamma_{F1}$  dualize to  $\hat{\Gamma}_{F1}$ , as can be read directly from the T-duality transformation rule of the action.

where again  $A$  is a general field of the supergravity background and we are assuming that the fermion  $\epsilon$  is actually a killing spinor of the bosonic background.

In this letter we have provide a new form for the type IIa/b superstring actions, expanded up to second order in the fermions in general bosonic backgrounds. Although there is a paper [6] with similar calculations, we point out that in that work the component expansion of the superfields corresponds to the so-called gauge completion expansion [9]. This method is based on a laborious comparison order-by-order between component supersymmetric transformations and superspace coordinate transformations. On the contrary, the normal coordinate expansion [4] used here, gives a much more friendly framework with more powerful results (the action of the superstring is a good example). We were able to cast the action in a ‘‘covariant’’ form for supersymmetry,  $\kappa$ -symmetry and T-duality. Also, we would like to stress that we consider only up to second order terms in the fermion expansion, not because of a limitation of the method but it was a pure choice that we took for brevity. The normal coordinate expansion can be carried out up to 32nd order if you wish, since it is defined by an iterative method with very simple rules.

We have also given a generalization of the T-duality rules for these actions, that are in agreement with the rules of Hassan [2, 10] (derived using pure supergravity arguments) and could be confronted to those found in [6]. T-duality rules for the Green-Schwarz superstring in superspace formalism were studied in [11] (see also [12]). Nevertheless, these studies are given in terms of superfields and therefore the structure of the T-duality rules for the components is implicit. It will be very interesting to compare both approaches by a normal coordinate expansion on their results.

As a last comment, it seems that the superstring actions founded here are well suited for studies on quantum properties like the beta function, the extension of the action to include couplings to the dilaton, etc...

## Acknowledgments

We thank J. Gomis, M. Grisar, D. Marolf and D. Zanon for useful discussions. This work was supported by INFN, MURST and by the European Commission RTN program HPRN-CT-2000-00131, in association with the University of Torino.

## 2 Appendix T-duality rules for background fields

Here we follow the Hassan formalism of [2] for T-duality: since we wish to apply T-duality along the 9th direction, let us introduce the following useful objects ( $\tilde{m}, \tilde{n} = 0, \dots, 8$ )

$$\begin{aligned}
 \Omega &= \frac{1}{\sqrt{g_{99}}} \Gamma^{11} \Gamma_9 \Rightarrow \Omega^2 = -1 \\
 E_{mn} &= g_{mn} + B_{mn} \\
 (Q_{\pm})^m_n &= \begin{pmatrix} \mp g_{99} & \mp (g \mp B)_{9\tilde{n}} \\ 0 & \mathbf{1}_9 \end{pmatrix} \\
 (Q_{\pm}^{-1})^m_n &= \begin{pmatrix} \mp g_{99}^{-1} & -(g \mp B)_{9\tilde{n}} \\ 0 & \mathbf{1}_9 \end{pmatrix}.
 \end{aligned} \tag{20}$$

The T-duality rules for  $E_{mn}$  are<sup>6</sup>,

$$\begin{aligned}
\tilde{E}_{\tilde{m}\tilde{n}} &= E_{\tilde{m}\tilde{n}} - E_{\tilde{m}9}g_{99}^{-1}E_{9\tilde{n}} \\
\tilde{E}_{\tilde{m}9} &= E_{\tilde{m}9}g_{99}^{-1} \\
\tilde{E}_{9\tilde{m}} &= -E_{9\tilde{m}}g_{99}^{-1} \\
\tilde{E}_{99} &= g_{99}^{-1} .
\end{aligned} \tag{21}$$

For the transformation of the vielbein and the spinors, we will use the Hassan conventions to avoid ambiguities<sup>7</sup>. The transformation rules for the vielbein are

$$\tilde{e}_a^m \equiv e_{(-)a}^m = (Q_-)^m{}_n e_a^n \Rightarrow \tilde{e}_m^a \equiv e_{(-)m}^a = (Q_-^{-1})^n{}_m e_n^a . \tag{22}$$

We will also need the alternative transformed vielbein

$$e_{(+a)}^m = (Q_+)^m{}_n e_a^n = \Lambda^b{}_a e_{(-)b}^m \Rightarrow e_{(+m)}^a = (Q_+^{-1})^n{}_m e_n^a = e_{(-)m}^b \Lambda_b^a . \tag{23}$$

Therefore going from IIa to IIa, we have:

$$\begin{aligned}
y_+ &= y_1 \Rightarrow \bar{y}_+ = \bar{y}_1 \\
y_- &= -\Omega y_2 \Rightarrow \bar{y}_- = \bar{y}_2 \Omega \\
D_m^{(0)} y_+ &= (Q_+^{-1})^n{}_m (\hat{D}_n^{(0)} y_1) \\
D_m^{(0)} y_- &= -\Omega (Q_-^{-1})^n{}_m (\hat{D}_n^{(0)} y_2) \\
W_m y_+ &= -\Omega (Q_-^{-1})^n{}_m (\hat{W}_n y_1) \\
W_m y_- &= (Q_+^{-1})^n{}_m (\hat{W}_n y_2)
\end{aligned}$$

Conversely, going from IIB to IIA we have

$$\begin{aligned}
y_1 &= y_+ \Rightarrow \bar{y}_1 = \bar{y}_+ \\
y_2 &= \Omega y_- \Rightarrow \bar{y}_2 = -\bar{y}_- \Omega \\
\hat{D}_m^{(0)} y_1 &= (Q_+^{-1})^n{}_m (D_n^{(0)} y_+) \\
\hat{D}_m^{(0)} y_2 &= \Omega (Q_-^{-1})^n{}_m (D_n^{(0)} y_-) \\
\hat{W}_m y_1 &= \Omega (Q_-^{-1})^n{}_m (W_n y_+) \\
\hat{W}_m y_- &= (Q_+^{-1})^n{}_m (W_n y_-)
\end{aligned}$$

## References

- [1] E. Bergshoeff, R. Kallosh, T. Ortin, D. Roest and A. Van Proeyen, *Class. Quant. Grav.* **18** (2001) 3359 [arXiv:hep-th/0103233].
- [2] S. F. Hassan, *Nucl.Phys.* B568(2000)145, [hep-th/9907152](#).
- [3] M. Grisar, M. Knutt, *Phys.Lett.* B500(2001)188, [hep-th/0011173](#) .

<sup>6</sup>Here and in the rest of this work, in ambiguous cases, we will put a tilde over the transformed fields

<sup>7</sup>We recalled that there are two possible choices  $e_{(\pm)a}^m$  for the transformed vielbein, related by a Lorentz transformation  $\Lambda^b{}_a$ .

- [4] I. McArthur, *Class. Quant. Grav.* **1** (1984) 233.  
 J. Atick and A. Dhar, *Nucl. Phys.* B284 (1987) 131.  
 M.T. Grisaru and D. Zanon, *Nucl. Phys.* B310 (1988) 57.  
 M.T. Grisaru, H. Nishino and D. Zanon, *Nucl. Phys.* B314 (1989) 363.  
 M.T. Grisaru, M.E. Knutt and W. Siegel, *Nucl. Phys.* B523 (1998) 663, (hep-th/9711120).  
 S.J. Gates, M.T. Grisaru, M.E. Knutt and W. Siegel, *Phys. Lett.* B421 (1998) 203,  
 (hep-th/9711151).
- [5] bibitemAlvarez:1994dn E. Alvarez, L. Alvarez-Gaume and Y. Lozano, *Nucl. Phys. Proc. Suppl.* **41** (1995) 1 [arXiv:hep-th/9410237].
- [6] M. Cvetič, H. Lu, C.N. Pope and K.S. Stelle, *Nucl. Phys.* B573 (2000) 149,  
 (hep-th/9907202).
- [7] E. Bergshoeff, E. Sezgin and P. K. Townsend, *Phys. Lett. B* **189** (1987) 75.  
 E. Bergshoeff, E. Sezgin and P. K. Townsend, *Annals Phys.* **185** (1988) 330.
- [8] E. Bergshoeff and P. K. Townsend, *Nucl. Phys. B* **490** (1997) 145 [arXiv:hep-th/9611173].
- [9] B. de Wit, K. Peeters and J. Plefka, *Nucl. Phys. B* **532** (1998) 99 [arXiv:hep-th/9803209].  
 A. Dasgupta, H. Nicolai and J. Plefka, *JHEP* 0005 (2000) 007, (hep-th/0003280).  
 J. Plefka, (hep-th/0009193).  
 K. Millar, W. Taylor and M. van Raamsdonk, (hep-th/0007157).
- [10] S. F. Hassan, *Nucl.Phys.Proc.Suppl.* 102(2001)72 , hep-th/0103149.  
 S. F. Hassan, *Nucl.Phys.* B583(2000)431, hep-th/9912236.
- [11] B. Kulik and R. Roiban, *JHEP* **0209** (2002) 007 [arXiv:hep-th/0012010].
- [12] Igor A. Bandos, Bernard Julia, hep-th/0303075.