

# Coleman's theorem on physical assumptions for no Goldstone bosons in two dimensions

Chigak Itoi

Department of Physics, Nihon University,  
Kanda-Surugadai, Chiyoda, Tokyo 101-8308, Japan

November 27, 2018

## Abstract

Thirty years ago, Coleman proved that no continuous symmetry is broken spontaneously in a two-dimensional relativistic quantum field theory. In his argument, however, it is difficult to understand the physical meaning of the assumption of no infrared divergence. I derive the same result directly from the cluster property of a local field regarded as a physically acceptable assumption.

*Introduction.* Coleman's theorem claims that the vacuum expectation value of a charged field must vanish in a two-dimensional relativistic quantum field theory with a continuous symmetry [1]. This is a remarkable nature of relativistic field theories, while in some non-relativistic systems in one space dimension a continuous symmetry is broken spontaneously at zero temperature. This theorem is quite useful so that his renowned paper has been cited many times for thirty years. In a few articles [2, 3], however, one finds criticism to Coleman's theorem based on some counter examples. Always, the assumption for no Goldstone bosons comes into question, in addition to an incorrect representation of Coleman's lemma. Here, I discuss a physical problem in the assumption of the theorem. Coleman assumed that a field theory with an infrared divergence in Wightman functions is ill-defined and therefore no massless scalar fields exist in two dimensions. Without this assumption one cannot show the theorem. The title of his paper "There are no Goldstone bosons in two dimensions" is not the result, but an assumption in

the theorem. Generally in physics, however, infrared divergence is important phenomenon. Therefore, the existence of infrared divergence should not be a criterion to exclude ill-defined theories. For example, the ferromagnetic quantum Heisenberg model in one space dimension has a divergent magnetic susceptibility at zero temperature and the  $SU(2)$  symmetry is broken spontaneously. In this case, the infrared divergence helps the ferromagnetic order to compete with the quantum fluctuation. Generally for a locally interacting lattice system with spontaneously broken continuous symmetry in one space dimension, Momoi proved that the corresponding susceptibility should diverge at zero temperature [4]. The infrared divergence plays a physically important role in this case. On the other hand, an infrared divergence is not so serious in some other cases. For example, one can use a massless scalar field in two dimensions to bosonize a fermion field or some others with different scaling dimensions. In this case, the infrared divergence can be easily removed from the physically important correlation functions. Therefore, the massless scalar field is not ill-defined even in two dimensions. The mathematical assumption to exclude massless scalar fields because of no infrared divergence in Coleman's argument is not understandable physically.

In this paper, I show impossibility of non-zero vacuum expectation value only for a local scalar field with a cluster property. This property is necessary for a local field above a healthy vacuum and sufficient for the theorem in any subtle case. Although the cluster property of a local field is proved in a primary representation of a  $C^*$ -algebra [5], it is a physical requirement for local fields from the viewpoint of physicists. The theorem proved on this assumption is indeed sufficiently useful for physical systems. It is neither surprising nor problematic to obtain a non-zero vacuum expectation value of a scalar field with no cluster property in two dimensions. All examples [2, 3] with a non-zero vacuum expectation value of scalar fields are not true counter examples to the theorem from this viewpoint. In the following, I present my results along Coleman's style.

*Assumptions.* A charge operator  $Q$  for the continuous symmetry and a charged scalar field  $\phi(x)$  satisfy a commutation relation

$$[Q, \phi(x)] = \phi(x). \tag{1}$$

The charge  $Q \equiv \int dx_1 j_0(x_0, x_1)$  is defined from the current  $j_\mu(x)$  which obeys  $\partial^\mu j_\mu(x) = 0$ . Despite only an abelian symmetry considered here for

simplicity, the present argument can be extended to a non-abelian symmetry with charges  $Q_i$  and for fields  $\phi_j(x)$  satisfying a homogeneous relation  $[Q_i, \phi_j(x)] = C_{ij}^k \phi_k(x)$ . I assume also the relativistic covariance of the theory. I will show that the vacuum expectation value of the charged local field must vanish if the correlation functions of this field satisfy the cluster property. The cluster property of the two-point function requires

$$\langle 0 | \delta\phi(x)^\dagger \delta\phi(y) | 0 \rangle \rightarrow 0, \quad (2)$$

as a space-like separation  $(x - y)^2 \rightarrow -\infty$ , where the deviation is defined by  $\delta\phi(x) \equiv \phi(x) - \langle 0 | \phi(x) | 0 \rangle$ . For example in a free field representation in unperturbed or perturbed conformal field theories, a compact boson  $\varphi(x)$  itself does not have the cluster property. On the other hand, a physical field operator with a positive conformal dimension, say  $e^{2\pi i\varphi/R}$  with a compact boson radius  $R$  or a current  $\partial_\mu\varphi(x)$ , has the cluster property. The infrared divergence can be easily removed from correlation functions of the physical operators by some suitable rescaling. Hereafter, we assume that the physically meaningless divergence is removed. The Fourier transformed two point functions are

$$F(k) = \int d^2x e^{ikx} \langle 0 | \delta\phi(x)^\dagger \delta\phi(0) | 0 \rangle = \rho_0(k^2) \theta(k_0), \quad (3)$$

$$F_\mu(k) = \int d^2x e^{ikx} \langle 0 | j_\mu(x) \delta\phi(0) | 0 \rangle = (\sigma k_\mu \delta(k^2) + \epsilon_{\mu\nu} k^\nu \rho_1(k^2)) \theta(k_0), \quad (4)$$

$$F_{\mu\nu}(k) = \int d^2x e^{ikx} \langle 0 | j_\mu(x) j_\nu(0) | 0 \rangle = (k_\mu k_\nu - \eta_{\mu\nu} k^2) \rho_2(k^2) \theta(k_0). \quad (5)$$

by using the relativistic covariance, the positivity of energy, the permutation symmetry and the current conservation. I assume the spectral condition  $\rho_i(k^2) = 0$  for  $k^2 < 0$  ( $i = 0, 1, 2$ ), and the positive semi-definiteness

$$\rho_0(k^2) \geq 0, \quad \rho_2(k^2) \geq 0, \quad (6)$$

and also I assume their integrability over  $0 < k^2 < \infty$ . Note that the vacuum expectation value is proportional to the coefficient  $\sigma$

$$\langle 0 | \phi(0) | 0 \rangle = \langle 0 | \int dx_1 [j_0(x_0, x_1), \delta\phi(0)] | 0 \rangle = \frac{\sigma}{4\pi}. \quad (7)$$

**Lemma.** *Let  $H(\kappa^2)$  be a positive semi-definite function of a single variable  $\kappa^2 \geq 0$  with a bound  $H(\kappa^2) \leq CG(\kappa^2)$  by some constant  $C > 0$  and a function*

$$G(\kappa^2) \equiv \int_{-\infty}^{\infty} dp \frac{\cos p}{\sqrt{p^2 + \kappa^2}}. \quad (8)$$

For the distribution function  $\rho_0(k^2)$  in Eq.(3) of a field  $\phi(x)$  with the cluster property (2), the following limit vanishes

$$\lim_{\lambda \rightarrow \infty} \int_0^\infty \frac{d\kappa^2}{\lambda} \rho_0\left(\frac{\kappa^2}{\lambda}\right) H(\kappa^2) = 0. \quad (9)$$

*Proof of Lemma.* A spectral representation of the Fourier transformed two-point function is obtained by changing the integration variable from  $k_0$  to  $m^2 = k_0^2 - k_1^2$ . In this representation, the cluster property (2) implies

$$\lim_{|x_1| \rightarrow \infty} \int_0^\infty dm^2 \rho_0(m^2) \int_{-\infty}^\infty dk_1 \frac{\cos k_1 x_1}{\sqrt{k_1^2 + m^2}} = 0. \quad (10)$$

By rescaling the integration variables, this condition is rewritten into

$$\lim_{|x_1| \rightarrow \infty} \int_0^\infty \frac{d\kappa^2}{|x_1|^2} \rho_0\left(\frac{\kappa^2}{|x_1|^2}\right) G(\kappa^2) = 0, \quad (11)$$

By the bound of  $G(\kappa^2)$  and the positive semi-definiteness of each function, the right hand side of Eq.(9) is bounded by

$$\int_0^\infty \frac{d\kappa^2}{\lambda} \rho_0\left(\frac{\kappa^2}{\lambda}\right) H(\kappa^2) \leq C \int_0^\infty \frac{d\kappa^2}{\lambda} \rho_0\left(\frac{\kappa^2}{\lambda}\right) G(\kappa^2). \quad (12)$$

The right hand side vanishes in the limit with the correspondence  $\lambda = |x_1|^2$  in Eq.(11). Thus the lemma has been proved.

**Theorem.** *The vacuum expectation value of a charged scalar field with the cluster property vanishes in a two-dimensional relativistic quantum field theory with a conserved charge with respect to the continuous symmetry.*

*Proof of Theorem.* I evaluate terms in Coleman's correlation inequality

$$\left[ \int d^2k F(k) |h_\lambda(k)|^2 \right] \left[ \int d^2k F_{00}(k) |h_\lambda(k)|^2 \right] \geq \left| \int d^2k F_0(k) |h_\lambda(k)|^2 \right|^2, \quad (13)$$

where I choose a test function

$$h_\lambda(k_0, k_1) \equiv f(\lambda(k_0^2 - k_1^2))g(k_1^2)$$

by a bounded positive semi-definite function  $f(\kappa^2)$  and  $g(\kappa^2)$  with a compact support. The support of  $f(\kappa^2)$  includes the origin and that of  $g(\kappa^2)$  does not. For example, one of the simplest choice is  $f(\kappa^2) = \theta(a^2 - \kappa^2)$  and  $g(k_1^2) = \theta(k_1^2 - b^2)\theta(c^2 - k_1^2)$  with positive numbers  $a < b < c$ . This test function differs from the original one, but has the same important property. The support of  $h_\lambda(k)$  lies on the light cone in the large  $\lambda$  limit. Now, I evaluate the first term in the inequality (13)

$$\begin{aligned} \int d^2k F(k) |h_\lambda(k)|^2 &= \int_0^\infty dm^2 \rho_0(m^2) |f(\lambda m^2)|^2 \int_{-\infty}^\infty dk_1 \frac{|g(k_1^2)|^2}{2\sqrt{k_1^2 + m^2}} \\ &\leq \int_0^\infty \frac{d\kappa^2}{\lambda} \rho_0\left(\frac{\kappa^2}{\lambda}\right) |f(\kappa^2)|^2 \int_{-\infty}^\infty dk_1 \frac{|g(k_1^2)|^2}{2|k_1|}. \end{aligned}$$

Since the support of the function  $g(k_1^2)$  does not include the origin, the factor  $\int dk_1 |g(k_1^2)|^2 / |k_1|$  is finite. The bounded positive semi-definite function  $|f(\kappa^2)|^2$  with a compact support is bounded by  $CG(\kappa^2)$  with some positive constant  $C$ , then the first term in the inequality (13) vanishes in the limit by the lemma

$$\lim_{\lambda \rightarrow \infty} \int d^2k F(k) |h_\lambda(k)|^2 = 0. \quad (14)$$

Next, I evaluate the second term in the left hand side of (13)

$$\int d^2k F_{00}(k) |h_\lambda(k)|^2 = \int_0^\infty dm^2 \rho_2(m^2) |f(\lambda m^2)|^2 \int_{-\infty}^\infty dk_1 \frac{k_1^2 |g(k_1^2)|^2}{\sqrt{k_1^2 + m^2}} \quad (15)$$

which is also bounded from above because of the boundedness of the integrand and the integrability of  $\rho_2(m^2)$ . Finally, I show that the right hand side of (13) is proportional to  $|\sigma|^2$  with a non-zero coefficient. The  $\delta$ -functional singularity in Eq.(4) gives

$$\int d^2k F_0(k) |h_\lambda(k)|^2 = \sigma f(0) \int_{-\infty}^\infty dk_1 |g(k_1^2)|^2. \quad (16)$$

Therefore, Coleman's correlation inequality (13) in the large  $\lambda$  limit prohibits the non-zero vacuum expectation value  $\sigma/4\pi$  of the charged local field with the cluster property. The theorem has been demonstrated in more physical situation.

*Note.* In this paper, the theorem is shown by a test function with a compact support including the light cone in a momentum space as in Coleman's proof. This corresponds to setting a test state extended over the whole space-time. To prove the theorem with a test state on a space-like surface as in the famous Mermin-Wagner theorem [6], Bogoliubov's inequality is no use at zero temperature. One can use more complicated Shastry's inequality, which was employed to prove no antiferromagnetic order in the antiferromagnetic quantum Heisenberg chain [7]. Also simpler Coleman's inequality works with a test function  $h_\lambda(k) = g(\lambda k_1^2)$ , where the  $k_0$ -independence implies no support compactness and a test state on a space-like surface. In either case, one needs the lemma under a relaxed condition on  $H(\kappa^2)$  when the bound by  $G(\kappa^2)$  is available only in some neighborhood of the origin. This version of lemma allows one to prove the theorem with the test state on a space-like surface.

*Acknowledgements* The author would like to thank T. Nihei and M. Yamanaka for reading the manuscript and helpful comments.

## References

- [1] Coleman, S. : There are no Goldstone bosons in two dimensions. *Commun. Math. Phys.* **31**, 259 (1973).
- [2] Nakanishi, N. : Free massless scalar field in two-dimensional space-time. *Prog. Theor. Phys.* **57**, 269 (1977).
- [3] Faber, M. and Ivanov, A. N. : Comments on Coleman's paper "There are no Goldstone bosons in two dimensions". hep-th/0204237.
- [4] Momoi, T. : Quantum fluctuations in quantum lattice systems with continuous symmetry. *J. Stat. Phys.* **85**, 193 (1996).
- [5] Haag, R. : *Local Quantum Physics* (2nd. ed.), Springer-Verlag (1996).
- [6] Marmin, N. D. and Wagner, H. : Absence of ferromagnetism or antiferromagnetism in one- or two-dimensional isotropic Heisenberg model. *Phys. Rev. Lett.* **17**, 1133 (1966).
- [7] Shastry, B. S. : Bounds for correlation functions of Heisenberg antiferromagnet. *J. Phys. A: Math. Gen.* **25**, L249 (1992).