

# Comments on BPS Bound State Decay

Anatoly Dymarsky<sup>†</sup> and Dmitry Melnikov<sup>‡</sup>

*Department of Physics at Moscow State University  
Vorobjevy Gory, 119899 Moscow, Russia*

*Institute for Theoretical and Experimental Physics  
B.Chermushkinskaya 25, 117259 Moscow, Russia*

March, 2003

## Abstract

In N=2 SYM theory with matter non-conservation of the S-charge of a BPS state under monodromies leads to so-called BPS state “decay”. A mechanism for such a behavior on the semiclassical level could be established through consideration of soliton-fermion bound state. Solutions to classical equations of motion allow to observe the BPS state “decay” *in vivo*.

---

## 1 Introduction

In the work of Seiberg and Witten [1] concerning N=2 SYM with matter in fundamental representation of SU(2) the following formula for central charge is established:

$$Z = n_m a_D + n_e a + \sum_{i=1}^{N_f} \frac{m_i S_i}{\sqrt{2}} \quad (1)$$

---

<sup>†</sup>dymarsky@gate.itep.ru

<sup>‡</sup>melnikov@gate.itep.ru

where besides the magnetic and electric charges an abelian global charge appears. This charge, called the S-charge or the spectral asymmetry, arises due to global  $SU(N_f)$  symmetry which is generally broken down to  $U(1)^{N_f}$  by the hypermultiplet mass term in the action. The S-charge is carried only by the fields from hypermultiplet and on equal footing with  $n_e$  and  $n_m$  it parameterizes the BPS spectrum of the N=2 theory.

The moduli space possesses a certain number of singularities. In this paper we are going to consider those singularities that correspond to massless quarks. From the central charge formula (1) it follows that  $Z$  vanishes for the states with

$$n_m = 0, \quad n_e = \pm 1, \quad S = \pm 1$$

at a special point  $a_0 = \pm m/\sqrt{2}$  on the moduli space.

The complex coupling  $\tau$  has a logarithmic singularity at this point

$$\tau(a) = -\frac{i}{2\pi} \ln(a - a_0), \tag{2}$$

$$a_D = c - \frac{i}{2\pi} (a - a_0) \ln(a - a_0).$$

Considering  $a$  and  $a_D$  as the functions on the complex  $a$  plane and investigating their monodromies around the point  $a_0$  we find

$$a \rightarrow a, \quad a_D \rightarrow a + a_D - \frac{m}{\sqrt{2}}.$$

Since the central charge should be invariant, the charges of a quantum state consisted of a monopole of magnetic charge  $n_m$  and quark with charges  $n_e$  and  $S$  should be shifted by this rotation:

$$(n_m, n_e, S) \rightarrow (n_m, n_e - n_m, S + n_m)$$

By multiple action of the monodromy in the presence of a magnetic charge  $n_m \neq 0$  in principle we could make the value of the S-charge arbitrarily large.

On the other hand for the quark-monopole bound state the value of the S-charge is bounded, since the only fields carrying S-charge are fermion modes, that obey the Pauli exclusive principle. Therefore starting from one particle state and encircling the singularity we come to two particle state.

In the limit  $m \gg \Lambda$  the singularity becomes semiclassical and hence the simple dynamical analysis of the theory near the singularity is appropriate. Below we consider the fermion equations of motion in the background of monopole fields. We find explicitly a normalizable fermion mode corresponding to the bound state, which carries S-charge. It is from the analysis of the properties of the classical solution that the mechanism of the BPS state decay becomes explicit.

Somewhat similar considerations were made by Henningson in [2], where the conditions for the existence of normalizable solution were investigated. The latter result is consistent with simple analysis of Marginal Stability from BPS Mass Formula performed in [3].

The notion of BPS state decay was originally introduced in the works by Bilal and Ferrari [4], where this phenomenon was first derived. Also it was the S-charge, that was calculated in simpler models under the name of spectral asymmetry in the works of [5]. Recently the new derivation of this charge based on holomorphicity was proposed by Ferrari [6]. In the present paper we rederive the result of [2] and present an obvious picture which describes classically the phenomenon of the BPS state decay.

## 2 Derivation of the fermion mode

### 2.1 Solution to Dirac equation

Consider  $N = 2$  SYM theory with  $SU(2)$  gauge group and matter hypermultiplets in fundamental representation.<sup>§</sup> This theory contains monopoles in its spectrum. There are also solutions that correspond to matter fermions in the external monopole field. The question is whether these solutions can be normalizable. In other words, whether there exist soliton-fermion bound states to which we refer as to one particle states. As we shall see below, such normalizable solutions exist only in some region of the moduli space. The point of singularity for the massless quark happens to belong to the boundary of this region, providing a natural explanation for the bound state decay described above.

The fermionic components of the hypermultiplet satisfy the Dirac equation

$$i\gamma^\mu D_\mu \Psi - (\hat{m} + \sqrt{2}\hat{A})\Psi = 0, \quad (3)$$

$$\hat{m} + \sqrt{2}\hat{A} = \Re(m + \sqrt{2}A) + i\gamma^5 \Im(m + \sqrt{2}A).$$

where the Dirac spinor  $\Psi$  is composed of two Weyl spinors as follows:

$$\Psi = \begin{pmatrix} \psi_q \\ \psi_{\bar{q}} \end{pmatrix}. \quad (4)$$

These two Weyl spinors belong to fundamental and antifundamental  $N=1$  chiral multiplets.

Working in the Gauss gauge  $A_0 = 0$ , substitute  $D_0$  by the energy eigenvalue  $iE$ . Now notice that the hamiltonian commutes with the operator  $\Gamma = \gamma^0\gamma^5$  in the case  $\Im(m) + \sqrt{2}\Im(A) = 0$ . Thus general solution to (3) could be found as the sum of  $\Gamma$  eigenfunctions  $\Psi = \Psi^+ + \Psi^-$ .

Substituting  $\Psi^+$  instead of  $\Psi$  we split the equation (3):

$$[\sigma^k D_k - \Re(m + \sqrt{2}A)]\psi^+ = 0, \quad (5)$$

$$[E - \Im(m + \sqrt{2}A)]\psi^+ = 0.$$

Here  $\psi^+$  is a chiral Weyl component of  $\Psi$ .

In the consideration above we treat  $m$  as a real constant in contrast with scalar field  $A$  which is generally complex. However it would be more useful for the future computations to keep  $A$  real. Since there is a  $U(1)_R$  symmetry acting on the fields, we could choose such a  $U(1)$  rotation

$$A \rightarrow e^{2i\alpha} A,$$

$$m \rightarrow e^{2i\alpha} m, \quad (6)$$

$$\Psi \rightarrow e^{-i\alpha\gamma^5} \Psi,$$

---

<sup>§</sup>Here we restrict ourselves to the case of  $N_f = 1$ . For more detailed review of  $N = 2$  actions see for example [7].

under which complexity of  $A$  flows into complexity of  $m$ . Although this symmetry is in general anomalous, this is not the case in the semiclassical region. Then solution to the second equation of (5) implies that the bound-state mass spectrum satisfies

$$E = \Im m(\mathcal{M}),$$

where  $\mathcal{M}$  now defines a complex mass parameter.

In the case of real scalar field  $A$  we could use the standard radially symmetric solution for 't Hooft-Polyakov monopole [8] in the BPS limit<sup>¶</sup>:

$$\begin{aligned} \sqrt{2}A^i &= an^i(1 - F(r)), \\ A_i^a &= \epsilon^{aij} \frac{n^j}{r}(1 - H(r)), \end{aligned} \tag{7}$$

where  $n^i$  is a unit vector,  $F$  and  $H$  are known functions of radius  $r$ . For future convenience we chose the normalization of the scalar field  $A$  in (7) slightly different from that in [1]. In our conventions it has asymptotic  $A \rightarrow a/\sqrt{2}$ . Furthermore Seiberg and Witten renormalize the charges of the BPS states in such a way that  $a \rightarrow a/2$ . So our  $a$  will be different from that of Seiberg and Witten by the factor of  $(2\sqrt{2})^{-1}$ .

Substituting (7) into (5) we find a solution in the form

$$(\psi^+)_a^\alpha = \delta_a^\alpha \chi_0 \xi + n^i (\sigma^i)^\alpha_\beta \delta_a^\beta \eta_0 \zeta, \tag{8}$$

where  $\alpha$  and  $a$  are spinor and color indices respectively. Functions  $\chi_0$  and  $\eta_0$ , solving the equations at zero  $m$  are defined as follows:<sup>||</sup>

$$\begin{aligned} \chi_0 &= \frac{1}{\sqrt{\rho \operatorname{sh} \rho}} \operatorname{th} \frac{\rho}{2}, \\ \eta_0 &= \frac{1}{\sqrt{\rho \operatorname{sh} \rho}} \operatorname{cth} \frac{\rho}{2}. \end{aligned} \tag{9}$$

Functions  $\xi$  and  $\zeta$  satisfy the system of the first order differential equations

$$\begin{aligned} \xi' &= r_0 m \zeta \operatorname{cth}^2 \left( \frac{\rho}{2} \right), \\ \zeta' &= r_0 m \xi \operatorname{th}^2 \left( \frac{\rho}{2} \right), \end{aligned} \tag{10}$$

which implies the pair of second order equations

$$\begin{aligned} \xi'' + \frac{2\xi'}{\operatorname{sh} \rho} - r_0^2 m^2 \xi &= 0, \\ \zeta'' - \frac{2\zeta'}{\operatorname{sh} \rho} - r_0^2 m^2 \zeta &= 0. \end{aligned} \tag{11}$$

---

<sup>¶</sup>We neglect the back reaction of matter fields to monopole fields assuming that in semiclassical limit the ratio of corresponding masses is small.

<sup>||</sup>Here we define  $\rho = \frac{r}{r_0}$ ,  $r_0^{-1} = a$ .

Explicit solutions to this equations could be found in terms of hypergeometric functions. We shall find normalizable solutions after an analysis of asymptotic behavior of the general solutions to (5).

The solution presented above is one of positive chirality. It could be shown as well that for negative chirality the similar solution could not be made normalizable. It is also natural to assume [2] that there are no other discrete spectrum solutions to (3).

## 2.2 Analysis of the Solution

In this subsection we investigate the asymptotic of the solution to (3) and rederive the conditions under which this solution exists [2]. Using the asymptotic we also derive the explicit solution in terms of hypergeometric functions.

Introduce a pair of functions  $\chi, \eta$  that will characterize the asymptotic behavior of solution (8):

$$\chi = \chi_0 \xi, \quad \eta = \eta_0 \zeta. \quad (12)$$

Since we are interested in soliton-fermion bound states, the fermion mode we are looking for should be a normalizable one, i.e. it should decrease fast enough at infinity and be regular at the origin, namely

$$\chi(0) = \text{const}, \quad \eta(0) = 0$$

In the  $\rho \rightarrow 0$  limit the fermion mode behaves as the solution to the equations

$$\begin{aligned} \xi'' + \frac{2\xi'}{\rho} - r_0^2 m^2 \xi &= 0, \\ \zeta'' - \frac{2\zeta'}{\rho} - r_0^2 m^2 \zeta &= 0. \end{aligned} \quad (13)$$

The regular solutions to (13) are as follows:

$$\xi = C \frac{\text{sh}(r_0 m \rho)}{\rho}, \quad (14)$$

$$\zeta = C(r_0 m \rho - 1)e^{r_0 m \rho} + C(r_0 m \rho + 1)e^{-r_0 m \rho},$$

with the coefficients fixed by asymptotical behavior of (10).

Assume that we have a solution that is regular at the origin. From the asymptotic (14) it follows that the first derivative of this solution is zero and the second derivative is positive at this point. Furthermore, the second derivative in exact equations (11) is positive wherever first derivative vanishes. Therefore such solutions could have only minima and cannot be normalizable. This means that solution to (11) is either regular at the origin and divergent at infinity or regular at infinity and divergent at the origin. However if we look at (9), we shall see that under certain conditions we could construct a normalizable zero-mode. The regular at the origin and increasing at infinity solution to (11) has the asymptotic

$$\xi, \zeta \sim e^{r_0 m \rho}. \quad (15)$$

It follows from (9) that the functions  $\chi_0, \eta_0$  have the asymptotic

$$\chi_0, \eta_0 \sim \frac{e^{-\rho}}{\sqrt{\rho}}, \quad (16)$$

and that for the functions  $\chi, \eta$

$$\chi, \eta \sim \sqrt{\frac{1}{\rho}} e^{(r_0 m - \frac{1}{2})\rho}. \quad (17)$$

We see that for the existence of normalizable zero mode the following conditions should be satisfied:

$$2r_0 m < 1, \quad \text{or} \quad \frac{2m}{a} < 1, \quad (18)$$

or  $a > m/\sqrt{2}$  in the standard normalization of [1]. This result was originally derived in [2], and then was confirmed by CMS considerations in [3] in the weak coupling limit.

Now turn to the equations (11). Changing variables from  $\rho$  to  $x = (\text{ch}\rho + 1)/2$  we obtain a pair of hypergeometric equations

$$\begin{aligned} x(x-1)y_1'' + (x + \frac{1}{2})y_1' - r_0^2 m^2 y_1 &= 0, \\ x(x-1)y_2'' + (x - \frac{3}{2})y_2' - r_0^2 m^2 y_2 &= 0. \end{aligned} \quad (19)$$

Solving (19) and expressing the solution in terms of original variable  $\rho$  we find the unknown functions  $\xi$  and  $\zeta$ :

$$\begin{aligned} \xi(\rho) &= CF\left(\frac{m}{a}, -\frac{m}{a}, -\frac{1}{2}, \text{ch}^2 \frac{\rho}{2}\right), \\ \zeta(\rho) &= \tilde{C}F\left(\frac{m}{a}, -\frac{m}{a}, \frac{3}{2}, \text{ch}^2 \frac{\rho}{2}\right). \end{aligned} \quad (20)$$

Here we took into account the asymptotic of solution determined above. The asymptotic of (10) also fixes the relation  $C = \tilde{C}$ .

### 2.3 BPS State Decay

From the considerations above we found that normalizable fermion mode with the energy defined by complex mass parameter  $E = \Im m \mathcal{M}$  exists only in the region of moduli space defined as\*\*

$$\Re e \mathcal{M} = m < \frac{|a|}{2}. \quad (21)$$

Now for complete understanding of the process we need one more thing, namely to find what region of the moduli space corresponds to the solutions to Dirac equation (3) from

---

\*\*In the fermion equations  $a$  was treated as  $v\text{ev}$  of real field  $A$ . To define this condition for all complex moduli plane we substitute the latter by absolute value of complex coordinate  $|a|$ .




Figure 1: The region of discrete spectrum (inclined semiplane) is continuously attached to the region of continuous spectrum (cone) by the line  $E = \Im m \mathcal{M}$  which goes through the point  $a = 2m$ .

continuous spectrum, i.e. non-bound BPS states. Considering asymptotic of (3) at infinity we obtain the system of linear differential equations, neglecting exponentially decaying non constant coefficients. The corresponding characteristic equation gives for fixed energy  $E$

$$E = \pm \left| m - \frac{a}{2} \right|. \quad (22)$$

This condition defines a cone in the  $E - a$  or equivalently in the  $E - \mathcal{M}$  parameter space with the singularity at  $a = 2m$  (Fig.1). Further we substitute the investigation on the complex  $a$  plane by that on the  $\mathcal{M}$  plane, which is equivalent due to  $U(1)$  symmetry. Continuous spectrum belongs to the interior of this cone. The region of discrete spectrum is an inclined plane  $E = \Im m \mathcal{M}$ , which is bounded due to the condition (21).

Now a closed contour on the complex  $\mathcal{M}$  corresponds to some 3 dimensional curve, which is restricted to belong either to inclined plane of discrete spectrum or to the interior of the continuous spectrum cone. Since the contour must be closed and smooth, this curve must have a spiral form. This means that encircling the singular point we cannot return to the initial value of energy. Furthermore the final point of a spiral curve belongs to the state from continuous spectrum, which means a decay of the initial bound state.

To clarify what happens when we encircle the singularity consider the dependence of energy  $E$  on the angle of rotation  $\varphi$  in the  $\mathcal{M}$  plane (Fig.2). We start at some real mass value  $m_0$  and hence at zero energy. At  $\varphi = \pi/2$  we reach the continuous spectrum. For  $\varphi$  from  $\pi/2$  to  $3\pi/2$  there is no any bound state, i.e. the initial bound state has decayed. After a full rotation  $\varphi = 2\pi$  the mode from continuous spectrum cannot descend to discrete spectrum and the only allowed values of energy will be  $E \geq |m_0 - a/2|$ . However if we move in the opposite direction starting from  $\varphi = 2\pi$  we shall also come to continuous spectrum at  $\varphi = 3\pi/2$ , but in the part of it which is below zero level. Since we move adiabatically and all the states for fermion below zero are occupied we conclude that after encircling the singularity one fermion mode runs to continuous spectrum and another one comes from the Dirac sea below.




Figure 2: The initial bound state decays after crossing the point of stability at  $\varphi = \pi/2$ . Another fermion mode replaces it rising from the continuous spectrum below.

### 3 Conclusions

Investigations of the properties of the classical solutions to (3) tells us that the adiabatic motion around the singularity is severely restricted in acceptable values of energy  $E$ . During this motion we necessarily cross the Curve of Marginal Stability for bound state. For an initial state this happens only once and that is why this state remains finally unbounded. On the other hand another fermion line comes from the continuous spectrum of Dirac sea below and forms a new bound state, thus confirming the predicted shift in the S-charge

$$S \rightarrow S + n_m.$$

We are grateful to K.Selivanov for initiating this work and various support and useful discussions at the different stages. We would also like to thank D.Gal'tsov, A.Gorsky and V.Ch.Zhukovsky for numerous stimulating discussions. A special thank should be said to S.Dubovsky, A.Vainshtein, A.Solovyov and K.Zarembo for clarifying some subtle questions. This work would not have appeared without support of A.Alexandrov, I.Gordeli, S.Klevtsov, G.Nozaдзе and V.Poberezhny. A.D. would like to thank the Cargèse 2002 ASI, where the part of work was done. Analogously D.M. would like to thank for hospitality Department of Theoretical Physics at Uppsala University. The work was supported in part by grants RFBR 01-02-17682, RFBR 00-02-06026, INTAS 00-334, by the Russian President's grant 00-15-99296 (A.D.), and by grants RFBR 01-01-00549, INTAS 00-561, the Russian President's grant 00-15-99296 (D.M.).

### References

- [1] N. Seiberg and E. Witten, Nucl. Phys. B **431** (1994) 484 [arXiv:hep-th/9408099].
- [2] M. Henningson, Nucl. Phys. B **461**, 101 (1996) [arXiv:hep-th/9510138].
- [3] A. Ritz, A. Vainshtein, Nucl. Phys. B **617**, 43 (2001) [arXiv:hep-th/0102121].

- [4] A. Bilal, F. Ferrari, Nucl. Phys. B 469 (1996) 387–402 [arXiv:hep-th/9602082];  
A. Bilal, F. Ferrari, Nucl. Phys. B 480 (1996) 589–619 [arXiv:hep-th/9605101];  
F. Ferrari, Nucl. Phys. B 501 (1997) 53–96 [arXiv:hep-th/9702166].
- [5] M. B. Paranjape and G. W. Semenoff, Phys. Lett. B **132** (1983) 369;  
A. J. Niemi, M. B. Paranjape and G. W. Semenoff, Phys. Rev. Lett. **53** (1984) 515.
- [6] F. Ferrari, Phys. Rev. Lett. 78 (1997) 795–798 [arXiv:hep-th/9609101].
- [7] L. Alvarez-Gaume and S. F. Hassan Fortsch.Phys. 45 (1997) 159-236  
[arXiv:hep-th/9701069];  
A. Bilal hep-th/0101055.
- [8] G. 't Hooft, Nucl. Phys. B **79** 276 (1974);  
A. M. Polyakov, JETP Lett. **20** (1974) 194.