

Note on Plane Wave Quantum Mechanics

Marcus Spradlin¹ and Anastasia Volovich²

¹ Department of Physics
Princeton University
Princeton, NJ 08544

² Kavli Institute for Theoretical Physics
University of California
Santa Barbara, CA 93106

Abstract

We study the quantum mechanics of BMN operators with two scalar impurities and arbitrarily many traces, at one loop and all genus. We prove an operator identity which partially elucidates the structure of this quantum mechanics, provides some support for a conjectured formula for the free all genus two-point functions, and demonstrates that a single $\mathcal{O}(g_2^2)$ contact term arises in the Hamiltonian as a result of transforming from the natural gauge theory basis to the string basis. We propose to identify the S-matrix of this quantum mechanics with the S-matrix of string theory in the plane-wave background.

Contents

1. Introduction	1
2. Definition of the Quantum Mechanics	3
2.1. Hilbert space	3
2.2. Inner product	4
2.3. Splitting-joining operator	4
2.4. Hamiltonian	5
3. Relation to Gauge Theory	6
3.1. Hilbert space	6
3.2. Inner product	7
3.3. Splitting-joining operator	7
3.4. Hamiltonian	7
4. The String Basis	8
4.1. Some identities	8
4.2. Hamiltonian in the string basis	9
4.3. The ‘supercharge’	10
4.4. The string field theory ‘prefactor’	10
5. Discussion and Speculations	11
5.1. A puzzle at two loops	11
5.2. The S-matrix	12
Appendix A. Proof of Lemma 1	13
A.1. Relation (1) on states of the first type	14
A.2. Relation (1) on states of the second type	15
A.3. Relation (2) on states of the second type	16
A.4. Relation (2) on states of the first type	17
Appendix B. Sums	18
Appendix C. More on Σ	20

1. Introduction

Recently there has been much interest in a particular limit of the AdS/CFT correspondence in which the $\text{AdS}_5 \times S^5$ background degenerates into a plane-wave and the free string theory becomes exactly solvable [1-33]. To reach this limit one focuses in the large N limit of $\mathcal{N} = 4$ $\text{SU}(N)$ super Yang-Mills theory on those operators with large R-charge J but finite $\Delta - J$, where Δ is the scaling dimension [1]. When J is taken to be of order \sqrt{N} , many quantities of physical interest are believed to be effectively perturbative in the parameters [1,3,5]

$$\lambda' \equiv \frac{g_{\text{YM}}^2 N}{J^2}, \quad g_2 \equiv \frac{J^2}{N}, \quad (1.1)$$

despite the fact that the 't Hooft coupling $\lambda = g_{\text{YM}}^2 N$ is going to infinity in this limit.

Recent efforts have successfully matched certain matrix elements of $\Delta - J$ in the gauge theory [12,15,17], after a suitable basis transformation (to be discussed extensively below), with matrix elements of the light-cone string field theory Hamiltonian [2,9], to first (or second) order in the string coupling and first order in λ' [21,22,27]. It has been emphasized in a number of papers that many interesting aspects of the gauge theory can be studied via a simple quantum mechanical model [8,15,18,30,33]: there is a space of states (gauge theory operators), an inner product (the free gauge theory two-point function), and a Hamiltonian (given by $H = \Delta - J$ in radial quantization).

In this note we study the simplest non-trivial version of this quantum mechanics [30]: that of BMN operators with two bosonic impurities that are orthogonal scalar fields, but have arbitrarily many traces. Everything in the bulk of the paper applies at one loop (i.e., $\mathcal{O}(\lambda')$), although we comment on higher loops in the final section. It is intended that this note can be read abstractly as a study of a particular quantum mechanics by those who are not necessarily familiar with the gauge theory, string field theory, or the string bit model. However, it is hoped that the results of this paper will serve to tie together these threads a little bit.

Of particular importance in the quantum mechanics is the splitting-joining operator Σ (so-called because it can increase or decrease the number of traces by one) [8]. It has been conjectured that the free two-point functions of BMN operators $\langle 1|2\rangle_{g_2}$ at finite g_2 are related to those at $g_2 = 0$ by the identity [18]

$$\langle 1|S^{-1}|2\rangle_{g_2} = \langle 1|2\rangle_{g_2=0}, \quad S = e^{g_2\Sigma}. \quad (1.2)$$

This formula has been confirmed by several calculations and refuted by none so far. If it turns out to be incorrect, then the results of this paper remain true, albeit difficult to interpret.

In light of (1.2), we identify $S^{-1/2}$ as the transformation between the ‘gauge theory’ basis and the ‘string’ basis [21]. That is, if $|k\rangle$ is a state which corresponds to a k -trace operator, then we identify $S^{-1/2}|k\rangle$ as a state with precisely k strings. Moreover if $|k\rangle$ is the gauge theory operator which corresponds via the BMN dictionary at $g_2 = 0$ to the string state $|\tilde{k}\rangle$, then we identify $|\tilde{k}\rangle = S^{-1/2}|k\rangle$ at finite g_2 . From the Hamiltonian $H = \Delta - J$ in the gauge theory basis we can construct the basis-transformed operator $\tilde{H} = S^{1/2}HS^{-1/2}$, which we call the string Hamiltonian.

In this note we prove an identity (Lemma 1, below) satisfied by Σ which implies that

$$e^{g_2 \Sigma} H = H^\dagger e^{g_2 \Sigma}. \quad (1.3)$$

This ensures that the string Hamiltonian \tilde{H} is hermitian with respect to the inner product (1.2). Obviously this is a basic requirement for identifying \tilde{H} with the Hamiltonian of light-cone string field theory in the plane-wave background. Another consequence of the identity is that while H in the gauge theory has manifestly no contact terms according to [30], the string Hamiltonian \tilde{H} has precisely one contact term at order g_2^2 (to first order in λ') [27]. The contact term is therefore an artifact of the change of basis. Finally, our formalism allows an analytic proof of the fact that $\Delta - J$ is positive definite at one loop. Of course, the supersymmetry algebra of the gauge theory requires this to be true, but it is certainly not obvious from the form of the generators given below, so it is satisfying to see that this can be proven analytically even after one has forgotten where the quantum mechanics comes from.

We conclude the paper with a discussion of some puzzles that appear at higher loops, and some comments on the S-matrix. In particular, we propose to identify the S-matrix obtained from this quantum mechanics with the S-matrix of string theory in the plane wave background.

2. Definition of the Quantum Mechanics

In this section we review the definition the quantum mechanics of BMN operators. To this end we (1) explain the Hilbert space, (2) provide an inner product, and (3) define some useful operators — in particular, the Hamiltonian. The reader may choose to think of this as an abstract quantum mechanical model. However, some insight into its structure is naturally gained by understanding precisely how these definitions arise from the gauge theory, as will be explained in the next section.

2.1. Hilbert space

The Hilbert space is spanned by two kinds of basis vectors. The first kind is

$$|n, r_0; r_1, \dots, r_k\rangle, \quad n \in \mathbb{Z}, \quad k \geq 0, \quad r_0, r_i \in [0, 1], \quad r_0 + \sum_{i=1}^k r_i = 1, \quad (2.1)$$

and the second kind is

$$|s_1; s_2; r_1, \dots, r_k\rangle, \quad k \geq 0, \quad s_1, s_2, r_i \in [0, 1], \quad s_1 + s_2 + \sum_{i=1}^k r_i = 1. \quad (2.2)$$

The order of the $\{r_i\}$ is not significant.

2.2. Inner product

We define the inner product $\langle | \rangle$ by

$$\begin{aligned} \langle m, r'_0; r'_1, \dots, r'_l | n, r_0; r_1, \dots, r_k \rangle &= \delta_{kl} \delta_{mn} \sum_{\pi \in S_k} \prod_{i=1}^k \delta(r'_i - r_{\pi(i)}), \\ \langle n, r_0; r_1, \dots, r_l | s_1; s_2; r_1, \dots, r_k \rangle &= 0, \\ \langle s'_1; s'_2; r'_1, \dots, r'_l | s_1; s_2; r_1, \dots, r_k \rangle &= \delta_{kl} \delta(s'_2 - s_2) \sum_{\pi \in S_k} \prod_{i=1}^k \delta(r'_i - r_{\pi(i)}). \end{aligned} \quad (2.3)$$

We will use the symbol \dagger to denote the adjoint with respect to this inner product.

2.3. Splitting-joining operator

We now define a hermitian operator $\Sigma = \Sigma_+ + \Sigma_-$ which plays a central role. The splitting operator Σ_+ is defined by¹

$$\begin{aligned} \Sigma_+ |n, r_0; r_1, \dots, r_k \rangle &= \frac{(-1)^n r_0^{3/2}}{\pi^2 n^2} \int_0^{r_0} ds \sin(\pi n \frac{r_0-s}{r_0}) \sin(\pi n \frac{s}{r_0}) |r_0 - s; s; r_1, \dots, r_k \rangle \\ &+ \int_0^{r_0} dr_{k+1} \sum_{m=-\infty}^{\infty} \sqrt{\frac{(r_0-r_{k+1})^3 r_{k+1}}{r_0}} \frac{\sin^2(\pi n \frac{r_0-r_{k+1}}{r_0})}{\pi^2 (m - n \frac{r_0-r_{k+1}}{r_0})^2} |m, r_0-r_{k+1}; r_1, \dots, r_{k+1} \rangle, \\ &+ \sum_{i=1}^k \int_0^{r_i} dr_{k+1} \sqrt{r_i (r_i - r_{k+1}) r_{k+1}} |n, r_0; r_1, \dots, r_i - r_{k+1}, \dots, r_{k+1} \rangle, \\ \Sigma_+ |s_1; s_2; r_1, \dots, r_k \rangle &= \int_0^{s_1} dr_{k+1} (s_1 - r_{k+1}) \sqrt{r_{k+1}} |s_1 - r_{k+1}; s_2; r_1, \dots, r_{k+1} \rangle \\ &+ \int_0^{s_2} dr_{k+1} (s_2 - r_{k+1}) \sqrt{r_{k+1}} |s_1; s_2 - r_{k+1}; r_1, \dots, r_{k+1} \rangle \\ &+ \sum_{i=1}^k \int_0^{r_i} dr_{k+1} \sqrt{r_i (r_i - r_{k+1}) r_{k+1}} |s_1; s_2; r_1, \dots, r_i - r_{k+1}, \dots, r_{k+1} \rangle, \end{aligned} \quad (2.6)$$

¹ A technical comment: note that

$$(-1)^n \sin(\pi n y) \sin(\pi n (1-y)) = -\sin^2(\pi n y) \quad \text{for } n \in \mathbb{Z}. \quad (2.4)$$

However, we have not used this identity to simplify the first line of Σ_+ because

$$\frac{(-1)^n \sin(\pi n y) \sin(\pi n (1-y))}{n^2} \quad \text{and} \quad -\frac{\sin^2(\pi n y)}{n^2} \quad (2.5)$$

disagree when $n = 0$. We will only use (2.4) when there is no such problem.

and the joining operator is $\Sigma_- = (\Sigma_+)^{\dagger}$. (The formula is presented in appendix C in order to avoid cluttering the text too much). The nomenclature will become obvious in the next section; for now one can note that Σ_+ (Σ_-) increases (decreases) the number of ‘traces’ by one. It will also become obvious, for example, why Σ_+ never takes a state of the second type into a state of the first type.

2.4. Hamiltonian

The free Hamiltonian is defined to be

$$H_0|n, r_0; r_1, \dots, r_k\rangle = \frac{n^2}{r_0^2}|n, r_0; r_1, \dots, r_k\rangle, \quad H_0|s_1; s_2; r_1, \dots, r_k\rangle = 0. \quad (2.7)$$

It will prove convenient to introduce an operator Q_0 which is a square root of the free Hamiltonian,

$$Q_0|n, r_0; r_1, \dots, r_k\rangle = \frac{n}{r_0}|n, r_0; r_1, \dots, r_k\rangle, \quad Q_0|s_1; s_2; r_1, \dots, r_k\rangle = 0. \quad (2.8)$$

Clearly both H_0 and Q_0 are hermitian. The full Hamiltonian H is given by

$$H = H_0 + g_2V, \quad V = H_+ + H_-, \quad (2.9)$$

where g_2 is the coupling constant and H_{\pm} are the interaction terms

$$H_{\pm} = Q_0[Q_0, \Sigma_{\pm}]. \quad (2.10)$$

Using the definitions above, it is easy to see that when acting on states of the first type,

$$\begin{aligned} & H_+|n, r_0; r_1, \dots, r_k\rangle \\ &= \int_0^{r_0} dr_{k+1} \sum_{m=-\infty}^{\infty} \sqrt{\frac{r_{k+1}}{r_0(r_0 - r_{k+1})}} \frac{m \sin^2(\pi n \frac{r_0 - r_{k+1}}{r_0})}{\pi^2(m - n \frac{r_0 - r_{k+1}}{r_0})} |m, r_0 - r_{k+1}; r_1, \dots, r_{k+1}\rangle, \\ & H_-|n, r_0; r_1, \dots, r_k\rangle \\ &= \sum_{i=1}^k \sum_{m=-\infty}^{\infty} \sqrt{\frac{r_i}{r_0(r_0 + r_i)}} \frac{m \sin^2(\pi m \frac{r_0}{r_0 + r_i})}{\pi^2(m - n \frac{r_0 + r_i}{r_0})} |m, r_0 + r_i; r_1, \dots, r_i, \dots, r_k\rangle, \end{aligned} \quad (2.11)$$

while on states of the second type

$$\begin{aligned} & H_+|s_1; s_2; r_1, \dots, r_k\rangle = 0, \\ & H_-|s_1; s_2; r_1, \dots, r_k\rangle = - \sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi m \frac{s_1}{s_1 + s_2})}{\pi^2 \sqrt{s_1 + s_2}} |m, s_1 + s_2; r_1, \dots, r_k\rangle. \end{aligned} \quad (2.12)$$

Note that H_+ is not the adjoint of H_- . Therefore V (and hence the Hamiltonian H) is not hermitian with respect to the inner product $\langle | \rangle$ defined in subsection 2.3. At this point all we can say is that

$$V = Q_0[Q_0, \Sigma], \quad V^\dagger = [\Sigma, Q_0]Q_0. \quad (2.13)$$

Let us note that the expression given in (2.11) differs from that of [30] in the following inconsequential ways: (1) their Hamiltonian has an additional factor of $4\pi^2$ (which can be absorbed into λ'), (2) the expressions (2.11) have slightly different factors in the square root, owing to a slightly different definition of the states (see next section), and (3) the arguments of the \sin^2 functions are slightly different, but equivalent (since m and n are integers). Finally, we remark that the authors of [30] had no need for the expressions (2.12) because they focused on diagonalizing H within the subspace of the first kind of state. Since acting with H will never produce states of the second type, it is consistent for their purposes to completely disregard the second component of the Hilbert space.

3. Relation to Gauge Theory

Here we summarize the relation between the definitions in the previous section and the BMN limit of the gauge theory. This section is provided for cultural enrichment; those readers content to study the structure of the quantum mechanics for its own sake may proceed to section 4.

3.1. Hilbert space

Recall that $\mathcal{N} = 4$ $SU(N)$ super Yang-Mills theory has 6 real scalar fields ϕ_i . The BMN operators can be constructed from three orthogonal complex combinations, which can be taken to be

$$\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2), \quad \psi = \frac{1}{\sqrt{2}}(\phi_3 + i\phi_4), \quad Z = \frac{1}{\sqrt{2}}(\phi_5 + i\phi_6). \quad (3.1)$$

The state-operator identification is then

$$\begin{aligned} |n, r_0; r_1, \dots, r_k\rangle &\leftrightarrow \int_0^{r_0} dx \frac{e^{2\pi i n x / r_0}}{\sqrt{r_0 r_1 \dots r_k}} \text{Tr}(\phi Z^{Jx} \psi Z^{J(r_0-x)}) \text{Tr}(Z^{J_1}) \dots \text{Tr}(Z^{J_k}), \\ |s_1; s_2; r_1, \dots, r_k\rangle &\leftrightarrow \frac{1}{\sqrt{r_1 \dots r_k}} \text{Tr}(\phi Z^{J s_1}) \text{Tr}(\psi Z^{J s_2}) \text{Tr}(Z^{J_1}) \dots \text{Tr}(Z^{J_k}), \end{aligned} \quad (3.2)$$

where $r_i = J_i/J$. We will use $O_{|a\rangle}$ to denote the operator corresponding to the state $|a\rangle$.

3.2. Inner product

The gauge theory inner product is related by the state-operator mapping to the two-point function in the free ($g_{\text{YM}} = 0$) theory according to the formula

$$\langle \overline{O}_{|1\rangle}(0) O_{|2\rangle}(x) \rangle \Big|_{\text{free}} = \frac{J^{-1} N^{J+2}}{(4\pi^2 x^2)^{J+2}} \langle 1|2 \rangle_{g_2}. \quad (3.3)$$

This formula may be viewed as the definition of $\langle 1|2 \rangle_{g_2}$.

The factor of J^{-1} in (3.3) can be motivated by checking this relation in the $g_2 \rightarrow 0$ limit. For operators of the first type one gets a factor of J^{k+1} (at large J) from contracting the fields, a factor of J^{-2} from converting the integrals in (3.2) to sums, and a factor of J^{-k} from converting k Kronecker delta-functions to the continuous delta functions in the first line of (2.3). For operators of the second type one gets a factor of J^k from contracting the Z fields and a factor of J^{-k-1} from converting $k+1$ Kronecker delta-functions.

3.3. Splitting-joining operator

The inner product we defined in subsection 2.2 corresponds to the gauge theory inner product (3.3) only at $g_2 = 0$. The splitting-joining operator Σ which we defined above gives the first order g_2 correction to the inner product according to the formula

$$\langle \overline{O}_{|1\rangle}(0) O_{|2\rangle}(x) \rangle \Big|_{\text{free}, \mathcal{O}(g_2)} = g_2 \frac{J^{-1} N^{J+2}}{(4\pi^2 x^2)^{J+2}} \langle 1|\Sigma|2 \rangle. \quad (3.4)$$

Equivalently: matrix elements of Σ may be computed in the gauge theory by calculating free, planar contractions between k -trace and $k+1$ -trace BMN operators.

This operator Σ has appeared in at least three different but equivalent guises in the plane wave literature:

- (1) It encodes free, planar three-point functions of BMN operators [5,12,17].
- (2) It is the permutation operator in the discretized string theory (bit model) [8,18].
- (3) In light-cone string field theory, Σ is the three-string vertex without prefactor [2,9].

3.4. Hamiltonian

The Hamiltonian of the quantum mechanical model corresponds in the gauge theory to $\Delta - J$, where Δ is the dilatation operator. At $g_2 = 0$ it is known that

$$\begin{aligned} (\Delta - J)|n, r_0; r_1, \dots, r_k \rangle &= 2\sqrt{1 + \lambda' n^2 / r_0^2} |n, r_0; r_1, \dots, r_k \rangle, \\ (\Delta - J)|s_1; s_2; r_1, \dots, r_k \rangle &= 0, \end{aligned} \quad (3.5)$$

and the expressions given for H in section 2 encapsulate the one-loop (i.e. $\mathcal{O}(\lambda')$) contribution to $\Delta - J$. The interaction terms $H_+ + H_-$ encode the first order (in g_2) elements of the one-loop anomalous dimension mixing matrix.

4. The String Basis

In this section we investigate some properties of the quantum mechanics. Of paramount importance is the relation

$$SH = H^\dagger S, \quad S = e^{g_2 \Sigma}, \quad (4.1)$$

to be proven below. Let us explain the significance of this result. We saw in section 2 that the Hamiltonian H is not hermitian with respect to the inner product $\langle | \rangle$. However, this is no cause for concern: while H is manifestly hermitian with respect to the gauge theory inner product $\langle | \rangle_{g_2}$ of (3.3), there is no reason for it to be hermitian with respect to $\langle | \rangle$ since the two inner products agree only at $g_2 = 0$!

It is believed that the gauge theory inner product at finite g_2 (defined by (3.3)) is given by the simple formula

$$\langle 1|2 \rangle_{g_2} = \langle 1|S|2 \rangle. \quad (4.2)$$

The relation (4.1) guarantees that H is hermitian with respect to (4.2). This provides a consistency check on the conjecture that (4.2) is actually the correct gauge theory inner product at finite g_2 .

4.1. Some identities

From the definition (2.13) we immediately obtain the relations

$$[H_0, \Sigma] = V - V^\dagger, \quad [Q_0, [Q_0, \Sigma]] = V + V^\dagger. \quad (4.3)$$

Next we present

Lemma 1.

$$[\Sigma, [\Sigma, Q_0]] = 0. \quad (4.4)$$

The lengthy proof of this result is given in appendix A. It would be interesting to understand the gauge theory origin of this lemma, which might be possible to derive as a consequence of supersymmetry.

Some immediate consequences of Lemma 1 which follow directly from the definition (2.13) include

$$[\Sigma, V + V^\dagger] = 0, \quad [\Sigma, [\Sigma, V]] = 0. \quad (4.5)$$

Next consider the formula

$$e^A B e^{-A} = B + [A, B] + \frac{1}{2!}[A, [A, B]] + \frac{1}{3!}[A, [A, [A, B]]] + \dots \quad (4.6)$$

which leads to

$$\begin{aligned} e^{\lambda g_2 \Sigma} H_0 e^{-\lambda g_2 \Sigma} &= H_0 + g_2 \lambda (V^\dagger - V) + \frac{g_2^2}{2} \lambda^2 [\Sigma, V^\dagger - V], \\ e^{\lambda g_2 \Sigma} g_2 V e^{-\lambda g_2 \Sigma} &= g_2 V + \frac{g_2^2}{2} \lambda [\Sigma, V - V^\dagger], \end{aligned} \quad (4.7)$$

for an arbitrary parameter λ . Remarkably, all higher order terms vanish as a consequence of (4.5). Combining (4.7) gives

$$e^{\lambda g_2 \Sigma} H e^{-\lambda g_2 \Sigma} = H - g_2 \lambda (V - V^\dagger) + \frac{g_2^2}{2} (\lambda - \lambda^2) [\Sigma, V - V^\dagger]. \quad (4.8)$$

For $\lambda = 1$ we find

$$S H S^{-1} = H^\dagger, \quad (4.9)$$

thereby establishing the desired relation (4.1).

4.2. Hamiltonian in the string basis

The basis which diagonalizes the inner product (4.2) is

$$|\tilde{a}\rangle = S^{-1/2} |a\rangle. \quad (4.10)$$

This basis has been identified as the basis of string states in the light-cone string field theory: if $|k\rangle$ corresponds to an operator with precisely k traces, then $S^{-1/2} |k\rangle$ corresponds to a state of precisely k strings.

It is convenient to define an operator \tilde{H} whose matrix elements in the gauge theory basis are the same as the matrix elements of the Hamiltonian H in the string basis:

$$\langle \tilde{a} | H | \tilde{b} \rangle_{g_2} = \langle a | \tilde{H} | b \rangle. \quad (4.11)$$

We will call \tilde{H} the ‘string Hamiltonian’, since its matrix elements are related to matrix elements of the light-cone string field theory Hamiltonian. Clearly, \tilde{H} is given by

$$\tilde{H} = S^{1/2} H S^{-1/2}. \quad (4.12)$$

Setting $\lambda = \frac{1}{2}$ in (4.8) leads to

$$\tilde{H} = H_0 + \frac{g_2}{2} (V + V^\dagger) + \frac{g_2^2}{8} [\Sigma, V - V^\dagger]. \quad (4.13)$$

Naturally, \tilde{H} is manifestly hermitian with respect to the inner product $\langle | \rangle$. It is remarkable that whereas the Hamiltonian in the gauge theory basis has no contact terms, the basis transformation introduces a single $\mathcal{O}(g_2^2)$ contact term.

4.3. The ‘supercharge’

Remarkably, it is easy to see that the string Hamiltonian \tilde{H} is a perfect square! In particular,

$$\tilde{H} = Q^\dagger Q, \quad Q = Q_0 + \frac{g_2}{2}[Q_0, \Sigma] = S^{-1/2}Q_0S^{1/2}. \quad (4.14)$$

The order g_2 term in (4.13) works out due to (4.3), while the g_2^2 term follows from (2.13) and

$$[\Sigma, Q_0[Q_0, \Sigma]] - [\Sigma, [\Sigma, Q_0]Q_0] - 2[\Sigma, Q_0][Q_0, \Sigma] = 0, \quad (4.15)$$

which is a consequence of (4.4). Note that it follows from Lemma 1 that

$$[\Sigma, [\Sigma, Q]] = 0. \quad (4.16)$$

Of course, the $\mathcal{N} = 4$ supersymmetry algebra requires that $\Delta - J$ is positive definite. However, it is nice to see that this can be analytically proven from the expressions (2.11), where this fact is not at all obvious.

One interesting open problem is to supersymmetrize this quantum mechanics by including fermionic impurities, in which case the appropriate generalization of Q would become an honest fermionic supercharge. It would be very interesting to see whether the fermionic extension of this quantum mechanics proceeds as in the string bit model [18], or whether the fermionic completion has a different flavor here.

4.4. The string field theory ‘prefactor’

Let us consider order g_2 matrix elements of \tilde{H} between two energy eigenstates — actually, two eigenstates of Q_0 with eigenvalues $\sqrt{E_1}, \sqrt{E_2}$. From (4.13) and (4.3) we immediately have

$$\langle 1|\tilde{H}|2\rangle = \frac{g_2}{2}\langle 1|[Q_0, [Q_0, \Sigma]]|2\rangle = \frac{g_2}{2}(\sqrt{E_1} - \sqrt{E_2})^2\langle 1|\Sigma|2\rangle. \quad (4.17)$$

This formula equates matrix elements of the string field theory Hamiltonian \tilde{H} to three-point functions $\langle 1|\Sigma|2\rangle$ in the gauge theory, with a certain ‘prefactor’. In the earliest literature on the subject, the prefactor was erroneously believed to be $E_1 - E_2$ instead of $\frac{1}{2}(\sqrt{E_1} - \sqrt{E_2})^2$. It would be very interesting to understand the generalization of this formula for more complicated processes, in particular those involving more than two impurities.

5. Discussion and Speculations

5.1. A puzzle at two loops

A study of the gauge theory at two-loops has recently been presented in [33]. The results led the authors to the conjecture that the Hamiltonian is given to all loops by

$$H_{\text{full}} = 2\sqrt{1 + \lambda'H} \quad (5.1)$$

(with firm calculational support only up to and including $\mathcal{O}(\lambda'^2)$), where H on the right-hand side is the one-loop Hamiltonian studied in the previous sections. The Hamiltonian in the string basis would then be

$$\tilde{H}_{\text{full}} = S^{1/2}H_{\text{full}}S^{-1/2} = 2\sqrt{1 + \lambda'\tilde{H}}. \quad (5.2)$$

It is not hard to derive from this a formula for the order g_2 contribution to the following matrix element, to all orders in λ' :

$$\begin{aligned} \frac{\langle n, 1 | \tilde{H}_{\text{full}} | m, y; 1 - y \rangle}{\langle n, 1 | \tilde{H} | m, y; 1 - y \rangle} &= \frac{\sqrt{1 + \lambda'n^2} - \sqrt{1 + \lambda'm^2/y^2}}{\frac{1}{2}\lambda'(n^2 - m^2/y^2)} + \mathcal{O}(g_2) \\ &= \left[1 - \frac{1}{4}\lambda'(n^2 + m^2/y^2) + \mathcal{O}(\lambda'^2) \right] + \mathcal{O}(g_2). \end{aligned} \quad (5.3)$$

On the other hand, this particular matrix element has also been studied extensively on the string field theory side of the BMN correspondence. In [25] it was shown that to all orders in λ' perturbation theory,

$$\begin{aligned} \frac{\langle n, 1 | P_{\text{all orders in } \lambda'}^- | m, y; 1 - y \rangle}{\langle n, 1 | P_{\text{first order in } \lambda'}^- | m, y; 1 - y \rangle} &= \frac{1}{\sqrt{1 + \lambda'n^2}\sqrt{1 + \lambda'm^2/y^2}} \\ &= 1 - \frac{1}{2}\lambda'(n^2 + m^2/y^2) + \mathcal{O}(\lambda'^2). \end{aligned} \quad (5.4)$$

The results (5.3) and (5.4) disagree by a factor of 2 at two loops (where the calculations of [33] are firm), and they disagree even more strongly at higher loops (where [33] merely conjectured). In particular, it is impossible to write down any function just of \tilde{H} that reproduces the result (5.4) to all orders in λ' . It is possible that this discrepancy involves an unallowed exchange of the order of limits between gauge theory and string field theory. This problem has manifested itself in [10], for example (where a “renormalization” by a finite amount occurs below (2.14)), and in [27], regarding the issue of intermediate states which do not conserve the number of impurities. It is an interesting open problem to understand precisely which observables we might expect to be able to study perturbatively on both sides of the duality.

5.2. The S-matrix

What is it that we would most like to know about the quantum mechanics studied in this paper; i.e. what is the ultimate goal of this course of research? We propose that the goal should be to calculate the non-relativistic S-matrix obtained from this quantum mechanics, which we identify with the S-matrix of string theory [24] in the plane wave background, after the appropriate basis transformation.²

Much of the literature on this subject has focused (quite successfully) on comparing matrix elements of \tilde{H} to matrix elements of the Hamiltonian P^- of light-cone string field theory in the plane wave background. The S-matrix proposal subsumes all of the supporting evidence (since it is a weaker proposal) and simultaneously satisfies the ardent skeptic who points out that only the S-matrix is a good observable in string theory (as matrix elements of the light-cone Hamiltonian are not coordinate invariant).

The authors of [30] set out to find the spectrum of H (which is identical to the spectrum of \tilde{H}). This would of course be very useful to know, but this might be a very difficult task in practice. The BMN correspondence suggests that H should be related somehow to the Hamiltonian of an interacting string theory, which likely has an exceedingly complicated spectrum! Indeed the authors of [30] encountered technical difficulties at genus two due to overlapping continuum states. However, one rarely studies string theory (or any quantum field theory) by attempting to diagonalize the Hamiltonian. Instead, the goal is usually to calculate the S-matrix.

Let us now recall some non-relativistic scattering theory. We write the string Hamiltonian as

$$\tilde{H} = H_0 + W, \quad W = \frac{g_2}{2}(V + V^\dagger) + \frac{g_2^2}{8}[\Sigma, V - V^\dagger]. \quad (5.5)$$

Then the transition matrix $T(z)$ can be obtained from the Born series

$$T(z) = W + WG_0(z)W + WG_0(z)WG_0(z)W + \dots, \quad (5.6)$$

² It is important to note that in general an S-matrix depends on how one chooses to divide the full Hamiltonian into a ‘free’ part and an ‘interacting’ part. In our case, the basis transformation $S^{1/2}$ does not commute with this division: the free and interacting parts of \tilde{H} are not respectively the transforms of the free and interacting parts of H . Therefore the S-matrices obtained from H and \tilde{H} are not unitarily related to each other (although it would be intriguing to see if there were some other, more complicated relationship between them).

where $G_0(z) = (z - H_0)^{-1}$ is the free propagator. As a function of the complex variable z , the operator $T(z)$ should have poles at the bound states of H and branch cuts along the continuous spectrum of H . The S-matrix is then given by

$$\langle 1|S|2\rangle = \langle 1|1 - 2\pi i\delta(E_1 - E_2)T(E_1 + i\epsilon)|2\rangle, \quad (5.7)$$

when $H_0|i\rangle = E_i|i\rangle$.

It is now possible (though technically complicated) to calculate the S-matrix to any desired order in g_2 . The divergences encountered in [30] at genus two would also occur if one tried to diagonalize the Hamiltonian of string theory in the plane wave background. According to our proposal, they should be regulated with a $+i\epsilon$ prescription (instead of a principal value) and interpreted as the usual branch cuts one finds in transition amplitudes when there is a continuum of intermediate scattering states.

Acknowledgements

It is a pleasure to thank D. Freedman, U. Gursoy, N. Itzhaki, R. Roiban, D. Vaman and H. Verlinde for helpful discussions. The work of M.S. was supported in part by DOE grant DE-FG02-91ER40671, and that of A.V. was supported in part by the National Science Foundation under Grant No. PHY99-07949.

Appendix A. Proof of Lemma 1

By separating $\Sigma = \Sigma_+ + \Sigma_-$ and defining $P_{\pm} = [Q_0, \Sigma_{\pm}]$, we can express (4.4) as three relations which must separately be satisfied:

$$\begin{aligned} (1) \quad & [\Sigma_-, P_-] = 0, \\ (2) \quad & [\Sigma_-, P_+] + [\Sigma_+, P_-] = 0, \\ (3) \quad & [\Sigma_+, P_+] = 0. \end{aligned} \quad (A.1)$$

Of course, (3) is equivalent to the adjoint of (1) so it will suffice to check only the relations (1) and (2). From the definitions in section 2 we have

$$\begin{aligned} P_+|n, r_0; r_1, \dots, r_k\rangle &= \sqrt{r_0} \int_0^{r_0} ds \frac{\sin^2(\pi n \frac{s}{r_0})}{\pi^2 n} |r_0 - s; s; r_1, \dots, r_k\rangle \\ &+ \int_0^{r_0} dr_{k+1} \sum_{m=-\infty}^{\infty} \sqrt{\frac{(r_0 - r_{k+1})r_{k+1}}{r_0}} \frac{\sin^2(\pi n \frac{r_0 - r_{k+1}}{r_0})}{\pi^2(m - n \frac{r_0 - r_{k+1}}{r_0})} |m, r_0 - r_{k+1}; r_1, \dots, r_{k+1}\rangle, \\ P_-|n, r_0; r_1, \dots, r_k\rangle &= \sum_{i=1}^k \sum_{m=-\infty}^{\infty} \sqrt{\frac{(r_0 + r_i)r_i}{r_0}} \frac{\sin^2(\pi m \frac{r_0}{r_0 + r_i})}{\pi^2(m - n \frac{r_0}{r_0 + r_i})} |m, r_0 + r_i; r_1, \dots, r_i, \dots, r_k\rangle \end{aligned} \quad (A.2)$$

on states of type one and

$$\begin{aligned}
P_+ |s_1; s_2; r_1, \dots, r_k\rangle &= 0, \\
P_- |s_1; s_2; r_1, \dots, r_k\rangle &= -\sqrt{s_1 + s_2} \sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi m \frac{s_1}{s_1+s_2})}{\pi^2 m} |m, s_1 + s_2; r_1, \dots, r_k\rangle
\end{aligned} \tag{A.3}$$

on states of the second type. We now consider separately the relations (A.1) on the two kinds of states.

A.1. Relation (1) on states of the first type

Acting with $\Sigma_- P_-$ on a state of type one gives

$$\begin{aligned}
\Sigma_- P_- |n, r_0; r_1, \dots, r_k\rangle &= \sum_{i \neq j} \sum_{m=-\infty}^{\infty} A_{ijm} |m, r_0 + r_i + r_j; r_1, \dots, r_i, \dots, r_j, \dots, r_k\rangle \\
&+ \sum_{i=1}^k \sum_{m=-\infty}^{\infty} \sqrt{\frac{(r_0 + r_i)r_i}{r_0}} \frac{\sin^2(\pi m \frac{r_0}{r_0+r_i})}{\pi^2(m - n \frac{r_0+r_i}{r_0})} \\
&\times \sum_{j \neq i} \sum_{l \neq j, i} \frac{1}{2} \sqrt{r_j r_l (r_j + r_l)} |m, r_0 + r_i; r_1, \dots, r_i, \dots, r_j, \dots, r_l, \dots, r_k, r_j + r_l\rangle,
\end{aligned} \tag{A.4}$$

where the coefficient A is given by

$$A_{ijm} = \sum_{p=-\infty}^{\infty} \sqrt{\frac{r_i(r_0 + r_i + r_j)^3 r_j}{r_0}} \frac{\sin^2(\pi p \frac{r_0}{r_0+r_i})}{\pi^2(p - n \frac{r_0+r_i}{r_0})} \frac{\sin^2(\pi m \frac{r_0+r_i}{r_0+r_i+r_j})}{\pi^2(m - p \frac{r_0+r_i+r_j}{r_0+r_i})^2}. \tag{A.5}$$

Similarly, acting with $P_- \Sigma_-$ gives

$$\begin{aligned}
P_- \Sigma_- |n, r_0; r_1, \dots, r_k\rangle &= \sum_{i \neq j} \sum_{m=-\infty}^{\infty} B_{ijm} |m, r_0 + r_i + r_j; r_1, \dots, r_i, \dots, r_j, \dots, r_k\rangle \\
&+ \frac{1}{2} \sum_{j \neq l} \sqrt{r_j r_l (r_j + r_l)} P_- |n, r_0; r_1, \dots, r_j, \dots, r_l, \dots, r_k, r_j + r_l\rangle,
\end{aligned} \tag{A.6}$$

with coefficient

$$B_{ijm} = \sum_{p=-\infty}^{\infty} \sqrt{\frac{(r_0 + r_i)^2 r_i (r_0 + r_i + r_j) r_j}{r_0}} \frac{\sin^2(\pi p \frac{r_0}{r_0+r_i})}{\pi^2(p - n \frac{r_0+r_i}{r_0})^2} \frac{\sin^2(\pi m \frac{r_0+r_i}{r_0+r_i+r_j})}{\pi^2(m - p \frac{r_0+r_i+r_j}{r_0+r_i})}. \tag{A.7}$$

Let us first study the A_{ijm} and B_{ijm} terms. If we define

$$y = \frac{r_0}{r_0 + r_i}, \quad a = n/y, \quad b = m \frac{r_0 + r_i}{r_0 + r_i + r_j} \tag{A.8}$$

then we can combine A and B into the formula

$$A_{ijm} - B_{ijm} = \frac{1}{\pi^4} \sqrt{\frac{r_i r_j}{r_0(r_0 + r_i + r_j)}} \times \left[(r_0 + r_i)^2 \sin^2\left(\pi m \frac{r_0 + r_i}{r_0 + r_i + r_j}\right) \sum_{p=-\infty}^{\infty} \frac{\sin^2(\pi p y)}{(p-a)(p-b)} \left(\frac{1}{p-a} + \frac{1}{p-b}\right) \right]. \quad (\text{A.9})$$

The sum over p can be performed with the help of (B.4), and we find after much simplification

$$A_{(ij)m} - B_{(ij)m} = \frac{r_i + r_j}{2\pi^2} \sqrt{\frac{r_i r_j (r_0 + r_i + r_j)}{r_0}} \frac{\sin^2\left(\pi m \frac{r_0}{r_0 + r_i + r_j}\right)}{m - n \frac{r_0 + r_i + r_j}{r_0}}. \quad (\text{A.10})$$

Notice that we have symmetrized in the i and j indices. This step is allowed (indeed, forced upon us) because of the manifest $i \leftrightarrow j$ symmetry of the states these coefficients multiply.

Now consider the second term in (A.4) and the second term in (A.6). It is easy to see that they are essentially the same. However, the latter has an additional term when P_- acts on the last trace $(r_j + r_l)$, giving the term

$$\begin{aligned} \dots + \frac{1}{2} \sum_{j \neq l} \sqrt{r_j r_l (r_j + r_l)} \sum_{m=-\infty}^{\infty} \sqrt{\frac{(r_0 + r_j + r_l)(r_j + r_l)}{r_0}} \\ \times \frac{\sin^2\left(\pi m \frac{r_0}{r_0 + r_j + r_l}\right)}{\pi^2 \left(m - n \frac{r_0 + r_j + r_l}{r_0}\right)} |m, r_0 + r_j + r_l; r_1, \dots, \mathfrak{r}_j, \dots, \mathfrak{r}_l, \dots, r_k\rangle. \end{aligned} \quad (\text{A.11})$$

This additional term is precisely the same as the one whose coefficient is given by (A.10)! This completes the proof that

$$[\Sigma_-, P_-] |n, r_0; r_1, \dots, r_k\rangle = 0. \quad (\text{A.12})$$

A.2. Relation (1) on states of the second type

Acting with $\Sigma_- P_-$ on a state of type two gives something of the form

$$\begin{aligned} \Sigma_- P_- |s_1; s_2; r_1, \dots, r_k\rangle &= \sum_{i=1}^k \sum_{m=-\infty}^{\infty} C_{im} |m, s_1 + s_2 + r_i; r_1, \dots, \mathfrak{r}_i, \dots, r_k\rangle \\ &- \sqrt{s_1 + s_2} \sum_{m=-\infty}^{\infty} \frac{\sin^2\left(\pi m \frac{s_1}{s_1 + s_2}\right)}{\pi^2 m} \\ &\times \frac{1}{2} \sum_{i \neq j} \sqrt{r_i r_j (r_i + r_j)} |m, s_1 + s_2; r_1, \dots, \mathfrak{r}_i, \dots, \mathfrak{r}_j, \dots, r_k, r_i + r_j\rangle. \end{aligned} \quad (\text{A.13})$$

with

$$C_{im} = -\sqrt{(s_1 + s_2 + r_i)^3 r_i} \sum_{p=-\infty}^{\infty} \frac{\sin^2(\pi p \frac{s_1}{s_1+s_2})}{\pi^2 p} \frac{\sin^2(\pi m \frac{s_1+s_2}{s_1+s_2+r_i})}{\pi^2 (m - p \frac{s_1+s_2+r_i}{s_1+s_2})^2}. \quad (\text{A.14})$$

Similarly, from $P_- \Sigma_-$ we get an expression similar to (A.13). The second term is identical, but the first term has the coefficient

$$D_{im} = \sqrt{(s_1 + s_2 + r_i) r_i} \left[-s_1 \frac{\sin^2(\pi m \frac{s_2}{s_1+s_2+r_i})}{\pi^2 m} - s_2 \frac{\sin^2(\pi m \frac{s_1}{s_1+s_2+r_i})}{\pi^2 m} + (s_1 + s_2) \sum_{p=-\infty}^{\infty} \frac{(-1)^p \sin(\pi p \frac{s_1}{s_1+s_2}) \sin(\pi p \frac{s_2}{s_1+s_2})}{\pi^2 p^2} \frac{\sin^2(\pi m \frac{s_1+s_2}{s_1+s_2+r_i})}{\pi^2 (m - p \frac{s_1+s_2+r_i}{s_1+s_2})} \right]. \quad (\text{A.15})$$

Applying the formulas in appendix B leads eventually to

$$C_{im} = D_{im}, \quad (\text{A.16})$$

thereby establishing that

$$[\Sigma_-, P_-] |s_1; s_2; r_1, \dots, r_k\rangle = 0. \quad (\text{A.17})$$

A.3. Relation (2) on states of the second type

Since P_+ annihilates states of the second type, we have simply

$$\begin{aligned} [\Sigma_-, P_+] |s_1; s_2; r_1, \dots, r_k\rangle &= \int_0^{s_1+s_2} dr_{k+1} \sum_{m=-\infty}^{\infty} E_m |m; s_1 + s_2 - r_{k+1}; r_1, \dots, r_{k+1}\rangle \\ &+ (s_1 + s_2)^2 \sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi m \frac{s_1}{s_1+s_2})}{\pi^2 m^2} \int_0^{s_1+s_2} ds \frac{\sin^2(\pi m \frac{s}{s_1+s_2})}{\pi^2 m} |s_1 + s_2 - s; s; r_1, \dots, r_k\rangle. \end{aligned} \quad (\text{A.18})$$

The second term vanishes because the summand is odd in m , and in the first term appears the coefficient

$$E_m = (s_1 + s_2) \sqrt{(s_1 + s_2 - r_{k+1}) r_{k+1}} \sum_{p=-\infty}^{\infty} \frac{\sin^2(\pi p \frac{s_1}{s_1+s_2})}{\pi^2 p^2} \frac{\sin^2(\pi p \frac{s_1+s_2-r_{k+1}}{s_1+s_2})}{\pi^2 (m - p \frac{s_1+s_2-r_{k+1}}{s_1+s_2})}. \quad (\text{A.19})$$

Next we consider $[\Sigma_+, P_-] |s_1; s_2; r_1, \dots, r_k\rangle$. Again there appears a term involving states of the second kind which vanishes due to an odd summand. The remaining terms of the

first type take the same form as in (A.18), but with a much more complicated coefficient

$$\begin{aligned}
F_m = & -\sqrt{(s_1 + s_2 - r_{k+1})^3 r_{k+1}} \sum_{p=-\infty}^{\infty} \frac{\sin^2(\pi p \frac{s_1}{s_1+s_2})}{\pi^2 p} \frac{\sin^2(\pi p \frac{s_1+s_2-r_{k+1}}{s_1+s_2})}{\pi^2 (m - p \frac{s_1+s_2-r_{k+1}}{s_1+s_2})^2} \\
& + (s_1 - r_{k+1}) \sqrt{(s_1 + s_2 - r_{k+1}) r_{k+1}} \frac{\sin^2(\pi m \frac{s_1-r_{k+1}}{s_1+s_2-r_{k+1}})}{\pi^2 m} \theta(s_1 - r_{k+1}) \\
& + (s_2 - r_{k+1}) \sqrt{(s_1 + s_2 - r_{k+1}) r_{k+1}} \frac{\sin^2(\pi m \frac{s_2-r_{k+1}}{s_1+s_2-r_{k+1}})}{\pi^2 m} \theta(s_2 - r_{k+1}).
\end{aligned} \tag{A.20}$$

We are attempting to show that

$$([\Sigma_-, P_+] + [\Sigma_+, P_-]) |s_1; s_2; r_1, \dots, r_k\rangle = 0. \tag{A.21}$$

This requires $E_m + F_m = 0$, which can be written as

$$\begin{aligned}
\sum_{p=-\infty}^{\infty} \frac{\sin^2(\pi p \frac{s_1}{s_1+s_2}) \sin^2(\pi p \frac{s_1+s_2-r_{k+1}}{s_1+s_2})}{p(p-a)} \left[\frac{1}{p} + \frac{1}{p-a} \right] &= \frac{\pi^2 (s_1 + s_2 - r_{k+1})}{m (s_1 + s_2)^2} \\
&\times \left[(s_1 - r_{k+1}) \sin^2(\pi m \frac{s_1-r_{k+1}}{s_1+s_2-r_{k+1}}) \theta(s_1 - r_{k+1}) + (s_1 \leftrightarrow s_2) \right],
\end{aligned} \tag{A.22}$$

where $a = m \frac{s_1+s_2}{s_1+s_2-r_{k+1}}$. This highly nontrivial identity is a consequence of (B.6), but requires some explanation since the constraint $1 \geq y_1 + y_2 \geq y_1 \geq y_2 \geq 0$ in (B.6) is crucial. The integral in (A.18) splits into three regions. In each region a different choice of y_1 and y_2 is necessary in order to satisfy the constraint. Taking $s_2 \geq s_1$ without loss of generality, the appropriate choices are

$$\begin{aligned}
r_{k+1} \in [0, s_1] : & \quad y_1 = \frac{s_1}{s_1 + s_2}, \quad y_2 = \frac{r_{k+1}}{s_1 + s_2}, \\
r_{k+1} \in [s_1, s_2] : & \quad y_1 = \frac{s_1 + s_2 - r_{k+1}}{s_1 + s_2}, \quad y_2 = \frac{s_1}{s_1 + s_2}, \\
r_{k+1} \in [s_2, s_1 + s_2] : & \quad y_1 = \frac{s_1}{s_1 + s_2}, \quad y_2 = \frac{s_1 + s_2 - r_{k+1}}{s_1 + s_2}.
\end{aligned} \tag{A.23}$$

After much simplification, we find in each case that (B.6) reproduces the right-hand side of (A.22).

A.4. Relation (2) on states of the first type

When we act with $[\Sigma_-, P_+] + [\Sigma_+, P_-]$ on a state of the first type we get many terms. Let us first collect four terms T_1, \dots, T_4 which are of the second type. From $\Sigma_- P_+$ we get

$$\begin{aligned}
T_1 = \frac{\sqrt{r_0}}{\pi^2 n} \int_0^{r_0} ds \sin^2(\pi n \frac{s}{r_0}) \sum_{i=1}^k \left[(r_0 - s) \sqrt{r_i} |r_0 - s + r_i; s; r_1, \dots, \mathfrak{r}_i, \dots, r_k\rangle \right. \\
\left. + s \sqrt{r_i} |r_0 - s; s + r_i; r_1, \dots, \mathfrak{r}_i, \dots, r_k\rangle \right]
\end{aligned} \tag{A.24}$$

and

$$T_2 = \sqrt{r_0} \int_0^{r_0} ds \frac{\sin^2(\pi n \frac{s}{r_0})}{\pi^2 n} \times \frac{1}{2} \sum_{i \neq j} \sqrt{r_i r_j (r_i + r_j)} |r_0 - s; s; r_1, \dots, \mathfrak{r}_i, \dots, \mathfrak{r}_j, \dots, r_k, r_i + r_j\rangle. \quad (\text{A.25})$$

Next from $(\Sigma_+ P_- - P_+ \Sigma_-)$ we get two terms which combine nicely into

$$T_3 = -\frac{1}{\pi^4} \sum_{i=1}^k \sqrt{\frac{(r_0 + r_i)^4 r_i}{r_0}} \sum_{m=-\infty}^{\infty} \int_0^{r_0+r_i} ds \frac{\sin^2(\pi m \frac{r_0}{r_0+r_i}) \sin^2(\pi m \frac{s}{r_0+r_i})}{m(m - n \frac{r_0+r_i}{r_0})} \times \left[\frac{1}{m} + \frac{1}{m - n \frac{r_0+r_i}{r_0}} \right] |r_0 + r_i - s; s; r_1, \dots, \mathfrak{r}_i, \dots, r_k\rangle, \quad (\text{A.26})$$

and from $-P_+ \Sigma_-$ we get an additional term which is exactly

$$T_4 = -T_2. \quad (\text{A.27})$$

We can simplify T_3 by using the sum (B.4) with $a = 0$, $b = n \frac{r_0+r_i}{r_0}$, but we must consider the cases $s < r_0$ and $s > r_0$ separately, using $y_1 = \frac{r_0}{r_0+r_i}$, $y_2 = \frac{s}{r_0+r_i}$ in the former and $y_1 = 1 - \frac{r_0}{r_0+r_i}$, $y_2 = 1 - \frac{s}{r_0+r_i}$ in the latter. After considerable simplification, we that

$$T_3 = -\frac{\sqrt{r_0}}{\pi^2 n} \sum_{i=1}^k \sqrt{r_i} \int_0^{r_0+r_i} ds \left[(r_0 - s) \sin^2(\pi n \frac{s}{r_0}) \theta(r_0 - s) + (s - r_i) \sin^2(\pi n \frac{s-r_i}{r_0}) \theta(s - r_0) \right] |r_0 + r_i - s; s; r_1, \dots, \mathfrak{r}_i, \dots, r_k\rangle. \quad (\text{A.28})$$

After shifting the variable of integration in the second term it becomes manifest that $T_3 = -T_1$. Adding up all of these terms therefore gives $T_1 + T_2 + T_3 + T_4 = 0!$

It remains to collect the large number of terms of the first type which arise. After several pages of algebra, these can be shown to all cancel using all of the various ingredients used in the previous stages of the proof. We do not show all of the steps since the formulas are exceedingly long and no new tricks are required.

Appendix B. Sums

Throughout this appendix, a, b, c denote non-integer (though otherwise arbitrary complex) numbers, and $y \in [0, 1]$. We start with the basic sum

$$\sum_{m=-\infty}^{\infty} \frac{e^{2\pi i m y}}{(m-a)(m-b)(m-c)} = f(a, b, c) + f(b, c, a) + f(c, a, b), \quad (\text{B.1})$$

$$f(a, b, c) = -\frac{\pi \csc(\pi a)}{(a-b)(a-c)} e^{\pi i a(2y-1)}.$$

which is easily evaluated by standard contour techniques (see for example [10]). It follows that

$$\sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi my)}{(m-a)(m-b)(m-c)} = g(a, b, c) + g(b, c, a) + g(c, a, b), \quad (\text{B.2})$$

$$g(a, b, c) = \frac{\pi \csc(\pi a) \sin(\pi ay) \sin(\pi a(1-y))}{(a-b)(a-c)}.$$

By taking appropriate limits of this result one can obtain a formula for the sum

$$F(y; a, b) \equiv \sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi my)}{(m-a)(m-b)} \left[\frac{1}{m-a} + \frac{1}{m-b} \right], \quad (\text{B.3})$$

which appears throughout appendix A. The expression for $F(y; a, b)$ is rather lengthy so we will not write it down, but a particularly useful special case is

$$F(y; a = n/y, b) = \frac{\pi^2}{(a-b)} \frac{\sin(\pi by)}{\sin(\pi b)} \left[2y \cos(\pi b(1-y)) - \frac{\sin(\pi by)}{\sin(\pi b)} \right], \quad (\text{B.4})$$

valid when n is an integer.

Next, for $1 \geq y_1 + y_2 \geq y_1 \geq y_2 \geq 0$ we obtain from (B.1) the more complicated result

$$\sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi my_1) \sin^2(\pi my_2)}{(m-a)(m-b)(m-c)} = h(a, b, c) + h(b, c, a) + h(c, a, b),$$

$$h(a, b, c) = \frac{\pi}{4(a-b)(a-c)} \frac{\sin(\pi ay_2)}{\sin(\pi a)} \left[2 \sin(\pi a(1-y_2)) \right. \quad (\text{B.5})$$

$$\left. + \sin(\pi a(1-2y_1+y_2)) + \sin(\pi a(2y_1+y_2-1)) \right].$$

Finally, taking the appropriate limits of the preceding equation gives the remarkable identity

$$\sum_{m=-\infty}^{\infty} \frac{\sin^2(\pi my_1) \sin^2(\pi my_2)}{m(m-a)} \left[\frac{1}{m} + \frac{1}{m-a} \right] = \frac{\pi^2 \sin^2(\pi ay_1) \sin^2(\pi ay_2)}{a \sin^2(\pi a)}$$

$$+ \frac{\pi^2 \sin(\pi ay_2)}{2a \sin(\pi a)} \left[(y_1 + y_2) \cos(\pi a(1-2y_1-y_2)) \right. \quad (\text{B.6})$$

$$\left. - (y_1 - y_2) \cos(\pi a(1-2y_1+y_2)) - 2y_2 \cos(\pi a(y_2-1)) \right].$$

Appendix C. More on Σ

Here we present the joining operator, which is the adjoint of (2.6):

$$\begin{aligned}
& \Sigma_- |n, r_0; r_1, \dots, r_k\rangle \\
&= \sum_{i=1}^k \sum_{m=-\infty}^{\infty} \sqrt{\frac{(r_0 + r_i)^3 r_i}{r_0}} \frac{\sin^2(\pi m \frac{r_0}{r_0 + r_i})}{\pi^2 (m - n \frac{r_0 + r_i}{r_0})^2} |m, r_0 + r_i; r_1, \dots, r_i, \dots, r_k\rangle \\
&+ \frac{1}{2} \sum_{i \neq j} \sqrt{r_i r_j (r_i + r_j)} |n, r_0; r_1, \dots, r_i, \dots, r_j, \dots, r_k, r_i + r_j\rangle, \\
& \Sigma_- |s_1; s_2; r_1, \dots, r_k\rangle = \tag{C.1} \\
& \sum_{m=-\infty}^{\infty} \frac{(-1)^m (s_1 + s_2)^{3/2}}{\pi^2 m^2} \sin(\pi m \frac{s_1}{s_1 + s_2}) \sin(\pi m \frac{s_2}{s_1 + s_2}) |m, s_1 + s_2; r_1, \dots, r_k\rangle \\
&+ \sum_{i=1}^k \sqrt{r_i} \left[s_1 |s_1 + r_i; s_2; r_1, \dots, r_i, \dots, r_k\rangle + s_2 |s_1; s_2 + r_i; r_1, \dots, r_i, \dots, r_k\rangle \right] \\
&+ \frac{1}{2} \sum_{i \neq j} \sqrt{r_i r_j (r_i + r_j)} |n, r_0; r_1, \dots, r_i, \dots, r_j, \dots, r_k, r_i + r_j\rangle.
\end{aligned}$$

References

- [1] D. Berenstein, J. M. Maldacena and H. Nastase, “Strings in flat space and pp waves from $N = 4$ super Yang Mills,” *JHEP* **0204**, 013 (2002) [arXiv:hep-th/0202021].
- [2] M. Spradlin and A. Volovich, “Superstring interactions in a pp-wave background,” *Phys. Rev. D* **66**, 086004 (2002) [arXiv:hep-th/0204146].
- [3] C. Kristjansen, J. Plefka, G. W. Semenoff and M. Staudacher, “A new double-scaling limit of $N = 4$ super Yang-Mills theory and PP-wave strings,” *Nucl. Phys. B* **643**, 3 (2002) [arXiv:hep-th/0205033].
- [4] D. J. Gross, A. Mikhailov and R. Roiban, “Operators with large R charge in $N = 4$ Yang-Mills theory,” *Annals Phys.* **301**, 31 (2002) [arXiv:hep-th/0205066].
- [5] N. R. Constable, D. Z. Freedman, M. Headrick, S. Minwalla, L. Motl, A. Postnikov and W. Skiba, “PP-wave string interactions from perturbative Yang-Mills theory,” *JHEP* **0207**, 017 (2002) [arXiv:hep-th/0205089].
- [6] Y. Kiem, Y. Kim, S. Lee and J. Park, “pp-wave / Yang-Mills correspondence: An explicit check,” *Nucl. Phys. B* **642**, 389 (2002) [arXiv:hep-th/0205279].
- [7] C. S. Chu, V. V. Khoze and G. Travaglini, “Three-point functions in $N = 4$ Yang-Mills theory and pp-waves,” *JHEP* **0206**, 011 (2002) [arXiv:hep-th/0206005].
- [8] H. Verlinde, “Bits, matrices and $1/N$,” arXiv:hep-th/0206059.
- [9] M. Spradlin and A. Volovich, “Superstring interactions in a pp-wave background II,” *JHEP* **0301**, 036 (2003) [arXiv:hep-th/0206073].
- [10] I. R. Klebanov, M. Spradlin and A. Volovich, “New effects in gauge theory from pp-wave superstrings,” *Phys. Lett. B* **548**, 111 (2002) [arXiv:hep-th/0206221].
- [11] U. Gursoy, “Vector operators in the BMN correspondence,” arXiv:hep-th/0208041.
- [12] N. Beisert, C. Kristjansen, J. Plefka, G. W. Semenoff and M. Staudacher, “BMN correlators and operator mixing in $N = 4$ super Yang-Mills theory,” *Nucl. Phys. B* **650**, 125 (2003) [arXiv:hep-th/0208178].
- [13] J. H. Schwarz, “Comments on superstring interactions in a plane-wave background,” *JHEP* **0209**, 058 (2002) [arXiv:hep-th/0208179].
- [14] A. Pankiewicz, “More comments on superstring interactions in the pp-wave background,” *JHEP* **0209**, 056 (2002) [arXiv:hep-th/0208209].
- [15] D. J. Gross, A. Mikhailov and R. Roiban, “A calculation of the plane wave string Hamiltonian from $N = 4$ super-Yang-Mills theory,” arXiv:hep-th/0208231.
- [16] J. G. Zhou, “pp-wave string interactions from string bit model,” *Phys. Rev. D* **67**, 026010 (2003) [arXiv:hep-th/0208232].
- [17] N. R. Constable, D. Z. Freedman, M. Headrick and S. Minwalla, “Operator mixing and the BMN correspondence,” *JHEP* **0210**, 068 (2002) [arXiv:hep-th/0209002].
- [18] D. Vaman and H. Verlinde, “Bit strings from $N = 4$ gauge theory,” arXiv:hep-th/0209215.

- [19] B. Eynard and C. Kristjansen, “BMN correlators by loop equations,” JHEP **0210**, 027 (2002) [arXiv:hep-th/0209244].
- [20] R. A. Janik, “BMN operators and string field theory,” Phys. Lett. B **549**, 237 (2002) [arXiv:hep-th/0209263].
- [21] J. Pearson, M. Spradlin, D. Vaman, H. Verlinde and A. Volovich, “Tracing the string: BMN correspondence at finite J^2/N ,” arXiv:hep-th/0210102.
- [22] J. Gomis, S. Moriyama and J. Park, “SYM description of SFT Hamiltonian in a pp-wave background,” arXiv:hep-th/0210153.
- [23] A. Pankiewicz and B. Stefański, jr., “pp-wave light-cone superstring field theory,” Nucl. Phys. B **657**, 79 (2003) [arXiv:hep-th/0210246].
- [24] D. Bak and M. M. Sheikh-Jabbari, “Strong evidence in favor of the existence of S-matrix for strings in plane waves,” JHEP **0302**, 019 (2003) [arXiv:hep-th/0211073].
- [25] Y. H. He, J. H. Schwarz, M. Spradlin and A. Volovich, “Explicit formulas for Neumann coefficients in the plane-wave geometry,” arXiv:hep-th/0211198.
- [26] Y. Kiem, Y. Kim, J. Park and C. Ryou, “Chiral primary cubic interactions from pp-wave supergravity,” JHEP **0301**, 026 (2003) [arXiv:hep-th/0211217].
- [27] R. Roiban, M. Spradlin and A. Volovich, “On light-cone SFT contact terms in a plane wave,” arXiv:hep-th/0211220.
- [28] U. Gursoy, “Predictions for pp-wave string amplitudes from perturbative SYM,” arXiv:hep-th/0212118.
- [29] J. A. Minahan and K. Zarembo, “The Bethe-ansatz for $N = 4$ super Yang-Mills,” JHEP **0303**, 013 (2003) [arXiv:hep-th/0212208].
- [30] N. Beisert, C. Kristjansen, J. Plefka and M. Staudacher, “BMN gauge theory as a quantum mechanical system,” Phys. Lett. B **558**, 229 (2003) [arXiv:hep-th/0212269].
- [31] J. Gomis, S. Moriyama and J. Park, “SYM description of pp-wave string interactions: Singlet sector and arbitrary impurities,” arXiv:hep-th/0301250.
- [32] K. Bobkov, “Graviton-scalar interaction in the pp-wave background,” arXiv:hep-th/0303007.
- [33] N. Beisert, C. Kristjansen and M. Staudacher, “The dilatation operator of $N = 4$ super Yang-Mills theory,” arXiv:hep-th/0303060.