

Anomalies in M-theory on singular  $G_2$ -manifoldsAdel Bilal<sup>1</sup> and Steffen Metzger<sup>1,2</sup><sup>1</sup> CNRS - Laboratoire de Physique Théorique, Ecole Normale Supérieure  
24 rue Lhomond, 75231 Paris Cedex 05, France<sup>2</sup> Sektion Physik, Ludwig-Maximilians-Universität  
München, Germany

e-mail: adel.bilal@lpt.ens.fr, metzger@physique.ens.fr

## Abstract

When M-theory is compactified on  $G_2$ -holonomy manifolds with conical singularities, charged chiral fermions are present and the low-energy four-dimensional theory is potentially anomalous. We reconsider the issue of anomaly cancellation, first studied by Witten. We propose a mechanism to cancel both abelian and non-abelian gauge anomalies locally, i.e. separately for each conical singularity. It is similar in spirit to the one used to cancel the normal bundle anomaly in the presence of ve-branes. It involves smoothly cutting off all fields close to the conical singularities, resulting in an anomalous variation of the 3-form  $C$  and of the non-abelian gauge fields present if there are also ADE singularities. Local cancellation of the mixed gauge-gravitational anomalies seems to require gravitational corrections beyond the Green-Schwarz term.

# 1 Introduction

M-theory compactified on a smooth 7-manifold  $X$  of  $G_2$ -holonomy gives rise to four-dimensional  $N = 1$  supergravity coupled to  $b_2(X)$  abelian vector multiplets and  $b_3(X)$  neutral chiral multiplets [1]. The theory contains no charged chiral fermions and there are no non-abelian gauge symmetries. Both phenomena are generated when  $X$  possesses conical and ADE-singularities [2, 3, 4, 5]. In this case the low-energy four-dimensional theory is potentially anomalous, but it was argued in ref. [6] that all anomalies cancel against various "inflow" terms.

The basic examples of conical singularities are taken from the asymptotics of the well-known non-compact  $G_2$ -manifolds [7]. Of course, Joyce's construction [8] gives compact  $G_2$ -manifolds, but no explicit example with conical singularities is known.

On the other hand, there are generalisations of  $G_2$ -holonomy manifolds which naturally are compact and have conical singularities. Mathematically they have so-called weak  $G_2$ -holonomy, and in many cases we can write down the metric explicitly [9]. Physically, they correspond to turning on a background value for the supergravity four-form field strength  $G = dC$ , thus creating a non-vanishing energy-momentum tensor which increases the curvature of  $X$  and makes it compact. When done appropriately one still has  $N = 1$  supergravity, but now in  $AdS_4$ . Doing quantum field theory, in particular loop calculations in  $AdS$  spaces is highly non-trivial, but one can still study anomalies and their cancellation, since they are topological in nature.

The aim of this note is to reconsider the anomaly cancellation mechanism for singular  $G_2$ -manifolds outlined in [6], but insisting on local cancellation, i.e. separately for each conical singularity. In particular, one has to be careful about the correct interpretation when using Stoke's theorem to rewrite bulk integrals as a sum of boundary terms. This is a general feature of anomaly cancellation through inflow from the bulk as soon as one has several "boundary" components. We show how local anomaly cancellation can be properly achieved by appropriate modifications of certain low energy effective interactions like e.g. the Chern-Simons term of eleven-dimensional supergravity, much in the same way as required for the cancellation of the normal bundle anomaly in the presence of D-branes [10]. The basic feature of these modifications is to smoothly cut off all the fields when a conical singularity is approached. If there are ADE singularities

which generate non-abelian gauge fields, this cut-off procedure induces corresponding modifications of the additional interactions present in this case. To study the mixed gauge-gravitational anomalies we also cut off the fluctuations of the geometry. Since we study quantum theory in a given background, only the fluctuations around this background are cut off, not the background itself. In any case, the relevant interactions  $S_i$  then naturally split into a "bulk" part  $S_i^{(1)}$  and a sum of terms  $S_i^{(2; \alpha)}$  localised at the various singularities  $P_\alpha$ . While the  $S_i^{(1)}$  are invariant, the variation of each  $S_i^{(2; \alpha)}$  cancels the corresponding anomaly at  $P_\alpha$  locally. This method to achieve local cancellation is rather general and powerful. Exactly the same mechanism can also be applied to discuss local gauge anomaly cancellation on weak  $G_2$ -holonomy manifolds with conical singularities as constructed in [9]. However, we also meet a surprise: the anomalous variation of the appropriately modified Green-Schwarz term matches the mixed gauge-gravitational anomaly locally, but it comes with the wrong sign. We are forced to conclude that local cancellation of this anomaly requires further gravitational corrections of higher order.

This paper is organized as follows: in section 2, we review the geometrical setup and the anomalies due to the chiral fermions present at the singularities. We remind the reader how global anomaly cancellation was shown in [6] and explain why local cancellation still remained to be proven. In section 3, we introduce our procedure of cutting off the fields close to the singularities and show how this leads to local cancellation of the gauge anomaly in the abelian case. We also give a preliminary discussion of the mixed gauge-gravitational anomaly. Section 4 deals with the non-abelian case where the cut-off procedure is more complicated due to the non-linearities. We show how the  $SU(N)^3$  and mixed  $U(1)_1 G^2$  anomalies indeed are all cancelled locally. Finally, we complete the discussion of the mixed gauge-gravitational anomaly. We conclude in section 5. In an appendix we briefly describe the compact weak  $G_2$ -manifolds with two conical singularities constructed in [9]. They provide useful explicit examples to have in mind throughout the main text.

## 2 Global anomaly cancellation for abelian gauge fields

Anomalies that arise upon compactification of M-theory on  $G_2$ -manifolds with conical singularities were first analysed by Witten [6]. The well-known non-compact metrics [7] are asymptotically, for large  $r$ , a cone on a compact six-manifold  $Y$  with  $Y = S^3 \times S^3$ ,  $Y = CP^3$  or  $Y = SU(3)/U(1)^2$ . The metrics on these manifolds all depend on some scale which we call  $r_0$ , and the conical limit is  $r_0 \rightarrow r$ . Of course, there is no singularity since, for small  $r \gg r_0$ , these metrics are perfectly regular. Mathematically, it is only in the limit  $r_0 \rightarrow 0$  that a conical singularity develops. However, if  $r_0$  is as small as the eleven-dimensional Planck length (or less) then, from the long-wavelength limit of supergravity, the manifold looks as if it had a conical singularity. Said differently, the curvature is of order  $\frac{1}{r_0^2}$  and supergravity ceases to be a valid approximation. It was argued [4] that generic singularities of compact  $G_2$ -manifolds are also conical.

In the vicinity of a conical singularity  $P$  we can always introduce a local coordinate  $r$  such that the metric can be written as

$$ds_X^2 = dr^2 + r^2 ds_Y^2 \quad (2.1)$$

with  $ds_Y^2$  the metric on the compact six-manifold  $Y$ . A necessary condition for  $ds_X^2$  to have  $G_2$ -holonomy is Ricci flatness. This in turn implies that  $Y$  is an Einstein space with  $R_{ab}^Y = 5g_{ab}$ . In fact,  $Y$  has weak  $SU(3)$ -holonomy. Furthermore, the Riemann tensors of  $X$  and  $Y$  are related as  $R_{cd}^{X ab} = \frac{1}{r^2} R_{cd}^{Y ab} + \frac{a}{c} \frac{b}{d} - \frac{a}{d} \frac{b}{c}$ ,  $a, b, c, d = 1, \dots, 6$ , and there are curvature invariants of  $X$  that diverge as  $r \rightarrow 0$ . It was argued in [3, 4, 5] that at each such singularity  $P$  there is a set  $T$  of four-dimensional chiral supermultiplets  $\psi_i$ ;  $2 \leq |T| \leq 3$ . (This set may be empty as is the case for  $Y = S^3 \times S^3$ .) They carry charges with respect to the abelian gauge group  $U(1)^k$  that arises from the Kaluza-Klein reduction of the three-form  $C$ . Note, however, that they need not be charged with respect to all  $U(1)$  gauge fields.

These charged chiral multiplets give rise to a gauge anomaly at a given singularity  $P$  characterised by the standard gauge anomaly polynomial

$$I^{\text{gauge}} = \frac{1}{6(2\pi)^2} \sum_{i=1}^k X_i^{X^k} \sum_{i=1}^3 q_i^i F_i \quad (2.2)$$

where  $\mathcal{X}^k$  labels the various chiral multiplets present at  $P$  and  $q^i$  is the charge of  $\mathcal{X}^k$  under the  $i^{\text{th}}$   $U(1)$ . The same chiral multiplets also give rise, at each singularity, to a mixed gauge-gravitational anomaly characterised by

$$I^{\text{mixed}} = \frac{1}{24} \sum_{i=1}^3 \sum_k q^i F_i \wedge p_1^0 \wedge \mathcal{X}^k \quad (2.3)$$

where  $p_1^0 = \frac{1}{8} \text{tr} R \wedge R$  is the first Pontryagin class of the four-dimensional spacetime  $M_4$ . In ref. [6], it was argued that these anomalies are cancelled locally, i.e. separately for each singularity, by an appropriate non-invariance of the Chern-Simons and Green-Schwarz terms of eleven-dimensional supergravity:

$$S_{\text{CS}} = \frac{1}{12} \int_{M_4} C \wedge G \wedge G = \frac{1}{6} \frac{T_2^3}{(\ell_p)^2} \int_{M_4} C \wedge G \wedge G; \quad (2.4)$$

$$S_{\text{GS}} = \frac{T_2}{2} \int_{M_4} C \wedge X_8; \quad (2.5)$$

where, for convenience, we replaced  $\ell_p^2$  by the membrane tension  $T_2$ , via the usual relation  $T_2^3 = \frac{(2\pi)^2}{\ell_p^2}$ . Here  $G = dC$  and  $X_8$  is the standard gravitational eight-form to be given below. With our conventions, the  $C_{MNP}$ -eld has dimension 0, so that the 3-form  $C$  and the 4-form  $G$  both have dimension 3. In particular,  $T_2 C$  is dimensionless.

Explicitly, the Kaluza-Klein reduction of  $C$  is

$$T_2 C = c + \omega_n \wedge \omega_n + A_i \wedge \omega_i + \omega_k \wedge \omega_k + \dots \quad (2.6)$$

where  $\omega_n$ ,  $\omega_i$  and  $\omega_k$  are harmonic 1-, 2- and 3-forms on  $X$ , and  $c$ ,  $\omega_n$ ,  $A_i$  and  $\omega_k$  are massless 3-, 2-, 1-form and scalar elds on  $M_4$ . The dots stand for contributions of massive elds. In particular, one gets a four-dimensional abelian gauge eld  $A_i = A_i dx^i$  for every harmonic 2-form  $\omega_i$  on  $X$ . Indeed, the gauge symmetry  $C \rightarrow C + d\lambda$  with  $T_2 \lambda = \omega_i \omega_i + \dots$  corresponds to a  $U(1)^k$  gauge transformation  $A_i \rightarrow A_i + d\lambda_i$ . Note that the standard dimension for a gauge eld  $A$  is 1, so that the one-forms  $A_i$  have dimension 0 and hence the  $\omega_i$  are dimensionless. The Kaluza-Klein reduction of  $C$  implies a similar reduction for  $G = dC$  which in particular contains a term  $F^i \wedge \omega_i$ .

Due to the conical singularities, one has to be a bit more precise about which class of harmonic 2-forms one is interested in. Inspection of the kinetic term  $\int_{M_4} dC \wedge dC$  shows that one needs square-integrable harmonic forms on  $X$ , i.e. forms satisfying

$\int_X \omega_i \wedge \omega_i < 1$ , in order to get massless 4-dimensional fields with finite kinetic terms. In particular, square-integrability requires an appropriate  $r$  dependence as  $r \rightarrow 0$ . As long as we are in a neighbourhood  $N$  of the conical singularity  $P$  where (2.1) holds, we can adapt the results of ref. [9]: every  $L^2$ -harmonic  $p$ -form  $\omega_p^{(Y)}$  on  $Y$  with  $p \leq 3$  trivially extends to a harmonic  $p$ -form  $\omega_p^{(X)} = \omega_p^{(Y)}$  on  $X$  such that  $\int_N \omega_p^{(X)} \wedge \omega_p^{(X)}$  is convergent. The same obviously is true for the Hodge duals  $\omega_p^{(X)}$  that are harmonic  $q$ -forms on  $X$  with  $q \leq 4$ . In order to decide whether these forms are  $L^2$  on  $X$ , i.e. whether  $\int_X \omega_p^{(X)} \wedge \omega_p^{(X)}$  converges at all singularities, one needs more information about the global structure of  $X$ , which we are lacking. However, for the examples of weak  $G_2$  cohomogeneity-one metrics with two conical singularities constructed in [9] this global information is available and it was shown that the harmonic forms considered above are indeed  $L^2$  and they are the only ones, so that  $b^p(X) = b^p(Y)$  for  $p \leq 3$ . Also, since for a compact Einstein space of positive curvature like  $Y$  the first Betti number always vanishes, in these examples one then has  $b^1(X) = 0$ . In the present case, we will simply need to assume that the  $L^2$ -harmonic  $p$ -forms on  $X$  for  $p \leq 3$  are given, in the vicinity of  $P$ , by the trivial extensions onto  $X$  of the  $L^2$ -harmonic  $p$ -forms on  $Y$ . In particular, then  $k = b^2(X) = b^2(Y)$  and  $b^1(X) = 0$ . (Of course, for compact smooth  $G_2$ -holonomy manifolds  $b^1(X)$  always vanishes.) Hence, the gauge group is  $U(1)^k$  and the terms  $\omega_n \wedge \omega_n$  are absent in (2.6).

In [6] it is argued that supergravity ceases to be valid close to the singularities and  $S_{CS}$  and  $S_{GS}$  should be taken as integrals over  $M_4 \setminus X^0$  only, where  $X^0$  is  $X$  with small neighbourhoods of the singularities excised. Then the boundary of  $X^0$  is  $\partial X^0 = [Y]$ , and the variation of  $S_{CS}$  can be rewritten as a sum of boundary terms [6]:

$$\delta S_{CS} = \int_{M_4 \setminus X^0} d \omega \wedge G \wedge G = \int_{M_4 \setminus Y} \omega \wedge G \wedge G : \quad (2.7)$$

Upon doing the KK reduction this yields

$$\delta S_{CS} = \int_{M_4} \omega_i F_j \wedge F_k - \int_Y \omega_i \wedge \omega_j \wedge \omega_k : \quad (2.8)$$

Next one uses the relation

$$\int_{2T} \omega^i \omega^j \omega^k = \int_Y \omega_i \wedge \omega_j \wedge \omega_k - d_{ijk}^{(1)} ; \quad (2.9)$$

found to be true for the three standard  $Y$  considered. For the example of  $Y = \mathbb{C}P^3$  there is a single harmonic 2-form  $!$ , given in terms of the Kähler form  $K$  as  $! = \frac{K}{2}$  and normalised such that  $\int_{\mathbb{C}P^3} ! \wedge ! \wedge ! = 1$ . This matches with the existence of a single multiplet with  $q = 1$ . Note that the orientation of  $Y$  is important. In the examples discussed in the appendix one has e.g.  $Y_1 = \mathbb{C}P^3$  and  $Y_2 = -\mathbb{C}P^3$ , so that one must have  $q_1 = 1$  and  $q_2 = -1$ . Given eq. (2.9), it was concluded in [6] that eq. (2.8) cancels the gauge anomaly of the chiral multiplets (2.2).

This cannot be the full story, however. In eq. (2.7) one uses Stokes' theorem to rewrite a bulk integral as a sum of boundary contributions. While mathematically perfectly correct, it is not necessarily meaningful to assign a physical interpretation to the boundary contributions individually.<sup>1</sup> These remarks suggest that the above argument (2.7) – (2.9) is insufficient to show the local character of the anomaly cancellation, separately at each singularity. Indeed, as it stands, the KK reduction of the integrand of the l.h.s. of eq. (2.7) does not give the desired contribution. For a  $U(1)^k$  gauge transformation with  $T_2 = \sum_i ! \wedge !$  the only piece contained in  $d \wedge G \wedge G = d \wedge dC \wedge dC$  which is a 4-form on  $M_4$  and only involves the massless fields is  $d_i \wedge F_j \wedge d_k$ . In particular, the desired piece  $d_i \wedge F_j \wedge F_k$  is a 5-form on  $M_4$  and cannot contribute. We conclude that eq. (2.7) is a somewhat artificial rewriting of zero, at least for the terms of interest to us, and that equations (2.8) and (2.9) only prove global anomaly cancellation, i.e. cancellation after summing the contributions of all singularities  $P$ . Indeed, global cancellation of the anomaly is the statement that  $\int_P I^{\text{gauge}} = 0$ . As remarked in [6], this is a simple consequence of eq. (2.9) and  $\int_Y \sum_i ! \wedge ! \wedge ! \wedge ! \wedge ! = \int_X d(\sum_i ! \wedge ! \wedge ! \wedge ! \wedge !) = 0$ . However, local cancellation still remains to be proven. As we will show next, it will require a modification of  $S_{CS}$ , much as when D-branes are present [10].

---

<sup>1</sup>As a trivial example consider integrating 0 over an interval  $[a; b]$ . If  $c(x)$  is any constant function we have  $\int_a^b 0 = \int_a^b dc = c(b) - c(a)$ . Obviously, there is no meaning in assigning a value  $c$  to the upper boundary and  $-c$  to the lower one.

### 3 Local anomaly cancellation for abelian gauge fields

#### 3.1 The modified fields

In the treatment of ref. [10] of the D3-brane anomaly a small neighbourhood of the D3-brane is cut out creating a boundary (analogous to  $M_4 \times Y$ ). Then the anomalous Bianchi identity  $dG = \frac{1}{2} \frac{F_2^3}{(2)^2} R$  ( $W_6$ ) is smeared out around this boundary and the Chern-Simons term gets an anomalous variation localised on this smeared out region. Alternatively, this could be viewed as due to a two-form field  $B$  living close to the boundary and transforming as  $\delta B = \dots$ . The CS-term is given by  $S_{CS} = \int \frac{1}{6} \frac{F_2^3}{(2)^2} R \mathcal{C} \wedge \mathcal{G} \wedge \mathcal{G}$  with appropriately modified  $\mathcal{C}$  and  $\mathcal{G}$  (which coincide with  $C$  and  $G = dC$  away from the D3-brane and its neighbourhood) such that  $S_{CS}$  is non-vanishing and cancels the left-over normal bundle anomaly.

Now we show that a similar treatment works for conical singularities. We first concentrate on the neighbourhood of a given conical singularity  $P$  with a metric locally given by  $ds_X^2 = dr^2 + r^2 ds_Y^2$ . The local radial coordinate obviously is  $r > 0$ , the singularity being at  $r = 0$ . As mentioned above, there are curvature invariants of  $X$  that diverge as  $r \rightarrow 0$ . In particular, supergravity cannot be valid down to  $r = 0$ . Rather than cutting off the manifold at some  $r = r > 0$ , we cut off the fields which can be done in a smooth way. However, we keep fixed the geometry, and in particular the metric and curvature on  $X$ . Said differently, we cut off all fields that represent the quantum fluctuations but keep the background fields (in particular the background geometry) as before. Introduce a smooth but finite regulator  $\epsilon$  and the regularised step function  $\theta_\epsilon(r - r)$  such that

$$\begin{aligned} \theta_\epsilon(r - r) &= 0 & \text{if } 0 < r - r < \epsilon &; \\ \theta_\epsilon(r - r) &= 1 & \text{if } r - r > \epsilon &; \end{aligned} \tag{3.1}$$

with  $\theta_\epsilon$  a non-decreasing smooth function between  $r - \epsilon$  and  $r + \epsilon$ . (Outside the neighbourhood where the local coordinate  $r$  is defined,  $\theta_\epsilon$  obviously equals 1.) We define the corresponding regularised Chern-Simons action as<sup>2</sup>

$$S_{CS} = \int \frac{1}{6} \frac{F_2^3}{(2)^2} R \mathcal{C} \wedge \mathcal{G} \wedge \mathcal{G} \tag{3.2}$$

---

<sup>2</sup>We write  $\delta$  rather than  $\delta$  since the latter symbol already denotes the gauge variation of a quantity.



Of course, if  $X$  has several conical singularities (see the appendix for examples),  $\mu$  must have the appropriate behaviour (3.1) at each singularity  $P$ . It can be constructed as the product of the individual  $\mu$ 's and then  $\mu$  becomes the sum of the individual  $\mu$ 's:

$$\mu = \sum_X \mu_X \quad (3.3)$$

When evaluating integrals one has to be careful since e.g.  $r^2 \notin \mathcal{D}$ , although  $r^2$  would be just as good a definition of a regularised step function. We write  $r^2 \theta$  which means that, in an integral, one can replace  $r^2$  by  $r^2 \theta$  when multiplied by a form that varies slowly between  $r$  and  $r+$ . However, one has e.g.  $\int r^2 = \int r^2 d^4x = \frac{1}{3} d^3x$ ,  $\int \frac{1}{3} d^3x = \frac{1}{3}$  where a crucial  $\frac{1}{3}$  has appeared. Then, for any ten-form  $\omega_{(10)}$ , not containing  $\mu$ 's or  $\mu$ 's, we have

$$\int_{M_4 \times X} \omega_{(10)} = \int_{M_4 \times Y} \frac{1}{n+1} \omega_{(10)} \quad (3.4)$$

It is always understood that the regulator is removed,  $\epsilon \rightarrow 0$ , after the integration.

Now we cut off the fields with this so that all fields vanish if  $r < r$  for some  $r$ . Starting from  $C$  and  $G = dC$  we define

$$\hat{C} = C \theta; \quad \hat{G} = G \theta \quad (3.5)$$

Then the gauge-invariant kinetic term for the  $C$ -field is constructed with  $\hat{G}$ :

$$S_{kin} = \frac{1}{4} \int_{r > r} \hat{G} \wedge \hat{G} = \frac{1}{4} \int_{r > r} dC \wedge dC \quad (3.6)$$

and the  $A_i$  resulting from the KK reduction of  $C$  still are massless gauge fields. To construct a satisfactory version of the Chern-Simons term, we first note that, of course,  $\hat{G} \notin d\hat{C}$  and  $d\hat{G} = G \notin 0$ . However, we want a modified  $G$ -field which vanishes for  $r < r$ , is closed everywhere and is gauge invariant. Closedness is achieved by subtracting from  $\hat{G}$  a term  $C \wedge \theta$ , but this no longer is gauge invariant under  $C = d$ .

In order to maintain gauge invariance we add another two-form field  $B$ , that effectively only lives on the subspace  $r < r < r+$ , with

$$B = \theta \omega \quad (3.7)$$

In the limit  $\epsilon \rightarrow 0$ ,  $B$  really is a ten-dimensional field, although we treat it as an "auxiliary" field that has no kinetic term. Of course, a gauge-invariant kinetic term

could be added as  $\int_{M_4 \times Y} \mathcal{C} \wedge (\mathcal{C} - dB)$  but it is irrelevant for our present purpose. In any case

$$\mathcal{E} = G \wedge (\mathcal{C} - dB) \quad (3.8)$$

satisfies all requirements:

$$d\mathcal{E} = 0; \quad \mathcal{E} = 0; \quad \mathcal{E} = 0 \text{ for } r < r_0 : \quad (3.9)$$

We have

$$\mathcal{E} = d\mathcal{C} \quad (3.10)$$

with

$$\mathcal{C} = C + B \wedge \quad (3.11)$$

and

$$\mathcal{E} = dC + \wedge = d(\quad) : \quad (3.12)$$

### 3.2 The $U(1)^3$ anomaly

All this is similar in spirit to ref. [10], and we propose that  $S_{CS}$  should be replaced by

$$\mathcal{S}_{CS} = \frac{1}{6} \frac{T_2^3}{(2\pi)^2} \int_{M_4 \times X} \mathcal{C} \wedge \mathcal{E} \wedge \mathcal{E} : \quad (3.13)$$

We may view the differences  $\mathcal{C} - C$  and  $\mathcal{E} - G$  as gravitational corrections in an effective low-energy description of M-theory. Actually, further gravitational terms of higher order can and do appear. They are irrelevant to the present discussion of pure gauge anomalies but, as we will see below, they are expected to play a role when studying mixed gauge-gravitational anomalies.

Note that in order to discuss local anomaly cancellation, i.e. cancellation singularity by singularity, we are not allowed to integrate by parts, i.e. use Stoke's theorem. More precisely, we must avoid partial integration in the  $r$ -direction since, as remarked above, this could shift contributions between the different singularities. However, once an expression is reduced to an integral over a given  $M_4 \times Y$ , corresponding to a given singularity, one may freely integrate by parts on  $M_4 \times Y$ , as usual. In particular consider any smooth  $p$ - and  $(9-p)$ -forms  $'$  and  $d'$  not containing  $r$ . Then

$$\begin{aligned} \int_{M_4 \times X} d' \wedge \quad^n &= \frac{1}{n+1} \int_{M_4 \times Y} d' \wedge \quad^n \\ &= \int_{M_4 \times Y} \quad^{p+1} \frac{1}{n+1} \wedge d' = \int_{M_4 \times X} \quad^{p+1} \wedge d' \quad^n : \quad (3.14) \end{aligned}$$

We see that whenever an integral contains a  $\int$  we are allowed to "integrate by parts", but the derivative  $d$  does not act on the  $\omega$ 's.

Writing out  $\mathcal{S}$  and  $\mathcal{C}$  explicitly, the modified Chern-Simons term reads

$$\mathcal{S}_{CS} = \frac{1}{6} \frac{T_2^3}{(\ell)^2} \int_{M_4 \times X} C \wedge G \wedge G^3 + (B \wedge G \wedge G - 2dB \wedge C \wedge G)^2$$

$$\mathcal{S}_{CS}^{(1)} + \sum_X \mathcal{S}_{CS}^{(2i)}; \quad (3.15)$$

where we used  $\int_P = \int$ , see eq. (3.3). In the limit  $\ell \rightarrow 0$ , the first term  $\mathcal{S}_{CS}^{(1)}$  reproduces the usual bulk term, but only for  $r \gg \ell$ , while the term  $\mathcal{S}_{CS}^{(2i)}$ , due to the presence of  $\int$ , each are localised on the ten-manifolds  $M_4 \times Y$  close to the singularities  $P$ . Although they look similar, they do not arise as boundary terms. We have

$$\mathcal{S}_{CS}^{(2i)} = \frac{1}{18} \frac{T_2^3}{(\ell)^2} \int_{M_4 \times Y} (B \wedge G \wedge G - 2dB \wedge C \wedge G)$$

$$= \frac{1}{6} \frac{T_2^3}{(\ell)^2} \int_{M_4 \times Y} B \wedge G \wedge G; \quad (3.16)$$

This result illustrates again eq. (3.14). An "anomalous" variation of each  $\mathcal{S}_{CS}^{(2i)}$  then arises since  $B = \ell \rightarrow 0$ :

$$\delta \mathcal{S}_{CS}^{(2i)} = \frac{1}{6} \frac{T_2^3}{(\ell)^2} \int_{M_4 \times Y} \delta B \wedge G \wedge G; \quad (3.17)$$

Of course, there is also the "usual" variation of  $\mathcal{S}_{CS}^{(1)}$ :

$$\delta \mathcal{S}_{CS}^{(1)} = \frac{1}{6} \frac{T_2^3}{(\ell)^2} \int_{M_4 \times X} dC \wedge G \wedge G^3; \quad (3.18)$$

Globally, this equals  $\int_P \mathcal{S}_{CS}^{(2i)}$  as could easily be seen when integrating by parts. Indeed, globally  $\mathcal{S}_{CS}$  is invariant. This is alright, since we know from [6] that globally, i.e. when summed over the singularities, there are no anomalies to be cancelled.

Next, we will see what happens upon Kaluza-Klein reduction. We will keep all massless fields, not only the gauge fields. In agreement with the above discussion we assume that  $b^1(X) = 0$ . As before, let  $\omega_i$  be a basis of  $L^2$ -harmonic 2-forms and  $\omega_k$  of  $L^2$ -harmonic 3-forms on  $X$ . Then

$$\begin{aligned} T_2 C &= c + A_i \omega_i + \omega_k \omega_k + \dots \\ T_2 G &= dc + F_i \omega_i + d\omega_k \omega_k \dots \\ T_2 B &= b + f_i \omega_i + \dots; \end{aligned} \quad (3.19)$$

$f_i$  and  $k$  are massless scalar fields similar to axions, while  $c$  and  $A_i$  are 3-form and 2-form fields on  $M_4$  respectively. The dots indicate contributions of  $m$  massive fields. Under a "gauge" transformation with

$$T_2 = \dots + \dots + \dots \quad (3.20)$$

one has for the 4-dimensional fields

$$c = d \dots; \quad A_i = d \dots; \quad k = 0; \quad \dots = \dots; \quad f = \dots \quad (3.21)$$

Then in the "bulk"-term  $\mathcal{S}_{CS}^{(1)}$ , the only non-vanishing contribution of the massless fields is

$$\mathcal{S}_{CS}^{(1)} = \frac{1}{6(2\pi)^2} \int_{M_4} \epsilon_{ijkl} F_i \wedge F_j \wedge F_k \wedge F_l + \dots \quad (3.22)$$

Obviously, this is gauge-invariant. For the terms  $\mathcal{S}_{CS}^{(2)}$ , localised near the singularities  $P$ , we get

$$\mathcal{S}_{CS}^{(2)} = \frac{1}{6(2\pi)^2} \int_{M_4} \epsilon_{ijkl} f_i F_j \wedge F_k \wedge F_l + \dots \quad (3.23)$$

Under a  $U(1)^k$  gauge transformation with  $\alpha = 0$  but  $\beta_i \neq 0$ , the second term in this expression is invariant, but the first one is not.

We finally conclude that under a  $U(1)^k$ -gauge transformation with  $\beta_i \neq 0$  (but  $\alpha = 0$ ) we have

$$\mathcal{S}_{CS}^{(1)} = \dots + \dots$$

$$\mathcal{S}_{CS}^{(2)} = \frac{1}{6(2\pi)^2} \int_{M_4} \epsilon_{ijkl} \beta_i F_j \wedge F_k \wedge F_l + \dots \quad (3.24)$$

Using the relation (2.9) it is then obvious that, separately at each singularity, this precisely cancels the gauge anomaly obtained from (2.2) via the descent equations. Hence, anomaly cancellation indeed occurs locally.

Before we go on, a remark is in order. To cancel the four-dimensional gauge anomalies we modified the eleven-dimensional Chern-Simons term, including a new interaction with the ten-dimensional non-dynamical  $B$ -field. This was natural and necessary to

have an invariant  $\mathcal{G}$ -eld. As a result, the gauge variation of the KK reduction of  $\mathcal{S}_{CS}^{(2;1)}$  no longer vanishes and was seen to cancel the four-dimensional gauge anomalies. This might look as if we had found a four-dimensional counterterm  $\int_{M_4}^R f_i F_j \wedge F_k$  to cancel the anomaly. Now, a relevant anomaly cannot be cancelled by the variation of a local four-dimensional counterterm of the gauge fields. The point is, of course, that this not only contains the gauge fields but also the axion-like fields  $f_i$  that arose from the non-dynamical B-field and it is the non-invariance of the  $f_i$  that leads to anomaly cancellation. This is quite different from adding a four-dimensional counterterm.

It is also interesting to note that under transformations with  $\delta = \delta_{CS}$  we get non-vanishing  $\mathcal{S}_{CS}^{(2;1)}$ , with no corresponding "fermion anomaly" to be cancelled locally. Of course, globally these variations vanish, but locally they do not. However, this is not harmful as it would be for gauge anomalies, since anyway, these transformations only affect fields that do not propagate on  $M_4$ .

### 3.3 The mixed gauge-gravitational anomaly

The mixed gauge-gravitational anomaly (2.3) should be cancelled similarly through local anomaly "inflow" from an appropriately modified Green-Schwarz term. This now involves the gauge fields and the gravitational fields. Which fields should be cut off at the singularities? Our general philosophy is to keep fixed the background fields and in particular the background geometry, but to cut off the fluctuations around this background. If we call  $\omega_0$  the spin-connection of the background geometry, and  $\omega = \omega_0 + \delta\omega$  the one of the full geometry including the fluctuations around  $\omega_0$ , then one should cut off only  $\delta\omega$  so that  $\omega = \omega_0 + \delta\omega \in \mathfrak{so}(3,1) + \mathfrak{e}$ . In principle one should then work with the gravitational 8-form  $\tilde{X}_8$  computed with this  $\delta\omega$ , and start with

$$\mathcal{S}_{GS} = \frac{T_2}{2} \int \mathcal{C} \wedge \tilde{X}_8 : \quad (3.25)$$

Dealing correctly with the cut-off spin connection requires some machinery which we will only introduce in the next section where we deal with non-abelian gauge fields. However, there we will also see that the two terms  $\int^R \mathcal{C} \text{tr} F^2$  and  $\int^R \mathcal{C} \text{tr} \mathbb{F}^2$  (with  $F$  a non-abelian field strength and  $\mathbb{F}$  its cut-off version) lead to exactly the same anomaly inflow for the mixed  $U(1)_1 G^2$  anomaly. Hence we expect that, in the same way,  $\int^R \mathcal{C} X_8$

and  $\int_{M_8} \mathbb{C}^{\mathbb{R}} \wedge X_8$  may also lead to the same anomaly in flow for the mixed  $U(1)_I$ -gravitational anomaly. We will explicitly verify this in section 4.4. Here we consider<sup>3</sup>

$$\mathbb{S}_{GS}^0 = \frac{T_2}{2} \int_{M_8} \mathbb{C} \wedge X_8 : \quad (3.26)$$

As usual,  $X_8$  is given by  $X_8 = \frac{1}{24(\ell_p)^3} \frac{1}{8} \text{tr} R^4 - \frac{1}{32} (\text{tr} R^2)^2$ , and  $R$  is evaluated with  $\omega = \omega_0 + \delta\omega$ . In principle, one should consider arbitrary fluctuations that do not necessarily preserve the product structure of the manifold,  $M_4 \times X$ , but, for simplicity, we will assume they do.<sup>4</sup> Then one can rewrite  $X_8$  in terms of the first Pontryagin classes  $p_1^0 = \frac{1}{8} \text{tr} R^2$  and  $p_1^{00} = \frac{1}{8} \text{tr} R^2$  of  $M_4$  and  $X$  respectively as  $X_8 = \frac{1}{48} p_1^0 \wedge p_1^{00}$ . Note that this is second and higher order in the fluctuations since the background geometry of  $M_4$  is flat at  $R^4$ . Hence

$$\mathbb{S}_{GS}^0 = \frac{T_2}{96} \int_{M_4 \times X} \mathbb{C} \wedge p_1^0 \wedge p_1^{00} : \quad (3.27)$$

Inserting the Kaluza-Klein decomposition of  $\mathbb{C}$  we get again a "bulk" part and a sum of contributions localised close to the singularities:

$$\begin{aligned} \mathbb{S}_{GS}^0 &= \mathbb{S}_{GS}^{0(1)} + \int_X \mathbb{S}_{GS}^{0(2; \cdot)} ; \\ \mathbb{S}_{GS}^{0(1)} &= \frac{1}{96} \int_{M_4 \times X} p_1^0 \wedge p_1^{00} ; \\ \mathbb{S}_{GS}^{0(2; \cdot)} &= \frac{1}{24} \int_{M_4} f_i p_1^0 \frac{1}{4} \int_Y \omega_i \wedge p_1^{00} : \end{aligned} \quad (3.28)$$

In the last integral over  $Y$ ,  $p_1^{00}$  now is the first Pontryagin class of  $Y$ . This follows easily from the properties of the characteristic classes for the geometry at hand.<sup>5</sup> Now one

<sup>3</sup>Alternative forms of the Green-Schwarz term would be  $\frac{T_2}{2} \int_{M_8} \mathbb{C}^{\mathbb{R}} \wedge X_7$  or  $\frac{T_2}{2} \int_{M_8} \mathbb{C}^{\mathbb{R}} \wedge X_7$ . Although globally equivalent to (3.25), respectively (3.26), a priori they could lead to different local variations. Nevertheless, we have checked that the final result always equals (3.31).

<sup>4</sup>As always, dangerous anomalies are associated with the massless modes. Hence, we only need to consider fluctuations of the metric that are massless. Massless fluctuations that do not preserve the product structure would arise e.g. if  $X$  had non-trivial Killing vectors. However, we know that for  $G_2$ -manifolds this is not the case.

<sup>5</sup>Explicitly, this can be seen as follows. On  $X$  one has  $\text{tr} R^6 = \frac{1}{6} \text{tr} R^6 + 2 \text{tr} R^7 \wedge R^7$ . But for the cones  $\text{tr} R^7 = 0$ . Furthermore, the curvature 2-forms on  $X$  and  $Y$  are related by  $R_X = R_Y \otimes e + \omega$  with  $e$  the 6-beins on  $Y$ . Since with the relevant geometries we have  $R_Y \wedge e \wedge e = 0$  one sees that  $\text{tr} R^6 = \text{tr} R^6$  and hence  $p_1^0(X) = p_1^0(Y)$ . Of course, eq. (3.28) is unchanged by the fluctuations of the geometry on  $X$  since  $\int_X$  has topologically invariant integrals, and so does  $\int_Y \omega_i \wedge p_1^{00}$ .

uses another relation which relates the charges of the chiral fermions to the geometric properties of  $Y$  namely

$$\frac{1}{4} \int_Y \text{tr} \wedge^2 p_1^0(Y) = \sum_{2T} q_i^i : \quad (3.29)$$

The only of the three examples for which this relation is non-trivial is  $Y = \mathbb{C}P^3$  where  $p_1^0 = 4! \wedge !$  and one correctly gets  $\int \text{tr} \wedge^2 ! \wedge ! = q = 1$ . Then

$$\mathbb{S}_{GS}^{0(2;)} = \sum_{2T} \frac{q_i^i}{24} \int_{M_4} f_i p_1^0 : \quad (3.30)$$

Finally, we conclude that under a  $U(1)_i$  gauge transformation  $\mathbb{S}_{GS}^{(1)}$  is invariant while

$$\mathbb{S}_{GS}^{0(2;)} = \sum_{2T} \frac{q_i^i}{24} \int_{M_4} i p_1^0 : \quad (3.31)$$

This is exactly what we need to cancel the mixed gauge-gravitational anomaly due to the chiral fermions associated with (2.3), except that (3.31) has the wrong sign!

We have carefully checked the signs. For the original Green-Schwarz term, written either as  $\frac{T_2}{2} \int C \wedge X_8$  or as  $\frac{T_2}{2} \int G \wedge X_7$  with  $dX_7 = X_8$  one can find both signs in the literature depending on the sign convention used for  $C$ . However, with the convention we use, where the coefficient in  $S_{CS}$  in front of  $CGG$  is  $\frac{1}{6} \frac{T_2^3}{(2)^2}$ , the correct sign for the GS-term is  $+\frac{T_2}{2} \int C \wedge X_8$ . In particular, this relative sign between both terms is a necessary condition for the cancellation of the normal bundle anomaly of the five-brane.

If, in the present setting, the contributions to the mixed gauge-gravitational anomaly from the fermions and from the modified Green-Schwarz term do not cancel but add up, we are forced to conclude that we are still missing some other contribution to this anomaly. Indeed, it is quite conceivable that we need to include higher-order gravitational corrections in the Chern-Simons term or elsewhere. It could be possible that, on top of the modifications  $G \rightarrow \mathbb{G}$  we already performed, one has to include terms like  $(1 + \dots) \text{tr} R \wedge R$  or  $\int_{3L} !$  into  $\mathbb{G}$ . Such terms vanish in the "bulk", but would give a non-vanishing  $d\mathbb{G}$  near the singularities and give contributions in  $S_{CS}$  similar to (3.31). However, since we have found no unambiguous way to fix the coefficients of such terms, we will not explore this possibility any further.

## 4 Anomally cancellation for non-abelian gauge fields

If  $X$  has ADE singularities non-abelian gauge fields are generated. Since ADE singularities have codimension four, the set of singular points is a three-dimensional submanifold  $Q$ . Such geometries have been discussed extensively in the literature see e.g. [2, 4, 6]. The interesting situation is when  $Q$  itself has a conical singularity. In the neighbourhood of such a singularity  $P$ , we may still think of  $X$  as a cone on  $Y$ , but now  $Y$  is an ADE orbifold. Let  $U$  be the two-dimensional singularity (fixed-point) set of  $Y$ . Then locally  $Q$  is a cone on  $U$ .

On the seven-dimensional spacetime  $M_4 \times Q$  there live non-abelian ADE gauge fields  $A$  with curvature  $F = dA + A^2$ . After KK reduction they give rise to four-dimensional ADE gauge fields and field strengths which we call again  $A$  and  $F$ . In addition, on  $M_4$ , we may still have abelian gauge fields  $A_i$  with field strength  $F_i = dA_i$ , which arise from the KK reduction of the C-eld. The four-dimensional chiral supermultiplets present at the singularities  $P$  now couple to the gauge fields of the non-abelian group  $G$  and are charged with respect to the abelian  $A_i$ . Then there are potentially  $U(1)^3, U(1)G^2$  and  $G^3$  anomalies. The first are cancelled as described above by inflow from the  $\mathbb{C} \wedge \mathbb{C} \wedge \mathbb{C}$  term. The  $G^3$  anomaly is present only for  $G = SU(N)$ .

### 4.1 Consistent versus covariant anomalies

In the non-abelian case anomalies can manifest themselves in two different ways, as consistent or covariant anomalies [11]. As is well-known from the early days of the triangle anomaly in four dimensions, if the regularisation of the one-loop diagram respects Bose symmetry in the external gauge fields one gets the consistent anomaly. Alternatively one may preserve gauge invariance (current conservation) for two of the three external gauge fields (including contributions of square and pentagon diagrams), with all non-invariance only in the third eld. This leads to the covariant form of the anomaly. For one chiral fermion of unit charge this (integrated) covariant anomaly is given by

$$A_{\text{covariant}}^{SU(N)^3} = \frac{1}{2(2\pi)^2} \int_{M_4} \text{tr} F^2 : \quad (4.1)$$



Similarly, the (integrated) consistent anomaly for one chiral fermion of unit charge is

$$A_{\text{consistent}}^{\text{SU}(N)^3} = \frac{1}{6(2\pi)^2} \int_{M_4} \text{tr} \, dA dA + \frac{1}{2} A^3 \quad : \quad (4.2)$$

The consistent anomaly is related via the descent equations<sup>6</sup> to the invariant six-form  $\frac{1}{6(2\pi)^2} \text{tr} F^3$ . Obviously, since the covariant anomaly cannot be obtained this way, it is not possible to find a local counterterm of the gauge fields such that  $A_{\text{covariant}} = A_{\text{consistent}} + \dots$ . However, on the level of the corresponding currents  $J$  one can find [11] a local  $X$  such that  $J_{\text{covariant}} = J_{\text{consistent}} + X$ .

Clearly, if the consistent anomalies cancel when summed over the contributions of all chiral fermions, the same is true for the covariant anomalies, and vice versa. Here, however, we want to cancel a non-vanishing anomaly due to chiral fermions (originating from a given conical singularity) by an appropriate anomaly inflow from a higher-dimensional action, i.e. by some  $S$ . Such a setup respects Bose symmetry between all gauge fields and it is clear that we must cancel the consistent anomaly, not the covariant one. Indeed, when invoking anomaly inflow, one wants to show that the resulting total effective action is invariant. But any non-invariance of part of the effective action must satisfy the Wess-Zumino consistency conditions [12] and hence be the consistent anomaly. There has been some discussion in the literature about consistent versus covariant inflow [13, 14, 15]: in all cases there is a consistent anomaly due to fermions to be cancelled by an inflow.<sup>7</sup>

Similarly, the mixed  $U(1)_i G^2$  anomaly can also be expressed in a covariant or consistent form. The consistent form derives from the invariant 6-form  $\frac{1}{6(2\pi)^2} 3 q^i F_i \text{tr} F^2$  via the descent equations. It can manifest itself as  $\frac{1}{2(2\pi)^2} q^i \text{tr} F^2$  or  $\frac{1}{2(2\pi)^2} q^i F_i \text{tr} \, dA$  or any combination of these two with total weight one:

$$A_{\text{consistent}}^{U(1)_i G^2} = \frac{q^i}{2(2\pi)^2} \int_{M_4} \text{tr} F^2 + (1 - \dots) F_i \text{tr} \, dA \quad : \quad (4.3)$$

<sup>6</sup>  $\text{tr} F^3 = d!_5$ ,  $!_5 = d!_4$ , with  $!_5 = \text{tr} \, A F^2 - \frac{1}{2} A^3 F + \frac{1}{10} A^5 = \text{tr} \, A dA dA + \frac{3}{2} A^3 dA + \frac{3}{5} A^5$  and  $!_4 = \text{tr} \, dA F - \frac{1}{2} A^3 = \text{tr} \, dA dA + \frac{1}{2} A^3$ .

<sup>7</sup> In these papers a first inflow computation [13] gave a covariant anomaly inflow in discrepancy with the consistent anomaly due to the fermions. It was then argued [14] that a careful computation of the inflow actually gives two pieces for the current, the old covariant one, and a new one converting the consistent fermion current into a covariant one. However, a more fruitful interpretation is to observe that the new inflow contribution is exactly what was needed to convert the old covariant inflow into a consistent inflow.

The parameter  $\alpha$  can be changed by the addition of a local four-dimensional counter-term  $\int_{M_4} \text{tr} F^2$  (where  $\text{tr} F^2 = \text{tr} F^2$ ). Note that the mixed  $U(1)_i G^2$  anomaly is present for any  $G$ , not only  $SU(N)$ . As for the pure  $SU(N)^3$  anomaly, the mixed anomaly can also be expressed in a covariant form, but we will not need it here.<sup>8</sup>

## 4.2 The $SU(N)^3$ anomaly

In ref. [6] it was argued that these anomalies can be cancelled by the non-invariance of certain interactions, namely  $\int_{M_4 Q} K \wedge \text{tr} F^2$  for the  $SU(N)^3$  anomaly and  $\int_{M_4 Q} C \wedge \text{tr} F^2$  for the mixed one. Here  $K$  is the curvature of a certain line bundle with a connection induced by the metric on  $X$ .  $K$  is a 2-form on  $Q$  and must obey  $dK = 0$  except at the singularities of  $Q$  where one could get  $\delta$ -function contributions. Thus it makes sense to define

$$n = \int_Q \frac{K}{2} : \quad (4.5)$$

Actually,  $\frac{K}{2}$  is the first Chern class of the line bundle and hence the  $n$  are integers. To show that the variation of  $S_1$  and  $S_2$  cancel the fermion anomalies, ref. [6] again integrates by parts on  $Q$ . According to our discussion above this only proves that anomalies cancel globally. To achieve local cancellation we should find interactions localised close to the singularities, such that their variations individually cancel the fermion anomalies at each singularity. This will again involve cutting off the gauge fields using  $\delta$ , but things will be slightly more complicated due to the non-linear structure of the non-abelian fields.

To see how we should cut off the non-abelian gauge fields we recall the important points of the abelian case: 1) all (fluctuating) fields should vanish close enough to the conical singularities and equal the usual ones "sufficiently far away" from these

<sup>8</sup>As for the  $SU(N)^3$  anomaly, the covariant form arises if, in the triangle diagram, one maintains  $U(1)$  or  $G$  gauge invariance at two of the vertices and then checks the gauge variations at the third vertex. If one probes for  $U(1)$  invariance, there are non-abelian gauge fields at the two other vertices, while when probing  $G$ -invariance there are one abelian and one non-abelian gauge field at the other vertices, yielding a relative combinatorial factor 2. Hence the covariant mixed anomaly is

$$A_{\text{covariant}}^{U(1)_i G^2} = \frac{q^i}{2(2\pi)^2} \int_{M_4} \text{tr} F^2 + 2F_i \text{tr} F : \quad (4.4)$$

This is similar to (4.3) with  $\alpha = \frac{1}{3}$ , but the coefficient is again 3 times larger and, of course, we have the covariant  $F_i \text{tr} F$  instead of  $F_i \text{tr} dA$ .

singularities, and 2) the modified field strengths should have the same properties as the unmodified ones. The first requirement allows fields to be a combination of terms involving  $n$  or  $n-1$  with  $n \geq 1$ . In the abelian case only  $n = 1$  occurred, see eqs (3.8) and (3.11). The field strength  $\mathcal{G}$  obeyed  $d\mathcal{G} = 0$  and  $\mathcal{G} = 0$  just as  $dG = 0$  and  $G = 0$ . Hence, it satisfied also the second requirement. This was guaranteed because the relation between  $\mathcal{G}$  and  $\mathcal{C}$  was the same as the one between  $G$  and  $C$ , namely  $\mathcal{G} = d\mathcal{C}$ , while the gauge transformation was  $\mathcal{C} = d(\ )$  with an explicit  $\epsilon$  to make sure the transformed field also satisfies the first requirement.

Now we want to apply both requirements to the non-abelian case. The gauge field  $A$  and field strength  $F$  (defined on the 7-manifold  $M_4 \times Q$ ) should be replaced by cut-off fields  $\mathcal{A}$  and  $\mathcal{F}$ . It is then clear that the second requirement will be satisfied if

$$\mathcal{A} = de + [\mathcal{A}; e] \quad (4.6)$$

and

$$\mathcal{F} = d\mathcal{A} + \mathcal{A}^2 : \quad (4.7)$$

From the first requirement then

$$e = \epsilon : \quad (4.8)$$

(Here,  $\epsilon$  is a Lie algebra-valued smooth function on  $M_4 \times Q$ .) In particular these equations guarantee that

$$\mathcal{F} = [\mathcal{F}; e] ; \quad d\mathcal{F} + [\mathcal{A}; \mathcal{F}] = 0 ; \quad (4.9)$$

as usual. The difference with the abelian case is that the non-linear structure (4.6) together with (4.8) imply that  $\mathcal{A}$  cannot simply be of the form  $a + f$ , but instead is

$$\mathcal{A} = \sum_{n=1}^k a_n + f_n : \quad (4.10)$$

The  $a_n$  are smooth 1-form fields on  $M_4 \times Q$ , while the  $f_n$  are smooth scalar fields, also on  $M_4 \times Q$  but effectively only on  $M_4 \times U$ . The latter are analogous to the  $B$ -field of the abelian case. Note that "in the bulk", i.e. for  $r > r_+$  where  $\epsilon = 1$  and  $\epsilon = 0$ , we have

$$\mathcal{A}_{\text{bulk}} = \sum_{n=1}^k a_n : \quad (4.11)$$

The gauge transformation (4.6) implies

$$\begin{aligned} a_i &= d & ; & & f &= \\ a_n &= [a_{n-1}; ] & ; & & f_n &= [f_{n-1}; ] & ; & n \geq 2 : \end{aligned} \quad (4.12)$$

In particular, it follows that the "bulk"-eld  $\mathbb{A}_{\text{bulk}}$  transforms as an ordinary gauge eld,

$$\mathbb{A}_{\text{bulk}} = d + \mathbb{A}_{\text{bulk}} ; \quad (4.13)$$

as it should. To simplify the notations below, we introduce

$$\begin{aligned} a &= a(\cdot) = \sum_{n=1}^X a_n \\ f &= f(\cdot) = \sum_{n=1}^X f_n \end{aligned} \quad (4.14)$$

so that

$$\mathbb{A} = a + f ; \quad (4.15)$$

Let furthermore  $a^0 = \sum_{n=1}^P a_n$ , as well as  $\hat{d}a = \sum_{n=1}^P (da_n)$  and  $\hat{d}f = \sum_{n=1}^P (df_n)$ . Of course,  $\hat{d}$  behaves as an exterior derivative and, in particular,  $\hat{d}^2 = 0$ . Then  $da = \hat{d}a$  and

$$d\mathbb{A} = \hat{d}a + (\hat{d}f) ; \quad (4.16)$$

Finally, we are in a position to show that the consistent fermion anomaly is cancelled by the non-invariance of the following interaction<sup>9</sup>

$$\mathcal{S}_1 = \frac{1}{6(2)^2} \int_{M_4 \times Q} \frac{K}{2} \wedge !_5(\mathbb{A}) \quad (4.17)$$

where  $!_5(\mathbb{A})$  is the standard Chern-Simons 5-form with  $\mathbb{A}$  replacing  $A$ , namely

$$!_5(\mathbb{A}) = \text{tr} \left( \mathbb{A} d\mathbb{A} d\mathbb{A} + \frac{3}{2} \mathbb{A}^3 d\mathbb{A} + \frac{3}{5} \mathbb{A}^5 \right) ; \quad (4.18)$$

This is a sum of a "bulk" term not containing  $\omega$  and a term linear in  $\omega = \sum_{i=1}^P \omega_i$ , so that we can again write

$$\mathcal{S}_1 = \mathcal{S}_1^{(1)} + \sum_{i=1}^X \mathcal{S}_1^{(2; i)} ; \quad (4.19)$$

<sup>9</sup>Actually, we should start with an  $\mathcal{S}_1$  where also  $K$  is replaced by  $\mathbb{K} = K_0 + d\omega$  with  $K_0$  corresponding to the background geometry and  $d\omega$  taking into account fluctuations around this background. Here  $\omega$  is the appropriately cut-off spin connection on the line bundle:  $\omega = \sum_{i=1}^P \omega_i$ . It is easy to include this  $d\omega$ -term into the computation which follows and show that it does not contribute to  $\mathcal{S}_1$ .

where the  $\mathcal{S}_1^{(2; \cdot)}$  reduce to integrals over  $M_4 \times U$ . Although it is straightforward to explicitly compute the  $\mathcal{S}_1^{(2; \cdot)}$ , the resulting expressions are not very illuminating. However, their gauge variations turn out to be simple, and this is why we will first compute the variations  $\delta \mathcal{S}_1^{(2; \cdot)}$ , and then reduce them to integrals over  $M_4 \times U$ .

Since  $\mathcal{F}$  satisfies the standard relation (4.6) we know that

$$\int_5(\mathcal{F}) = d \int_4(e^{\mathcal{F}}) \quad (4.20)$$

with

$$\int_4(e^{\mathcal{F}}) = \text{tr} e^{\mathcal{F}} d\mathcal{F} + \frac{1}{2}\mathcal{F}^3; \quad (4.21)$$

so that

$$\mathcal{S}_1 = \frac{1}{6(2-\kappa)^2} \int_{M_4 \times Q} \frac{\kappa}{2} \wedge d \int_4(e^{\mathcal{F}}); \quad (4.22)$$

The next step is to explicitly evaluate the integrand, substituting  $e = \exp(\mathcal{F})$  and (4.15) and (4.16) for  $\mathcal{F}$  and  $d\mathcal{F}$ . We get

$$\begin{aligned} \mathcal{S}_1^{(1)} &= \frac{1}{6(2-\kappa)^2} \int_{M_4 \times Q} \frac{\kappa}{2} \wedge \text{tr} \left( \hat{d}a \hat{d}a + \frac{1}{2} \hat{d}a^3 \right) \\ \mathcal{S}_1^{(2; \cdot)} &= \frac{1}{6(2-\kappa)^2} \int_{M_4 \times Q} \frac{\kappa}{2} \wedge \text{tr} \left( \hat{d}a \hat{d}a - d \hat{d}a a^0 + a^0 \hat{d}a \right) + d \hat{d}(\hat{d}a f + f \hat{d}a) \\ &\quad + \frac{1}{2} \hat{d}a^3 - \frac{1}{2} d(a^0 a^2 + a a^0 a + a^2 a^0) \\ &\quad + \frac{1}{2} d \hat{d}(a^2 f + a f a + f a^2); \end{aligned} \quad (4.23)$$

(Of course,  $\hat{d}a^3$  is shorthand for  $\hat{d}a a^2 - a \hat{d}a a + a^2 \hat{d}a$ , etc.)

To go further, we perform the Kaluza-Klein reduction. Each 1-form field  $a_n$  on  $M_4 \times Q$  becomes a 1-form field on  $M_4$  which we also denote by  $a_n$ , and a scalar field  $f_n$  on  $M_4$  for every harmonic 1-form  $\omega_1$  on  $Q$ , plus massive modes. The scalars  $f_n$  simply become scalars on  $M_4$  (again denoted  $f_n$ ), plus massive modes. Since  $\kappa \wedge \omega_1$  already is a 3-form on  $Q$ , the  $\omega_1$  cannot contribute in  $\mathcal{S}_1^{(2; \cdot)}$ , while in  $\mathcal{S}_1^{(1)}$  we must pick out the part linear in  $\omega_1$ . It is  $\text{tr} \left( \hat{d} \omega_1 \hat{d}a + \hat{d}a \omega_1 + \frac{1}{2} a^2 \omega_1 - \frac{1}{2} a \omega_1 a + \frac{1}{2} \omega_1 a^2 \right) \omega_1$ , where now  $\hat{d} \omega_1$  is the exterior derivative on  $M_4$ . Clearly, after integration over  $M_4$  this term vanishes:

$$\mathcal{S}_1^{(1)} = 0; \quad (4.24)$$

It remains to evaluate  $\mathcal{S}_1^{(2;)}$  with  $a_n$  and  $f_n$  now 1- and 0-forms on  $M_4$ . To see how to perform the integrals over  $Q$ , consider e.g. the terms that only involve two a-elds:

$$\begin{aligned}
 & \int_{M_4 \times Q} \frac{K}{2} \wedge \text{tr} \left( \hat{d}a \hat{d}a + d\hat{a} a^0 + a^0 \hat{d}a \right) \\
 = & \int_{M_4 \times Q} \frac{K}{2} \wedge \text{tr} \sum_{n,m=1}^{\infty} \left( da_n da_m + d\hat{a}_m a_n + d\hat{a}_n a_m \right) \wedge \dots \\
 = & \int_{M_4 \times U} \frac{K}{2} \wedge \text{tr} \sum_{n,m=1}^{\infty} \frac{1}{n+m+1} \left( da_n da_m + d\hat{a}_m a_n + d\hat{a}_n a_m \right) :
 \end{aligned} \tag{4.25}$$

At this point the  $Q$ -integral is reduced to a sum of integrals over  $M_4 \times U$  and now we can safely integrate by parts. The three terms then all are  $da_n da_m$  and the coefficients add up as  $\frac{1}{n+m+1} (1+m+n) = 1$ . Using (4.5) we get  $n \int_{M_4} \text{tr} dA dA$ , where now

$$A = \sum_{n=1}^{\infty} a_n \tag{4.26}$$

is the Kaluza-Klein reduction of the "bulk" eld  $\hat{A}_{\text{bulk}}$  encountered before in (4.11). Similarly one sees that the terms involving  $f$  do not contribute,<sup>10</sup> while the terms involving three a-elds add up to give  $n \int_{M_4} \text{tr} \frac{1}{2} dA^3$ . We conclude that

$$\mathcal{S}_1^{(2;)} = \frac{n}{6(2\pi)^2} \int_{M_4} \text{tr} dA dA + \frac{1}{2} A^3 : \tag{4.27}$$

Provided the  $n$  coincide with the number of charged chiral multiplets present at the singularity  $P$ , as suggested in [6], the non-invariance of the interaction (4.17) cancels the  $SU(N)^3$  anomaly locally, separately at each singularity.

Quite remarkably, the final result is simple with all contributions of the different  $a_n$  adding up to reproduce  $\frac{1}{4} \left( \sum_n a_n \right)^3 = \frac{1}{4} (A)^3$ . Alternatively, one might have first expanded  $\mathcal{S}_1$ . Then  $\mathcal{S}_1^{(2;)}$  would have reduced to an integral over  $M_4 \times U$  with the integral of  $\frac{K}{2}$  over  $U$  just giving  $n$ . The result would have been a four-dimensional action involving infinitely many elds  $a_n$  and  $f_n$ . While the gauge transformations of each term individually are complicated, we know that they sum up to give (4.27).

<sup>10</sup>Of course, we could have "integrated by parts" according to the rule (3.14) directly in  $\mathcal{S}_1^{(2;)}$  in (4.23), showing immediately that the  $f$ -elds do not contribute.

### 4.3 The $U(1)_i G^2$ anomaly

It remains to discuss the cancellation of the mixed  $U(1)_i G^2$  anomaly (4.3). Clearly, the variation of an interaction  $\int_{M_4} \mathbb{C} \wedge \text{tr} F^2$  can cancel the consistent anomaly (4.3) with  $\mathbb{C} = 1$ . However, following our general philosophy, we should really start with

$$\mathbb{S}_2 = \frac{T_2}{2(\ell_p)^2} \int_{M_4} \mathbb{C} \wedge \text{tr} F^2 : \quad (4.28)$$

Note that in the bulk this coincides with the standard interaction  $\int_{M_4} \mathbb{C} \wedge \text{tr} F^2$ . Since  $\text{tr} F^2$  was designed to be gauge invariant under (4.6) only  $\mathbb{C}$  contributes to the gauge variation of  $\mathbb{S}_2$ . Again, we write  $\mathbb{S}_2 = \mathbb{S}_2^{(1)} + \mathbb{S}_2^{(2; i)}$ . Inserting  $\mathbb{C} = d\hat{a} + \hat{a}^2$  from eq. (3.12) and  $F = \hat{d}a + a^2 + (\hat{d}f + af - fa - \hat{a})$  from (4.15) and (4.16) we get

$$\begin{aligned} \mathbb{S}_2^{(1)} &= \frac{T_2}{2(\ell_p)^2} \int_{M_4} d \text{tr} (\hat{d}a \hat{d}a + 2a^2 \hat{d}a) \\ \mathbb{S}_2^{(2; i)} &= \frac{T_2}{2(\ell_p)^2} \int_{M_4} \text{tr} (\hat{d}a \hat{d}a + 2a^2 \hat{d}a) \\ &\quad + 2d \text{tr} \hat{d}(f \hat{d}a + fa^2) - a^0 (\hat{d}a + a^2)^i : \quad (4.29) \end{aligned}$$

When we perform the Kaluza-Klein reduction,  $T_2 \rightarrow \int_{S^1} \hat{a}_i$  and  $a_n \rightarrow a_n + \frac{1}{n} \hat{a}_i$ , in  $\mathbb{S}_2^{(1)}$  only terms linear in  $\hat{a}_i$  can contribute, but they vanish after partial integration over  $M_4$ , just as for  $\mathbb{S}_1^{(1)}$ . Also as before, in  $\mathbb{S}_2^{(2; i)}$ ,  $\frac{1}{n} \hat{a}_i$  cannot contribute, while the terms containing  $f$  again vanish after partial integration over  $M_4$ . The remaining terms combine to yield

$$\mathbb{S}_2^{(2; i)} = \frac{1}{2(\ell_p)^2} \int_U \hat{a}_i \text{tr} (dA dA + 2A^2 dA) : \quad (4.30)$$

Provided

$$\int_U \hat{a}_i = \sum_{2\pi} q_i^i ; \quad (4.31)$$

this exactly cancels the mixed  $U(1)_i G^2$  anomaly locally. Note that the variation of  $\mathbb{S}_2^0 = \frac{T_2}{2(\ell_p)^2} \int_{M_4} \mathbb{C} \wedge \text{tr} F^2$  would have produced exactly the same result.

### 4.4 The mixed gauge-gravitational anomaly once more

Now we dispose of the necessary machinery to show that the variation of the modified Green-Schwarz term (3.25) with cut-off  $\hat{X}_8$  leads to the same local anomaly contribution as the variation of (3.26) using the ordinary  $X_8$ .

To begin with, we replace the spin connection  $\omega = \omega_0 + \omega$  by its cut-off version

$$\omega = \omega_0 + \omega \quad (4.32)$$

$\omega_0$  represents the fixed background  $R^4 \times X$  and the fluctuations. For the time being we make no assumption about  $\omega$ , but later on we will again restrict to fluctuations that preserve the product structure of the manifold. Again we write

$$\omega = \omega^{(0)} + \omega^{(1)} + \dots \quad (4.33)$$

with  $\omega^{(n)} = \sum_{n=1}^p \omega_n^{(n)}$  and  $\omega^{(n)} = \sum_{n=1}^p \omega_n^{(n)}$ . Of course,

$$\sum_{n=1}^p \omega_n^{(n)} = \omega \quad (4.34)$$

since in the bulk, where  $\omega = 1$  and  $\omega = 0$ , we want  $\omega$  to coincide with  $\omega_0$ .

We require that under a local Lorentz transformation with parameter  $\epsilon_L$  one has

$$\omega = d\epsilon_L + [\omega; \epsilon_L] \quad ; \quad \epsilon_L = \epsilon_L \quad (4.35)$$

This ensures that

$$\mathbb{R} = d\omega + \omega^2 \quad (4.36)$$

transforms covariantly:  $\mathbb{R} = [\mathbb{R}; \epsilon_L]$  and

$$\text{tr} \mathbb{R}^n = 0 \quad (4.37)$$

Comparing powers of  $\omega$  and  $\mathbb{R}$  in eq. (4.35) shows that the background is not transformed,  $\omega_0 = 0$ , as expected, and  $\omega_1 = D_0 \epsilon_L$ ,  $\omega_1 = \epsilon_L$  and, for  $n \geq 2$ ,

$\omega_n = [\omega_{n-1}; \epsilon_L]$ ,  $\omega_n = [\omega_{n-1}; \epsilon_L]$ , where  $D_0$  is the covariant derivative with  $\omega_0$ . Again we define  $\hat{\omega}^{(0)} = \sum_{n=1}^p \omega_n^{(n)}$  and  $\hat{\omega}^{(n)} = \sum_{n=1}^p (\omega_n^{(n)})$ , and idem for  $\hat{\mathbb{R}}$ . Then  $d\omega = \hat{d} \omega^{(0)} + \hat{d} \omega^{(1)}$  and for the curvature we find

$$\mathbb{R}(\omega) = \hat{\mathbb{R}}(\omega) + \dots \quad (4.38)$$

with

$$\begin{aligned} \hat{\mathbb{R}}(\omega) &= \mathbb{R}_0 + \hat{d} \omega^{(0)} + \omega^{(0)} + \omega^{(0)2} \\ \hat{\mathbb{R}}(\omega) &= \hat{d} \omega^{(1)} + [\omega^{(0)}; \omega^{(1)}] + \dots \end{aligned} \quad (4.39)$$



Two useful identities which follow from the Bianchi identity  $d\mathbb{R} = [\mathbb{R}; \epsilon]$  are

$$\begin{aligned} \hat{d}\hat{R} &= [\hat{R}; \epsilon]_0 + \dots \\ \hat{d} &= (\dots) (\dots) + [\hat{R}; \epsilon] \hat{d}^0 (\dots) (\dots) (\dots) : (4.40) \end{aligned}$$

Finally, the cut-off gravitational 8-form  $\mathbb{X}_8$  then is given by

$$\begin{aligned} \mathbb{X}_8 &= \frac{1}{192(2)^3} \text{tr}\hat{R}^4 - \frac{1}{4} (\text{tr}\hat{R}^2)^2 \\ &= \frac{1}{192(2)^3} \text{tr}\hat{R}^4 + 4 \text{tr}\hat{R}^3 - \frac{1}{4} (\text{tr}\hat{R}^2)^2 - \text{tr}\hat{R}^2 \text{tr}\hat{R} : (4.41) \end{aligned}$$

By construction,  $\mathbb{X}_8$ , as well as each of the four terms individually, is invariant under local Lorentz transformations. Hence, using  $\mathbb{C} = C + B$  and  $\epsilon = \epsilon^P$ , we find

$$\begin{aligned} \mathbb{S}_{GS} &= \mathbb{S}_{GS}^{(1)} + \mathbb{S}_{GS}^{(2;\epsilon)} \\ \mathbb{S}_{GS}^{(1)} &= \frac{T_2}{192(2)^4} \int_C \text{tr}\hat{R}^4 - \frac{1}{4} (\text{tr}\hat{R}^2)^2 \\ \mathbb{S}_{GS}^{(2;\epsilon)} &= \frac{T_2}{192(2)^4} \int_B \text{tr}\hat{R}^4 + 4C \text{tr}\hat{R}^3 - \frac{1}{4} B (\text{tr}\hat{R}^2)^2 - C \text{tr}\hat{R}^2 \text{tr}\hat{R} : (4.42) \end{aligned}$$

While  $\mathbb{S}_{GS}^{(1)}$  involves an integral over all of  $M_4 \times X$ , each  $\mathbb{S}_{GS}^{(2;\epsilon)}$  reduces to an integral over  $M_4 \times Y$ . We could perfectly well do this reduction first and then compute the gauge variation of each  $\mathbb{S}_{GS}^{(2;\epsilon)}$ . However, as for the Yang-Mills case, the computations are more compact if we first take the variation and then evaluate the integral. Using  $C = d$ ,  $B = \dots$  and the invariance of the gravitational terms we find

$$\begin{aligned} \mathbb{S}_{GS}^{(1)} &= \frac{T_2}{192(2)^4} \int d \text{tr}\hat{R}^4 - \frac{1}{4} (\text{tr}\hat{R}^2)^2 \\ \mathbb{S}_{GS}^{(2;\epsilon)} &= \frac{T_2}{192(2)^4} \int \text{tr}\hat{R}^4 + 4d \text{tr}\hat{R}^3 - \frac{1}{4} (\text{tr}\hat{R}^2)^2 - d \text{tr}\hat{R}^2 \text{tr}\hat{R} : (4.44) \end{aligned}$$

We will discuss  $\mathbb{S}_{GS}^{(1)}$  later on. As familiar by now, thanks to the presence of  $\epsilon$ , each  $\mathbb{S}_{GS}^{(2;\epsilon)}$  is reduced to an integral over  $M_4 \times Y$  with every  $\epsilon^k$  contributing a factor  $\frac{1}{k+1}$ . On  $M_4 \times Y$ , the derivative  $\hat{d}$  acts as an ordinary derivative  $d$ , and we are allowed to integrate by parts. As explained in eq. (3.14) above, it is easy to see that exactly

the same result is obtained if one first replaces  $d$  by  $\hat{d}$  and integrates all  $\hat{d}$  by parts directly in (4.44), remembering that  $\hat{d}$  only acts on the  $\omega_0, \omega_n$  and  $\omega_{n+1}$ , but not on the  $\omega_n$ . Using this observation, we rewrite

$$\mathbb{S}_{GS}^{(2;)} = \frac{T_2}{192(2)^4} \int \text{tr} \hat{R}^4 - 4 \hat{d}(\text{tr} \hat{R}^3) - \frac{1}{4} (\text{tr} \hat{R}^2)^2 + \hat{d}(\text{tr} \hat{R}^2 \text{tr} \hat{R}) \quad ; \quad (4.45)$$

Now use the identities (4.40) to show that  $\hat{d} \text{tr} \hat{R}^2 = 0$  and

$$\hat{d}(\text{tr} \hat{R}^1) = \text{tr} \hat{R}^1 \hat{d} \omega_0 + (\omega_0 + \omega_n) \omega_0 + \omega_0 (\omega_0 + \omega_n) = \frac{1}{1+1} \frac{\partial}{\partial} \text{tr} \hat{R}^{1+1} \quad (4.46)$$

so that

$$\mathbb{S}_{GS}^{(2;)} = \frac{T_2}{192(2)^4} \int \left( 1 + \frac{\partial}{\partial} \right) \text{tr} \hat{R}^4 - \frac{1}{4} (\text{tr} \hat{R}^2)^2 \quad ; \quad (4.47)$$

Next, we expand the integrand in powers of  $\omega_n$ . Writing  $\hat{R} = \sum_{n=0}^P \hat{R}_n \omega_n$  with  $\hat{R}_0 = R_0$  and  $\hat{R}_n = \hat{d} \omega_n + \omega_0 \omega_n + \omega_n \omega_0 + \sum_{r=1}^P \omega_r \omega_{n-r}$  for  $n \geq 1$ , we see that the 1 in the parenthesis in (4.47) contributes a factor  $\frac{1}{n+m+k+1+1}$  to the integral over  $r$  while the  $\frac{\partial}{\partial}$  contributes a factor  $\frac{n+m+k+1}{n+m+k+1+1}$ , both adding up to 1. As a result, we get

$$\mathbb{S}_{GS}^{(2;)} = \frac{T_2}{192(2)^4} \int_{M_4 \times Y} \text{tr} R^4 - \frac{1}{4} (\text{tr} R^2)^2 - \frac{T_2}{2} \int_{M_4 \times Y} X_8(R) \quad ; \quad (4.48)$$

where now

$$R = \sum_{n=0}^{\hat{X}} \hat{R}_n \quad ; \quad (4.49)$$

Clearly,  $R$  is the value of the curvature "in the bulk" of  $M_4 \times X$  (or its appropriate pullback onto  $M_4 \times Y$ ), and corresponds to the fluctuating geometry  $\omega = \omega_0 + \omega_n$  without cutting off anything. Hence we see that, in the end, this rather sophisticated treatment reproduces the same result as the more naive  $\mathbb{S}_{GS}^0 = \frac{T_2}{2} \int \mathbb{C} \wedge X_8$  of eq (3.26). It is clear from our analysis that this same simplification occurs for any invariant quantity made from combinations of  $\text{tr} R^1$  or  $\text{tr} R^k$ .

It remains to discuss  $\mathbb{S}_{GS}^{(1)}$ . With  $\omega = \omega_i \omega_i$ , the integral will be non-vanishing only if  $\text{tr} \hat{R}^4 - \frac{1}{4} (\text{tr} \hat{R}^2)^2$  is a 3-form on  $M_4$  and a 5-form on  $X$ . If the fluctuations of the metric preserve the product structure  $M_4 \times X$ , this is clearly impossible, and we conclude

$$\mathbb{S}_{GS}^{(1)} = 0 \quad ; \quad (4.50)$$

For more general fluctuations, however, it is less clear what happens and we will not pursue this issue further.

## 5 Conclusions

We have reconsidered the anomaly cancellation mechanism on  $G_2$ -holonomy manifolds with conical singularities, first outlined in [6]. It turned out that we needed to modify the eleven-dimensional Chern-Simons and Green-Schwarz terms, and similarly the interactions  $S_1$  and  $S_2$  present on ADE singularities, by (smoothly) cutting off the fields close to the conical singularities. This induces anomalous variations of the cut-off 3-form field  $\mathbb{C}$  and of the cut-off non-abelian gauge field  $\mathbb{F}$ . These anomalous variations are localized in the regions close to the conical singularities where the cut-off is done. This implies that the corresponding non-invariance of the action is also localized there and we get one  $S^{(i)}$  term for each conical singularity  $P_i$ . Each of these terms then exactly cancels the various anomalies that are present at these singularities due to the charged chiral fermions living there. Thus anomaly cancellation indeed occurs locally, i.e. separately for each conical singularity. For the mixed gauge-gravitational anomaly we met a surprise. The inflow terms would cancel the anomaly but for their sign. We argued that further gravitational correction terms must be present to achieve complete local cancellation of these mixed anomalies.

Throughout the whole discussion it is always assumed that the  $G_2$ -holonomy manifold is compact, although the explicit examples of conical singularities are actually taken from the known non-compact  $G_2$ -holonomy manifolds, assuming that conical singularities on compact  $G_2$ -manifolds have the same structure. As mentioned in the introduction, there exist close relatives of  $G_2$ -holonomy manifolds which are weak  $G_2$ -holonomy manifolds. In this case, it is quite easy to construct compact examples with conical singularities and explicitly known metrics. This is done in [9] and will be briefly recalled in the appendix. The conical singularities are exactly as assumed in the present paper, namely for  $r \neq 0$  they are cones on some  $Y$  with the same  $Y$ 's as considered here. This implies that the whole discussion of chiral fermions present at the singularities and of the anomaly cancellation of the present paper directly carries over to these weak  $G_2$ -holonomy manifolds.

## Acknowledgements

Stefan Metzger gratefully acknowledges support by the Gottlieb Daimler- und Karl Benz-Stiftung. We would like to thank Luis Alvarez-Gaume, Jean-Pierre D'Erendinger, Jean Iliopoulos, Ruben Minasian, Ivo Sachs, Julius Wess and Jean Zinn-Justin for helpful discussions.

## 6 Appendix

Here we will briefly recall the geometry of the singular weak  $G_2$ -holonomy manifolds  $X$  constructed in [9]. Although they have weak  $G_2$ -holonomy rather than  $G_2$ -holonomy, they are the prototype of the compact manifolds with conical singularities one has in mind throughout the present paper.

In [9] it was shown that for every non-compact  $G_2$ -manifold that is asymptotic (for large  $r$ ) to a cone on  $Y$ , there is an associated compact weak  $G_2$ -manifold with its metric given by

$$ds_X^2 = dr^2 + R \sin^2 \frac{r}{R} ds_Y^2 ; \quad 0 \leq r \leq R ; \quad (\text{A } 1)$$

It has two conical singularities. The first one, at  $r = 0$ , is a cone on  $Y$ , while the second one, at  $r = R$ , is a cone on  $\bar{Y}$ . Here  $\bar{Y}$  equals  $Y$  but with its orientation reversed. This reversal of orientation simply occurs since we define  $\bar{Y}$  always such that the normal vector points away from the singularity. Hence:

$$Y_1 = Y ; \quad Y_2 = \bar{Y} ; \quad (\text{A } 2)$$

For these examples we have all the necessary global information, and it was shown in [9] that the square-integrable harmonic  $p$ -forms on  $X$ , for  $p \geq 3$ , are the trivial extensions of the square-integrable harmonic  $p$ -forms on  $Y$ . In particular, we have  $b^1(X) = b^1(Y) = 0$ ,  $b^2(X) = b^2(Y)$  and  $b^3(X) = b^3(Y)$ .

According to the general cut-off procedure described in section 3.1, for these examples one introduces two local coordinates  $r_1 = r$  and  $r_2 = R - r$ . It follows that

$= \omega_1 + \omega_2$ , where, in the limit of vanishing regularisation,

$$\begin{aligned} \omega_1 &= \int_{M_4 \times Y} (\eta_1 - \epsilon) d\eta_1 = \int_{M_4 \times Y} (r - \epsilon) dr \\ \omega_2 &= \int_{M_4 \times Y} (\eta_2 - \epsilon) d\eta_2 = \int_{M_4 \times Y} (r - (R - \epsilon)) dr : \end{aligned} \quad (\text{A } 3)$$

Then for a smooth 10-form one has

$$\int_{M_4 \times X} Z = \int_{M_4 \times Y} Z \Big|_{r=\epsilon} - \int_{M_4 \times Y} Z \Big|_{r=R-\epsilon} = \int_{M_4 \times Y_1} Z + \int_{M_4 \times Y_2} Z : \quad (\text{A } 4)$$

## References

- [1] G. Papadopoulos and P.K. Townsend, Compactification of D = 11 supergravity, Phys. Lett. B 357 (1995) 300, hep-th/9506159.
- [2] B.S. Acharya, M-theory, Joyce orbifolds and super Yang-Mills, Adv. Theor. Math. Phys. 3, 227, hep-th/9812205.
- [3] M. Cvetič, G. Shiu and A.M. Uranga, Chiral Four-Dimensional N = 1 Supersymmetric Type IIA Orientifolds from Intersecting D 6-Branes, Nucl. Phys. B 615 (2001) 3, hep-th/0107166.
- [4] M. Atiyah and E. Witten, M-theory dynamics on a manifold with G<sub>2</sub>-holonomy, Adv. Theor. Math. Phys. 6 (2003) 1, hep-th/0107177.
- [5] B.S. Acharya and E. Witten, Chiral Fermions from Manifolds of G<sub>2</sub> Holonomy, hep-th/0109152.
- [6] E. Witten, Anomaly cancellation on G<sub>2</sub>-manifolds, hep-th/0108165.
- [7] R. Bryant and S. Salamon, On the construction of complete metrics with exceptional holonomy, Duke Math. Journal 58 (1989) 829;  
G.W. Gibbons, D.N. Page and C.N. Pope, Einstein metrics on S<sup>3</sup>; R<sup>3</sup> and R<sup>4</sup> bundles, Comm. Math. Phys. 127 (1990) 529.

- [8] D. D. Joyce, Compact Riemannian 7-manifolds with holonomy  $G_2$  I, *J. Di. Geom.* 43 (1996) 291, and idem II, *J. Di. Geom.* 43 (1996) 329.
- [9] A. Bilal and S. Metzger, Compact weak  $G_2$ -manifolds with conical singularities, [hep-th/0302021](#).
- [10] D. Freed, J.A. Harvey, R. Minasian and G. Moore, Gravitational anomaly cancellation for M-theory vebranes, *Adv. Theor. Math. Phys.* 2 (1998) 31, [hep-th/9803205](#).
- [11] W. A. Bardeen and B. Zumino, Consistent and covariant anomalies in gauge and gravitational theories, *Nucl. Phys. B* 244 (1984) 421.
- [12] J. Wess and B. Zumino, Consequences of Anomalous Ward Identities, *Phys. Lett. B* 37 (1971) 95.
- [13] C. G. Callan and J.A. Harvey, Anomalies and Fermion Zero Modes on Strings and Domain Walls, *Nucl. Phys. B* 250 (1985) 427.
- [14] S. G. Nakulich, Axionic strings, *Nucl. Phys. B* 296 (1988) 937.
- [15] J.A. Harvey and O. Ruchayskiy, The local Structure of Anomaly Inflow, *JHEP* 0106, 044 (2001), [hep-th/0007037](#).