VARIATIONAL DERIVATION OF RELATIVISTIC FERMION-ANTIFERMION WAVE EQUATIONS IN QED

Andrei G. Terekidi and Jurij W. Darewych

Department of Physics and Astronomy, York University, Toronto, Ontario, M3J 1P3, Canada

Abstract We present a variational method for deriving relativistic two-fermion wave equations in a Hamiltonian formulation of QED. A reformulation of QED is performed, in which covariant Green functions are used to solve for the electromagnetic field in terms of the fermion fields. The resulting modified Hamiltonian contains the photon propagator directly. The reformulation permits one to use a simple Fock-space variational trial state to derive relativistic fermion-antifermion wave equations from the corresponding quantum field theory. We verify that the energy eigenvalues obtained from the wave equation agree with known results for positronium.

1. Introduction

The description of relativistic bound and quasi-bound (i.e. unstable) few body systems continues to be an active area of research. The traditional method of treating relativistic bound states in quantum field theory (QFT) is by means of the Bethe-Salpeter (BS) equation. However, this approach has a number of difficulties, including the appearance of relative-time coordinates and negative-energy solutions. In practice, the interaction kernels (potentials) in the BS equation are obtained from covariant perturbation theory, which may be of questionable validity for strongly coupled systems. In addition, the BS formalism is difficult to implement for systems of more than two particles.

An alternative approach might be the variational method, which is non-perturbative in principle. The variational method has not been widely used in quantum field theory, in contrast to non-relativistic systems describable by the Schrödinger theory, in part because of the difficulty of constructing realistic yet tractable trial states.

It has been pointed out in previous publications [1,2] that various models in QFT, including QED, can be reformulated, using mediating-field Green functions, into a form particularly convenient for variational calculations. This approach was applied recently to the study of relativistic two-body states in the scalar Yukawa (Wick-Cutkosky) theory [3,4,5]. In the present paper we shall implement this approach to the realistic QED theory, where comparison with experimentally verified results are possible. In particular, we shall use the reformulated QED Hamiltonian to derive a relativistic fermion-antifermion wave equation and discuss its solution.

The reformulation of the QED is presented in section 2, while the Hamiltonian and equal time quantization are given in section 3. In section 4 we use the variational principle, with simple Fockspace trial states to derive the relativistic fermion-antifermion equations, and present their "partial wave" decomposition for all possible J^{PC} states. The relativistic radial equations are presented in

section 5, while their non-relativistic and semi relativistic limits are given in 6. In section 7 the energy eigenvalues are shown to yield the correct the fine and hyperfine structure for all states. Concluding remarks are given in section 8.

2. Reformulation of field equations and Lagrangian

The Lagrangian of QED is $(\hbar = c = 1)$

$$
\mathcal{L} = \overline{\psi}(x) \left(i \gamma^{\mu} \partial_{\mu} - m - e \gamma^{\mu} A_{\mu}(x) \right) \psi(x) \n- \frac{1}{4} \left(\partial_{\alpha} A_{\beta}(x) - \partial_{\beta} A_{\alpha}(x) \right) \left(\partial^{\alpha} A^{\beta}(x) - \partial^{\beta} A^{\alpha}(x) \right).
$$
\n(1)

The corresponding Euler-Lagrange equations of motion are the coupled Dirac-Maxwell equations,

$$
(i\gamma^{\mu}\partial_{\mu} - m)\psi(x) = e\gamma^{\mu}A_{\mu}(x)\psi(x), \qquad (2)
$$

and

$$
\partial_{\mu}\partial^{\mu}A^{\nu}(x) - \partial^{\nu}\partial_{\mu}A^{\mu}(x) = j^{\nu}(x),\tag{3}
$$

where

$$
j^{\nu}(x) = e\overline{\psi}(x)\gamma^{\nu}\psi(x). \tag{4}
$$

The equations $(2)-(3)$ can be decoupled in part by using the well-known formal solution [6,7] of the Maxwell equation (3), namely

$$
A_{\mu}(x) = A_{\mu}^{0}(x) + \int d^{4}x' D_{\mu\nu}(x - x')j^{\nu}(x'), \qquad (5)
$$

where $D_{\mu\nu}(x-x')$ is a Green function (or photon propagator in QFT terminology), defined by

$$
\partial_{\alpha}\partial^{\alpha}D_{\mu\nu}(x-x') - \partial_{\mu}\partial^{\alpha}D_{\alpha\nu}(x-x') = g_{\mu\nu}\delta^{4}(x-x'),\tag{6}
$$

and $A^0_\mu(x)$ is a solution of the homogeneous (or "free field") equation (3) with $j^\mu(x) = 0$.

We recall, in passing, that equation (6) does not define the covariant Green function $D_{\mu\nu}(x-x')$ uniquely. For one thing, one can always add a solution of the homogeneous equation (eq. (6) with $g_{\mu\nu} \to 0$). This allows for a certain freedom in the choice of $D_{\mu\nu}$, as is discussed in standard texts (e.g. ref. $[6,7]$). In practice, the solution of eq. (6) , like that of eq. (3) , requires a choice of gauge. However, we do not need to specify one at this stage.

Substitution of the formal solution (5) into equation (2) yields the "partly reduced" equations,

$$
\left(i\gamma^{\mu}\partial_{\mu} - m\right)\psi(x) = e\gamma^{\mu}\left(A_{\mu}^{0}(x) + \int d^{4}x' D_{\mu\nu}(x - x')j^{\nu}(x')\right)\psi(x),\tag{7}
$$

which is a nonlinear Dirac equation. To our knowledge no exact (analytic or numeric) solution of equation (7) for classical fields have been reported in the literature. However, approximate solutions have been discussed by various authors, particularly Barut and his co-workers (see ref. [8,9] and citations therein). In any case, our interest here is in the quantized field theory.

The partially reduced equation (7) is derivable from the stationary action principle

$$
\delta S\left[\psi\right] = \delta \int d^4x \mathcal{L}_R = 0\tag{8}
$$

with the Lagrangian density

$$
\mathcal{L}_R = \overline{\psi}(x) \left(i \gamma^\mu \partial_\mu - m - e \gamma_\mu A_0^\mu(x) \right) \psi(x) - \frac{1}{2} \int d^4 x' j^\mu(x') D_{\mu\nu}(x - x') j^\nu(x) \tag{9}
$$

provided that the Green function is symmetric in the sense that

$$
D_{\mu\nu}(x - x') = D_{\mu\nu}(x' - x) \quad \text{and} \quad D_{\mu\nu}(x - x') = D_{\nu\mu}(x - x'). \tag{10}
$$

One can proceed to do conventional covariant perturbation theory using the reformulated QED Lagrangian (9). The interaction part of (9) has a somewhat modified structure from that of the usual formulation of QED. Thus, there are two interaction terms. The last term of (9) is a "current-current" interaction which contains the photon propagator sandwiched between the fermionic currents. As such, it corresponds to Feynman diagrams without external photon lines. The term containing A_0^{μ} $\frac{\mu}{0}$ corresponds to diagrams that cannot be generated by the term containing $D_{\mu\nu}$, including diagrams involving external photon lines (care would have to be taken not to double count physical effects). However, we shall not pursue covariant perturbation theory in this work. Rather, we shall consider a variational approach that allows one to derive relativistic few-fermion equations, and to study their bound and scattering solutions.

3. Hamiltonian of the quantized theory in the equal-time formalism

We consider this theory in the quantized, equal-time formalism. To this end we write down the Hamiltonian density corresponding to the Lagrangian (9), with the term for the free A_0^{μ} $\frac{\mu}{0}(x)$ field suppressed since it will not contribute to the results presented in this paper. The relevant expression is:

$$
\mathcal{H}_R = \mathcal{H}_0 + \mathcal{H}_I,\tag{11}
$$

where

$$
\mathcal{H}_0 = \psi^{\dagger}(x) \left(-i \overrightarrow{\alpha} \cdot \nabla + m\beta \right) \psi(x), \tag{12}
$$

$$
\mathcal{H}_I = \frac{1}{2} \int d^4 x' j^\mu(x') D_{\mu\nu}(x - x') j^\nu(x), \tag{13}
$$

We construct a quantized theory by the imposition of anticommutation rules for the fermion fields, namely

$$
\left\{\psi_{\alpha}(\mathbf{x},t),\psi_{\beta}^{\dagger}(\mathbf{y},t)\right\} = \delta_{\alpha\beta}\delta^{3}\left(\mathbf{x}-\mathbf{y}\right),\tag{14}
$$

while all other vanish. In addition, if A_0^{μ} $\mu_0^{\mu} \neq 0$, there would be the usual commutation rules for the A_0^{μ} $_0^{\mu}$ field, and commutation of the A_0^{μ} $\frac{\mu}{0}$ field operators with the ψ field operators.

To specify our notation, we quote the usual Fourier decomposition of the field operators, namely

$$
\psi(x) = \sum_{s} \int \frac{d^3 p}{\left(2\pi\right)^{3/2}} \left(\frac{m}{\omega_p}\right)^{1/2} \left[b_{\mathbf{p}s} u\left(\mathbf{p}, s\right) e^{-ip\cdot x} + d_{\mathbf{p}s}^{\dagger} v\left(\mathbf{p}, s\right) e^{ip\cdot x}\right],\tag{15}
$$

with $p = p^{\mu} = (\omega_p, \mathbf{p})$, and $\omega_p = \sqrt{m^2 + \mathbf{p}^2}$. Dirac spinors u and v for free particles of mass m, where $(\gamma^{\mu}p_{\mu}-m)u(\mathbf{p},s)=0$, $(\gamma^{\mu}p_{\mu}+m)v(\mathbf{p},s)=0$, are normalized such that

$$
u^{\dagger}(\mathbf{p},s) u(\mathbf{p},\sigma) = v^{\dagger}(\mathbf{p},s) v(\mathbf{p},\sigma) = \frac{\omega_p}{m} \delta_{s\sigma}, \qquad (16)
$$

$$
\dagger(\mathbf{p},s)v(\mathbf{p},\sigma) = v^{\dagger}(\mathbf{p},s)u(\mathbf{p},\sigma) = 0.
$$
 (17)

The creation and annihilation operators b^{\dagger} , b of the (free) fermions of mass m, and d^{\dagger} , d for the corresponding antifermions, satisfy the usual anticommutation relations. The non-vanishing ones are

$$
\left\{b_{\mathbf{p}s}, b_{\mathbf{q}\sigma}^{\dagger}\right\} = \left\{d_{\mathbf{p}s}, d_{\mathbf{q}\sigma}^{\dagger}\right\} = \delta_{s\sigma}\delta^3\left(\mathbf{p} - \mathbf{q}\right). \tag{18}
$$

4. Variational principle and fermion-antifermion trial states

u

Unfortunately we do not know how to obtain exact eigenstates of the Hamiltonian (11). Therefore we shall resort to a variational approximation, based on the variational principle

$$
\delta \left\langle \psi \right| \hat{H} - E \left| \psi \right\rangle_{t=0} = 0. \tag{19}
$$

For a fermion-antifermion system , the simplest Fock-space trial state that can be written down in the rest frame is

$$
|\psi_T\rangle = \sum_{s_1 s_2} \int d^3 \mathbf{p} F_{s_1 s_2}(\mathbf{p}) b_{\mathbf{p} s_1}^{\dagger} d_{-\mathbf{p} s_2}^{\dagger} |0\rangle ,
$$
 (20)

where $F_{s_1s_2}$ are four adjustable functions. We use this trial state to evaluate the matrix elements needed to implement the variational principle (19), namely

$$
\langle \psi_T | : \hat{H}_0 - E : | \psi_T \rangle = \sum_{s_1 s_2} \int d^3 \mathbf{p} F_{s_1 s_2}^*(\mathbf{p}) F_{s_1 s_2}(\mathbf{p}) (2\omega_p - E)
$$
 (21)

and

$$
\langle \psi_T | : \hat{H}_I : | \psi_T \rangle = \frac{e^2 m^2}{(2\pi)^3} \sum_{s_1's_2's_1s_2} \int \frac{d^3 \mathbf{p} d^3 \mathbf{p}'}{\omega_p \omega_{p'}} F_{s_1s_2}^*(\mathbf{p}) F_{s_1's_2'}(\mathbf{p}') \times
$$

$$
\times \left(\begin{array}{c} -\overline{u} \left(\mathbf{p}, s_1 \right) \gamma^{\mu} u \left(\mathbf{p}', s_1' \right) D_{\mu\nu} (p - p') \overline{v} \left(-\mathbf{p}', s_2' \right) \gamma^{\nu} v \left(-\mathbf{p}, s_2 \right) \\ +\overline{u} \left(\mathbf{p}, s_1 \right) \gamma^{\mu} v \left(-\mathbf{p}, s_2 \right) D_{\mu\nu} (p + p') \overline{v} \left(-\mathbf{p}', s_2' \right) \gamma^{\nu} u \left(\mathbf{p}', s_1' \right) \end{array} \right), \tag{22}
$$

where $p = (\omega_p, \mathbf{p}), p' = (\omega_{p'}, \mathbf{p'}),$ with $\mathbf{p} + \mathbf{p'} = 0$ (i.e. $p + p' = (2\omega_p, 0)$) in the rest frame, and

$$
D_{\mu\nu}(x - x') = \int \frac{d^4k}{(2\pi)^4} D_{\mu\nu}(k) e^{-ik \cdot (x - x')}.
$$
 (23)

We have normal-order the entire Hamiltonian, since this circumvents the need for mass renormalization which would otherwise arise. Not that there is difficulty with handling mass renormalization in the present formalism (as shown in various earlier papers; see, for example, ref. [10] and citations therein). It is simply that we are not interested in mass renormalization here, since it has no effect on the two-body bound state energies that we obtain in this paper. Furthermore, the approximate trial state (20), which we use in this work, is incapable of sampling loop effects. Thus, the normal-ordering of the entire Hamiltonian does not "sweep under the carpet" loop effects, since none arise at the present level of approximation, that is with the trial state $|\psi_T\rangle$ specified in eq. (20) .

The variational principle (19) leads to the following equation

$$
\sum_{s_1s_2} \int d^3 \mathbf{p} \left(2\omega_p - E\right) F_{s_1s_2}(\mathbf{p}) \delta F_{s_1s_2}^*(\mathbf{p})
$$

$$
-\frac{m^2}{\left(2\pi\right)^3} \sum_{\sigma_1 \sigma_2 s_1 s_2} \int \frac{d^3 \mathbf{p} d^3 \mathbf{q}}{\omega_p \omega_q} F_{\sigma_1 \sigma_2}(\mathbf{q}) \left(-i\right) \mathcal{M}_{s_1s_2\sigma_1 \sigma_2}(\mathbf{p}, \mathbf{q}) \delta F_{s_1s_2}^*(\mathbf{p}) = 0, \tag{24}
$$

where $\mathcal{M}_{s_1s_2\sigma_1\sigma_2}(\mathbf{p},\mathbf{q})$ is an invariant "matrix element", which contains two terms:

$$
\mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} \left(\mathbf{p}, \mathbf{q} \right) = \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2}^{ope} \left(\mathbf{p}, \mathbf{q} \right) + \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2}^{ann} \left(\mathbf{p}, \mathbf{q} \right), \tag{25}
$$

where

$$
\mathcal{M}_{s_1s_2\sigma_1\sigma_2}^{ope}(\mathbf{p},\mathbf{q}) = -\overline{u}(\mathbf{p},s_1) \left(-ie\gamma^{\mu}\right) u(\mathbf{q},\sigma_1) i D_{\mu\nu}(p-q) \overline{v}(-\mathbf{q},\sigma_2) \left(-ie\gamma^{\nu}\right) v(-\mathbf{p},s_2),\tag{26}
$$

$$
\mathcal{M}_{s_1s_2\sigma_1\sigma_2}^{ann}(\mathbf{p},\mathbf{q}) = \overline{u}(\mathbf{p},s_1) \left(-ie\gamma^{\mu}\right)v\left(-\mathbf{p},s_2\right)iD_{\mu\nu}\left(p+q\right)\overline{v}\left(-\mathbf{q},\sigma_2\right)\left(-ie\gamma^{\nu}\right)u\left(\mathbf{q},\sigma_1\right),\tag{27}
$$

correspond to the usual one-photon exchange and virtual annihilation Feynman diagrams.

At this point it is worthwhile to make a few comments about our equation (24) and to compare its general features with other two-fermion equations, particularly field-theory based approaches. Firstly we note that the present variational derivation leads to momentum-space Salpeter-like equations, with at most four independent components $F_{s_1s_2}(\mathbf{p})$. The equations have only positiveenergy solutions, as is evident from eq. (24) with the interaction turned off, in which case only $E = 2\omega_p > 0$ is obtained. This is in contrast to the BS equation, which is a 16-component equation and contains both positive, negative and mixed energy solutions.

The interaction kernels, represented by the covariant M -matrices, result from the variational derivation, that is, they are not put in by hands. This is in contrast to two fermion equations, which are not derived from a underlying quantum field theory, such as various two-body generalizations of the one-body Dirac equation. There are many such equations on the market, for example the eight component two-fermion equation of Pilkuhn [11]. In these treatments QFT effects, such as the virtual annihilation interaction (eq. (27)) do not arise naturally but need to be added in.

The fact that only the lowest order ("tree level") diagrams appear in our eq. (24) is a reflection of the fact that we have used the simplest possible variational ansatz (20). Even so, it is important to note that, because of the reformulation discussed in section 2 and 3, their derivation does not require additional Fock-space terms in the variational state (20) as is the case in traditional (nonreformulated) treatments (e.g. [12]-[14]).

In the non-relativistic limit, the functions $F_{s_1s_2}$ can be written as

$$
F_{s_1s_2}(\mathbf{p}) = F(\mathbf{p})\Lambda_{s_1s_2},\tag{28}
$$

where the non-zero elements of Λ_{ij} for total spin singlet $(S = 0)$ states are $\Lambda_{12} = -\Lambda_{21} = \frac{1}{\sqrt{2}}$ $\frac{1}{2}$, while for the spin triplet $(S = 1)$ states the non-zero elements are $\Lambda_{11} = 1$ for $m_s = +1, \Lambda_{12} = \Lambda_{21} = \frac{1}{\sqrt{2}}$ 2 for $m_s = 0$, and $\Lambda_{22} = 1$ for $m_s = -1$. We use the notation that the subscripts 1 and 2 of Λ correspond to $m_s = 1/2$ and $m_s = -1/2$ (or \uparrow and \downarrow) respectively. Substituting (28) into (24), the variational procedure, after multiplying the result by $\Lambda_{s_1s_2}$ and summing over s_1 and s_2 , gives the equation

$$
(2\omega_p - E)F(\mathbf{p}) = \frac{1}{(2\pi)^3} \int d^3 \mathbf{q} \mathcal{K}(\mathbf{p}, \mathbf{q}) F(\mathbf{q}),
$$
\n(29)

where

$$
\mathcal{K}(\mathbf{p}, \mathbf{q}) = -i \frac{m^2}{\omega_p \omega_q} \sum_{s_1 s_2 \sigma_1 \sigma_2} \Lambda_{s_1 s_2} \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{p}, \mathbf{q}) \Lambda_{\sigma_1 \sigma_2}.
$$
 (30)

To lowest-order in $|p|/m$ (i.e. in the non-relativistic limit), the kernel (30) reduces to $K =$ $e^2/|\mathbf{p}-\mathbf{q}|^2$, and so (29) reduces to the (momentum-space) Schrödinger equation

$$
\left(\frac{\mathbf{p}^2}{2\mu} - \varepsilon\right) F(\mathbf{p}) = \frac{1}{(2\pi)^3} \int d^3 \mathbf{q} \frac{e^2}{|\mathbf{p} - \mathbf{q}|^2} F(\mathbf{q}),\tag{31}
$$

where $\varepsilon = E - 2m$ and $\mu = m/2$. This verifies that the relativistic two-fermion equation (24) has the expected non-relativistic limit.

In the relativistic case we do not complete the variational procedure in (24) at this stage to get equations for the four adjustable functions $F_{s_1s_2}$, because they are not independent in general. Indeed we require that the trial state be an eigenstate of the total angular momentum operator (in relativistic form), its projection, parity and charge conjugation, namely that

$$
\begin{bmatrix}\n\hat{\mathbf{J}}^2 \\
\hat{J}_3 \\
\hat{\mathcal{P}} \\
\hat{C}\n\end{bmatrix} |\psi_T\rangle = \begin{bmatrix}\nJ(J+1) \\
m_J \\
P \\
C\n\end{bmatrix} |\psi_T\rangle, \qquad (32)
$$

where $m_j = J, J-1, ..., -J$ as usual. We present explicit forms for the operators $\hat{\mathbf{J}}^2, \hat{J}_3$ in Appendix A. The form for $\hat{\mathbf{J}}^2$, eq. (109), in particular, is not readily found in standard texts and reference books.

The functions $F_{s_1s_2}(\mathbf{p})$ can be written in the general form

$$
F_{s_1s_2}(\mathbf{p}) = \sum_{\ell_{s_1s_2}} \sum_{m_{s_1s_2}} f_{s_1s_2}^{\ell_{s_1s_2}m_{s_1s_2}}(p) Y_{\ell_{s_1s_2}}^{m_{s_1s_2}}(\hat{\mathbf{p}}),
$$
\n(33)

where $Y_{\ell_{\text{max}}}^{m_{s_1s_2}}$ $\zeta_{s_1s_2}^{ms_1s_2}(\hat{\bf p})$ are the usual spherical harmonics. Here and henceforth we will use the notation $p = |\mathbf{p}|$ etc. (four-vectors will be written as p^{μ}). The orbital indexes $\ell_{s_1s_2}$ and $m_{s_1s_2}$ depend on the spin indexes s_1 and s_2 and are specified by equations (32). The radial coefficients $f_{s_1s_2}^{\ell_{s_1s_2}m_{s_1s_2}}(p)$ in the expansion (33) also depend on the spin variables.

Substitution of (33) into (20) and then into (32) leads to two categories of relations among the adjustable functions, as shown in Appendices A and B. It follows that, for trial states of the form (20), the total spin of the system is a good quantum number, and the states of the system separate into singlet states with the total spin $S = 0$ (parastates) and into triplet states with $S = 1$ (orthostates). We should point out that this phenomenon is characteristic of the fermion antifermion systems, which are charge conjugation eigenstates, and does not arise for systems like μ^+e^- .

The singlet states

In this case $\ell_{s_1s_2} \equiv \ell = J$, $m_{11} = m_{22} = 0$ and $m_{12} = m_{21} = m_J$. The nonzero components of $F_{s_1s_2}(\mathbf{p})$ are $F_{\uparrow\downarrow}(\mathbf{p}) \equiv F_{12}(\mathbf{p}), F_{\downarrow\uparrow}(\mathbf{p}) \equiv F_{21}(\mathbf{p})$ and have the form

$$
F_{s_1s_2}(\mathbf{p}) = f_{s_1s_2}^{(sgl)J}(p)Y_J^{m_{s_1s_2}}(\hat{\mathbf{p}}),
$$
\n(34)

where the relations between $f_{12}^{(sgl)J}(p)$ and $f_{21}^{(sgl)J}(p)$ involve the Clebsch-Gordan (C-G) coefficients $C_{Jm}^{(sgl)Jm_{s_1s_2}}$ $\lim_{J_{mJ}}$ that is

$$
f_{s_1s_2}^{(sgl)J}(p) = C_{Jm_J}^{(sgl)Jm_{s_1s_2}} f^J(p),
$$
\n(35)

as it shown in Appendix A. We see that the spin and radial variables separate for the singlet states in the sense that the factors $f_{s_1s_2}^{(sgl)J}(p)$ have a common radial function $f^{J}(p)$. Thus, for the singlet states we obtain

$$
F_{s_1s_2}(\mathbf{p}) = C_{Jm_J}^{(sgl)Jm_{s_1s_2}} f^J(p) Y_J^{m_J}(\hat{\mathbf{p}}).
$$
\n(36)

The C-G coefficients $C_{Jm}^{(sgl)Jm_{s_1s_2}}$ $J_{m_J}^{(sgl)Jm_{s_1s_2}}$ have a simple form: $C_{Jm_J}^{(sgl)Jm_{11}}$ $\frac{J_{mJ}}{J_{mJ}}=C_{Jm_{J}}^{(sgl)Jm_{22}}$ $J_{m_J}^{(sgl)Jm_{22}} = 0, C_{Jm_J}^{(sgl)Jm_{12}}$ $\frac{Jm_J}{Jm_J} =$ $-C_{J m_J}^{(sgl) J m_{21}}$ $J_{m_J}^{(sg_l)Jm_{21}} = 1$ (see Appendix A). Therefore for the singlet states we can write expression (20) in the explicit form

$$
|\psi_T\rangle = \int d^3 \mathbf{p} f^J(p) Y_J^{m_J}(\hat{\mathbf{p}}) \left(b_{\mathbf{p}\uparrow}^\dagger d_{-\mathbf{p}\downarrow}^\dagger - b_{\mathbf{p}\downarrow}^\dagger d_{-\mathbf{p}\uparrow}^\dagger \right) |0\rangle. \tag{37}
$$

These states are characterized by the quantum numbers J, m_J parity $P = (-1)^{J+1}$ and charge conjugation $C = (-1)^J$. As we can see, the quantum numbers ℓ (orbital angular momentum), and total spin S are good quantum numbers for the singlet states as well. The spectroscopical notation is 1J_J .

The triplet states

The solution of the system (32) for $S = 1$ leads to two cases (Appendix A), namely $\ell_{s_1 s_2} \equiv \ell = J$, for which

$$
F_{s_1s_2}(\mathbf{p}) = f_{s_1s_2}^{(tr)J}(p) Y_J^{m_{s_1s_2}}(\hat{\mathbf{p}}),
$$
\n(38)

and $\ell_{s_1s_2} \equiv \ell = J \mp 1$, for which

$$
F_{s_1s_2}(\mathbf{p}) = f_{s_1s_2}^{J-1}(p)Y_{J-1}^{m_{s_1s_2}}(\hat{\mathbf{p}}) + f_{s_1s_2}^{J+1}(p)Y_{J+1}^{m_{s_1s_2}}(\hat{\mathbf{p}}),
$$
\n(39)

where

$$
m_{11} = m_J - 1, \quad m_{12} = m_{21} = m_J, \quad m_{22} = m_J + 1. \tag{40}
$$

The expressions for $f_{s_1s_2}^{\ell}(p)$ in both cases involve the C-G coefficients $C_{Jm_J}^{(tr)\ell m_s}$ $J_{m_J}^{(tr)em_s}$ for $S=1$ listed in Appendix A, that is

$$
f_{s_1s_2}^{(tr)\ell}(p) = C_{Jm_J}^{(tr)\ell m_s} f^{\ell}(p),\tag{41}
$$

where the index m_s is defined as

$$
m_s = +1
$$
, when $m_{s_1s_2} = m_{11}$,
\n $m_s = 0$, when $m_{s_1s_2} = m_{12} = m_{21}$,
\n $m_s = -1$, when $m_{s_1s_2} = m_{22}$. (42)

Thus, for the triplet states with $\ell = J$

$$
F_{s_1s_2}(\mathbf{p}) = C_{Jm_J}^{(tr)Jm_s} f^J(p) Y_J^{m_{s_1s_2}}(\hat{\mathbf{p}}).
$$
\n(43)

These functions correspond to states, which can be characterized by the quantum numbers J, m_J , parity $P = (-1)^{J+1}$ and charge conjugation $C = (-1)^{J+1}$. The orbital angular momentum ℓ , as well as the total spin $S = 1$, are good quantum numbers in this case. The spectroscopic notation for these states is $^{3}J_{J}$.

For the triplet states with $\ell = J \mp 1$ we obtain the result

$$
F_{s_1s_2}(\mathbf{p}) = C_{Jm_J}^{(tr)(J-1)m_s} f^{J-1}(p) Y_{J-1}^{m_{s_1s_2}}(\hat{\mathbf{p}}) + C_{Jm_J}^{(tr)(J+1)m_s} f^{J+1}(p) Y_{J+1}^{m_{s_1s_2}}(\hat{\mathbf{p}}),\tag{44}
$$

which involves two radial functions $f^{J-1}(p)$ and $f^{J+1}(p)$ corresponding to $\ell = J-1$ and $\ell = J+1$. This means that ℓ is not a good quantum number. Such states are characterized by quantum numbers J, m_J , $P = (-1)^J$, charge conjugation $C = (-1)^J$ and spin $S = 1$. In spectroscopic notation, these states are a mixture of $(3)(J-1)$ _J and $(3)(J+1)$ _J states.

The requirement that the states be charge conjugation eigenstates (the last equation of (32)) is intimately tied to the conservation of total spin. Indeed, a linear combination of singlet and triplet states like

$$
F_{s_1s_2}(\mathbf{p}) = C_1 f_{s_1s_2}^{(sgl)J}(p) Y_J^{m_{s_1s_2}}(\hat{\mathbf{p}}) + C_2 f_{s_1s_2}^{(tr)J}(p) Y_J^{m_{s_1s_2}}(\hat{\mathbf{p}}),
$$
\n(45)

satisfies the first three equations of (32) . However, it is unacceptable for describing a fermionantifermion system because the first and the second terms in (45) have different charge conjugation. For a system of two particles of different mass (such as μ^+e^-) charge conjugation is not applicable, so that the total spin would not be conserved.

5. The relativistic radial equations and application to positronium-like systems

We return to equation (24) and replace the functions $F_{s_1s_2}(p)$ by the expression (36) for singlet states and by (43) and (44) for triplet states. The variational procedure then leads to the following results:

For the singlet states $\ell = J$, $P = (-1)^{J+1}$, $C = (-1)^J$, the radial equations are

$$
(2\omega_p - E) f^{J}(p) = \frac{m^2}{(2\pi)^3} \int \frac{q^2 dq}{\omega_p \omega_q} \mathcal{K}^{(sgl)}(p,q) f^{J}(q), \qquad (46)
$$

where the kernel

$$
\mathcal{K}^{(sgl)}(p,q) = -i \sum_{s_1s_2\sigma_1\sigma_2m_J} \int d\hat{\mathbf{p}} d\hat{\mathbf{q}} C_{Jm_J}^{(sgl)s_1s_2\sigma_1\sigma_2} \mathcal{M}_{s_1s_2\sigma_1\sigma_2}(\mathbf{p}, \mathbf{q}) Y_J^{m_J*}(\hat{\mathbf{p}}) Y_J^{m_J}(\hat{\mathbf{q}}), \tag{47}
$$

is defined by the invariant M -matrix and the coefficients

$$
C_{Jm_J}^{(sgl)s_1s_2\sigma_1\sigma_2} \equiv C_{Jm_J}^{(sgl)Jm_{\sigma}} C_{Jm_J}^{(sgl)Jm_s} / \sum_{\nu_1\nu_2m_J} \left(C_{Jm_J}^{(sgl)Jm_{\nu}} \right)^2.
$$
 (48)

Here we have summed over m_J , because of the $(2J + 1)$ -fold energy degeneracy.

For the triplet states, we obtain different equations for the $\ell = J$, and $\ell = J \mp 1$ cases. Thus for the states with $\ell = J$, $P = (-1)^{J+1}$, $C = (-1)^{J+1}$ the result is

$$
(2\omega_p - E) f^{J}(p) = \frac{m^2}{(2\pi)^3} \int \frac{q^2 dq}{\omega_p \omega_q} \mathcal{K}^{(tr)}(p,q) f^{J}(q), \qquad (49)
$$

where the kernel $K^{(tr)}$ is formally like that of (47), namely,

$$
\mathcal{K}^{(tr)}(p,q) = -i \sum_{s_1 s_2 \sigma_1 \sigma_2 m_J} C^{(tr)s_1 s_2 \sigma_1 \sigma_2}_{Jm_J} \int d\hat{\mathbf{p}} d\hat{\mathbf{q}} \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{p}, \mathbf{q}) Y_J^{m_{s_1 s_2}}(\hat{\mathbf{p}}) Y_J^{m_{\sigma_1 \sigma_2}}(\hat{\mathbf{q}}). \tag{50}
$$

However it involves different C-G coefficients, namely

$$
C_{Jm_J}^{(tr)s_1s_2\sigma_1\sigma_2} = C_{Jm_J}^{(tr)Jm_{\sigma}} C_{Jm_J}^{(tr)Jm_s} / \sum_{\nu_1\nu_2m_J} \left(C_{Jm_J}^{(tr)Jm_{\nu}}\right)^2.
$$
 (51)

For the triplet states with $\ell = J \mp 1$, we have two independent radial functions $f^{J-1}(p)$ and $f^{J+1}(p)$. Thus the variational equation (24) leads to a system of coupled equations for $f^{J-1}(p)$ and $f^{J+1}(p)$. It is convenient to write them in matrix form,

$$
(2\omega_p - E) \mathbb{F}(p) = \frac{m^2}{(2\pi)^3} \int \frac{q^2 dq}{\omega_p \omega_q} \mathbb{K}(p, q) \mathbb{F}(q) ,
$$
 (52)

where

$$
\mathbb{F}(p) = \left[\begin{array}{c} f^{J-1}(p) \\ f^{J+1}(p) \end{array} \right],\tag{53}
$$

and

$$
\mathbb{K}(p,q) = \left[\begin{array}{cc} \mathcal{K}_{11}(p,q) & \mathcal{K}_{12}(p,q) \\ \mathcal{K}_{21}(p,q) & \mathcal{K}_{22}(p,q) \end{array} \right]. \tag{54}
$$

The kernels K_{ij} are similar in form to (47) and (50), that is

$$
\mathcal{K}_{ij}(p,q) = -i \sum_{\sigma_1 \sigma_2 s_1 s_2 m_J} C_{Jm_J ij}^{s_1 s_2 \sigma_1 \sigma_2} \int d\hat{\mathbf{p}} d\hat{\mathbf{q}} \, \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{p}, \mathbf{q}) Y_{\ell_j}^{m_{\sigma_1 \sigma_2}} (\hat{\mathbf{q}}) Y_{\ell_i}^{m_{s_1 s_2}}^* (\hat{\mathbf{p}}). \tag{55}
$$

However the coefficients $C_{Jm_J ij}^{s_1 s_2 \sigma_1 \sigma_2}$ are defined by expression

$$
C_{Jm_Jij}^{s_1s_2\sigma_1\sigma_2} = C_{Jm_J}^{(tr)\ell_j m_\sigma} C_{Jm_J}^{(tr)\ell_i m_s} / \sum_{s_1s_2m_J} \left(C_{Jm_J}^{(tr)\ell_i m_s} \right)^2, \tag{56}
$$

where $\ell_1 = J - 1$, $\ell_2 = J + 1$ and m_S is as defined in Eq. (42). The system (52) reduses to a single equation for $J = 0$ since $f^{J-1}(p) = 0$ in that case.

Our equation (24) , or its radial components (46) , (49) , (52) , contain the relativistic two-body kinematics (kinetic energy, recoil effects) exactly, but the dynamics are included approximately due to the limited nature of our trial state (20). This limitation is reflected in the fact that the interaction kernels of our equations contain only "tree-level" Feynman diagrams. Nevertheless our equations (46), (49), (52) have no negative-energy solutions, in contrast to the BS equation. They are variationally derived, hence the energy eigenvalues obtained from them will give meaningful values for any strength of the coupling.

To our knowledge, it is not possible to obtain analytic solutions of the relativistic radial momentum-space equations (46), (49) and (52). Thus one must resort to numerical or other approximation methods. Numerical solutions of such equations are discussed, for example, in [10], while a variational approximation has been employed in [5]. However, in this paper we will concentrate on perturbative $O(\alpha^4)$ solutions, since it is important to verify that our equations yield the correct fine structure for systems like positronium.

Our equations will yield energies which are incomplete beyond $O(\alpha^4)$, because our variational trial state (20), as mentioned, reflects only "tree-level" Feynman diagrams, that is no radiative corrections are incorporated. One could, of course, augment them by the addition of invariant matrix elements corresponding to higher-order Feynman diagrams (including radiative corrections) to the existing M -matrices in the kernels of our equations, as is done in the BS formalism. Indeed, such an approach has been used in a similar, though not variational, treatment of positronium by Zhang and Koniuk [15]. These authors show that the inclusion of invariant matrix elements corresponding to single-loop diagrams yields positronium energy eigenstates which are accurate to $O(\alpha^5, \alpha^5 \ln \alpha)$. However such augmentation of the kernels "by hand" would be contrary to the spirit of the present variational treatment, and we shall not pursue it in this work.

6. Semi-relativistic expansions and the non-relativistic limit

For perturbative solutions of our radial equations, it is necessary to work out expansions of the relevant expressions to first order beyond the non-relativistic limit. This shall be summarized in the present section. We perform the calculation in the Coulomb gauge, in which the photon propagator has the form [16]

$$
D_{00}(\mathbf{k}) = \frac{1}{\mathbf{k}^2}, \ D_{0j}(\mathbf{k}) = 0, \ D_{ij}(k^{\mu}) = \frac{1}{k^{\mu}k_{\mu}} \left(\delta_{ij} - \frac{k_{i}k_{j}}{\mathbf{k}^2}\right),
$$
 (57)

where $k^{\mu} = (\omega_p - \omega_q, \mathbf{p} - \mathbf{q}).$

To expand the amplitudes M of (26) and (27) to one order of $(p/m)^2$ beyond the non-relativistic limit, we take the free-particle spinors to be

$$
u(\mathbf{p},i) = \begin{bmatrix} \left(1 + \frac{\mathbf{p}^2}{8m^2}\right) \\ \frac{\left(\overrightarrow{\sigma} \cdot \mathbf{p}\right)}{2m} \end{bmatrix} \varphi_i, \qquad v(\mathbf{p},i) = \begin{bmatrix} \frac{\left(\overrightarrow{\sigma} \cdot \mathbf{p}\right)}{2m} \\ \left(1 + \frac{\mathbf{p}^2}{8m^2}\right) \end{bmatrix} \chi_i,
$$
\n(58)

as discussed in Appendix C. In this approximation the photon propagator takes on the form

$$
D_{00}\left(\mathbf{p}-\mathbf{q}\right)=\frac{1}{\left(\mathbf{p}-\mathbf{q}\right)^{2}},\quad D_{ij}\left(\mathbf{p}-\mathbf{q}\right)\simeq-\frac{1}{\left(\mathbf{p}-\mathbf{q}\right)^{2}}\left(\delta_{ij}-\frac{\left(p-q\right)_{i}\left(p-q\right)_{j}}{\left(\mathbf{p}-\mathbf{q}\right)^{2}}\right).
$$
(59)

Corresponding calculations give for the orbital part of the M-matrix

$$
\mathcal{M}_{s_1s_2\sigma_1\sigma_2}^{ope (orb)}(\mathbf{p}, \mathbf{q}) = ie^2 \left\{ \frac{1}{(\mathbf{p}-\mathbf{q})^2} + \frac{1}{m^2} \left(\frac{1}{4} + \frac{\mathbf{q} \cdot \mathbf{p}}{(\mathbf{p}-\mathbf{q})^2} + \frac{(\mathbf{p} \times \mathbf{q})^2}{(\mathbf{p}-\mathbf{q})^4} \right) \right\} \delta_{s_1\sigma_1} \delta_{s_2\sigma_2}.
$$
 (60)

The terms of the expansion linear in spin correspond to the spin-orbit interaction:

$$
\mathcal{M}_{s_1s_2\sigma_1\sigma_2}^{ope(s-o)}(\mathbf{p},\mathbf{q}) = \frac{3e^2}{4m^2} \varphi_{s_1}^{\dagger} \chi_{\sigma_2}^{\dagger} \frac{\left(\overrightarrow{\sigma}^{(+)} - \overrightarrow{\sigma}^{(-)}\right) \cdot (\mathbf{p} \times \mathbf{q})}{\left(\mathbf{p} - \mathbf{q}\right)^2} \varphi_{\sigma_1} \chi_{s_2}.
$$
(61)

Here $\vec{\sigma}^{(+)}$ and $\vec{\sigma}^{(-)}$ are positron and electron spin matrices respectively, defined as follow: $\vec{\sigma}^{(+)}\varphi_{\sigma_1}$ Here $\vec{\sigma}^{(+)}$ and $\vec{\sigma}^{(-)}$ are positron and electron spin matrices respectively, defined as follow: $\vec{\sigma}^{(+)}\varphi_{\sigma_1}\chi_{s_2} = (\vec{\sigma}^{(+)}\varphi_{\sigma_1})\chi_{s_2}, \vec{\sigma}^{(-)}\varphi_{\sigma_1}\chi_{s_2} = \varphi_{\sigma_1}(\vec{\sigma}^{(-)}\chi_{s_2})$. The quadratic spi terms are

$$
\mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2}^{ope(s-s)}(\mathbf{p}, \mathbf{q}) = \frac{ie^2}{4m^2} \varphi_{s_1}^{\dagger} \chi_{\sigma_2}^{\dagger} \left\{ -\frac{\left(\overrightarrow{\sigma}^{(+)} \cdot (\mathbf{p} - \mathbf{q})\right) \left(\overrightarrow{\sigma}^{(-)} \cdot (\mathbf{p} - \mathbf{q})\right)}{(\mathbf{p} - \mathbf{q})^2} + \overrightarrow{\sigma}^{(+)} \cdot \overrightarrow{\sigma}^{(-)} \right\} \varphi_{\sigma_1} \chi_{s_2}.
$$
 (62)

Lastly, the virtual annihilation contribution is given by

$$
\mathcal{M}_{s_1s_2\sigma_1\sigma_2}^{ann}(p,q) = -\frac{ie^2}{4m^2} \varphi_{s_1}^{\dagger} \chi_{\sigma_2}^{\dagger} \left\{ \overrightarrow{\sigma}^{(+)} \cdot \overrightarrow{\sigma}^{(-)} \right\} \varphi_{\sigma_1} \chi_{s_2},\tag{63}
$$

where we have excluded a divergent term, which appears in the Coulomb gauge calculation. This divergence is an artefact of the Coulomb gauge. It does not arise, for example, in the Lorentz gauge, where only the expression (63) is obtained. However the Lorentz gauge is not convenient for obtaining all other $O(\alpha^4)$ corrections because it contains spurious degrees of freedom (longitudinal polarization) of the photon.

We have used expressions (60)-(63) to obtain the corresponding radial kernels. Details of the calculations can be found in Appendix D. We use the notation $z = (p^2 + q^2)/2pq$, and $Q_{\lambda}(z)$ is the Legendre function of the second kind [17]. The contributions of the various terms to the kernel are as follows:

Singlet states with $\ell = J \ (J \ge 0), P = (-1)^{J+1}, C = (-1)^{J}$

Orbital term

$$
\mathcal{K}^{(sgl)(o)}(p,q) = \frac{2\pi e^2}{pq} Q_J(z) + \frac{\pi e^2}{m^2} \left(-\frac{J-3}{2} \left(\frac{p}{q} + \frac{q}{p} \right) Q_J(z) + (J+1) Q_{J+1}(z) - 2\delta_{J,0} \right). \tag{64}
$$

Spin-orbit interaction

$$
\mathcal{K}^{(sgl)(s-o)}(p,q) = 0. \tag{65}
$$

Spin-spin interaction

$$
\mathcal{K}^{(sgl)(s-s)}(p,q) = \frac{2\pi e^2}{m^2} \delta_{J,0}.
$$
\n(66)

Triplet states with $\ell = J \ (J \ge 1), P = (-1)^{J+1}, C = (-1)^{J+1}$

Orbital term

$$
K^{(tr)(o)}(p,q) = \frac{2\pi e^2}{pq} Q_J(z) + \frac{\pi e^2}{m^2} \left(-\frac{J-3}{2} \left(\frac{p}{q} + \frac{q}{p} \right) Q_J(z) + (J+1) Q_{J+1}(z) \right).
$$
 (67)

Spin-orbit interaction

$$
K^{(tr)(s-o)}(p,q) = -\frac{3\pi e^2}{m^2} \frac{1}{2J+1} \left\{ Q_{J+1}(z) - Q_{J-1}(z) \right\}.
$$
 (68)

Spin-spin interaction

$$
\mathcal{K}^{(tr)(s-s)}(p,q) = \frac{\pi e^2}{2m^2} \left(\frac{p}{q} + \frac{q}{p}\right) Q_J(z) - \frac{\pi e^2}{m^2} \frac{1}{2J+1} \left\{ JQ_{J+1}(z) + (J+1)Q_{J-1}(z) \right\}.
$$
 (69)

Triplet states with $\ell = J - 1$ $(J \ge 1)$, $\ell = J + 1$ $(J \ge 0)$, $P = (-1)^J$, $C = (-1)^J$

The off-diagonal elements of the kernel matrix (Eqs.. $(52)-(54)$), K_{12} and K_{21} which are responsible for mixing of states with $\ell = J - 1$ and $\ell = J + 1$, get a non-zero contribution from the spin-spin interactions only:

$$
\mathcal{K}_{12}(p,q) = \mathcal{K}_{21}(p,q) = \frac{\pi e^2}{5m^2} \frac{\sqrt{J(J+1)}}{(2J+1)} \left(\frac{p}{q} Q_{J+1}(z) + \frac{q}{p} Q_{J-1}(z) - 2Q_J(z) \right). \tag{70}
$$

The contributions to the diagonal elements of the kernel matrix are the following:

Orbital terms

$$
\mathcal{K}_{11}^{(o)}(p,q) = \frac{2\pi e^2}{pq} Q_{J-1}(z) \n+ \frac{\pi e^2}{m^2} \left(-\frac{J-4}{2} \left(\frac{p}{q} + \frac{q}{p} \right) Q_{J-1}(z) + JQ_J(z) - 2\delta_{J-1,0} \right),
$$
\n(71)

$$
\mathcal{K}_{22}^{(o)}(p,q) = \frac{2\pi e^2}{pq} Q_{J+1}(z) \n+ \frac{\pi e^2}{m^2} \left(-\frac{J-2}{2} \left(\frac{p}{q} + \frac{q}{p} \right) Q_{J+1} + (J+2) Q_{J+2} \right).
$$
\n(72)

Spin-orbit interaction

$$
\mathcal{K}_{11}^{(s-o)}(p,q) = \frac{3\pi e^2}{m^2} \frac{J-1}{2J-1} \left(Q_J(z) - Q_{J-2}(z) \right),\tag{73}
$$

$$
\mathcal{K}_{22}^{(s-o)}(p,q) = -\frac{3\pi e^2}{m^2 c^2} \frac{J+2}{2J+3} \left(Q_{J+2}(z) - Q_J(z) \right). \tag{74}
$$

Spin-spin interaction

$$
\mathcal{K}_{11}^{(s-s)}(p,q) = \frac{\pi e^2}{2m^2} \frac{1}{2J+1} \left(\left(\frac{p}{q} + \frac{q}{p} \right) Q_{J-1}(z) - 2Q_J(z) \right),\tag{75}
$$

$$
\mathcal{K}_{22}^{(s-s)}(p,q) = \frac{\pi e^2}{2m^2} \frac{1}{2J+3} \left(\left(\frac{p}{q} + \frac{q}{p} \right) Q_{J+1}(z) - 2Q_{J+2}(z) \right). \tag{76}
$$

Annihilation term

$$
\mathcal{K}^{ann}(p,q) = -\frac{2\pi e^2}{m^2} \delta_{J-1,0}.
$$
\n(77)

We note that in the non-relativistic limit only the first terms of the orbital part of the kernels survive. They have the common form $2\pi i e^2 Q_{\ell}(z)/pq$, hence all radial equations reduce to the form

$$
(2\omega_p - E) f^{\ell}(p) = \frac{m^2 e^2}{\pi \omega_p p} \int_0^\infty dq \frac{q}{\omega_q} Q_\ell(z) f^{\ell}(q). \tag{78}
$$

Recalling, also, that

$$
\omega_p = \sqrt{m^2 + \mathbf{p}^2} \simeq m \left(1 + \frac{1}{2} \left(\frac{\mathbf{p}}{m} \right)^2 \right),\tag{79}
$$

we obtain, in the non relativistic limit, the momentum-space Schrödinger radial equations

$$
\left(\frac{\mathbf{p}^2}{2\mu} - \varepsilon\right) f^{\ell}(p) = \frac{\alpha}{\pi} \frac{1}{p} \int_0^\infty dq \, q \, Q_J(z) f^{\ell}(q),\tag{80}
$$

where $\alpha = e^2/4\pi$, $\mu = \frac{m}{2}$ $\frac{n}{2}, \varepsilon = E - 2m.$

7. Energy levels. Fine and hyperfine structure

The relativistic energy eigenvalues $E_{n,J}$ can be calculated from the expression

$$
E \int_0^{\infty} dp \, p^2 f^J(p) f^J(p) = \int_0^{\infty} dp \, p^2 \, 2\omega_p f^J(p) f^J(p) - \frac{m^2}{(2\pi)^3} \int_0^{\infty} \frac{dp p^2}{\omega_p} \int_0^{\infty} dq \frac{q^2}{\omega_q} \mathcal{K}^{(sgl,tr)}(p,q) f^J(p) f^J(q)
$$
(81)

for the singlet and $\ell = J$ triplet states.

For the $\ell = J \mp 1$ triplet states the corresponding result is (see equation (52))

$$
E \int_0^\infty dp \, p^2 \mathbb{F}^\dagger(p) \mathbb{F}(p) = \int_0^\infty dp \, p^2 \, 2\omega_p \mathbb{F}^\dagger(p) \mathbb{F}(p) - \frac{m^2}{(2\pi)^3} \int_0^\infty \frac{dpp^2}{\omega_p} \int_0^\infty dq \frac{q^2}{\omega_q} \mathbb{K}(p,q) \mathbb{F}^\dagger(p) \mathbb{F}(q).
$$
 (82)

To obtain results for E to $O(\alpha^4)$ we use the forms of the kernels expanded to $O(p^2/m^2)$ (eqs. (64)-(77)) and replace $f^{\ell}(p)$ by their non-relativistic (Schrödinger) form (see (147), Appendix D). The most important integrals that we used for calculating (81) and (82), are given in Appendix D. In Appendix E we show that the contribution of kernels K_{12} and K_{21} in (82), is zero at $O(\alpha^4)$. Thus, the energy corrections for the triplet states with $\ell = J - 1$ and $\ell = J + 1$ can be calculated independently.

The results will be presented in the form $\Delta \varepsilon = E - 2m + \alpha^2 m/2n^2$.

Singlet states $(\ell = J \ (J \ge 0), \ P = (-1)^{J+1}, \ C = (-1)^{J})$ The kinetic energy corrections

$$
\Delta \varepsilon_K^{(sgl)} = -\frac{\alpha^4 m}{8} \left(\frac{1}{2J+1} \frac{1}{n^3} - \frac{3}{8} \frac{1}{n^4} \right). \tag{83}
$$

The potential energy corrections

$$
\Delta \varepsilon_P^{(sgl)(o)} = -\frac{\alpha^4 m}{8} \left(\left(\frac{3}{2J+1} - 2\delta_{J,0} \right) \frac{1}{n^3} - \frac{1}{n^4} \right),\tag{84}
$$

$$
\Delta \varepsilon_P^{(sgl)(s-o)} = 0,\t\t(85)
$$

$$
\Delta \varepsilon_P^{(sgl)(s-s)} = -\frac{\alpha^4 m}{4} \frac{\delta_{J,0}}{n^3}.
$$
\n(86)

The total energy corrections

$$
\Delta \varepsilon^{(sgl)} = -\frac{\alpha^4 m}{8} \left(\frac{4}{2J+1} \frac{1}{n^3} - \frac{11}{8n^4} \right). \tag{87}
$$

Triplet states $(\ell = J \ (J \geq 1), \ P = (-1)^{J+1}, \ C = (-1)^{J+1})$ The kinetic energy corrections

$$
\Delta \varepsilon_K^{(tr)} = -\frac{\alpha^4 m}{8} \left(\frac{1}{2J+1} \frac{1}{n^3} - \frac{3}{8} \frac{1}{n^4} \right). \tag{88}
$$

The potential energy corrections

$$
\Delta \varepsilon_P^{(tr)(o)} = -\frac{\alpha^4 m}{8} \left(\left(\frac{3}{2J+1} - 2\delta_{J,0} \right) \frac{1}{n^3} - \frac{1}{n^4} \right),\tag{89}
$$

$$
\Delta \varepsilon_P^{(tr)(s-o)} = -\frac{\alpha^4 m}{8} \frac{3}{J(J+1)(2J+1)} \frac{1}{n^3},\tag{90}
$$

$$
\Delta \varepsilon_P^{(s-s)} = \frac{\alpha^4 m}{8} \frac{1}{J\left(J+1\right)\left(2J+1\right)} \frac{1}{n^3}.\tag{91}
$$

The total energy corrections

$$
\Delta \varepsilon^{(tr)} = -\frac{\alpha^4 m}{8} \left(\left(\frac{4}{2J+1} + \frac{2}{J(J+1)(2J+1)} \right) \frac{1}{n^3} - \frac{11}{8} \frac{1}{n^4} \right). \tag{92}
$$

Triplet states $(\ell = J - 1 \ (J \ge 1), \ P = (-1)^J, \ C = (-1)^J)$ The kinetic energy corrections

$$
\Delta \varepsilon_K^{(tr)(J-1)} = -\frac{\alpha^4 m}{8} \left(\frac{1}{2J - 1} \frac{1}{n^3} - \frac{3}{8} \frac{1}{n^4} \right). \tag{93}
$$

The potential energy corrections

$$
\Delta \varepsilon_P^{(tr)(o)(J-1)} = -\frac{\alpha^4 m}{8} \left[\left(\frac{3}{2J-1} - 2\delta_{J,1} \right) \frac{1}{n^3} - \frac{1}{n^4} \right],\tag{94}
$$

$$
\Delta \varepsilon_P^{tr(s-o)(J-1)} = \frac{\alpha^4 m}{8} \frac{3(1-\delta_{J,1})}{J(2J-1)} \frac{1}{n^3},\tag{95}
$$

$$
\Delta \varepsilon_P^{tr(s-s)(J-1)} = -\frac{\alpha^4 m}{8} \frac{1 - \delta_{J,1}}{J(2J+1)(2J-1)} \frac{1}{n^3},\tag{96}
$$

$$
\Delta \varepsilon^{(ann)} = \frac{\alpha^4 m}{4} \frac{1}{n^3} \delta_{J,1}.
$$
\n(97)

The total energy corrections

$$
\Delta \varepsilon^{tr(J-1)} = -\frac{\alpha^4 m}{8} \left(\left(\frac{4}{2J-1} - \frac{2(3J+1)(1-\delta_{J,1})}{J(2J+1)(2J-1)} - 4\delta_{J,1} \right) \frac{1}{n^3} - \frac{11}{8} \frac{1}{n^4} \right). \tag{98}
$$

Triplet states $(\ell = J + 1 \; (J \ge 0), \; P = (-1)^J, \; C = (-1)^J)$ The kinetic energy corrections

$$
\Delta \varepsilon_K^{(tr)(J+1)} = -\frac{\alpha^4 m}{8} \left(\frac{1}{2J+3} \frac{1}{n^3} - \frac{3}{8} \frac{1}{n^4} \right). \tag{99}
$$

The potential energy corrections

$$
\Delta \varepsilon_P^{(tr)(o)(J+1)} = -\frac{\alpha^4 m}{8} \left[\frac{3}{2J+3} \frac{1}{n^3} - \frac{1}{n^4} \right],\tag{100}
$$

$$
\Delta \varepsilon_P^{tr(s-o)(J+1)} = -\frac{\alpha^4 m}{8} \frac{3}{(J+1)(2J+3)} \frac{1}{n^3},\tag{101}
$$

$$
\Delta \varepsilon_P^{tr(s-s)(J+1)} = -\frac{\alpha^4 m}{8} \frac{1}{(J+1)(2J+3)(2J+1)} \frac{1}{n^3}.
$$
\n(102)

The total energy corrections

$$
\Delta \varepsilon^{tr(J+1)} = -\frac{\alpha^4 m}{8} \left(\frac{2}{2J+3} \left(2 + \frac{3J+2}{(J+1)(2J+1)} \right) \frac{1}{n^3} - \frac{11}{8} \frac{1}{n^4} \right). \tag{103}
$$

These results are in agreement with the well-known positronium fine structure results [18], [19].

8. Concluding remarks

We have considered a reformulation of electrodynamics, in which covariant Green functions are used to solve the field equations for the mediating electromagnetic field in terms of the fermion field. This leads to a reformulated Hamiltonian with an interaction term in which the photon propagator appears sandwiched between fermionic currents.

The variational method within a Hamiltonian formalism of quantum field theory is used to determine approximate eigensolutions for bound relativistic fermion-antifermion states. The reformulation enables us to use the simplest possible trial state to derive a relativistic momentum-space Salpeter-like equation for a positronium-like system. The invariant M matrices corresponding to one-photon exchange and virtual annihilation Feynman diagrams arise directly in the interaction kernel of this equation.

The trial states are chosen to be eigenstates of the total angular momentum operator $\hat{\mathbf{J}}^2$ and \hat{J}_3 , along with parity and charge conjugation. A general relativistic reduction of the wave equations to radial form is given. For given J there is a single radial equation for total spin zero singlet states, but for spin triplet states there are, in general two coupled equations. We show how the classification of states follows naturally from the system of eigenvalue equations obtained with our trial state.

It is not possible, as far as we know, to obtain analytic solutions of our relativistic radial equations nor the resulting eigenvalues of the particle-antiparticle system described. However, it is possible to obtain $O(\alpha^4)$ corrections analytically for all states using perturbation theory. The results agree with well known results for positronium, obtained on the basis of the Bethe-Salpeter equation [19], which lends credence to the validity of our variationally derived equations.

The method presented here can be generalized to include effects higher order in alpha by using dressed propagators in place of the bare propagators. This shall be the subject of a forthcoming work.

Acknowledgments

The financial support of the Natural Sciences and Engineering Research Council of Canada for this work is gratefully acknowledged.

Appendix A. Total angular momentum operator in relativistic form

The total angular momentum operator is defined by expression

$$
\hat{\mathbf{J}} = \int d^3 \mathbf{x} : \psi^\dagger(x) \, (\hat{\mathbf{L}} + \hat{\mathbf{S}}) \psi(x) ; \tag{104}
$$

where $\hat{\mathbf{L}} = \hat{\mathbf{x}} \times \hat{\mathbf{p}}$ and $\hat{\mathbf{S}} = \frac{1}{2} \overrightarrow{\sigma}$ are the orbital angular momentum and spin operators. We use the standard representation for the Pauli matrices

$$
\widehat{\overrightarrow{\sigma}} = \left[\begin{array}{cc} \overrightarrow{\sigma} & 0 \\ 0 & \overrightarrow{\sigma} \end{array} \right],\tag{105}
$$

$$
\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.
$$
 (106)

Using the field operator $\psi(x)$ in the form (15), after tedious calculations we obtain

$$
\widehat{J}_1 = \int d^3 \mathbf{q} \left(\begin{array}{c} \widehat{L}_{q1} \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\downarrow} + d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) \\ + \frac{1}{2} \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\downarrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\uparrow} + d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) \right),
$$
\n
$$
\widehat{J}_2 = \int d^3 \mathbf{q} \left(\begin{array}{c} \widehat{L}_{q2} \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\downarrow} + d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) \\ + \frac{i}{2} \left(-b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\downarrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\uparrow} - d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\downarrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\uparrow} \right) \end{array} \right), \tag{107}
$$
\n
$$
\widehat{J}_3 = \int d^3 \mathbf{q} \left(\begin{array}{c} \widehat{L}_{q3} \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\downarrow} + d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\uparrow} \right) \end{array} \right),
$$

Here $\widehat{\mathbf{L}}_q$ is the orbital angular momentum operator in momentum representation:

$$
(\widehat{\mathbf{L}}_q)_i \equiv \widehat{L}_{qi} = -i \left(\mathbf{q} \times \nabla_q \right)_i. \tag{108}
$$

Note that these expressions are valid for any t , since the time-dependent phase factors of the form $e^{i\omega_q t}$ cancel out.

For the operator $\mathbf{\hat{J}}^2 = \hat{J}_1^2 + \hat{J}_2^2 + \hat{J}_3^2$ we have

$$
\hat{\mathbf{J}}^{2} = \int d^{3} \mathbf{q} \left(\begin{array}{c} \left(\widehat{\mathbf{L}}_{q}^{2} + \frac{3}{4} \right) \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\downarrow} + d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) \\ + \widehat{L}_{q3} \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\uparrow} + \widehat{L}_{q-} d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\downarrow} + \widehat{L}_{q+} d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\uparrow} \right) + \\ + \widehat{L}_{q3} \left(b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} b_{\mathbf{q}\downarrow} + d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + b_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) \\ + \frac{1}{2} \left(b_{q'\uparrow}^{\dagger} b_{q'\uparrow} d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) + \frac{1}{2} \left(b_{q'\uparrow}^{\dagger} b_{q'\uparrow} d_{\mathbf{q}\uparrow}^{\dagger} d_{\mathbf{q}\uparrow} + b_{\mathbf{q}\uparrow}^{\dagger} b_{\mathbf{q}\uparrow} + d_{\mathbf{q}\downarrow}^{\dagger} d_{\mathbf{q}\downarrow} \right) \\ + \frac{1}{2} \int d^{3} \mathbf{q}' d^{3} \mathbf{q} \left(\begin{array}{c} \frac{\partial}{\partial q} \left(b_{\mathbf{q}\uparrow} + b_{\mathbf{q
$$

where

$$
\widehat{L}_{q+} = \widehat{L}_{q1} + i\widehat{L}_{q2}, \qquad \widehat{L}_{q-} = \widehat{L}_{q1} - i\widehat{L}_{q2}.
$$
\n(110)

The requirement that the trial state (20) be an eigenstate of \hat{J}^2 and \hat{J}_z leads to the system of equations

$$
(\hat{L}_3 + 1) F_{11} = m_J F_{11},
$$

\n
$$
\hat{L}_3 F_{12} = m_J F_{12},
$$

\n
$$
\hat{L}_3 F_{21} = m_J F_{21},
$$

\n
$$
(\hat{L}_3 - 1) F_{22} = m_J F_{22},
$$
\n(111)

$$
\left(J(J+1) - \hat{\mathbf{L}}^2 - 2 - 2\hat{L}_3\right) F_{11} = \hat{L}_-(F_{12} + F_{21}),
$$
\n
$$
\left(J(J+1) - \hat{\mathbf{L}}^2 - 1\right) F_{12} = F_{21} + \hat{L}_+ F_{11} + \hat{L}_- F_{22},
$$
\n
$$
\left(J(J+1) - \hat{\mathbf{L}}^2 - 1\right) F_{21} = F_{12} + \hat{L}_+ F_{11} + \hat{L}_- F_{22},
$$
\n
$$
\left(J(J+1) - \hat{\mathbf{L}}^2 - 2 + 2\hat{L}_3\right) F_{22} = \hat{L}_+ (F_{12} + F_{21}).
$$
\n(112)

Substitution of the expressions (33) for $F_{s_1s_2}$ and use of eq. (111) gives

$$
m_{12} = m_{21} = m_J, \qquad m_{11} = m_J - 1, \qquad m_{22} = m_J + 1,\tag{113}
$$

$$
\ell_{11} = \ell_{22} = \ell_{12} = \ell_{21} = \ell,\tag{114}
$$

and

$$
(J(J+1) - \ell(\ell+1) - 2m_J) f_{11}^{\ell}(p) = \sqrt{(\ell - m_J + 1)(\ell + m_J)} f_{12}^{\ell}(p)
$$

\n
$$
+ \sqrt{(\ell - m_J + 1)(\ell + m_J)} f_{21}^{\ell}(p),
$$

\n
$$
(J(J+1) - \ell(\ell+1) - 1) f_{12}^{\ell}(p) = f_{21}^{\ell}(p)
$$

\n
$$
+ \sqrt{(\ell + m_J)(\ell - m_J + 1)} f_{11}^{\ell}(p)
$$

\n
$$
+ \sqrt{(\ell - m_J)(\ell + m_J + 1)} f_{22}^{\ell}(p),
$$

\n
$$
(J(J+1) - \ell(\ell+1) - 1) f_{21}^{\ell}(p) = f_{12}^{\ell}(p)
$$

\n
$$
+ \sqrt{(\ell + m_J)(\ell - m_J + 1)} f_{11}^{\ell}(p)
$$

\n
$$
+ \sqrt{(\ell - m_J)(\ell + m_J + 1)} f_{22}^{\ell}(p),
$$

\n
$$
(J(J+1) - \ell(\ell+1) + 2m_J) f_{22}^{\ell}(p) = \sqrt{(\ell + m_J + 1)(\ell - m_J)} f_{12}^{\ell}(p)
$$

\n
$$
+ \sqrt{(\ell + m_J + 1)(\ell - m_J)} f_{21}^{\ell}(p).
$$

\n(115)

The singlet states correspond to the solution $f_{11}^{\ell}(p) = f_{22}^{\ell}(p) = 0$, $f_{12}^{\ell}(p) = -f_{21}^{\ell}(p)$ of this system with $\ell = J \ (J \geq 0)$.

For the triplet states the solutions are $f_{12}^{\ell}(p) = f_{21}^{\ell}(p) \equiv f^{\ell}(p)$, and, for $\ell = J - 1 \ (J \ge 1)$:

$$
(J - m_J) f_{11}^{J-1}(p) = \sqrt{(J - m_J) (J + m_J - 1)} f^{J-1}(p), \qquad (116)
$$

$$
(J+m_J) f_{22}^{J-1}(p) = \sqrt{(J+m_J)(J-m_J-1)} f^{J-1}(p), \qquad (117)
$$

for $\ell = J \ (J \geq 1)$:

$$
m_J f_{11}^J(p) = -\sqrt{(J+m_J)(J-m_J+1)} f^J(p), \qquad (118)
$$

$$
m_J f_{22}^J(p) = \sqrt{(J - m_J)(J + m_J + 1)} f^J(p), \qquad (119)
$$

for $\ell = J + 1 \ (J \ge 0)$:

$$
(J+1+mJ) f11J+1(p) = -\sqrt{(J-mJ+2)(J+mJ+1)} fJ+1(p),
$$
\n(120)

$$
(J+1-m_J) f_{22}^{J+1}(p) = -\sqrt{(J-m_J+1)(J+m_J+2)} f^{J+1}(p). \tag{121}
$$

It is convenient to introduce the table of coefficients $C_{Jm}^{(tr)\ell m_s}$ $\frac{Jm_J}{Jm_J}$:

These coefficients coincide with the usual Clebsch-Gordan coefficients for $S = 1$ except for a factor 2 in the denominator, which we absorb into the normalization constant.

Appendix B. Parity and charge conjugation

We consider the application of the parity operator to the trial state (20) :

$$
\widehat{\mathcal{P}}|\psi_T\rangle = \sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_1s_2}(\mathbf{p}) \widehat{\mathcal{P}} b_{ps_1}^{\dagger} d_{-ps_2}^{\dagger} |0\rangle \n= \sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_1s_2}(\mathbf{p}) \widehat{\mathcal{P}} b_{ps_1}^{\dagger} \widehat{\mathcal{P}}^{-1} \widehat{\mathcal{P}} d_{-ps_2}^{\dagger} \widehat{\mathcal{P}}^{-1} \widehat{\mathcal{P}} |0\rangle.
$$
\n(122)

Making use of the properties

$$
\widehat{\mathcal{P}}b_{ps_1}^{\dagger}\widehat{\mathcal{P}}^{-1} = \eta^P b_{-ps_1}^{\dagger}, \qquad \widehat{\mathcal{P}}d_{-ps_2}^{\dagger}\widehat{\mathcal{P}}^{-1} = -\eta^P d_{ps_2}^{\dagger}, \qquad \widehat{\mathcal{P}}\left|0\right\rangle = \left|0\right\rangle,\tag{123}
$$

where η^P is the intrinsic parity $((\eta^P)^2 = 1)$, it follows that

$$
\hat{\mathcal{P}}|\psi_{T}\rangle = \sum_{s_{1}s_{2}} \int d^{3} \mathbf{p} F_{s_{1}s_{2}}(\mathbf{p}) \hat{\mathcal{P}} b_{ps_{1}}^{\dagger} d_{-ps_{2}}^{\dagger} |0\rangle = -\sum_{s_{1}s_{2}} \int d^{3} \mathbf{p} F_{s_{1}s_{2}}(-\mathbf{p}) b_{ps_{1}}^{\dagger} d_{-ps_{2}}^{\dagger} |0\rangle
$$

$$
= P \sum_{s_{1}s_{2}} \int d^{3} \mathbf{p} F_{s_{1}s_{2}}(\mathbf{p}) b_{ps_{1}}^{\dagger} d_{-ps_{2}}^{\dagger} |0\rangle , \qquad (124)
$$

where the parity eigenvalue P depends on the symmetry of $F_{s_1s_2}(p)$ in different states: For the singlet states $(\ell = J)$ we get from (36) $F_{s_1 s_2}(-p) = (-1)^J F_{s_1 s_2}(p)$, so that $P = (-1)^{J+1}$. For the triplet states with $\ell = J$ we get from (38) $F_{s_1 s_2}(-p) = (-1)^J F_{s_1 s_2}(p)$, hence $P = (-1)^{J+1}$. For the triplet states with $\ell = J \pm 1$ we get from (39) $F_{s_1s_2}(-p) = (-1)^{J+1} F_{s_1s_2}(p)$, therefore $P = (-1)^J$.

Charge conjugation is associated with the interchange of the particle and antiparticle. Applying the charge conjugation operator to the trial state (20) we get

$$
\widehat{\mathcal{C}}|\psi_T\rangle = \sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_1s_2}(\mathbf{p}) \widehat{\mathcal{C}} b_{ps_1}^\dagger d_{-ps_2}^\dagger |0\rangle \tag{125}
$$

$$
= \sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_1s_2}(\mathbf{p}) \widehat{\mathcal{C}} b_{ps_1}^{\dagger} \widehat{\mathcal{C}}^{-1} \widehat{\mathcal{C}} d_{-ps_2}^{\dagger} \widehat{\mathcal{C}}^{-1} \widehat{\mathcal{C}} |0\rangle. \tag{126}
$$

Using the relations

$$
\widehat{C}b_{ps_1}^{\dagger}\widehat{C}^{-1} = \eta^C d_{ps_1}^{\dagger}, \quad \widehat{C}d_{-ps_2}^{\dagger}\widehat{C}^{-1} = \eta^C b_{-ps_2}^{\dagger}, \quad \widehat{C}|0\rangle = |0\rangle, \tag{127}
$$

where $(\eta^C)^2 = 1$, we obtain

$$
\widehat{\mathcal{C}}|\psi_T\rangle = \sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_1s_2}(\mathbf{p}) \widehat{\mathcal{C}} b_{ps_1}^\dagger d_{-ps_2}^\dagger |0\rangle = -\sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_2s_1}(\mathbf{p}) b_{ps_1}^\dagger d_{-ps_2}^\dagger |0\rangle
$$

$$
= C \sum_{s_1s_2} \int d^3 \mathbf{p} F_{s_1s_2}(\mathbf{p}) b_{ps_1}^\dagger d_{-ps_2}^\dagger |0\rangle , \qquad (128)
$$

where the charge conjugation quantum number C depends on the symmetry of $F_{s_1s_2}(p)$ in different states:

For the singlet states $(\ell = J)$ we get from (36) $F_{s_1s_2}(-p) = (-1)^{J+1} F_{s_1s_2}(p)$, hence $C = (-1)^J$. For the triplet states with $\ell = J$ we get from (38) $F_{s_1s_2}(-p) = (-1)^J F_{s_1s_2}(p)$, therefore $C =$ $(-1)^{J+1}$.

For the triplet states with $\ell = J \pm 1$ we get from (39) $F_{s_1s_2}(-p) = (-1)^{J+1} F_{s_1s_2}(p)$, so that $C = (-1)^J$.

Appendix C. Expansion of the spinors

We recall the form of the particle spinors:

$$
u(\mathbf{p}, i) = N_{\mathbf{p}} \begin{bmatrix} 1 \\ \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{\omega_p + m} \end{bmatrix} \varphi_i,
$$
 (129)

where

$$
\varphi_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \varphi_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad N_{\mathbf{p}} = \sqrt{\frac{\omega_p + m}{2m}}.
$$
 (130)

The antiparticle or "positron" representation for the $v_i(p)$ spinors has the form

$$
v(\mathbf{p},i) = N_{\mathbf{p}} \begin{bmatrix} \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{\omega_p + m} \\ 1 \end{bmatrix} \chi_i,
$$
\n(131)

where

$$
\chi_1 = \left[\begin{array}{c} 0 \\ 1 \end{array} \right], \quad \chi_2 = -\left[\begin{array}{c} 1 \\ 0 \end{array} \right]. \tag{132}
$$

The normalization is

$$
\overline{u}(\mathbf{p},i)u(\mathbf{p},j) = \delta_{ij}, \qquad \overline{v}(\mathbf{p},i)v(\mathbf{p},j) = -\delta_{ij}.
$$
\n(133)

Expanding in powers of p/m and keeping the lowest non-trivial order terms,

$$
\frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{\omega_p + m} \simeq \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{2m},\tag{134}
$$

$$
N_{\mathbf{p}} = \sqrt{\frac{\omega_p + m}{2m}} \simeq 1 + \frac{\mathbf{p}^2}{8m^2},\tag{135}
$$

we obtain the result

$$
u(\mathbf{p},i) \simeq \left(1 + \frac{\mathbf{p}^2}{8m^2}\right) \left[\begin{array}{c} 1\\ \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{2m} \end{array}\right] \varphi_i = \left[\begin{array}{c} \left(1 + \frac{\mathbf{p}^2}{8m^2}\right)\\ \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{2m} \end{array}\right] \varphi_i, \tag{136}
$$

$$
v(\mathbf{p},i) \simeq \left(1 + \frac{\mathbf{p}^2}{8m^2}\right) \left[\begin{array}{c} \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{2m} \\ 1 \end{array}\right] \chi_i = \left[\begin{array}{c} \frac{(\overrightarrow{\sigma} \cdot \mathbf{p})}{2m} \\ \left(1 + \frac{\mathbf{p}^2}{8m^2}\right) \end{array}\right] \chi_i. \tag{137}
$$

Appendix D. Some useful identities and integrals

The following identity is useful for evaluating the M matrices:

$$
\frac{\left(\left(\mathbf{p}-\mathbf{q}\right)\cdot\mathbf{p}\right)^{2}}{\left(\mathbf{p}-\mathbf{q}\right)^{4}} = \frac{\mathbf{p}^{2}}{\left(\mathbf{p}-\mathbf{q}\right)^{2}} - \frac{\left(\mathbf{p}\times\mathbf{q}\right)^{2}}{\left(\mathbf{p}-\mathbf{q}\right)^{4}}.
$$
\n(138)

The angular integration in (47), (50), (55) involves the following integrals

$$
\int d\hat{\mathbf{p}} d\hat{\mathbf{q}} F(\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}) Y_{J'}^{m'_{J}}(\hat{\mathbf{q}}) Y_{J}^{m_{J}*}(\hat{\mathbf{p}}) = 2\pi \delta_{J'J} \delta_{m'_{J}m_{J}} \int d(\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}) F(\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}) P_{J}(\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}), \qquad (139)
$$

$$
\int d\left(\hat{\mathbf{p}}\cdot\hat{\mathbf{q}}\right) \frac{\hat{\mathbf{p}}\cdot\hat{\mathbf{q}}}{\left(\mathbf{p}-\mathbf{q}\right)^2} P_J\left(\hat{\mathbf{p}}\cdot\hat{\mathbf{q}}\right) = \frac{1}{|\mathbf{p}||\mathbf{q}|} \left(\frac{J+1}{2J+1} Q_{J+1}\left(z\right) + \frac{J}{2J+1} Q_{J-1}\left(z\right)\right),\tag{140}
$$

$$
\int d(\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}) \frac{(\mathbf{p} \times \mathbf{q})^2}{(\mathbf{p} - \mathbf{q})^4} P_J(\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}) = \frac{(J+1)(J+2)}{2(2J+1)} Q_{J+1}(z) - \frac{J(J-1)}{2(2J+1)} Q_{J-1}(z), \quad (141)
$$

where \mathcal{F} ($\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}$) is an arbitrary function of $\hat{\mathbf{p}} \cdot \hat{\mathbf{q}}$, $P_J(x)$ is the Legendre polynomial, and $Q_J(z)$ is the Legendre function of the second kind of order J.

The following integrals are needed for the calculation of the relativistic energy corrections.

$$
\int_0^\infty \int_0^\infty dp \, dq \, p^2 q^2 f^J(p) f^J(q) = 2\pi \left(\frac{\alpha \mu}{n}\right)^3 \delta_{J,0},\tag{142}
$$

$$
\int_0^\infty \int_0^\infty dp \, dq \, p q f^J(p) f^J(q) Q_J(z_1) = \frac{\pi \alpha \mu}{n^2},\tag{143}
$$

$$
\int_0^\infty \int_0^\infty dp \, dq \, p^2 q^2 f^J(p) f^J(q) Q_J(z_1) =
$$
\n
$$
\int_0^\infty \int_0^\infty dp \, dq \, p^3 q f^J(p) f^J(q) Q_J(z_1) = \pi \left(\frac{\alpha \mu}{n}\right)^3 \left(\frac{4}{2J+1} - \frac{1}{n}\right),
$$
\n(144)

$$
\int_0^\infty \int_0^\infty dp \, dq \, p^2 q^2 f^J(p) f^J(q) Q_{J-1}(z_1) = \pi \left(\frac{\alpha \mu}{n}\right)^3 \left(\frac{2}{J} - \frac{1}{n}\right),\tag{145}
$$

$$
\int_0^\infty \int_0^\infty dp \, dq \, p^2 q^2 f^J(p) f^J(q) Q_{J+1}(z_1) = \pi \left(\frac{\alpha \mu}{n}\right)^3 \left(\frac{2}{J+1} - \frac{1}{n}\right). \tag{146}
$$

Here f^J is the nonrelativistic hydrogen-like radial wave function in momentum space [19]

$$
f^{J}(p) \equiv f_{n}^{J}(p) = \left(\frac{2}{\pi} \frac{(n-J-1)!}{(n+J)!}\right)^{1/2} \frac{n^{J+2} p^{J} 2^{2(J+1)} J!}{(n^{2} p^{2} + 1)^{J+2}} \mathcal{G}_{n-J-1}^{J+1}\left(\frac{n^{2} p^{2} - 1}{n^{2} p^{2} + 1}\right),\tag{147}
$$

where G_{n-J}^{J+1} $_{n-J-1}^{J+1}(x)$ are Gegenbauer functions.

Appendix E. \mathcal{K}_{12} , \mathcal{K}_{21} kernels for $l = J \mp 1$ states

The contribution of the kernel K_{12} to the energy correction is

$$
\int dp \, dq \, p^2 q^2 \mathcal{K}_{12}(p,q) \, f^{J-1}(p) f^{J+1}(q),\tag{148}
$$

where

$$
\mathcal{K}_{12}(p,q) = \sum_{\sigma_1 \sigma_2 s_1 s_2} C_{Jm_J 12}^{s_1 s_2 \sigma_1 \sigma_2} \int d\hat{\mathbf{p}} d\hat{\mathbf{q}} \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2}^{ope(s-s)} (\mathbf{p}, \mathbf{q}) Y_{J+1}^{m_{\sigma_1 \sigma_2}}(\hat{\mathbf{q}}) Y_{J-1}^{m_{s_1 s_2}}(\hat{\mathbf{p}}). \tag{149}
$$

This requires the following integral

$$
\sum_{\sigma_1 \sigma_2 s_1 s_2} C_{Jm_J 12}^{s_1 s_2 \sigma_1 \sigma_2} \int d^3 \mathbf{p} \, d^3 \mathbf{q} \, f^{J-1}(p) Y_{J-1}^{m_{s_1 s_2}}(\hat{\mathbf{p}}) \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2}^{ope(s-s)}(\mathbf{p}, \mathbf{q}) \, f^{J+1}(q) Y_{J+1}^{m_{\sigma_1 \sigma_2}}(\hat{\mathbf{q}}). \tag{150}
$$

We calculate this form in coordinate space. The Fourier transform of $M_{s_1s_2\sigma_1\sigma_2}$ (**p**, **q**) is

$$
\mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{p}, \mathbf{q}) = \int d^3 \mathbf{r} \, d^3 \mathbf{r}' \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{r}, \mathbf{r}') \, e^{-i(\mathbf{p} - \mathbf{q}) \cdot (\mathbf{r} - \mathbf{r}')} , \tag{151}
$$

where the $M_{s_1s_2\sigma_1\sigma_2}(\mathbf{r},\mathbf{r}')$ matrix is a local operator in general [16], that is

$$
\mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{r}, \mathbf{r}') = \mathcal{M}_{s_1 s_2 \sigma_1 \sigma_2} (\mathbf{r}) \, \delta (\mathbf{r} - \mathbf{r}'). \tag{152}
$$

We apply this transformation to the $M_{s_1s_2\sigma_1\sigma_2}^{ope(s-s)}(\mathbf{p},\mathbf{q})$ matrix (see eq. (62)). Because of the angular integration in (149), only the first term in (62) survives. The Fourier transformation of that term is

$$
\frac{\left(\overrightarrow{\sigma}^{(+)}\cdot\left(\mathbf{p}-\mathbf{q}\right)\right)\left(\overrightarrow{\sigma}^{(-)}\cdot\left(\mathbf{p}-\mathbf{q}\right)\right)}{4m^2\left(\mathbf{p}-\mathbf{q}\right)^2} \rightarrow 3\frac{\left(\overrightarrow{\sigma}^{(+)}\cdot\mathbf{r}\right)\left(\overrightarrow{\sigma}^{(-)}\cdot\mathbf{r}\right)}{16\pi m^2 r^5}.
$$
\n(153)

Furthermore,

$$
\int d^3 \mathbf{p} f^{J-1}(p) Y_{J-1}^{m_{s_1 s_2}*}(\hat{\mathbf{p}}) e^{-i \mathbf{p} \cdot \mathbf{r}} = R_n^{J-1}(r) Y_{J-1}^{m_{s_1 s_2}*}(\hat{\mathbf{r}}), \tag{154}
$$

$$
\int d^3 \mathbf{q} f^{J+1}(q) Y_{J+1}^{m_{s_1 s_2}*}(\hat{\mathbf{q}}) e^{-i \mathbf{q} \cdot \mathbf{r}} = R_n^{J+1}(r) Y_{J+1}^{m_{s_1 s_2}}(\hat{\mathbf{r}}), \tag{155}
$$

where

$$
R_n^{\ell}(r) = -\frac{2}{n^2} \sqrt{\frac{(n-\ell-1)!}{((n+\ell)!)^3}} e^{-r/n} \left(\frac{2r}{n}\right)^{\ell} L_{n+\ell}^{2\ell+1}\left(\frac{2r}{n}\right).
$$
 (156)

The associated Laguerre function L^{μ}_{λ} $\mu_{\lambda}(\rho)$ is related to the confluent hypergeometric function by

$$
L^{\mu}_{\lambda}(\rho) = (-1)^{\mu} \frac{(\lambda!)^2}{\mu! (\lambda - \mu)!} F(-\lambda + \mu, \mu + 1; \rho).
$$
 (157)

The generating function for the Laguerre function is

$$
U_{\mu}(\rho, u) \equiv (-1)^{\mu} \frac{u^{\mu}}{(1-u)^{\mu+1}} \exp\left(-\frac{u\rho}{1-u}\right) = \sum_{\lambda=\mu}^{\infty} \frac{L_{\lambda}^{\mu}(\rho)}{\lambda!} u^{\lambda},\tag{158}
$$

hence

$$
\sum_{\sigma_{1}\sigma_{2}s_{1}s_{2}} C_{Jm_{J}12}^{s_{1}s_{2}\sigma_{1}\sigma_{2}} \int d^{3} \mathbf{p} d^{3} \mathbf{q} f^{J-1}(p) Y_{J-1}^{m_{s_{1}s_{2}}*}(\hat{\mathbf{p}}) \mathcal{M}_{s_{1}s_{2}\sigma_{1}\sigma_{2}}^{ope(s-s)}(\mathbf{p}, \mathbf{q}) f^{J+1}(q) Y_{J+1}^{m_{\sigma_{1}\sigma_{2}}}(\hat{\mathbf{q}})
$$
\n
$$
= \sum_{\sigma_{1}\sigma_{2}s_{1}s_{2}} C_{Jm_{J}12}^{s_{1}s_{2}\sigma_{1}\sigma_{2}} \int d^{3} \mathbf{r} R_{n}^{J-1}(r) Y_{J-1}^{m_{s_{1}s_{2}}*}(\hat{\mathbf{r}})
$$
\n
$$
\times \left(3\alpha \frac{(\overrightarrow{\sigma}^{(+)}\cdot\mathbf{r})(\overrightarrow{\sigma}^{(-)}\cdot\mathbf{r})}{16\pi m^{2}r^{5}}\right) R_{n}^{J+1}(r) Y_{J+1}^{m_{s_{1}s_{2}}}(\hat{\mathbf{r}})
$$
\n
$$
= \frac{3\alpha}{16\pi m^{2}} \int dr \, r^{2} \frac{1}{r^{3}} R_{n}^{J-1}(r) R_{n}^{J+1}(r) \times
$$
\n
$$
\times \sum_{\sigma_{1}\sigma_{2}s_{1}s_{2}} C_{Jm_{J}12}^{s_{1}s_{2}\sigma_{1}\sigma_{2}} \int d\hat{\mathbf{r}} Y_{J-1}^{m_{s_{1}s_{2}}*}(\hat{\mathbf{r}}) (\overrightarrow{\sigma}^{(+)}\cdot\hat{\mathbf{r}}) (\overrightarrow{\sigma}^{(-)}\cdot\hat{\mathbf{r}}) Y_{J+1}^{m_{s_{1}s_{2}}}(\hat{\mathbf{r}}).
$$
\n(159)

It follows that

$$
\sum_{\sigma_1 \sigma_2 s_1 s_2} C_{Jm_J 12}^{s_1 s_2 \sigma_1 \sigma_2} \int d\hat{\mathbf{r}} Y_{J-1}^{m_{s_1 s_2}}(\hat{\mathbf{r}}) \left(\vec{\sigma}^{(+)} \cdot \hat{\mathbf{r}}\right) \left(\vec{\sigma}^{(-)} \cdot \hat{\mathbf{r}}\right) Y_{J+1}^{m_{s_1 s_2}}(\hat{\mathbf{r}}) = \frac{1}{15} \frac{\sqrt{J(J+1)}}{2J+1},\tag{160}
$$

but

$$
\int_0^\infty dr \, r^2 \frac{1}{r^3} R_n^{J-1}(r) \, R_n^{J+1}(r) = 0. \tag{161}
$$

The last expression can be proved in the following way. Let us consider the more general case

$$
\int_0^\infty dr \, r^{\beta+2} R_n^{\ell}(r) \, R_n^{\ell'}(r) \,. \tag{162}
$$

The generating function for $R_n^{\ell}(r)$ is

$$
G_{n\ell}(r,u) = -\frac{2}{n^2} \sqrt{\frac{(n-\ell-1)!}{((n+\ell)!)^3}} e^{-r/n} \left(\frac{2r}{n}\right)^{\ell} (-1)^{2\ell+1} \frac{u^{2\ell+1}}{(1-u)^{2\ell+2}} \exp\left\{-\frac{u}{1-u} \frac{2r}{n}\right\}.
$$
 (163)

Then we consider the expression

$$
\int_{0}^{\infty} dr r^{\beta+2} G_{n\ell}(r, u) G_{n\ell'}(r, v)
$$
\n
$$
= \int_{0}^{\infty} dr r^{\beta+2} \frac{4}{n^4} \sqrt{\frac{(n-\ell-1)!(n-\ell'-1)!}{((n+\ell)!)^3 ((n+\ell')!)^3}} e^{-2r/n} \left(\frac{2r}{n}\right)^{\ell+\ell'} \times
$$
\n
$$
\times \frac{u^{2\ell+1}v^{2\ell'+1}}{(1-u)^{2\ell+2}(1-v)^{2\ell'+2}} \exp\left\{-\left(\frac{u}{1-u}+\frac{v}{1-v}\right)\frac{2r}{n}\right\}
$$
\n
$$
= \frac{4}{n^4} \sqrt{\frac{(n-\ell-1)!(n-\ell'-1)!}{((n+\ell)!)^3 ((n+\ell')!)^3}} \frac{u^{2\ell+1}v^{2\ell'+1}}{(1-u)^{2\ell+2}(1-v)^{2\ell'+2}} \times
$$
\n
$$
\times \int_{0}^{\infty} dr \left(\frac{2r}{n}\right)^{\beta+2+\ell+\ell'} \exp\left\{-\left(1+\frac{u}{1-u}+\frac{v}{1-v}\right)\frac{2r}{n}\right\}.
$$
\nand

It is well known that

$$
\int_0^\infty d\rho \rho^\beta e^{-\rho} = \Gamma(\beta + 1),\tag{165}
$$

therefore

$$
\int_0^\infty dr \left(\frac{2r}{n}\right)^{\beta+2+\ell+\ell'} \exp\left\{-\left(1+\frac{u}{1-u}+\frac{v}{1-v}\right)\frac{2r}{n}\right\}
$$

$$
= \left(\frac{n}{2}\right)^{\beta+3} \left(\frac{(1-u)(1-v)}{1-uv}\right)^{\beta+3+\ell+\ell'} \Gamma\left(\beta+3+\ell+\ell'\right) \tag{166}
$$

and

$$
\int_0^{\infty} dr r^{\beta+2} G_{n\ell}(r, u) G_{n\ell'}(r, v)
$$
\n
$$
= \frac{2^{-\beta-1}}{n^{-\beta+1}} \sqrt{\frac{(n-\ell-1)!(n-\ell'-1)!}{((n+\ell)!)^3 ((n+\ell')!)^3}} \times \frac{u^{2\ell+1} v^{2\ell'+1} (1-u)^{\beta+1-\ell+\ell'} (1-v)^{\beta+1+\ell-\ell'}}{(1-uv)^{\beta+3+\ell+\ell'}} \Gamma(\beta+3+\ell+\ell').
$$
\n(167)

We expand this expression in a series,

$$
\int_0^{\infty} dr r^{\beta+2} G_{n\ell}(r, u) G_{n\ell'}(r, v) = \sum_{\eta \eta'} C_{\eta \eta'}(n, \beta, \ell, \ell') u^{\eta} u^{\eta'}.
$$
 (168)

It is not difficult to show [20], that the coefficient $C_{n+\ell,n+\ell'}$ represents the integral

$$
C_{n+\ell,n+\ell'}\left(n,\beta,\ell,\ell'\right) = \int_0^\infty dr r^{\beta+2} R_n^{\ell}(r) R_n^{\ell'}(r). \tag{169}
$$

Simple but tedious calculations show that this coefficient is zero for $\beta = -3$, $\ell = J - 1$, $\ell' = J + 1$. Thus the kernel K_{12} does not contribute to the energy corrections to $O(\alpha^4)$. The same result is obtained for the kernel K_{21} .

References

1. J. W. Darewych, Annales Fond. L. de Broglie (Paris) 23, 15 (1998).

2. J. W. Darewych, in Causality and Locality in Modern Physics, G Hunter et al. (eds.), Kluwer, Dordrecht, p. 333, (1998).

3. J. W. Darewych, Can. J. Phys. 76, 523 (1998).

4. M. Barham and J. W. Darewych, J. Phys. A 31, 3481 (1998).

5. B. Ding and J. Darewych, J. Phys. G 26, 907 (2000).

6. J. D. Jackson, Classical Electrodynamics (John Wiley, New York, 1975).

7. A. O. Barut, Electrodynamics and Classical Theory of Fields and Particles (Dover, new York, 1980).

8. W. T. Grandy, Relativistic Quantum Mechanics of Leptons and Fields (Kluwer, Dordrecht, 1991).

9. A. O. Barut, in Geometrical and Algebraic Aspects of Nonlinear Field Theory, edited by S. De Filippo, M. Marinaro, G. Marmo and G. Vilasi, (Elsevier New York, 1989), p. 37.

10. J. W. Darewych and L. Di Leo, J. Phys. A 29, 6817 (1996).

11. H. Pilkuhn, J. Phys. B 28, 4421 (1995), H. Pilkuhn, Relativistic Quantum Mechanics (Springer 2003).

12. J. W. Darewych and M. Horbatsch, J. Phys. B 22, 973 (1989); 23, 337 (1990).

13. W. Dykshoorn and R. Koniuk, Phys. Rev. A 41, 64 (1990).

14. T. Zhang and R. Koniuk, Can. J. Phys. 70, 683 (1992).

15. T. Zhang and R. Koniuk, Can. J. Phys. 70, 670 (1992).

16. V. B. Berestetskii, E. M. Lifshitz, and L. P. Pitaevski, Quantum Electrodynamics (Pergamon, New York, 1982).

17. G. Arfken and H. Weber, Mathematical Methods for Physicists (Academic Press, 2001).

18. M. A. Stroscio, Positronium: A Review Of The Theory, Physics Reports C 22, 5-215 (1975).

19. H. A. Bethe and E. E. Salpeter, Quantum Mechanics of One and Two-Electron Atoms (Springer-Verlag/Academic, New York, 1957).

20. M. Mizushima, Quantum Mechanics of Atomic Spectra and Atomic Structure (W. A. Benjamin, Inc., New York, 1970).