

The Gravity dual of the Non-Perturbative $N = 2$ SUSY Yang-Mills Theory

Satchidananda Naik
Harish-chandra Research Institute
Chhatnag Road, Jhusi
Allahabad-211 019, INDIA

December 25, 2018

Abstract

The anomalous Ward identity is derived for $N = 2$ SUSY Yang-Mills theories, which is resulted out of Wrapping of D_5 branes on Supersymmetric two cycles. From the Ward identity One obtains the Witten-Dijkgraaf-Verlinde-Verlinde equation and hence can solve for the pre-potential. This way one avoids the problem of enhancon which maligns the non-perturbative behaviour of the Yang-Mills theory resulted out of Wrapped branes.

1. Introduction

Recently the gauge theory/gravity duality, which is commonly known as AdS/CFT duality is extended to non-conformal pure $\mathcal{N} = 1$ or $\mathcal{N} = 2$ supersymmetric Yang-Mills (SYM) theories [1]. So far the perturbative behaviour of the $\mathcal{N} = 2$ SYM is produced by this duality. The conventional folklore is that the instantons which are responsible for the non-perturbative part of the pre-potential are suppressed in the large N limit (since the gauge gravity duality is valid only in the large N limit). Also the non-perturbative strong coupling behaviour of SYM concerns the dilaton which is plagued with the singularity of "enhanced" [2]. Here we establish the anomalous super conformal Ward identity for $\mathcal{N} = 2$ SYM in the gravity dual picture. Further the super conformal Ward identity is written as Witten-Dijkgraaf-Verlinde-Verlinde (WDVV)[3] equation from which one can obtain the exact pre-potential.

2. The Strategy

We start with type IIB little string theory or a collection of a large number of NS_5 brane in the vanishing string coupling limit which gives rise to $D = 6$ SYM [4]. Then we dimensionally reduce two of its spatial world volume in such a way that we retain $\mathcal{N} = 2$ SYM in the low energy limit. The NS_5 brane has $SO(4)$ R-symmetry as the normal bundle. When one identifies the $U(1)$ subgroup of the $SO(4)$ R-symmetry with the $U(1)$ spin connection of the two cycle which is compactified, one gets a covariant constant spinor and SUSY is retained, which is commonly known as twisting [1]. This is called wrapping of NS_5 brane on a supersymmetric two cycles. If the compact space is a two-sphere, then there will be no extra hyper-multiplet and in the low energy limit i.e. in the scale much lower than the radius of the sphere we will get pure $\mathcal{N} = 2$ SYM. Thus it amounts to consider a gauged $D = 7$ supergravity solution and then lift it to get the solutions in ten dimensions. We use here the results of [5, 6] classical solutions of $D = 7$ gauged supergravity which is amenable to ten dimensional string theory.

$$ds_{10}^2 = e^\Phi \left[dx_{1,3}^2 + \frac{z}{\lambda^2} (d\theta^2 + \sin^2 \theta d\varphi^2) + \frac{1}{\lambda^2} e^{2x} dz^2 + \frac{1}{\lambda^2} \left(d\theta_1^2 + \frac{e^{-x}}{f(x)} \cos^2 \theta_1 (d\theta_2 + \cos \theta d\varphi)^2 + \frac{e^x}{f(x)} \sin^2 \theta_1 d\theta_3^2 \right) \right], \quad (1)$$

where the dilaton is

$$e^{2\Phi} = e^{2z} \left[1 - \sin^2 \theta_1 \frac{1 + c e^{-2z}}{2z} \right] \quad (2)$$

and

$$f(x) = e^x \cos^2 \theta_1 + e^{-x} \sin^2 \theta_1, \quad (3)$$

also

$$e^{-2x} = 1 - \frac{1 + c e^{-2z}}{2z} \quad (4)$$

where λ is the gauge coupling constant of seven dimensional gauged supergravity and c is a parameter as the integration constant of the classical solution. For $c \geq -1$ the range of the radial variable is $z_0 \leq z \leq \infty$ where z_0 is the solution for $e^{-2x(z_0)} = 0$. Here θ and φ are the angles of compact two-sphere with radius of compactification as $\frac{z}{\lambda^2}$ and θ_1, θ_2 and θ_3 are angles of transverse three-sphere. The conservation of the RR-charge on the transverse sphere S_3 fixes $\frac{1}{\lambda^2} = N g_s \alpha'$ for N number of D_5 branes with string coupling g_s . The D_5 brane action is given by

$$S = -\tau_5 \int d^6 \xi e^{-\Phi} \sqrt{-\det (G + 2\pi \alpha' F)} + \tau_5 \int \left(\sum_n C^{(n)} \wedge e^{2\pi \alpha' F} \right) \quad (5)$$

where F is the world volume gauge field and τ_5 is the brane tension. . The BPS condition is fixed from the condition of the vanishing of the potential between two branes which gives θ_1 to be $\frac{\pi}{2}$. This condition makes the transverse boundary of the D brane to be a two dimensional space consisting z and θ_3 which will eventually the moduli space of $\mathcal{N} = 2$ SYM.

We want to establish here the anomalous super conformal Ward identity. In the presence of gravity the trace anomaly

$$\langle \theta_\mu^\mu \rangle = \frac{1}{2} \frac{\beta(g)}{g^3} (F_{\mu\nu}^a)^2 + \frac{c(g^2)}{16\pi^2} (W_{\mu\nu\rho\sigma})^2 - \frac{a(g^2)}{16\pi^2} (\tilde{R}_{\mu\nu\rho\sigma})^2 \quad (6)$$

where $\beta(g)$ is the beta function of SYM, $a(g)$ and $c(g)$ are central functions near the criticality, $W_{\mu\nu\rho\sigma}$ is the Weyl tensor and $\tilde{R}_{\mu\nu\rho\sigma}$ is the dual of the curvature tensor. However this relation can be extracted from the two point functions of the energy momentum tensors[7]. In Ref.[8], it is shown how to extract these functions from the absorption cross-section of soft dilatons or gravitons by the D branes. The probability of absorption is taken as ratio of the flux near z_0 to the in coming flux at very large z . This gives

$$\sigma = \frac{N^4}{128\pi^3} (z - z_0)^2 \omega^3. \quad (7)$$

where ω is the frequency of the soft gluon. Here we see the presence of $(z - z_0)^2$ which if we write as in Ref.[6] $e^z = \rho$ we get a term $\left(\log \frac{\rho}{\rho_0}\right)^2$ in the cross-section signaling the asymptotic freedom or logarithmic coupling. Also we see the presence of enhancon when z_0 is zero. This gives $\beta(g) = -\frac{N}{8\pi^2} g^3$. Similarly one can also calculate from $U(1)$ R-current the chiral anomaly. Combining trace anomaly, chiral

anomaly and supertrace anomaly one can write the Ward identity as [9]

$$2\mathcal{F} - \mathcal{F}(\mathcal{A})' \mathcal{A} = \frac{N}{8\pi^2} \langle tr \psi^2 \rangle, \quad (8)$$

where \mathcal{F} is the effective potential or the pre-potential, \mathcal{A} is the chiral multiplet coupled to vector multiplet in the $\mathcal{N} = 2$ SYM and $tr \psi^2$ is the anomaly multiplet for example $tr F^2$ will correspond to θ_μ^μ . Here the bosonic component of \mathcal{A} corresponds to $z - i\theta_3$ or in radial coordinate $ve^{i\gamma}$. In the broken phase the branes will be distributed on a circle or \mathcal{A}_i the eigen values of \mathcal{A} which are $U(N)$ matrices will be distributed on a circle. The second part of eq.(8) will read as $\sum_i \frac{\partial \mathcal{F}}{\partial \mathcal{A}_i} \mathcal{A}_i$. This equation one can in principle solve in the large N limit and obtain the exact pre-potential in this limit [10].

References

- [1] J. Maldacena, Adv. Theor. Math. Phys. 2 (1998) 231, J.Maldacena and C. Nunez, Phys. Rev.Lett. 86,(2001)588.
- [2] C.V.Johnson,A.Peet and J.Polchinski, Phys. Rev. D 61(2000)086001
- [3] E. Witten, Nuc.Phys. B 340(1990)281, R. Dijkgraaf,E.Verlinde and H. Verlinde, Nuc.Phys. B 352(1991)59, B. Dubrovin, Lect. notes. hep-th-9407018.
- [4] N. Seiberg, Phys. Lett. B 408(1997)98
- [5] J. Gauntlet,N.Kim,D.Martelli , D. Waldram, Phys.Rev. D 64 (2001) 106008, F.Bigazzi,A.Cotrone, A. Zaffaroni, Phys.Lett.B 519(2001)269.
- [6] M.Bertolini,P. Di Vecchia, M. Frau, A.Lerda, R. Marotta, I. Pesando, JHEP 0102 (2001)014, P. Di Vecchia, A.Lerda and P. Merlatti, Nucl.Phys.B646 (2002) 43, P. Di Vecchia, hep-th-0212162.
- [7] D. Anselmi, JHEP 9805(1998)005, D. Anselmi,D.Z. Freedman, M. Grisaru, A. Johansen, Nucl. Phys. B 526(1998) 543.
- [8] S.S. Gubser and I.R. Klebanov, Phys.Lett. B 413(1997)41.
- [9] Marco Matone, Phys. Lett. B 357 (1995)342,G.Bonelli and M. Matone Phys. Rev. Lett. 77 (1996)4712, F.Fucito and G. Travaglini,Phys.Rev. D 55(1997)1099, N.Dorey,V.Khoze,M.Mattis, Phys.Lett. B390 (1997)205, P.S. Howe and P.C.West, Phys.Lett.B 400(1997)30.

- [10] M.R. Douglas and S. Shenker, Nucl.Phys.B447 (1995) 271, F. Ferrari, Nucl.Phys.B 612 (2001)151.