

Path Integral Techniques in Conformal Field Theory

André van Tonder

Department of Physics, Brown University
Box 1843, Providence, RI 02906
andre@het.brown.edu

December 1, 2004

Abstract

We present the theory of a two-dimensional conformal scalar field using path integral techniques. We derive the conformal anomaly using an adaptation of the method of Fujikawa, and compare the result with a derivation based on a Pauli-Villars measure, where the anomaly is shifted from the path integral measure to the energy-momentum trace. In the path integral approach the energy-momentum is a true coordinate-invariant tensor quantity, and we explain how it is related to the corresponding non-tensor object arising in the operator approach, obtaining an intuitive explanation within the context of the path integral approach for the anomalous transformation law and anomalous Ward identities of the latter. After carefully calculating nontrivial contact terms arising in certain energy-momentum products, we use these to provide a simple consistency check confirming the change of variables formula for the path integral and to review the relationship between the conformal anomaly and the energy-momentum two-point functions.

Keywords: Conformal Field Theory, Path Integral, Anomaly
Brown preprint: BROWN-HET-1429

1 Introduction

We present the theory of a two-dimensional conformal scalar field using path integral techniques.

While many presentations of conformal field theory use path integrals informally, most emphasize operator methods or the axiomatic approach for nontrivial calculations. Here we attempt a somewhat rigorous path-integral based introduction to a simple conformal theory, aimed at those who may be more familiar with the path integral approach to field theory, or who wish to understand the relationship between the different approaches. As we show, the relationship is in some respects rather subtle.

The path integral is particularly useful for the insight it provides into anomalies. We present a derivation of the conformal anomaly based on the method of Fujikawa, which exhibits the anomaly as a dependence of the path integral measure on the metric. We also provide a Pauli-Villars derivation of the anomaly. It turns out that the Pauli-Villars path integral measure is invariant under conformal transformations, and the anomaly is shifted to the expectation value of the trace of the energy-momentum tensor, which becomes nonzero.

A serious pitfall in relating the path integral and operator approaches is that equations obtained using path integrals, such as Ward identities, often become invalid upon naive replacement of bare quantities by the corresponding renormalized objects appearing in the operator formalism. Because this may seem to call into question some of the very properties of path integration, such as invariance under change of variables, that are used to justify the Ward identities in the first place, a more careful analysis is necessary. We resolve the problem by carefully defining the path integral measure so that the expectation values appearing in these identities are all finite. To do this, we use a Pauli-Villars regularization, which is a physically intuitive, coordinate invariant and nonperturbative regularization technique that is largely insensitive to the details of the continuum limit.

We point out that there is a subtle relationship between the energy-momentum tensor in the path-integral formalism and the corresponding quantity in the operator approach. In the path integral approach, the natural energy-momentum tensor is a true coordinate-invariant quantity. We show how it is related to the corresponding nontensor object found in the operator approach, obtaining an intuitive explanation for the anomalous behaviour of the latter within the context of the path integral. The difference in be-

haviour turns out to be due to a residual effect of the measure as encoded by the Pauli-Villars fields.

We confirm by explicit calculation that with the appropriate Pauli-Villars measure, the full energy-momentum tensor satisfies the classical Ward identities. We explain how these relate to the anomalous Ward identities found in the operator approach. We conclude with a simple consistency check confirming the validity of the change of variables formula for the path integral, and we review the relationship between the conformal anomaly and the energy-momentum two-point functions. Important in our derivations and checks are certain contact terms arising in expectation values of energy-momentum tensors. While these have been derived in the axiomatic approach, we provide an explicit first-principles calculation.

This paper reviews and clarifies known results in conformal field theory from a somewhat different point of view. Various of these results have not, to our knowledge, been calculated before using these methods.

2 The conformal anomaly

In this section we derive the conformal anomaly using an adaptation of the methods of Fujikawa [1, 2, 3], who showed that anomalies have a very intuitive origin as a non-invariance of the path integral measure under classical symmetries. In the next section, we provide an independent derivation of the anomaly using a Pauli-Villars definition of the path integral measure, where we show that the origin of the anomaly is shifted from the measure to the expectation value of the trace of the energy-momentum tensor.

We are interested in the dependence of the partition function

$$Z(g) \equiv \int [d\phi]_g e^{-S(g,\phi)}$$

on the background metric g_{ij} . Here $[d\phi]_g$ denotes the ∞ -form

$$[d\phi]_g \equiv \bigwedge_{n=0}^{\infty} da_n^g = da_0^g \wedge da_1^g \wedge \dots,$$

where the coefficients $a_n^g : \phi \rightarrow \mathbf{R}$ depend on the metric via the expansion

$$\phi(z) = \sum_{n=0}^{\infty} a_n^g \phi_n^g(z),$$

and where $\phi_n^g(z)$ denotes an orthonormal basis of field configurations satisfying

$$\int d^2x \sqrt{g} \phi_m^g(x) \phi_n^g(x) = \delta_{mn}. \quad (1)$$

To calculate the variation $\delta_g[d\phi]_g$ of the path integral measure with respect to deformations of the metric, we will need the variation

$$\delta_g a_m^g = \delta_g \int d^2x \sqrt{g} \phi_m^g \phi \quad (2)$$

$$= \int d^2x \sqrt{g} \left(\frac{1}{2} g^{ij} \delta g_{ij} \right) \phi_m^g \phi + \int d^2x \sqrt{g} (\delta \phi_m^g) \phi. \quad (3)$$

The metric dependence of ϕ_m^g is not uniquely fixed by the above orthonormality requirement. However, given a metric g , any two orthonormal bases are related by a unitary transformation that will leave the form $[d\phi]_g$ invariant. Therefore we can, without loss of generality, choose one particular metric dependence for ϕ_m^g compatible with orthonormality. To find such a choice, we vary both sides of (1) with respect to the metric, obtaining

$$0 = \int d^2x \sqrt{g} \left(\frac{1}{2} g^{ij} \delta g_{ij} \right) \phi_m^g \phi_n^g + \int d^2x \sqrt{g} (\delta_g \phi_m^g) \phi_n^g + \int d^2x \sqrt{g} \phi_m^g (\delta_g \phi_n^g).$$

A suitable choice for $\delta_g \phi_m^g$ is therefore

$$\delta_g \phi_m^g \equiv -\frac{1}{4} g^{ij} \delta g_{ij} \phi_m^g.$$

Inserting this in (3) gives

$$\begin{aligned} \delta_g a_m^g &= \int d^2x \sqrt{g} \left(\frac{1}{4} g^{ij} \delta g_{ij} \right) \phi_m^g \phi \\ &= \sum_n a_m^g \cdot \int d^2x \sqrt{g} \left(\frac{1}{4} g^{ij} \delta g_{ij} \right) \phi_m^g \phi_n^g \\ &\equiv \sum_n a_m^g C_{mn}, \end{aligned}$$

where

$$C_{mn} \equiv \int d^2x \sqrt{g} \phi_m^g \left(\frac{1}{4} g^{ij} \delta g_{ij} \right) \phi_n^g$$

We then find, using the normal rules for manipulating differential forms,

$$\delta_g[d\phi]_g \equiv \delta_g \left(\bigwedge_n da_n^g \right) \quad (4)$$

$$= \sum_m \cdots \wedge da_{m-1}^g \wedge \delta(da_m^g) \wedge da_{m+1}^g \wedge \cdots \quad (5)$$

$$= \sum_m \cdots \wedge da_{m-1}^g \wedge \left(\sum_n da_n^g C_{nm} \right) \wedge da_{m+1}^g \wedge \cdots \quad (6)$$

$$= \left(\sum_m C_{mm} \right) \bigwedge_n da_n^g \quad (7)$$

$$= (\text{Tr } C)_g [d\phi]_g, \quad (8)$$

where C is the operator

$$C \equiv \frac{1}{4} g^{ij} \delta g_{ij} = \frac{1}{2} \frac{1}{\sqrt{g}} \delta \sqrt{g} \equiv \delta \omega.$$

Notice that $\delta \sqrt{g}$ is the local change of volume. In other words, the path integral measure will be scale dependent. This is the origin of the conformal anomaly.

We now need to calculate the trace

$$(\text{Tr } C)_g = \int d^2x \sqrt{g} \delta \omega(x) A(x),$$

where the infinite sum

$$A(x) \equiv \sum_m \phi_m^g(x) \phi_m^g(x)$$

does not in general converge. A natural short-distance regularization, which can be taken as part of the definition of the path integral, is obtained by considering instead the limit as $\epsilon \rightarrow 0$ of the sum [4, 5]

$$\sum_m \phi_m^g(x) e^{\epsilon \Delta} \phi_m^g(x) = \langle x | e^{\epsilon \Delta} | x \rangle$$

where Δ denotes the Laplacian

$$\Delta \equiv \frac{1}{\sqrt{g}} \partial_i [\sqrt{g} g^{ij} \partial_j (\cdot)].$$

Here the position basis bras and kets are defined via $\langle x | y \rangle \equiv \delta(x-y)/\sqrt{g}$, to make $\langle x | \phi \rangle = \phi(x)$ with respect to the inner product $\langle \phi_1 | \phi_2 \rangle = \int \sqrt{g} \phi_1 \phi_2$.

As required for consistency with our previous remark that a unitary transformation leave $[d\phi]_g$ invariant, the regularized expression $\langle x | e^{\epsilon\Delta} | x \rangle$ depends only on the metric and not on the particular orthonormal basis ϕ_m^g that we have chosen. Also notice that if we insert an eigenbasis $|m\rangle$ of Δ , the above becomes

$$\langle x | e^{\epsilon\Delta} | x \rangle = \sum_m \langle m | x \rangle \langle x | m \rangle e^{\epsilon \lambda_m}$$

where the λ_m are the corresponding (negative) eigenvalues, making it clear that the regularization corresponds to a large-momentum suppression.

We therefore need to calculate the small t behaviour of the function

$$G(x, t|y, 0) \equiv \theta(t) \langle x | e^{t\Delta} | y \rangle,$$

where $\theta(t)$ denotes the step function, inserted to avoid the regime $t < 0$ where $\langle x | e^{t\Delta} | y \rangle$ diverges. It is easy to check that

$$(\partial_t - \Delta_x) G(x, t|y, 0) = \delta(t) \langle x | y \rangle = \frac{1}{\sqrt{g}} \delta(t) \delta(x - y).$$

In other words, the distribution $G(x, t|y, 0)$ solves the heat or diffusion equation given a point source at y at time $t = 0$, and is known as a *heat kernel*. Without loss of generality, we can choose coordinates so that $y = 0$. In two dimensions, we can further choose the coordinate system so that

$$\begin{aligned} g_{ij}(x) &= e^{2\omega} \delta_{ij}, \\ \omega(0) &= 0. \end{aligned}$$

so that the equation for $G(x, t|y, 0)$ becomes

$$(\partial_t - e^{-2\omega(x)} \Delta_0) G(x, t|0, 0) = \delta(t) \delta(x),$$

where $\Delta_0 \equiv \delta^{ij} \partial_i \partial_j$. Regarding this as an operator equation in the space $\mathcal{L}^2(\mathbf{R}^3)$, we write the solution as

$$G(x, t|0, 0) = \langle x, t | \left(\frac{1}{\partial_t - e^{-2\omega(x)} \Delta_0} \right) | 0, 0 \rangle.$$

Since we are interested in the limit as $t \rightarrow 0$, for which the diffusion becomes increasingly short-ranged, we will develop an expansion for $G(0, t|0, 0)$ in terms of derivatives of ω at the origin. Expanding around $x^i = 0$, we have

$$e^{-2\omega(x)} = 1 - 2(\partial_i \omega) x^i + [-(\partial_i \partial_j \omega) + 2(\partial_i \omega)(\partial_j \omega)] x^i x^j + \dots,$$

where the derivatives are all evaluated at $x^i = 0$. We can then write [4]

$$G = \frac{1}{A - B} = \frac{1}{A} + \frac{1}{A} B \frac{1}{A} + \frac{1}{A} B \frac{1}{A} B \frac{1}{A} + \dots$$

where $A \equiv \partial_t - \Delta_0$, so that $1/A$ is the flat space solution

$$\langle x, t | \left(\frac{1}{A} \right) | 0, 0 \rangle = \frac{1}{4\pi t} e^{-x^2/4t} \theta(t)$$

while

$$B \equiv \left\{ -2(\partial_i \omega) x^i + [-(\partial_i \partial_j \omega) + 2(\partial_i \omega)(\partial_j \omega)] x^i x^j + \dots \right\} \Delta_0$$

Inserting this expansion in the first-order contribution $A^{-1} B A^{-1}$ to $G(0, \epsilon|0, 0)$, the first term, with odd integrand proportional to x^i , vanishes. The next term in $A^{-1} B A^{-1}$ is proportional to

$$\begin{aligned} & \int_0^\epsilon dt \int d^2 x \frac{1}{4\pi(\epsilon-t)} e^{-x^2/4(\epsilon-t)} x^i x^j \Delta_0 \frac{1}{4\pi t} e^{-x^2/4t} \\ &= \int_0^\epsilon dt \int d^2 x \frac{1}{4\pi(\epsilon-t)} \frac{1}{4\pi t} e^{-x^2/4(\epsilon-t)} x^i x^j \left[-\frac{1}{t} + \frac{x^2}{4t^2} \right] e^{-x^2/4t} \\ &= \frac{1}{2} \delta^{ij} \int_0^\epsilon dt \int d^2 x \frac{1}{4\pi(\epsilon-t)} \frac{1}{4\pi t} e^{-x^2/4(\epsilon-t)} x^2 \left[-\frac{1}{t} + \frac{x^2}{4t^2} \right] e^{-x^2/4t} \\ &= \frac{1}{2} \delta^{ij} \int_0^\epsilon dt \int d^2 x \frac{1}{4\pi(\epsilon-t)} \frac{1}{4\pi t} x^2 \left[-\frac{1}{t} + \frac{x^2}{4t^2} \right] e^{-\epsilon x^2/4t(\epsilon-t)}. \end{aligned}$$

Writing the terms containing x^2 and x^4 as derivatives with respect to $\lambda \equiv \epsilon/4t(\epsilon-t)$ of the Gaussian integral

$$\int d^2 x e^{-\lambda x^2} = \frac{\pi}{\lambda},$$

this simplifies to

$$\frac{1}{2\pi} \delta^{ij} \int_0^\epsilon dt \left\{ -\frac{(\epsilon-t)}{\epsilon^2} + \frac{2(\epsilon-t)^2}{\epsilon^3} \right\} = \frac{1}{12\pi} \delta^{ij}$$

The next nonzero term in the contribution $A^{-1}BA^{-1}$ has integrand proportional to $x^i x^j x^k x^l$. This may be checked to be of order ϵ . Further terms are of even higher order in ϵ , so that the contribution to G from the term $A^{-1}BA^{-1}$ can be written

$$G_1(0, \epsilon|0, 0) = \frac{1}{12\pi} [-\Delta\omega + 2 \partial_i \omega \partial^i \omega] + o(\epsilon).$$

There is one additional contribution of order ϵ^0 to $G(0, \epsilon|0, 0)$. It comes from the second-order term $A^{-1}BA^{-1}BA^{-1}$ in the above expansion. It is

$$4 (\partial_i \omega) (\partial_j \omega) \int_0^\epsilon dt \int_0^t du \int d^2x \int d^2y \times \\ \times \frac{1}{4\pi(\epsilon-t)} e^{-x^2/4(\epsilon-t)} x^i \Delta_0^x \frac{1}{4\pi(t-u)} e^{-(x-y)^2/4(t-u)} y^j \Delta_0^y \frac{1}{4\pi u} e^{-y^2/4u}$$

By similar manipulations, this becomes

$$-\frac{1}{6\pi} \partial_i \omega \partial^i \omega,$$

so that

$$G_2(0, \epsilon|0, 0) = -\frac{1}{6\pi} \partial_i \omega \partial^i \omega + o(\epsilon)$$

Notice that this contributes a term that exactly cancels the term of the same form in $G^2(0, \epsilon|0, 0)$. We find the result

$$G(0, \epsilon|0, 0) = \frac{1}{4\pi\epsilon} - \frac{1}{12\pi} \Delta\omega + o(\epsilon) \\ = \frac{1}{4\pi\epsilon} + \frac{1}{24\pi} R + o(\epsilon),$$

where we have used

$$R = -2 e^{-2\omega} \Delta\omega$$

and $\omega(o) = 0$. Inserting our result into the formula for the variation of the measure, we obtain

$$\delta_g [d\phi]_g = \frac{1}{4\pi\epsilon} \int d^2x \sqrt{g} \delta\omega(x) + \frac{1}{24\pi} \int d^2x \sqrt{g} \delta\omega(x) R.$$

The first term diverges as $\epsilon \rightarrow 0$, but may be exactly canceled by adding the counterterm

$$\frac{1}{2\pi\epsilon} \int d^2x \sqrt{g}$$

to the original action. After doing this, we obtain our final result

$$\delta_g \int [d\phi]_g e^{-S(\phi,g)} = \left(\frac{1}{24\pi} \int d^2x \sqrt{g} \delta\omega(x) R \right) \int [d\phi]_g e^{-S(\phi,g)}. \quad (9)$$

3 A Pauli-Villars derivation of the anomaly

In the previous section the conformal anomaly was obtained from the dependence of the path integral measure on the metric. In the case of a Weyl transformation $g_{ij} \rightarrow e^{2\omega} g_{ij}$, we can absorb the anomalous dependence of the measure into the energy-momentum tensor by defining

$$\begin{aligned} \delta \int [d\phi]_g e^{-S(g,\phi)} &\equiv \frac{1}{4\pi} \int d^2x \sqrt{g} \delta g^{ij} \langle \tilde{T}_{ij} \rangle_g \\ &= \frac{1}{4\pi} \int d^2x \sqrt{g} (-2 \delta\omega) \langle \tilde{T}^i_i \rangle_g, \end{aligned}$$

where

$$T_{ij} \equiv -2\pi \left(\partial_i \phi \partial_j \phi - \frac{1}{2} g_{ij} g^{kl} \partial_k \phi \partial_l \phi \right)$$

and

$$\tilde{T}^i_i \equiv T^i_i - \frac{1}{12} R = -\frac{1}{12} R,$$

since T^i_i is identically zero. The curvature term was entirely due to the variation $\delta[d\phi]_g$ of the integration measure.

It is very instructive to derive the same result using a Pauli-Villars regularization [6, 7, 8]. This was first done by Vilenkin in [9]. For related work, including comparisons of the Pauli-Villars method to the ζ -function and point-splitting methods, see for example [10, 11, 12, 13].

With a Pauli-Villars definition of the path integral, we shall see that the integration measure will in fact be invariant under variations of g , and the necessary curvature term will instead be due to the fact that the Pauli-Villars fields are massive, so that $T^i_i \neq 0$, and we shall indeed find that

$$\langle T^i_i \rangle_g \rightarrow -\frac{1}{12} R$$

in the limit where the Pauli-Villars masses go to infinity.

In our following discussions, we will have need to carefully regulate the theory in both the infrared and ultraviolet. For our infrared regularization we give a small mass m to the field ϕ . The ultraviolet regularization will consist in adding auxiliary scalar Pauli-Villars fields χ_i with masses M_i and statistics chosen to make the path integral converge. In the end, we will be interested in the limits $m \rightarrow 0$ where the field ϕ becomes massless, and $M_i \rightarrow \infty$ where the Pauli-Villars fields become non-dynamical.

First consider the variation

$$\delta_g \left([d\phi]_g \wedge [d\chi_1]_g \wedge \cdots \wedge [d\chi_n]_g \right).$$

As in the previous section, each real scalar field χ_i will contribute a variation term of the same form as that contributed by ϕ . The contribution of Grassmann scalar fields χ_i will be of opposite sign. Indeed, the reader may easily check that with Grassmann integration rules, if $\delta\eta = \epsilon\eta$ where ϵ is a real parameter, then for the change of variables formula $\int d\eta' f(\eta') = \int d\eta f(\eta)$ to hold (equivalent to $\delta \int d\eta f(\eta) = 0$) one needs $\delta(d\eta) \equiv -\epsilon d\eta$, so that in formula (8) of our derivation of the previous section we get traces of opposite sign for Grassmann fields. Defining $c_i = 1$ for χ_i real commuting, and $c_i = -1$ for χ_i Grassmann anticommuting, we find

$$\begin{aligned} \delta_g \left([d\phi]_g \wedge [d\chi_1]_g \wedge \cdots \wedge [d\chi_n]_g \right) \\ = \left(1 + \sum_i c_i \right) \left(\frac{1}{4\pi\epsilon} \int d^2x \sqrt{g} \delta\omega(x) + \frac{1}{24\pi} \int d^2x \sqrt{g} \delta\omega(x) R \right) \times \\ \times \left([d\phi]_g \wedge [d\chi_1]_g \wedge \cdots \wedge [d\chi_n]_g \right) \end{aligned}$$

We therefore can choose our Pauli-Villars field statistics to cancel the variation of $[d\phi]_g$. In particular, if we choose

$$1 + \sum_i c_i = 0, \tag{10}$$

the variation of the measure is zero

$$\delta_g \left([d\phi]_g \wedge [d\chi_1]_g \wedge \cdots \wedge [d\chi_n]_g \right) = 0.$$

Since the measure is now invariant, the anomaly will have to be due entirely to the expectation value of the trace of the energy-momentum tensor

$$\langle T^i_i \rangle_g = 2\pi \left\langle m^2 \phi^2 + \sum_i M_i^2 \chi_i^2 \right\rangle_g.$$

Since the propagator of a Grassmann scalar has the opposite sign to that of ϕ , we find in flat space

$$\langle T^i_i \rangle_\delta = 2\pi \int \frac{d^2 p}{(2\pi)^2} \left\{ \frac{m^2}{p^2 + m^2} + \sum_i c_i \frac{M_i^2}{p^2 + M_i^2} \right\}, \quad (11)$$

where $c_i = 1$ for Grassmann fields and $c_i = -1$ for real fields. As long as we impose the condition

$$m^2 + \sum_i c_i M_i^2 = 0, \quad (12)$$

on the Pauli-Villars masses, the integral converges. The answer in this case is

$$\begin{aligned} & -\frac{1}{4\pi} \lim_{\Lambda \rightarrow \infty} \left(m^2 \ln \frac{\Lambda^2}{m^2} + \sum_i c_i M_i^2 \ln \frac{\Lambda^2}{M_i^2} \right) \\ & = \frac{1}{4\pi} \left(m^2 \ln \frac{m^2}{\mu^2} + \sum_i c_i M_i^2 \ln \frac{M_i^2}{\mu^2} \right), \end{aligned}$$

where the limit is finite due to the Pauli-Villars condition. Here μ is an arbitrary parameter with dimension of mass, irrelevant due to the first Pauli-Villars condition (10). As $m \rightarrow 0$, the first term in (11) falls away and we get

$$\frac{1}{4\pi} \sum_i c_i M_i^2 \ln \frac{M_i^2}{\mu^2},$$

This diverges in the limit $M_i \rightarrow \infty$ but can be exactly compensated by a counterterm of the form

$$\frac{1}{(4\pi)^2} \sum_i c_i M_i^2 \ln \frac{M_i^2}{\mu^2} \int d^2 x \sqrt{g}$$

in the original action. We could also avoid the need for the counterterm by imposing the additional Pauli-Villars condition

$$m^2 \ln \frac{m^2}{\mu^2} + \sum_i c_i M_i^2 \ln \frac{M_i^2}{\mu^2} = 0. \quad (13)$$

If we do this, note that to satisfy the three Pauli-Villars conditions (10), (12) and (13), while keeping the ability to take the masses $M_i \rightarrow \infty$, we will need at least five auxiliary fields. If we use exactly five, two of these should be commuting and three anticommuting.

We now consider the case of a curved space, where we need to calculate the limit

$$\langle T_i^i \rangle_g = 2\pi \lim_{\Lambda \rightarrow \infty} \left\{ m^2 G_{m,\Lambda}(x, x) + \sum_i M_i^2 G_{M_i,\Lambda}(x, x) \right\}.$$

Here $G_{m,\Lambda}(x, x)$ denotes the two-point function of a scalar field computed with large-momentum cutoff Λ . The Pauli-Villars conditions are precisely what are needed to make the limit $\Lambda \rightarrow \infty$ finite.

In a background g_{ij} , the propagator $G_M(x, x')$ satisfies

$$\partial_i [\sqrt{g} g^{ij} \partial_j G_M(x, x')] - \sqrt{g} M^2 G_M(x, x') = \delta(x - x').$$

Given x' , we can change coordinates so that x' lies at the origin. In two dimensions, we can furthermore choose the coordinate system so that

$$\begin{aligned} g_{ij}(x) &= e^{2\omega} \delta_{ij}, \\ \omega(0) &= 0. \end{aligned}$$

so that the equation for the propagator becomes

$$\begin{aligned} (e^{-2\omega(x)} \Delta - M^2) G_M(x, 0) &= e^{-2\omega(x)} \delta(x) \\ &= \delta(x), \end{aligned}$$

where $\Delta \equiv \delta^{ij} \partial_i \partial_j$. As we increase the Pauli-Villars mass M , the propagator becomes increasingly short-ranged, and we will develop an expansion for $G_M(x)$ in terms of derivatives of ω at the origin. Expanding around $x^i = 0$, we have

$$e^{-2\omega(x)} = 1 - 2(\partial_i \omega) x^i + [-(\partial_i \partial_j \omega) + 2(\partial_i \omega)(\partial_j \omega)] x^i x^j + \dots,$$

where the derivatives are all evaluated at $x^i = 0$. We can then write

$$G_M = \frac{1}{A-B} = \frac{1}{A} + \frac{1}{A} B \frac{1}{A} + \frac{1}{A} B \frac{1}{A} B \frac{1}{A} + \dots,$$

where $A \equiv \Delta - M^2$, so that $1/A$ is the flat space propagator, while

$$B \equiv \{2(\partial_i \omega) x^i + [(\partial_i \partial_j \omega) - 2(\partial_i \omega)(\partial_j \omega)] x^i x^j + \dots\} \Delta.$$

In the first-order contribution $A^{-1} B A^{-1}$ to $G_M(0, 0)$, the term with odd integrand proportional to x^i vanishes. The next term in $A^{-1} B A^{-1}$ is proportional to

$$\begin{aligned} & \int d^2 x x^i x^j \int \frac{d^2 k}{(2\pi)^2} \frac{e^{ikx}}{k^2 + M^2} \int \frac{d^2 p}{(2\pi)^2} \frac{p^2 e^{-ipx}}{p^2 + M^2} \\ &= - \int d^2 x \int \frac{d^2 k}{(2\pi)^2} \int \frac{d^2 p}{(2\pi)^2} \frac{1}{k^2 + M^2} \frac{p^2}{p^2 + M^2} \partial_{k_i} \partial_{k_j} e^{i(k-p)x} \\ &= -(2\pi)^2 \int \frac{d^2 k}{(2\pi)^2} \int \frac{d^2 p}{(2\pi)^2} \frac{1}{k^2 + M^2} \frac{p^2}{p^2 + M^2} \partial_{k_i} \partial_{k_j} \delta(k-p) \\ &= - \int \frac{d^2 k}{(2\pi)^2} \left\{ \frac{-2\delta^{ij}}{(k^2 + M^2)^2} + \frac{8k_i k_j}{(k^2 + M^2)^3} \right\} \frac{k^2}{k^2 + M^2} \\ &= -\delta^{ij} \int \frac{d^2 k}{(2\pi)^2} \left\{ \frac{-2}{(k^2 + M^2)^2} + \frac{4k^2}{(k^2 + M^2)^3} \right\} \frac{k^2}{k^2 + M^2} \\ &= -2\delta^{ij} \int \frac{d^2 k}{(2\pi)^2} \frac{k^2 (k^2 - M^2)}{(k^2 + M^2)^4} \\ &= -\frac{1}{12\pi} \frac{1}{M^2} \delta^{ij}. \end{aligned}$$

The next nonzero term in $A^{-1} B A^{-1}$ has integrand proportional to $x^i x^j x^k x^l$. This may be checked to be of order $1/M^4$. Further terms are of even higher order in $1/M$, so that the full first-order contribution is

$$G_{M,1}(0, 0) = -\frac{1}{12\pi} \frac{1}{M^2} [\Delta\omega - 2\partial_i \omega \partial^i \omega] + o\left(\frac{1}{M^4}\right),$$

which will contribute a term of zeroth order in $1/M^2$ to the expectation value $\langle T^i_i \rangle_g$.

There is one additional contribution of order $1/M^2$ to $G_M(0, 0)$. It comes from the second-order term $A^{-1}BA^{-1}BA^{-1}$ in the above expansion. It is

$$4 (\partial_i \omega) (\partial_j \omega) \int d^2x \int d^2y x^i y^j \times \\ \times \int \frac{d^2k}{(2\pi)^2} \frac{e^{ikx}}{k^2 + M^2} \int \frac{d^2p}{(2\pi)^2} \frac{p^2 e^{ip(y-x)}}{p^2 + M^2} \int \frac{d^2q}{(2\pi)^2} \frac{q^2 e^{-iqy}}{q^2 + M^2}.$$

By similar manipulations, this becomes

$$-8M^2 \delta^{ij} (\partial_i \omega) (\partial_j \omega) \int \frac{d^2k}{(2\pi)^2} \frac{k^4}{(k^2 + M^2)^5} = -\frac{1}{6\pi} \frac{1}{M^2} \partial_i \omega \partial^i \omega,$$

so that

$$G_{M,2}(0, 0) = -\frac{1}{6\pi} \frac{1}{M^2} \partial_i \omega \partial^i \omega + o\left(\frac{1}{M^4}\right)$$

Notice that this contributes a term that exactly cancels the term of the same form in $G_M^1(0, 0)$. We find

$$G_M^\Lambda(0, 0) = -\frac{1}{4\pi} \ln \frac{\Lambda^2}{M^2} - \frac{1}{12\pi} \frac{1}{M^2} \Delta \omega + o\left(\frac{1}{M^4}\right) \\ = -\frac{1}{4\pi} \ln \frac{\Lambda^2}{M^2} + \frac{1}{24\pi} \frac{1}{M^2} R + o\left(\frac{1}{M^4}\right).$$

The matter contribution can be expanded in m by writing the equation for the propagator as

$$(\Delta - e^{2\omega(x)} m^2) G_m(x, 0) = \delta(x),$$

so that

$$G_m = \frac{1}{A - B} = \frac{1}{A} + \frac{1}{A} B \frac{1}{A} + \frac{1}{A} B \frac{1}{A} B \frac{1}{A} + \dots$$

where $A \equiv \Delta - m^2$ and

$$B \equiv m^2 (1 - e^{2\omega(x)}) \equiv m^2 \gamma(x).$$

Therefore

$$\begin{aligned}
G_m^\Lambda(0,0) &= -\frac{1}{4\pi} \ln \frac{\Lambda^2}{m^2} \\
&+ m^2 \int d^2x \frac{1}{2\pi} K_0(m|x|) \gamma(x) \frac{1}{2\pi} K_0(m|x|) \\
&+ m^4 \int d^2x \int d^2y \frac{1}{2\pi} K_0(m|x|) \gamma(x) \frac{1}{2\pi} K_0(m|y-x|) \gamma(y) \frac{1}{2\pi} K_0(m|y|) \\
&+ \dots,
\end{aligned}$$

where the Bessel function decays exponentially for large x and

$$2\pi K_0(m|x|) = \int \frac{dk^2}{2\pi} \frac{e^{ikx}}{k^2 + m^2} \rightarrow -2\pi \ln(m|x|)$$

as $m|x| \rightarrow 0$, so that the integrals are well-defined for a large class of functions $\gamma(x)$. Since

$$\lim_{m \rightarrow 0} m \ln(m|x|) \rightarrow 0,$$

all terms but the first in the above series vanish as $m \rightarrow 0$ as long as $\gamma(x)$ is sufficiently well-behaved that we may exchange the integral and the limit. We find

$$G_m^\Lambda(0,0) \rightarrow -\frac{1}{4\pi} \ln \frac{\Lambda^2}{m^2}$$

as $m \rightarrow 0$. Including the Pauli-Villars contributions, we find

$$\langle T^i \rangle_g = -\frac{1}{2} \left(m^2 \ln \frac{\Lambda^2}{m^2} + \sum_i c_i M_i^2 \ln \frac{\Lambda^2}{M_i^2} \right) - \frac{1}{12} R + o\left(\frac{1}{M^2}\right),$$

where we have used the Pauli-Villars condition (10) to obtain the negative sign for the curvature term. As before, the terms in parentheses can be compensated by a counterterm or made zero by the condition (13) on the Pauli-Villars masses. After doing this, we find the result that we set out to prove

$$\langle T^i \rangle_g \rightarrow -\frac{1}{12} R$$

as the masses $M_i \rightarrow \infty$.

4 A study of T_{zz}

It is very instructive to observe the effect of the Pauli-Villars regularization on the expectation values of T_{zz} and $T_{\bar{z}\bar{z}}$. Writing $z \equiv x^1 + ix^2$, we have

$$\langle T_{zz} \rangle_g = -2\pi \lim_{\Lambda \rightarrow \infty} \left\{ -\partial_z^2 G_m^\Lambda(x, x) - \sum_i \partial_z^2 G_{M_i}^\Lambda(x, x) \right\}.$$

In general, the matter contribution $\langle T_{zz}^{m=0} \rangle$ may diverge in the cutoff parameter, and this divergence will be compensated by the zeroth-order contribution

$$\sum_i \langle T_{zz}^{M_i} \rangle_0 \equiv 2\pi \sum_i c_i \partial_z^2 \langle x | A^{-1} | x \rangle$$

of the Pauli-Villars fields, as we shall see in the explicit examples of the next section. With this in mind, we introduce a name for the combination

$$\hat{T}_{zz} \equiv T_{zz}^{m=0} + \lim_{M_i \rightarrow \infty} \sum_i \langle T_{zz}^{M_i} \rangle_0. \quad (14)$$

This may be regarded as defining a particular minimal subtraction renormalization prescription. As we shall see, it is this combination that will turn out to correspond to the renormalized energy-momentum operator occurring in the operator formalism. Obviously, it differs from the full energy-momentum tensor T_{zz} of our system.

The Pauli-Villars contributions may be expanded in $1/M^2$ as in our calculation of the trace anomaly in the previous section. We find a first-order contribution proportional to

$$\begin{aligned} & -2\pi \left[-(\partial_i \partial_j \omega) + 2(\partial_i \omega)(\partial_j \omega) \right] \times \\ & \times \int d^2x x^i x^j \partial_z K_0(M|x|) \partial_z \Delta K_0(M|x|). \end{aligned}$$

The term with integrand proportional to $\bar{z}\bar{z}$ vanishes

$$\begin{aligned} & \int d^2x \bar{z}\bar{z} \partial_z K_0(Mr) \partial_z \Delta K_0(Mr) \\ & = M^2 \int d^2x \bar{z}\bar{z}\bar{z}\bar{z} \partial_r K_0(Mr) \partial_r \Delta K_0(Mr) \\ & = 0, \end{aligned}$$

since the integrand is odd under $z \rightarrow e^{i\pi/4} z$. Similarly, the term with integrand proportional to $z\bar{z}$ vanishes, and we are left with the term proportional to

$$\begin{aligned}
& \int d^2x z^2 \partial_z K_0(Mr) \partial_z \Delta K_0(Mr) \\
&= - \int d^2x z^2 \int \frac{d^2k}{(2\pi)^2} \frac{\frac{1}{2} \bar{k} e^{\frac{1}{2} i (k\bar{z} + \bar{k}z)}}{k\bar{k} + M^2} \int \frac{d^2p}{(2\pi)^2} \frac{\frac{1}{2} \bar{p} (p\bar{p}) e^{-\frac{1}{2} i (p\bar{z} + \bar{p}z)}}{p\bar{p} + M^2} \\
&= (2\pi)^2 \int \frac{d^2k}{(2\pi)^2} \int \frac{d^2p}{(2\pi)^2} \frac{\bar{k}}{k\bar{k} + M^2} \frac{\bar{p} (p\bar{p})}{p\bar{p} + M^2} \partial_{\bar{p}}^2 \delta^2(k - p) \\
&= \int \frac{d^2k}{(2\pi)^2} \frac{\bar{k}}{k\bar{k} + M^2} \partial_{\bar{k}}^2 \left(\frac{\bar{k} (k\bar{k})}{k\bar{k} + M^2} \right),
\end{aligned}$$

which is of the form $\int d^2k f (\partial_{\bar{k}}^2 g)$. In the last step, we could just as well have used the identity

$$\partial_{\bar{p}}^2 \delta^2(k - p) = \partial_{\bar{k}}^2 \delta^2(k - p)$$

to write this in the form $\int d^2k (\partial_{\bar{k}}^2 f) g$. However, the reader may check by explicit calculation that

$$\int d^2k f (\partial_{\bar{k}}^2 g) \neq \int d^2k (\partial_{\bar{k}}^2 f) g,$$

an ambiguity due to naively exchanging the integral over x with those over k and p . In general, integrals may be exchanged only after a careful analysis of their uniformity of convergence [14]. Here we see an order dependence proportional to the quantities $f (\partial_{\bar{k}}^2 g)$ or $(\partial_{\bar{k}}^2 f) g$ integrated over the surface of the integration region. Since these surface integrands are of order $1/r$, they do contribute in two dimensions (the corresponding surface terms may be checked to be of order $1/r^2$ in our previous calculation of the trace anomaly, and therefore did not threaten the validity of that calculation). Fortunately we can avoid these uniformity issues by using the equation of motion for $K_0(mr)$ in position space first as follows:

$$\begin{aligned}
& \int d^2x z^2 \partial_z K_0(Mr) \partial_z \Delta K_0(Mr) \\
&= \int d^2x z^2 \partial_z K_0(Mr) \partial_z \left(-\delta^2(x) + M^2 K_0(Mr) \right) \\
&= M^2 \int d^2x z^2 \partial_z K_0(Mr) \partial_z K_0(Mr)
\end{aligned}$$

where we have used the behaviour $\partial_z K_0(Mr) = o(1/r)$ as $r \rightarrow 0$ to drop the term containing $\partial_z \delta^2(x)$. This then becomes

$$\begin{aligned}
& M^2 \int d^2x z^2 \int \frac{d^2k}{(2\pi)^2} \frac{\frac{1}{2} \bar{k} e^{\frac{1}{2} i (k\bar{z} + \bar{k}z)}}{k\bar{k} + M^2} \int \frac{d^2p}{(2\pi)^2} \frac{\frac{1}{2} \bar{p} e^{-\frac{1}{2} i (p\bar{z} + \bar{p}z)}}{p\bar{p} + M^2} \\
&= -M^2 \int \frac{d^2k}{(2\pi)^2} \frac{\bar{k}}{k\bar{k} + M^2} \partial_{\bar{k}}^2 \left(\frac{\bar{k}}{k\bar{k} + M^2} \right) \\
&= 2M^2 \int \frac{d^2k}{(2\pi)^2} \left\{ \frac{k\bar{k}}{(k\bar{k} + M^2)^2} - \frac{(k\bar{k})^2}{(k\bar{k} + M^2)^4} \right\} \\
&= \frac{1}{12\pi}.
\end{aligned}$$

The first-order contribution is therefore

$$\langle T_{zz}^M \rangle_1 = -\frac{1}{6} [-(\partial_z^2 \omega) + 2(\partial_z \omega)^2] + o\left(\frac{1}{M^2}\right).$$

The only other contribution of order 0 in $1/M^2$ is the second-order term

$$\begin{aligned}
& -2\pi \cdot 4 (\partial_z \omega)^2 \int d^2x \int d^2y zw \partial_z K_0(M|x|) \Delta K_0(M|y-x|) \partial_w \Delta K_0(M|y|) \\
&= -8\pi (\partial_z \omega)^2 \int d^2x \int d^2y zw \partial_z K_0(M|x|) (-\delta(y-x) + M^2 K_0(M|y-x|)) \times \\
&\quad \times \partial_w (-\delta(y) + M^2 K_0(M|y|)) \\
&= -8\pi (\partial_z \omega)^2 \left\{ M^4 \int d^2x \int d^2y zw \partial_z K_0(M|x|) K_0(M|y-x|) \partial_w K_0(M|y|) \right. \\
&\quad \left. - M^2 \int d^2x z^2 \partial_z K_0(M|x|) \partial_z K_0(M|x|) \right\},
\end{aligned}$$

where again we have manipulated the integrals in position space into a form where the surface terms in momentum space will vanish. This becomes

$$\begin{aligned}
& -8\pi (\partial_z \omega)^2 \left\{ -M^4 \int \frac{d^2k}{(2\pi)^2} \partial_{\bar{k}} \left(\frac{\bar{k}}{k\bar{k} + M^2} \right) \frac{1}{k\bar{k} + M^2} \partial_{\bar{k}} \left(\frac{\bar{k}}{k\bar{k} + M^2} \right) \right. \\
&\quad \left. + M^2 \int \frac{d^2k}{(2\pi)^2} \left[\partial_{\bar{k}} \left(\frac{\bar{k}}{k\bar{k} + M^2} \right) \right]^2 \right\} \\
&= \frac{1}{6} (\partial_z \omega)^2,
\end{aligned}$$

so that the second order contribution is

$$\langle T_{zz}^M \rangle_2 = \frac{1}{6} (\partial_z \omega)^2 + o\left(\frac{1}{M^2}\right).$$

As $M \rightarrow \infty$, we get for each Pauli-Villars field

$$\langle T_{zz}^{M_i} \rangle = \langle T_{zz}^{M_i} \rangle_0 + \frac{1}{6} c_i [(\partial_z^2 \omega) - (\partial_z \omega)^2].$$

Including the matter contribution and using the Pauli-Villars condition (3), we find

$$\langle T_{zz} \rangle = \langle \hat{T}_{zz} \rangle - \frac{1}{6} [(\partial_z^2 \omega) - (\partial_z \omega)^2] \quad (15)$$

$$\equiv \langle \hat{T}_{zz} \rangle - \frac{1}{12} t_{zz}. \quad (16)$$

Since $\Gamma_{zz}^z = 2 \partial_z \omega$, we may write t_{zz} in the form

$$t_{zz} = \partial_z \Gamma_{zz}^z - \frac{1}{2} (\Gamma_{zz}^z)^2$$

used by Eguchi and Ooguri in the context of the axiomatic approach to conformal field theory [15, 16]. One may verify that t_{zz} has the nontensor transformation law

$$\delta_v t_{zz} \equiv -\partial_z^3 v^z + v^i \partial_i t_{zz} - 2(\partial_z v^z) t_{zz}. \quad (17)$$

Since T_{zz} is a component of a true tensor, the term $\langle \hat{T}_{zz} \rangle$ in (16) is therefore not a tensor in isolation. From the transformation laws for T_{zz} and t_{zz} , it follows that $\langle \hat{T}_{zz} \rangle$ transforms as

$$\delta_v \langle \hat{T}_{zz} \rangle = -\frac{1}{12} \partial_z^3 v^z + v^i \partial_i \langle \hat{T}_{zz} \rangle - 2(\partial_z v^z) \langle \hat{T}_{zz} \rangle, \quad (18)$$

which coincides with the transformation law obtained in the operator product formalism after point-splitting renormalization [15, 16, 17, 18, 19, 20, 21]. The above result provides a straightforward explanation, in the context of the path integral formalism, for the familiar anomalous transformation law of \hat{T}_{zz} . In the path integral formalism, the more natural object is in fact the true tensor T_{zz} , which includes the full Pauli-Villars correction and, as we

shall see, satisfies simpler Ward identities. The non-covariant split (16) is rather less natural from this point of view.

Note that it would be incorrect to try to absorb the Pauli-Villars effects into an operator redefinition by replacing

$$T_{zz} \rightarrow \hat{T}_{zz} - \frac{1}{12} t_{zz}$$

in general calculations of expectation values. For example, with such a redefinition one would lose essential contact terms in the Ward identities that we consider in section 8.

The above transformation law for t_{zz} can be integrated to give, for $z' = f(z)$,

$$t_{z'z'} dz' \otimes dz' = t_{zz} dz \otimes dz - \{f, z\} dz \otimes dz,$$

where

$$\{f, z\} \equiv -6 \lim_{w \rightarrow z} \left(\frac{f'(w) f'(z)}{(f(w) - f(z))^2} - \frac{1}{(w - z)^2} \right),$$

is called the Schwarzian derivative. In this form, the transformation is easily seen to be precisely the difference between the renormalization subtractions needed in the operator product formalism for energy-momentum tensors defined in different coordinate systems, which provides a simple way to see that \hat{T}_{zz} corresponds to the usual operator formalism definitions.

Defining $t_{z\bar{z}} = t_{\bar{z}z} = 0$ and $t_{\bar{z}\bar{z}} = \partial_{\bar{z}} \Gamma_{\bar{z}\bar{z}}^{\bar{z}} - \frac{1}{2} (\Gamma_{\bar{z}\bar{z}}^{\bar{z}})^2$, we may summarize the results of this section as

$$\langle T_{ij} \rangle_g = \langle \hat{T}_{ij} \rangle_g - \frac{1}{12} t_{ij} - \frac{1}{24} g_{ij} R.$$

5 Examples

In this section we provide some example computations of energy-momentum tensor expectation values on a couple of simple manifolds.

First we consider the plane, for which

$$\langle T_{zz}^m \rangle_g = -\frac{1}{4} \int \frac{d^2 k}{(2\pi)^2} \frac{(k_1 - ik_2)^2}{k^2 + m^2}.$$

The numerator in the integrand is $k_1^2 - 2ik_1 k_2 - k_2^2$. The cross term is odd in k_1 and k_2 separately, while the sum of the first and the last term is odd under

$k_1 \leftrightarrow k_2$, so that the integral vanishes when integrated over the particular choice $0 \leq k \leq \Lambda$ of cutoff region in momentum space, suitable for manifolds for which k_1 and k_2 are continuous. With this cutoff region, we get the contribution

$$\langle T_{zz}^{m=0} \rangle = 0.$$

Similarly,

$$\langle T_{zz}^{M_i} \rangle = 0.$$

for each Pauli-Villars field, and we find

$$\langle T_{zz} \rangle = 0 = \langle \hat{T}_{zz} \rangle.$$

In section 3 we showed, using a Pauli-Villars definition of the path integral measure, that the expectation value

$$\langle T_{z\bar{z}} \rangle = \frac{1}{24} g_{z\bar{z}} R = 0$$

on the plane.

In a later section, we will also need the expectation value

$$\langle \partial_z \phi \partial_{\bar{z}} \phi \rangle = - \int \frac{d^2 k}{(2\pi)^2} \frac{k^2}{k^2 + m^2}.$$

This integrand is not odd, so that our simple argument for T_{zz} cannot be used. Computing this with a large-momentum cutoff Λ , we find

$$\langle \partial_z \phi \partial_{\bar{z}} \phi \rangle = -\frac{1}{2\pi} \Lambda^2 + \frac{m^2}{2\pi} \cdot \frac{1}{2} \ln \frac{\Lambda^2}{m^2}.$$

As we saw, both these terms can be exactly canceled by introducing Pauli-Villars fields with the appropriate statistics and mass conditions, and we obtain

$$\langle \partial_z \phi \partial_{\bar{z}} \phi \rangle = 0$$

on the plane.

Let us employ the Pauli-Villars formalism to perform a path-integral calculation of the expectation value $\langle T_{zz} \rangle$ on an infinite cylinder of circumference

L , a simple nontrivial manifold with a length scale that will give rise to a Casimir energy. We have, for a field of mass M ,

$$\begin{aligned}\langle T_{zz}^M \rangle &= -2\pi \cdot \frac{1}{4} \cdot \frac{1}{L} \cdot \sum_n \int \frac{dk_2}{2\pi} \frac{\left(\frac{2\pi n}{L}\right)^2 - k_2^2}{\left(\frac{2\pi n}{L}\right)^2 + k_2^2 + M^2} \\ &= -\frac{\pi}{2} \cdot \frac{1}{L} \sum_n \left\{ \sqrt{\left(\frac{2\pi n}{L}\right)^2 + M^2} - \frac{M^2/2}{\sqrt{\left(\frac{2\pi n}{L}\right)^2 + M^2}} \right\}.\end{aligned}$$

We have performed the integral over k_2 by closing the integration contour either above or below the real line. Although the resulting contour integral strictly diverges as we move the contour to infinity, this is one term in the full integrand containing both the matter and all Pauli-Villars, the integral of which will converge due to the Pauli-Villars conditions, validating the contour integral argument.

We first evaluate the matter contribution, for which $M = 0$. Inserting a convergence factor, the dependence on which will be canceled by the Pauli-Villars contributions, we find

$$\begin{aligned}\langle T_{zz}^{m=0} \rangle &= -\frac{\pi}{L} \sum_{n>0} \frac{2\pi n}{L} e^{-\epsilon 2\pi n/L} \\ &= -\frac{\pi}{L} \cdot \frac{2\pi}{L} \frac{e^{-\epsilon 2\pi/L}}{(1 - e^{-\epsilon 2\pi/L})^2}.\end{aligned}$$

Using the expansion

$$\frac{e^{-\nu}}{(1 - e^{-\nu})^2} = \frac{1}{\nu^2} - \frac{1}{12} + o(\nu),$$

we get

$$\langle T_{zz}^{m=0} \rangle = -\frac{1}{2\epsilon^2} + \frac{1}{24} \left(\frac{2\pi}{L}\right)^2 + o(\epsilon).$$

It remains to show that, as promised, the Pauli-Villars fields will cancel the dependence on ϵ as $\epsilon \rightarrow 0$. As $M \rightarrow \infty$, we can write each Pauli-Villars

contribution as a continuous integral as follows.

$$\begin{aligned}
\langle T_{zz}^M \rangle &= -\frac{1}{2} \cdot M^2 \cdot \frac{2\pi}{ML} \sum_{n \geq 0} \left\{ \sqrt{\left(\frac{2\pi n}{ML}\right)^2 + 1} - \frac{1/2}{\sqrt{\left(\frac{2\pi n}{ML}\right)^2 + 1}} \right\} e^{-\epsilon M 2\pi n / ML} \\
&\rightarrow -\frac{1}{2} \cdot M^2 \int_0^\infty dx \left\{ \sqrt{x^2 + 1} - \frac{1/2}{\sqrt{x^2 + 1}} \right\} e^{-\epsilon M x} \\
&= -\frac{1}{2} \cdot M^2 \cdot \frac{\sqrt{\pi}}{2} \left\{ \frac{2}{\epsilon M} \Gamma\left(\frac{3}{2}\right) [\mathbf{H}_1(\epsilon M) - N_1(\epsilon M)] \right. \\
&\quad \left. + \frac{1}{2} \Gamma\left(\frac{1}{2}\right) [\mathbf{H}_0(\epsilon M) - N_0(\epsilon M)] \right\},
\end{aligned}$$

where \mathbf{H}_ν denotes Struve functions and N_ν Bessel functions of the second kind [22]. Using the small z expansions

$$\begin{aligned}
\mathbf{H}_0(z) &= o(z), \\
\mathbf{H}_1(z) &= o(z^2), \\
\pi N_0(z) &= 2 \left(\ln \frac{z}{2} + \mathbf{C} \right) + o(z^2) \\
\pi N_1(z) &= -\frac{2}{z} - \frac{z}{2} + z \left(\ln \frac{z}{2} + \mathbf{C} \right) + o(z^3),
\end{aligned}$$

we find

$$\langle T_{zz}^M \rangle = -\frac{1}{2\epsilon^2} - \frac{1}{8} M^2 + o(\epsilon).$$

By the Pauli-Villars conditions $\sum c_i = 1$ and $\sum c_i M_i^2 = 0$ from (10) and (12), the first term cancels the ϵ dependence of the matter field and the second term falls away. Our final answer for the full expectation value becomes

$$\langle T_{zz} \rangle = \frac{1}{24} \left(\frac{2\pi}{L} \right)^2.$$

Computed the Pauli-Villars measure, this is finite without further renormalization.

6 Reparametrizations and the measure

Consider a family of reparametrizations $f_\lambda : M \rightarrow M$, $f_0 = \text{id}$ where λ is a real parameter, acting on a scalar field via push-forward

$$\phi(z) \rightarrow \phi^\lambda(z) \equiv \phi(f_{-\lambda}(z)).$$

Given a metric g on M , let ϕ_n be a basis of field configurations satisfying

$$\delta_{mn} = \int d^2x \sqrt{g} \phi_m(x) \phi_n(x)$$

and expand ϕ^λ as

$$\phi^\lambda(x) \equiv \sum_n a_n^\lambda \phi_n(x),$$

For each λ , the map $\phi \mapsto (a_0^\lambda, a_1^\lambda, \dots)$ may be regarded as a coordinate chart on the space of field configurations ϕ . Denoting the ∞ -form

$$\bigwedge_n da_n^\lambda \equiv da_0^\lambda \wedge da_1^\lambda \wedge \dots \equiv [d\phi^\lambda],$$

the change of variables theorem tells us that

$$\int [d\phi^\lambda] e^{-S(\phi^\lambda)} = \int [d\phi] e^{-S(\phi)}.$$

Note that this had better be true if the concept of integration is to make sense in a chart-independent way. In preparation for our derivation of the Ward identities in the next section, let us determine how the form $[d\phi^\lambda]$ depends on λ . We can project out the coefficient

$$a_m^\lambda = \int d^2x \sqrt{g} \phi_m(x) \phi^\lambda(x),$$

and calculate

$$\begin{aligned} \frac{d}{d\lambda} a_m^\lambda &= \int d^2x \sqrt{g} \phi_m(x) (-v^i) \partial_i \phi^\lambda(x) \\ &= - \sum_n a_n^\lambda \int d^2x \sqrt{g} \phi_m(x) v^i \partial_i \phi_n(x) \\ &\equiv \sum_n C_{mn} a_n^\lambda \end{aligned}$$

where v^i is the vector field whose flow gives f_λ , in other words

$$v^i(x) \equiv \left. \frac{df_\lambda^i(x)}{d\lambda} \right|_{\lambda=0}$$

The normal rules for manipulating differential forms now give

$$\frac{d}{d\lambda} [d\phi^\lambda] \equiv \frac{d}{d\lambda} \bigwedge_n da_n^\lambda = \left(\sum_m C_{mm} \right) \bigwedge_n da_n^\lambda = (\text{Tr } C) [d\phi^\lambda].$$

Now

$$\begin{aligned} C_{mm} &= - \int d^2x \sqrt{g} \phi_m(x) v^i \partial_i \phi_m(x) \\ &= \int d^2x \sqrt{g} (v^i \partial_i \phi_m(x)) \phi_m(x) \\ &\quad + \int d^2x \phi_m(x) \partial_i (\sqrt{g} v^i) \phi_m(x) \\ &= \frac{1}{2} \int d^2x \phi_m(x) \partial_i (\sqrt{g} v^i) \phi_m(x) \\ &= \frac{1}{2} \int d^2x \sqrt{g} \phi_m(x) (\nabla_i v^i) \phi_m(x) \end{aligned}$$

where ∇_i denotes the covariant derivative on the worldsheet with respect to the deformed metric $f_\lambda^* g$. Similar to the calculation of the Weyl anomaly, we get the trace of an operator $\nabla_i v^i$ representing a change in area, in this case along the flow of v . The calculation of the trace can be done via a heat kernel regularization and is identical to the calculation of the Weyl anomaly, with $\nabla_i v^i$ replacing $2\delta\omega$. We get

$$\frac{d}{d\lambda} [d\phi^\lambda] = \left(\frac{1}{8\pi\epsilon} \int d^2x \sqrt{g} \nabla_i v^i + \frac{1}{48\pi} \int d^2x \sqrt{g} (\nabla_i v^i) R_g \right) [d\phi^\lambda]$$

The integrand $\sqrt{g} \nabla_i v^i = \partial_i (\sqrt{g} v^i)$ in the first term is a total derivative, and will integrate to zero for suitable boundary conditions on v^i , which we will assume. A similar partial integration in the second term, and remembering that $\nabla_i = \partial_i$ on scalars, leads to

$$\frac{d}{d\lambda} [d\phi^\lambda] = \left(-\frac{1}{48\pi} \int d^2x \sqrt{g} v^i \nabla_i R_g \right) [d\phi^\lambda]. \quad (19)$$

In particular, on a flat manifold, the measure will be invariant under reparametrizations of ϕ . It is also interesting to note that the prefactor is independent of λ .

7 Ward identities

The basis for deriving the energy-momentum conservation Ward identities is the change of variables theorem [18]

$$\int [d\phi^\lambda] e^{-S(\phi^\lambda)} = \int [d\phi] e^{-S(\phi)}, \quad (20)$$

where

$$S(\phi^\lambda) = \frac{1}{2} \int d^2x \sqrt{g} g^{ij} \partial_i \phi^\lambda \partial_j \phi^\lambda$$

Notice that if f_λ is a conformal transformation, then S is independent of λ , so that classically we would have a symmetry. In the path integral formalism, the associated conserved Noether current follows from the variation of (20) with respect to more general, non-conformal deformations f_λ . Differentiating with respect to λ , the right hand side gives zero. For the left hand side, we have already calculated the variation of $[d\phi^\lambda]$. The variation of S is

$$\begin{aligned} \frac{dS}{d\lambda} &= \frac{1}{2} \cdot \frac{d}{d\lambda} \int d^2x \sqrt{g} g^{ij} \partial_i \phi^\lambda \partial_j \phi^\lambda \\ &= \frac{1}{2} \cdot \frac{d}{d\lambda} \int d^2x \sqrt{f_\lambda^* g} (f_\lambda^* g)^{ij} \partial_i \phi \partial_j \phi && \text{(by pullback)} \\ &= \frac{1}{4\pi} \int d^2x \sqrt{f_\lambda^* g} \delta_\lambda (f_\lambda^* g)^{ij} T_{ij}^{f_\lambda^* g}(\phi) && \text{(definition of } T) \\ &= \frac{1}{4\pi} \int d^2x \sqrt{f_\lambda^* g} (\nabla_{f_\lambda^* g}^i v^j + \nabla_{f_\lambda^* g}^j v^i) T_{ij}^{f_\lambda^* g}(\phi) \\ &= \frac{1}{4\pi} \int d^2x \sqrt{g} (\nabla^i v^j + \nabla^j v^i) T_{ij}^g(\phi^\lambda) && \text{(by push-forward)} \\ &= \frac{1}{2\pi} \int d^2x \sqrt{g} (\nabla^i v^j) T_{ij}(\phi^\lambda) && \text{(symmetry of } T) \\ &= \frac{1}{2\pi} \int d^2x \sqrt{g} (\nabla^i (v^j T_{ij}(\phi^\lambda)) - v^j \nabla^i T_{ij}(\phi^\lambda)) \\ &= -\frac{1}{2\pi} \int d^2x \sqrt{g} v^j \nabla^i T_{ij}(\phi^\lambda), \end{aligned}$$

where

$$T_{ij}^g(\phi^\lambda) \equiv -2\pi \left(\partial_i \phi^\lambda \partial_j \phi^\lambda - \frac{1}{2} g_{ij} g^{kl} \partial_k \phi^\lambda \partial_l \phi^\lambda \right).$$

and where we have used the property $\sqrt{g} \nabla^i J_i = \partial^i(\sqrt{g} J_i)$ of the covariant derivative to get a total derivative which we have dropped in the last line. Differentiation of equation (20) with respect to λ then gives

$$\begin{aligned} 0 &= \int d^2x \sqrt{g} \left\langle \frac{1}{2\pi} v^j \nabla^i T_{ij}(\phi^\lambda) - \frac{1}{48\pi} v^i \nabla_i R \right\rangle_\lambda \\ &= \frac{1}{2\pi} \int d^2x \sqrt{g} v^j \nabla^i \left\langle T_{ij}(\phi^\lambda) - \frac{1}{24} g_{ij} R \right\rangle_\lambda \end{aligned}$$

Since v^j is arbitrary, we get the conservation law

$$\nabla^i \left\langle T_{ij}(\phi^\lambda) - \frac{1}{24} g_{ij} R \right\rangle_\lambda = 0, \quad (21)$$

where the variation of the measure has contributed a curvature term not present in the classical conservation law. Remembering our definition of \tilde{T}_{ij} ,

$$\nabla^i \left\langle \tilde{T}_{ij}(\phi^\lambda) \right\rangle_\lambda = 0.$$

In this derivation, which assumed a single massless scalar field for which T_i^i is identically zero, the curvature term in the conservation law was due to the variation of the path integral measure under a reparametrization.

As it stands, though, this formula is not quite correct. In fact, as we have seen, the expectation values $\langle T_{ij} \rangle$ are not generally finite, and we needed to introduce Pauli-Villars fields to obtain finite values for $\langle T_{ij} \rangle$ in a coordinate-invariant way. Using (19), we find as in section 3 that the combined matter-Pauli-Villars path integral measure is invariant under reparametrizations, and by the above argument we then find

$$\nabla^i \langle T_{ij}(\phi^\lambda) \rangle_\lambda = 0. \quad (22)$$

The curvature dependence then comes from the fact that, with the Pauli-Villars regularization, we have seen that

$$\langle T_{ij}(\phi^\lambda) \rangle_\lambda = \left\langle \hat{T}_{ij}(\phi^\lambda) \right\rangle_\lambda - \frac{1}{12} t_{ij} - \frac{1}{24} g_{ij} R.$$

8 The Ward identity for T_{ij} , or, where is the anomaly?

To derive the Ward identity with an insertion of T_{ij} , we use the change of variables theorem in the form

$$\int [d\phi^\lambda] T_{ij}(\phi^\lambda)_x e^{-S(\phi^\lambda)} = \int [d\phi] T_{ij}(\phi)_x e^{-S(\phi)}. \quad (23)$$

Here $[d\phi^\lambda]$ denotes the full measure including Pauli-Villars fields, which is independent of λ as explained in section 7. Differentiating both sides with respect to λ , we therefore find the Ward identity

$$\langle \delta_{\mathbf{v}} T_{ij}(\phi^\lambda)_x \rangle_\lambda + \frac{1}{4\pi} \int d^2y h^{kl}(y) \langle T_{ij}(\phi^\lambda)_x T_{kl}(\phi^\lambda)_y \rangle_\lambda = 0, \quad (24)$$

where

$$\delta_{\mathbf{v}} T_{ij} = v^i \partial_i T_{ij} - (\partial_i v^m) T_{mj} - (\partial_j v^m) T_{im}$$

and

$$h^{kl} \equiv \nabla^k v^l - \nabla^l v^k.$$

It is important to note that the transformation law for T_{ij} appearing in the Ward identity is the classical one. This is indeed correct, and simply expresses the fact that the full energy-momentum tensor, including Pauli-Villars contributions, is a true coordinate invariant tensor quantity. As discussed in section 4, it is the quantity \hat{T}_{zz} introduced in (14) that satisfies an anomalous transformation law. We shall see by explicit calculation that the above Ward identity is indeed correct, but contains contact terms that can be compensated by a redefinition of T_{ij} to obtain the familiar anomalous Ward identity for \hat{T}_{zz} .

For simplicity, we restrict attention to the plane and consider

$$\begin{aligned} \langle \delta_{\mathbf{v}} T_{zz}(\phi^\lambda) \rangle_\lambda &= -\frac{1}{4\pi} \int d^2w h^{ww} \langle T_{zz}(\phi^\lambda) T_{ww}(\phi^\lambda) \rangle_\lambda \\ &\quad - 2 \cdot \frac{1}{4\pi} \int d^2w h^{w\bar{w}} \langle T_{zz}(\phi^\lambda) T_{w\bar{w}}(\phi^\lambda) \rangle_\lambda \\ &\quad - \frac{1}{4\pi} \int d^2w h^{\bar{w}\bar{w}} \langle T_{zz}(\phi^\lambda) T_{\bar{w}\bar{w}}(\phi^\lambda) \rangle_\lambda \end{aligned}$$

The first expectation value on the right hand side is easily calculated by a double contraction to be

$$\langle T_{zz}(\phi^\lambda) T_{ww}(\phi^\lambda) \rangle_\lambda = \frac{1}{2} \frac{1}{(z-w)^4}.$$

The expectation values in the second and third terms are not discussed in many standard treatments, but in fact contribute contact terms, in the absence of which the above identity would be untrue. The presence of the contact terms are noted in [19, 23, 24], although we have been unable to find a calculation from first principles as presented below.

The contact terms are nontrivial to calculate. Consider for example $\langle T_{zz} T_{\bar{w}\bar{w}} \rangle$. Naively taking appropriate derivatives of the double contraction for a massless field would give the square of the delta function, which does not exist as a well-defined distribution. Also troublesome is the expectation value $\langle T_{zz} T_{w\bar{w}} \rangle$. Since $T_{w\bar{w}}$ is identically zero in a massless theory, one might expect the answer to be zero. With our careful definition of the path integral, we shall see that this is only true up to a contact term.

Let us therefore calculate these expectation values more carefully using our regularized path integral. We realize an infrared regularization by introducing a mass m for the field ϕ , eventually to be taken to zero, while the ultraviolet regularization is taken care of, as before, by the Pauli-Villars auxiliary fields whose masses we eventually take to infinity.

We start by considering the expectation value $\langle T_{zz} T_{\bar{w}\bar{w}} \rangle$. Writing the contractions in terms of derivatives of the propagator

$$\langle \phi(x) \phi(0) \rangle = \int \frac{d^2 p}{(2\pi^2)} \frac{e^{-ipx}}{p^2 + m^2}$$

and Fourier transforming the result gives the familiar one-loop Feynman integral

$$\langle T_{zz}(x) T_{\bar{z}\bar{z}}(0) \rangle = \frac{2(2\pi)^2}{16} \int \frac{d^2 p}{(2\pi)^2} e^{-ip \cdot x} \int \frac{d^2 k}{(2\pi)^2} \left\{ F(m) - \sum_i c_i F(M_i) \right\}$$

where

$$F(m) \equiv \frac{k^2 (p-k)^2}{[k^2 + m^2] [(p-k)^2 + m^2]}. \quad (25)$$

In the absence of the Pauli-Villars field contributions, the integral over k would have both a quadratic and a logarithmic divergence. The regularization consists in choosing the coefficients c_i and masses M_i so as to make

the integral finite. Assuming this has been done, we can then write, using identities such as $k^2 = k^2 + m^2 - m^2$,

$$\begin{aligned}
& \int \frac{d^2k}{(2\pi)^2} \left\{ \frac{k^2 (p-k)^2}{[k^2 + m^2] [(p-k)^2 + m^2]} + P.V. \right\} \\
&= \int \frac{d^2k}{(2\pi)^2} \left\{ 1 - \frac{m^2}{(p-k)^2 + m^2} - \frac{m^2}{k^2 + m^2} \right. \\
&\quad \left. + \frac{m^2}{[k^2 + m^2] [(p-k)^2 + m^2]} + P.V. \right\} \\
&= \int \frac{d^2k}{(2\pi)^2} \left\{ 1 - \frac{2m^2}{k^2 + m^2} + \frac{m^4}{[k^2 + m^2] [(p-k)^2 + m^2]} + P.V. \right\},
\end{aligned}$$

where the shift of k in the last line is permitted since the integral converges. Integrating the first two terms between 0 and Λ , we get, for large Λ ,

$$\frac{1}{4\pi} \Lambda^2 + \frac{1}{2\pi} m^2 \ln \frac{\Lambda^2}{m^2} + \int \frac{d^2k}{(2\pi)^2} \frac{m^4}{[k^2 + m^4] [(p-k)^2 + m^2]} + P.V.$$

By the Pauli-Villars conditions (10), (12) and (13),

$$\begin{aligned}
1 + \sum_i c_i &= 0, \\
m^2 + \sum_i c_i M_i^2 &= 0, \\
m^2 \ln \frac{m^2}{\mu^2} + \sum_i c_i M_i^2 \ln \frac{M_i^2}{\mu^2} &= 0,
\end{aligned}$$

the first two terms cancel entirely [13]. What remains is the finite integral

$$\begin{aligned}
\langle T_{zz}(x) T_{\bar{z}\bar{z}}(0) \rangle &= \frac{2m^4 (2\pi)^2}{16} \int \frac{d^2p}{(2\pi)^2} e^{-ip \cdot x} \int \frac{d^2k}{(2\pi)^2} \frac{1}{[k^2 + m^2]} \frac{1}{[(p-k)^2 + m^2]} \\
&\quad + P.V.
\end{aligned}$$

Feynman's trick gives

$$\begin{aligned}
& \int \frac{d^2k}{(2\pi)^2} \frac{1}{k^2 + m^2} \frac{1}{(p-k)^2 + m^2} \\
&= \int \frac{d^2k}{(2\pi)^2} \int_0^1 dx \frac{1}{\{x(k^2 + m^2) + (1-x)[(p-k)^2 + m^2]\}^2} \\
&= \int \frac{d^2k}{(2\pi)^2} \int_0^1 dx \frac{1}{(k^2 + x(1-x)p^2 + m^2)^2},
\end{aligned}$$

where we have redefined $k - (x - 1)p \rightarrow k$ to complete the square in the last line. Performing the straightforward integration over k , we find

$$\frac{1}{4\pi} \int_0^1 dx \frac{1}{x(1-x)p^2 + m^2},$$

Changing variables [25] from x to

$$s \equiv \frac{m^2}{x(1-x)},$$

and including the prefactor, we find the spectral representation [24]

$$\begin{aligned} & \frac{1}{4\pi} \cdot \frac{2(2\pi)^2}{16} \int_{4m^2}^{\infty} \frac{ds}{s} \frac{2m^4}{\sqrt{1-4m^2/s}} \cdot \frac{1}{p^2+s} \\ &= \frac{1}{16} \cdot 4\pi \int_{2m}^{\infty} \frac{d\mu}{\mu} \frac{2m^4}{\sqrt{1-4m^2/\mu^2}} \cdot \frac{1}{p^2+\mu^2} \\ &= \frac{1}{16} \cdot \frac{\pi}{3} \int_{2m}^{\infty} \frac{d\mu}{\mu^5} \frac{24m^4}{\sqrt{1-4m^2/\mu^2}} \cdot \frac{1}{p^2+\mu^2} \\ &= \frac{1}{16} \cdot \frac{\pi}{3} \int d\mu c(\mu, m) \cdot \frac{\mu^4}{p^2+\mu^2} \end{aligned}$$

for the Fourier transform of the expectation value. Here the spectral function

$$c(\mu, m) \equiv \frac{24m^4}{\mu^5 \sqrt{1-4m^2/\mu^2}} \theta(\mu - 2m)$$

is dimensionless and has area equal to 1, independent of m . Contributions to the expectation value come from two-particle intermediate states propagating between 0 and x . The lowest of these has energy $2m$, which explains the lower bound on the integral.

To confirm that the area is one, we calculate [22]

$$\begin{aligned} 24m^4 \int_{2m}^{\infty} \frac{d\mu}{\mu^5} \frac{1}{\sqrt{1-4m^2/\mu^2}} &= \frac{3}{2} \int_1^{\infty} \frac{d\eta}{\eta^4} \frac{1}{\sqrt{\eta^2-1}} \\ &= \frac{3}{2} \cdot \frac{1}{2} \cdot B(2, \frac{1}{2}) \\ &= 1. \end{aligned}$$

As $m \rightarrow 0$, $c(\mu, m)$ develops a spike at $2m$ and goes to zero elsewhere. It follows that

$$c(\mu, m) \rightarrow \delta(\mu) \quad \text{as } m \rightarrow 0.$$

The Fourier transformed expectation value, including the Pauli-Villars contributions, is then

$$\begin{aligned} & \frac{1}{16} \cdot \frac{\pi}{3} \int d\mu \left\{ c(\mu, m) + \sum_i c_i c(\mu, M_i) \right\} \cdot \frac{\mu^4}{p^2 + \mu^2} \\ &= \frac{1}{16} \cdot \frac{\pi}{3} \int d\mu \left\{ c(\mu, m) + \sum_i c_i c(\mu, M_i) \right\} \cdot \left(\mu^2 - p^2 + \frac{p^4}{p^2 + \mu^2} \right) \end{aligned}$$

A change of variables from μ to η as above shows that the contribution

$$\int d\mu \left\{ c(\mu, m) + \sum_i c_i c(\mu, M_i) \right\} \mu^2$$

is proportional to $m^2 + \sum_i c_i M_i^2$, which is zero by the Pauli-Villars conditions. So is the contribution

$$\int d\mu \left\{ c(\mu, m) + \sum_i c_i c(\mu, M_i) \right\} p^2,$$

which is proportional to $1 + \sum_i c_i$ by the fact that $c(\mu, \cdot)$ has unit area. We are therefore left with

$$\frac{1}{16} \cdot \frac{\pi}{3} \int d\mu \left\{ c(\mu, m) + \sum_i c_i c(\mu, M_i) \right\} \cdot \frac{p^4}{p^2 + \mu^2}.$$

Here, as we take the Pauli-Villars masses to infinity, we find

$$\int_{2M_i} d\mu c(\mu, M_i) \cdot \frac{p^4}{p^2 + \mu^2} \rightarrow 0 \quad \text{as } M_i \rightarrow \infty$$

because of the lower bound on the integration and the unit area property of $c(\mu, \cdot)$ making the integrand of order $1/M_i^2$. All that remains is the matter contribution which, as we remove the infrared cutoff, is

$$\frac{1}{16} \cdot \frac{\pi}{3} \int d\mu c(\mu, m) \cdot \frac{p^4}{p^2 + \mu^2} \rightarrow \frac{1}{16} \cdot \frac{\pi}{3} \cdot p^2 \quad \text{as } m \rightarrow 0,$$

since in this limit $c(\mu, m) \rightarrow \delta(\mu)$. Fourier transforming, we find

$$\langle T_{zz}(x) T_{\bar{z}\bar{z}}(0) \rangle \rightarrow -\frac{\pi}{12} \partial_z \partial_{\bar{z}} \delta(x) \quad (26)$$

as $m \rightarrow 0$.

It is important to point out that, in addition to the above contribution, we would expect additional terms due to self-contractions. However, as discussed in section 5, these all vanish on the plane.

Next we calculate the expectation value

$$\langle T_{z\bar{z}}(z) T_{z\bar{z}}(0) \rangle.$$

Since the mass term breaks conformal invariance, $T_{z\bar{z}}$ is not zero. In fact

$$T_{z\bar{z}} = \frac{\pi}{2} m^2 \phi^2.$$

Although this indeed goes identically to zero as $m \rightarrow 0$, a contact term survives in the limit $m \rightarrow 0$. Indeed,

$$\langle T_{z\bar{z}}(x) T_{z\bar{z}}(0) \rangle = 2m^4 \left(\frac{\pi}{2}\right)^2 \int \frac{d^2 p}{(2\pi)^2} e^{-ip \cdot x} \int \frac{d^2 k}{(2\pi)^2} \left\{ \frac{1}{k^2 + m^2} \frac{1}{(p-k)^2 + m^2} + P.V. \right\},$$

which is the same expression we obtained above for $\langle T_{zz}(x) T_{\bar{z}\bar{z}}(0) \rangle$. Therefore

$$\langle T_{z\bar{z}}(x) T_{z\bar{z}}(0) \rangle \rightarrow -\frac{\pi}{12} \partial_z \partial_{\bar{z}} \delta(x) \quad (27)$$

as $m \rightarrow 0$.

The remaining expectation value $\langle T_{zz}(x) T_{z\bar{z}}(0) \rangle$ has Fourier transform (with a slight abuse of notation we denote $\bar{k} \equiv k_1 - ik_2$ but keep $k^2 = k_1^2 + k_2^2$)

$$\begin{aligned} & -\frac{1}{4} \cdot 2m^2 \cdot 2\pi \cdot \left(\frac{\pi}{2}\right) \int \frac{d^2 k}{(2\pi)^2} \left\{ \frac{\bar{k}(\bar{p} - \bar{k})}{[k^2 + m^2][(p-k)^2 + m^2]} + P.V. \right\} \\ & = -\frac{2m^2 (2\pi)^2}{16} \int \frac{d^2 k}{(2\pi)^2} \int_0^1 dx \left\{ \frac{[\bar{k} + (1-x)\bar{p}][\bar{p} - \bar{k} - (1-x)\bar{p}]}{[k^2 + x(1-x)p^2 + \mu^2]^2} + P.V. \right\} \\ & = -\frac{2m^2 (2\pi)^2}{16} \int \frac{d^2 k}{(2\pi)^2} \int_0^1 dx \left\{ \frac{x(1-x)\bar{p}^2}{[k^2 + x(1-x)p^2 + \mu^2]^2} + P.V. \right\}, \end{aligned}$$

where in the last line we have dropped odd integrands, including terms proportional to $k_1^2 - k_2^2$ and $k_1 k_2$. Performing the integration over k and changing variables from x to μ as before, we find

$$\begin{aligned} & -\frac{1}{16} \cdot \frac{\pi}{3} \int_{2m}^{\infty} \frac{d\mu}{\mu^5} \frac{24 m^4}{\sqrt{1 - 4m^2/\mu^2}} \cdot \frac{\mu^2 \bar{p}^2}{p^2 + \mu^2} + P.V. \\ &= -\frac{1}{16} \cdot \frac{\pi}{3} \int d\mu \left\{ c(\mu, m) \cdot \frac{\mu^2 \bar{p}^2}{p^2 + \mu^2} + P.V. \right\} \\ &= -\frac{1}{16} \cdot \frac{\pi}{3} \int d\mu \left\{ c(\mu, m) \cdot \bar{p}^2 - c(\mu, m) \cdot \frac{p^2 \bar{p}^2}{p^2 + \mu^2} + P.V. \right\}. \end{aligned}$$

As before, the first integrand will cancel due to the Pauli-Villars condition $1 + \sum_i c_i = 0$, while the second term will vanish for the Pauli-Villars fields in the limit of infinite mass. Again, all that remains is the matter contribution

$$-\frac{1}{16} \cdot \frac{\pi}{3} \int d\mu c(\mu, m) \cdot \frac{p^2 \bar{p}^2}{p^2 + \mu^2} \rightarrow -\frac{1}{16} \cdot \frac{\pi}{3} \bar{p}^2$$

as $m \rightarrow 0$. Fourier transforming, we get

$$\langle T_{zz}(x) T_{z\bar{z}}(0) \rangle \rightarrow \frac{\pi}{12} \partial_z^2 \delta(x) \quad (28)$$

Summarizing, we have

$$\langle T_{zz}(x) T_{zz}(0) \rangle = \frac{1}{2} \frac{1}{(z-w)^4}, \quad (29)$$

$$\langle T_{zz}(x) T_{\bar{z}\bar{z}}(0) \rangle = -\frac{\pi}{12} \partial_z \partial_{\bar{z}} \delta(x), \quad (30)$$

$$\langle T_{z\bar{z}}(x) T_{z\bar{z}}(0) \rangle = -\frac{\pi}{12} \partial_z \partial_{\bar{z}} \delta(x), \quad (31)$$

$$\langle T_{zz}(x) T_{z\bar{z}}(0) \rangle = \frac{\pi}{12} \partial_z^2 \delta(x). \quad (32)$$

The same formulae were obtained in the axiomatic approach to conformal field theory in [19]. A separate argument provided in [24] motivates them as follows: If we knew that conservation of energy-momentum held even in the limit of coinciding points, we could have expected the form of these correlation functions by inserting a spectral decomposition of the unit operator between the two T s. By conservation of T_{ij} , the correlator must then have the form

$$\langle T_{\mu\nu}(x) T_{\rho\sigma}(0) \rangle = \frac{\pi}{3} \int d\mu c(\mu) \int \frac{d^2 p}{(2\pi)^2} e^{ipx} \frac{(g_{\mu\nu} p^2 - p_\mu p_\nu)(g_{\rho\sigma} p^2 - p_\rho p_\sigma)}{p^2 + \mu^2},$$

which, noting that the Fourier transform of $1/z^4$ is $(\pi/24)\bar{p}^4/|p|^2$, indeed coincides with our results upon taking $c(\mu) \propto \delta(\mu)$, as expected for a massless theory. Our calculation confirms this explicitly. Further work on contact terms of energy-momentum tensors is reported in [26].

As a consistency check, even without performing the full calculation above, a simple algebraic argument applied to the original integrands, combined with a shift of variables, which is permitted in the presence of the Pauli-Villars fields, shows that, for example,

$$\partial_{\bar{z}} \langle T_{zz}(x) T_{z\bar{z}}(0) \rangle + \partial_z \langle T_{\bar{z}\bar{z}}(x) T_{z\bar{z}}(0) \rangle = 0.$$

In other words, the Pauli-Villars regularization does not break conservation of energy-momentum, even in the limit of coinciding points.

We are now finally ready to calculate

$$\begin{aligned} \langle \delta_v T_{zz}(\phi^\lambda) \rangle_\lambda &= -\frac{1}{4\pi} \int d^2w h^{ww} \langle T_{zz}(\phi^\lambda) T_{ww}(\phi^\lambda) \rangle_\lambda \\ &\quad - 2 \cdot \frac{1}{4\pi} \int d^2w h^{w\bar{w}} \langle T_{zz}(\phi^\lambda) T_{w\bar{w}}(\phi^\lambda) \rangle_\lambda \\ &\quad - \frac{1}{4\pi} \int d^2w h^{\bar{w}\bar{w}} \langle T_{zz}(\phi^\lambda) T_{\bar{w}\bar{w}}(\phi^\lambda) \rangle_\lambda \end{aligned}$$

Expressing the components of h in terms of v , for example, $h^{ww} = 2(\partial_{\bar{w}}v^w + \partial_{\bar{w}}v^w)$, discarding total derivative terms, and remembering that $\partial_{\bar{w}}(1/w) = \pi \delta(w)$, we find

$$\begin{aligned} \langle \delta_v T_{zz}(\phi^\lambda) \rangle_\lambda &= -\frac{1}{12} \partial_z^3 v^z \\ &\quad + \frac{1}{12} \partial_z^3 v^z + \frac{1}{12} \partial_z \partial_z^2 v^{\bar{z}} \\ &\quad - \frac{1}{12} \partial_{\bar{z}} \partial_z^2 v^{\bar{z}} \\ &= 0 \end{aligned}$$

In other words, the contact terms neatly cancel the anomalous contribution coming from the $1/(z-w)^4$ term in $\langle T_{zz}(\phi^\lambda) T_{ww}(\phi^\lambda) \rangle_\lambda$. Since the classical variation $\delta_v T_{zz}$ is linear in T_{zz} , we have

$$\langle \delta_v T_{zz}(\phi^\lambda) \rangle_\lambda = \delta_v \langle T_{zz}(\phi^\lambda) \rangle_\lambda = 0$$

on the plane, confirming the validity of the Ward identity for the bare tensor T_{zz} . In particular, it is important to note that the Ward identity for T_{ij} has no anomaly.

We now notice that if we define the quantity \hat{T}_{zz} to coincide with T_{zz}

$$\hat{T}_{zz} = T_{zz} \quad (33)$$

on the plane with trivial metric, and deform \hat{T}_{zz} according to the nontensor transformation law

$$\delta_v \hat{T}_{zz} \equiv -\frac{1}{12} \partial_z^3 v^z + v^i \partial_i \hat{T}_{zz} - 2(\partial_z v^z) \hat{T}_{zz}, \quad (34)$$

as we deform the metric along the flow of v , the extra term in the transformation law of \hat{T}_{zz} will exactly cancel the contributions coming from the contact terms (second and third lines) in the Ward identity. In terms of \hat{T}_{zz} , the Ward identity can therefore be expressed as

$$\begin{aligned} \left\langle \delta_v \hat{T}_{zz}(\phi^\lambda) \right\rangle_\lambda &= -\frac{1}{4\pi} \int d^2w h^{ww} \left\langle \hat{T}_{zz}(\phi^\lambda) \hat{T}_{ww}(\phi^\lambda) \right\rangle_\lambda \\ &= -\frac{1}{\pi} \int d^2w (\partial_{\bar{w}} v^w) \left\langle \hat{T}_{zz}(\phi^\lambda) \hat{T}_{ww}(\phi^\lambda) \right\rangle_\lambda, \end{aligned}$$

which is the form used in most treatments of conformal field theory. In particular, the energy-momentum tensor obtained from the common point-splitting renormalization of the operator product coincides with \hat{T}_{zz} , as can be seen from its transformation law.

It is possible to express the metric-dependence of \hat{T}_{zz} generated by the above transformation law directly as [15, 16]

$$\hat{T}_{zz} = T_{zz} + \frac{1}{12} t_{zz},$$

where

$$t_{zz} \equiv \partial_z \Gamma_{zz}^z - \frac{1}{2} (\Gamma_{zz}^z)^2.$$

Looking back to section 4, we see that \hat{T}_{zz} here coincides with the corresponding \hat{T}_{zz} of equation (16). Using the property

$$\partial_{\bar{z}} t_{zz} = -\frac{1}{2} g_{z\bar{z}} \partial_z R,$$

the redefined \hat{T}_{zz} is easily seen to still satisfy the conservation law

$$\langle \partial_{\bar{z}} \hat{T}_{zz} \rangle = 0$$

on a flat manifold.

9 Checking the change of variables theorem

Fundamental to path-integral derivations of conservation laws and Ward identities is the generalization (23)

$$\int [d\phi^\lambda] e^{-S(\phi^\lambda)} = \int [d\phi] e^{-S(\phi)} \quad (35)$$

of the change of variables theorem from finite to infinite dimensions. While this should be true if the concept of integration is to make sense in a chart-independent way, in the absence of a general proof it is something that should be verified case by case.

It is instructive to check this formula to second order around $\lambda = 0$ on the plane. As noted in section 7, the full measure, including Pauli-Villars fields, is independent of λ , so that differentiating the left hand side of (35) twice with respect to λ gives

$$\begin{aligned} \frac{d^2}{d\lambda^2} \Big|_{\lambda=0} \int [d\phi^\lambda] e^{-S(\phi^\lambda)} &= \frac{1}{4\pi} \int d^2x \langle \delta_v T_{ij}(\phi)_x \rangle \\ &+ \left(\frac{1}{4\pi} \right)^2 \int d^2x \int d^2y h^{ij}(x) h^{kl}(y) \langle T_{ij}(\phi)_x T_{kl}(\phi)_y \rangle, \end{aligned}$$

where

$$h^{ij} \equiv \delta_\lambda (f_\lambda^* g)^{ij} \Big|_{\lambda=0} = \partial^i v^j + \partial^j v^i.$$

For the change of variables formula to be valid to second order, this expression should be zero. But notice that this expression is just the integral over x of the Ward identity (24), laboriously verified in the previous section. We therefore find the required result

$$\frac{d^2}{d\lambda^2} \Big|_{\lambda=0} \int [d\phi^\lambda] e^{-S(\phi^\lambda)} = 0.$$

10 Relating the conformal anomaly and the Ward identity

The result of the previous section can also be interpreted as telling us that the second-order variation of the partition function is zero when we drag the metric along the flow of a vector field. Such a deformation does not change the curvature of an initially flat surface, and therefore, as expected, the Weyl anomaly did not contribute.

Let us now instead consider the change of

$$\int [d\phi]_g e^{-S(g,\phi)}$$

to second order under a deformation $h_{ij} \equiv \delta g_{ij}$ of the trivial metric $g_{ij} = \delta_{ij}$ on the plane, not necessarily generated by a vector field. Since in general this cannot be compensated by a change of variables, we do not expect the variation to be zero. In general, a second derivative will bring down up to two instances of the energy-momentum tensor from the exponent, so that we will have to calculate terms of the form $\langle T_{ij} T_{kl} \rangle$, for which we are forced to use the Pauli-Villars regularization of the previous sections. Since the Pauli-Villars measure $[d\phi]_g$ is invariant under variations of g , we can write

$$\begin{aligned} \delta_g^2 \int [d\phi]_g e^{-S(g,\phi)} &= \frac{1}{4\pi} \cdot \frac{1}{2} \int d^2x \sqrt{g} g_{kl} h^{kl} h^{ij} \langle T_{ij} \rangle \\ &+ \frac{1}{4\pi} \int d^2x \sqrt{g} h^{ij} \langle \delta_g T_{ij} \rangle \\ &+ \left(\frac{1}{4\pi} \right)^2 \int d^2x \sqrt{g} \int d^2y \sqrt{g} h^{ij}(x) h^{kl}(y) \langle T_{ij}(\phi)_x T_{kl}(\phi)_y \rangle. \end{aligned}$$

On the plane, $\sqrt{g} = 1$ and, as shown in section 5, $\langle T_{zz} \rangle$, $\langle T_{\bar{z}\bar{z}} \rangle$, and $\langle T_{z\bar{z}} \rangle$ are all zero with the Pauli-Villars measure, so that the first term vanishes. Remembering that T_{ij} depends on the metric, we calculate

$$\delta_g \langle T_{ij} \rangle = 2\pi \cdot \frac{1}{2} \delta_g (g_{ij} g^{kl}) \langle \partial_k \phi \partial_l \phi \rangle.$$

The expectation values $\langle \partial_k \phi \partial_l \phi \rangle$, are all linear combinations of $\langle T_{zz} \rangle$, $\langle T_{\bar{z}\bar{z}} \rangle$, $\langle T_{z\bar{z}} \rangle$ and $\langle \partial_z \phi \partial_{\bar{z}} \phi \rangle$, all of which vanish on the plane as shown in in section 5. So

$$\delta_g \langle T_{ij} \rangle = 0.$$

Let us now see how this Ward identity is related to the conformal anomaly discussed in sections 2 and 3. For simplicity we first consider a Weyl variation

$$\delta_\omega g_{ij} = 2 \delta\omega g_{ij}$$

of the flat metric. Then $h^{z\bar{z}} = h^{\bar{z}z} = -4 \delta\omega$, and the above formula becomes

$$\begin{aligned} \delta_\omega^2 \int [d\phi]_g e^{-S(g,\phi)} &= \left(\frac{1}{4\pi}\right)^2 \int d^2z \int d^2w 4 h^{z\bar{z}} h^{w\bar{w}} \langle T_{z\bar{z}} T_{w\bar{w}} \rangle \\ &= \left(\frac{1}{4\pi}\right)^2 \int d^2z \int d^2w 4 h^{z\bar{z}} h^{w\bar{w}} \left(-\frac{\pi}{12}\right) \partial_z \partial_{\bar{z}} \delta(z-w) \langle 1 \rangle \\ &= -\frac{1}{12\pi} \int d^2z \delta\omega \Delta \delta\omega \langle 1 \rangle \\ &= \frac{1}{24\pi} \int d^2x \delta\omega \delta R \langle 1 \rangle, \end{aligned}$$

where we used

$$R = -2 e^{-2\omega} \Delta\omega \quad \text{for} \quad g_{ij} = e^{2\omega} \delta_{ij}.$$

This formula concides precisely with the second variation of the formula (9) for the conformal anomaly

$$\delta_\omega \int [d\phi]_g e^{-S(\phi,g)} = \left(\frac{1}{24\pi} \int d^2x \sqrt{g} \delta\omega(x) R\right) \int [d\phi]_g e^{-S(\phi,g)}.$$

derived in two separate ways in sections 2 and 3.

For more generic deformations of the metric, we have

$$\begin{aligned} \delta_g^2 \int [d\phi]_g e^{-S(g,\phi)} &= \left(\frac{1}{4\pi}\right)^2 \int d^2x \sqrt{g} \int d^2y \sqrt{g} h^{ij}(x) h^{kl}(y) \langle T_{ij}(\phi)_x T_{kl}(\phi)_y \rangle \\ &= \left(\frac{1}{4\pi}\right)^2 \int d^2z \int d^2w \left\{ h^{zz} h^{ww} \left(\frac{1}{2}\right) \frac{1}{(z-w)^2} \right. \\ &\quad \left. + 2 h^{zz} h^{w\bar{w}} \left(\frac{\pi}{12}\right) \partial_z^2 \delta(z-w) \right. \\ &\quad \left. + h^{zz} h^{\bar{w}\bar{w}} \left(-\frac{\pi}{12}\right) \partial_z \partial_{\bar{z}} \delta(z-w) \right. \\ &\quad \left. + \dots \right\} \langle 1 \rangle, \end{aligned}$$

where we have inserted the contact term two-point functions derived before. To save space, we only wrote out the first three terms. Now, using

$$\pi \delta(z - w) = \partial_z \partial_{\bar{z}} \ln |z - w|^2,$$

and performing partial integrations, we find

$$\begin{aligned} & -\frac{1}{12} \left(\frac{1}{4\pi} \right)^2 \int d^2z \int d^2w \left(-\partial_z^2 h^{zz} - \partial_{\bar{z}}^2 h^{\bar{z}\bar{z}} + 2 \partial_z \partial_{\bar{z}} h^{z\bar{z}} \right) \times \\ & \quad \times \ln |z - w|^2 \left(-\partial_w^2 h^{ww} - \partial_{\bar{w}}^2 h^{\bar{w}\bar{w}} + 2 \partial_w \partial_{\bar{w}} h^{w\bar{w}} \right) \\ & = -\frac{1}{12} \left(\frac{1}{4\pi} \right)^2 \int d^2z \int d^2w \delta_g R(z) \ln |z - w|^2 \delta_g R(w). \end{aligned}$$

Notice that all the $\langle TT \rangle$ contact terms were necessary to obtain the correct curvature factors δR .

Our final result for the plane is [4, 18]

$$\delta_g^2 \int [d\phi]_g e^{-S(g,\phi)} = -\frac{1}{12} \left(\frac{1}{4\pi} \right)^2 \int d^2z \int d^2w \delta_g R(z) \ln |z - w|^2 \delta_g R(w) \langle 1 \rangle.$$

Notice that this result is entirely consistent with that of the previous section, since for a deformation of g by a vector field we have $\delta R = 0$, so that the right hand side vanishes.

11 Conclusion

In this article we provided a path integral presentation of the theory of a two-dimensional conformal scalar field.

We derived the conformal anomaly using an adaptation of the method of Fujikawa, which explains the anomaly as arising from a metric dependence of the path integral measure. We then re-derived the result using a Pauli-Villars path integral measure. Here the anomaly was shifted from the path integral measure, which became metric-independent, onto the trace of the energy-momentum tensor, which became nonzero.

In the path integral approach the energy-momentum tensor is a true coordinate-invariant quantity. We explained how this is related to the corresponding nontensor object found in the operator approach, obtaining an intuitive explanation for the anomalous behaviour of the latter within the

context of the path integral. The difference in behaviour was seen as due to a residual effect of the Pauli-Villars fields.

We confirmed by explicit calculation that the full energy-momentum tensor satisfies the classical Ward identities, and we explained how these relate to the anomalous Ward identities found in the operator approach. We concluded with a simple consistency check confirming the change of variables formula for the path integral, and we reviewed the relationship between the conformal anomaly and the energy-momentum two-point functions. Important in our derivations and checks were nontrivial contact terms arising in certain energy-momentum products.

It is our hope that this work may be helpful in sharpening the intuition of the reader regarding the origin, somewhat opaque in the operator approach, of some simple quantum or anomalous behaviours in conformal field theory. We have restricted attention to a scalar field, which provides a simple laboratory for showcasing various path integral techniques.

Acknowledgments

I would like to thank Dr. Miquel Dorca for very useful discussions. I would also like to thank Prof. Antal Jevicki and the Brown University Physics department for their support.

References

- [1] K. Fujikawa, *Path integral for gauge theories with fermions*, *Physical Review D* **21** No. 10 (1980), 2848-2858.
- [2] K. Fujikawa, *Path integral of relativistic strings*, *Physical Review D* **25** No. 10 (1982), 2584-2592.
- [3] M. Nakahara, *Geometry, Topology and Physics*, Institute of Physics Publishing, London, 1990.
- [4] P. Di Francesco, P. Mathieu and D. Sénéchal, *Conformal Field Theory*, Spinger-Verlag, New York (1996).

- [5] S. Albeverio, J. Jost, S. Paycha and S. Scarlatti, *A Mathematical Introduction to String Theory*, London Mathematical Society Lecture Note Series 225, Cambridge University Press, Cambridge (1997).
- [6] W. Pauli and F. Villars, *On the invariant regularization in relativistic quantum field theory*, *Reviews of Modern Physics* **21** (1949), 434-444.
- [7] A. Zee, *Quantum Field Theory in a Nutshell*, Princeton University Press, Princeton, 2003.
- [8] M.E. Peskin, D.V. Shroeder, *An Introduction to Quantum Field Theory* Perseus Books, Cambridge, Massachusetts, 1995.
- [9] A. Vilenkin, *Pauli-Villars regularization and trace anomalies*, *Il Nuovo Cimento* **44A** No. 3 (1978), 441-449.
- [10] N.D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, London (1982).
- [11] C.J.C. Burges, D.Z. Freedman, S. Davis and G.W. Gibbons, *Supersymmetry in anti-de-Sitter space*, *Annals of Physics* **167** (1986), 285-316.
- [12] R. Camporesi and A. Higuchi, *Stress-energy tensor in anti-de-Sitter spacetime*, *Physical Review D* **45** No. 10 (1992), 3591-3603.
- [13] D. Anselmi, *Covariant Pauli-Villars regularization of quantum gravity at the one loop order*, *Physical Review D* **48** (1993), 5751-5763.
- [14] E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis*, 4th ed., Cambridge University Press, Cambridge (1999).
- [15] T. Eguchi, *Conformal and current algebras on a general Riemann surface*, in Proceedings of the First Asia Pacific Workshop on High Energy Physics: *Conformal Field Theory, Anomalies and Superstrings*, B.E. Baaquie, editors ... [et al.], World Scientific, Singapore (1987).
- [16] T. Eguchi and H. Ooguri, *Conformal and current algebras on a general Riemann surface*, *Nuclear Physics* **B282** (1987), 308-328.
- [17] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, *Nuclear Physics* **B241** (1984), 333-380.

- [18] J. Polchinski, *String Theory*, Vol. 1, Cambridge University Press, Cambridge (1998).
- [19] K. Gawędzki, *Lectures on conformal field theory*, in *Quantum Fields and Strings: A Course for Mathematicians*, Vol. 2, P. Deligne, editors ... [et al.], American Mathematical Society, Providence (1999).
- [20] M. Henkel, *Conformal Invariance and Critical Phenomena*, Springer-Verlag, Berlin Heidelberg (1999).
- [21] A. Alvarez-Gaumé, C. Gomez, G. Moore, C. Vafa, *Strings in the operator formalism*, *Nuclear Physics* **B303** (1988), 455.
- [22] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, 5th ed., Alan Jeffrey, editor, Academic Press, San Diego (1994).
- [23] A. Cappelli and J.I. Latorre, *Perturbation theory of higher-spin conserved currents off criticality*, *Nuclear Physics* **B340** (1990), 659-691.
- [24] A. Cappelli, D. Friedan and J.I. Latorre, *c-Theorem and spectral representation*, *Nuclear Physics* **B352** (1991), 616-670.
- [25] L.S. Brown, *Quantum Field Theory*, Cambridge University Press, Cambridge (1992).
- [26] S. Forte and J. Latorre, *A proof of the irreversibility of renormalization group flows in four dimensions*, *Nuclear Physics* **B535** (1998), 709-728.