[hep-th/0503224](http://arxiv.org/abs/hep-th/0503224) TIT/HEP-536 TU-740 March, 2005

# Non(anti)commutative  $\mathcal{N} = (1, 1/2)$ Supersymmetric  $U(1)$  Gauge Theory

Takeo Araki<sup>1</sup>, Katsushi Ito<sup>2</sup> and Akihisa Ohtsuka<sup>2</sup>

<sup>1</sup> Department of Physics Tohoku University Sendai, 980-8578, Japan

<sup>2</sup>Department of Physics Tokyo Institute of Technology Tokyo, 152-8551, Japan

#### Abstract

We study a reduction of deformation parameters in non(anti)commutative  $\mathcal{N}=2$ harmonic superspace to those in non(anti)commutative  $\mathcal{N} = 1$  superspace. By this reduction we obtain the exact gauge and supersymmetry transformations in the Wess-Zumino gauge of non(anti)commutative  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory defined in the deformed harmonic superspace. We also find that the action with the first order correction in the deformation parameter reduces to the one in the  $\mathcal{N} = 1$  superspace by some field redefinition. We construct deformed  $\mathcal{N} = (1, 1/2)$  supersymmetry in  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in non(anti)commutative  $\mathcal{N}=1$  superspace.

### 1 Introduction

Supersymmetric field theories in deformed superspace[\[1,](#page-28-0) [2\]](#page-28-1) have been recently attracted much attentions in view of studying superstring effective field theories on the D-branes with graviphoton background [\[3,](#page-28-2) [4,](#page-28-3) [5\]](#page-29-0). Non(anti)commutative superspace is a deformed superspace with nonanticommutative Grassmann coordinates and the ∗-product. Field theories in non(anti)commutative superspace is constructed in terms of superfields and have been extensively studied  $[6][7][8]-[16]$  $[6][7][8]-[16]$  $[6][7][8]-[16]$  $[6][7][8]-[16]$  $[6][7][8]-[16]$ .

 $\mathcal{N}=1$  super Yang-Mills theory in the non(anti)commutative  $\mathcal{N}=1$  superspace has been formulated in [\[6\]](#page-29-1), where the deformation preserves the chiral structure of the theory. As in the commutative case, one can take the Wess-Zumino(WZ) gauge for vector superfields to write down the deformed action in terms of component fields. Since supersymmetry transformation cannot keep the WZ gauge, one may perform additional gauge transformation to recover the WZ gauge. In the nonanticommutative case, this gauge transformation induces the terms which depends on the deformation parameter C. Therefore, even if we redefine the component fields such that these fields transform canonically under the gauge transformation, the supersymmetry transformations receive the deformation due to non(anti)commutativity.

In a previous paper [\[7\]](#page-29-2), we wrote down the deformed action of  $\mathcal{N}=1$  supersymmetric  $U(N)$  gauge theory with matter fields in both fundamental and adjoint representations. We also examined invariance of the action under the  $\mathcal{N} = 1/2$  supersymmetry transformation. In particular, in the case of adjoint matter fields, we claimed that we could not find the (deformed) extended supersymmetry transformation and concluded that only the  $\mathcal{N} = 1/2$  supersymmetry is preserved. It was pointed out in [\[12\]](#page-30-1), however, that this conclusion had discrepancy with the general argument concerning the symmetry of the deformed  $\mathcal{N} = 2$  superspace whose Poisson structure constructed by the chiral supercharges  $Q^i_\alpha$  preserves at least the  $\mathcal{N} = (1, 0)$  supersymmetry of the theory generated by  $Q^1_{\alpha}$ ,  $Q^2_{\alpha}$ . Moreover, it is natural to expect that the action defined in the deformed harmonic superspace leads to the one in the deformed  $\mathcal{N}=1$  superspace by the reduction of deformation parameters of the harmonic superspace to the deformed  $\mathcal{N}=1$  superspace. In this case, it was also argued in [\[12\]](#page-30-1) that the deformation preserves the  $\mathcal{N} = (1, 1/2)$ 

supersymmetry generated by  $Q^1$ ,  $Q^2$  and  $\overline{Q}_2$ . The existence of  $\mathcal{N} = (1, 1/2)$  supersym-metry is partly supported by our recent work[\[13\]](#page-30-2), in which the deformed  $\mathcal{N} = (1,0)$ supersymmetry has been constructed in the  $\mathcal{N}=2$  supersymmetric  $U(1)$  gauge theory in non(anti)commutative harmonic superspace with generic (non-singlet) deformations. Thus we expect  $\mathcal{N} = (1, 1/2)$  supersymmetry in the  $\mathcal{N} = 2$  supersymmetric  $U(N)$  gauge theory in the non(anti)commutative  $\mathcal{N}=1$  superspace.

In this paper, we will construct deformed  $\mathcal{N} = (1, 1/2)$  supersymmetry in  $\mathcal{N} = 2$ supersymmetric  $U(1)$  gauge theory in non(anti)commutative  $\mathcal{N} = 1$  superspace. We compare this theory to the one in non(anti) commutative  $\mathcal{N}=2$  harmonic superspace obtained by the reduction to the deformed  $\mathcal{N} = 1$  superspace. By this reduction we find the exact gauge and supersymmetry transformations preserving the WZ gauge. We also find that these transformations and the  $O(C)$  action in the WZ gauge defined in the deformed harmonic superspace reduces to the one in the  $\mathcal{N}=1$  superspace by field redefinition.

This paper is organised as follows: In sect. 2, we review  $\mathcal{N}=2$  supersymmetric  $U(1)$ gauge theory in non(anti)commutative  $\mathcal{N} = 1$  superspace and non(anti)commutative  $\mathcal{N} = 2$  harmonic superspace with generic non-singlet deformation parameters C. For the latter superspace we construct  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in this superspace at  $O(C)$ . We then study the reduction of the deformation parameters in non(anti)commutative harmonic superspace to the deformed  $\mathcal{N}=1$  superspace. We find that the  $O(C)$  action reduces to the one defined in the deformed  $\mathcal{N} = 1$  superspace and argue invariance under  $\mathcal{N} = (1, 0)$  supersymmetry. In sect. 3, we investigate the reduced theory in more detail. We construct explicitly the exact gauge and  $\mathcal{N} = (1, 1/2)$ supersymmetry transformations of the component fields in the WZ gauge, where detailed calculations are explained in two appendices. After the field redefinition, we find deformed  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation which keeps the action in the deformed  $\mathcal{N}=1$  superspace invariant. Sect. 5 is devoted to discussion and conclusion. In appendix [A,](#page-19-0) we describe some useful reduction formulas of harmonic variables. The details of calculation of the exact  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation laws are presented in appendix [B.](#page-20-0)

## 2  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in non(anti)commutative superspaces

In this section we review  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in the deformed  $\mathcal{N} = 1$ superspace [\[6,](#page-29-1) [7\]](#page-29-2) and non(anti)commutative  $\mathcal{N} = 2$  harmonic superspace [\[12,](#page-30-1) [13,](#page-30-2) [14\]](#page-30-3).

## 2.1 The deformed  $\mathcal{N}=1$  superspace

Let  $(x^{\mu}, \theta^{\alpha}, \bar{\theta}^{\dot{\alpha}})$  be the supercoordinates of  $\mathcal{N}=1$  superspace[\[17\]](#page-30-4). Here  $\mu=0, 1, 2, 3$  are spacetime indices,  $\alpha, \dot{\alpha} = 1, 2$  the spinor indices. We study spacetime with the Euclidean signature so that chiral and antichiral fermions transform independently. We may call this  $\mathcal{N} = 1$  superspace as  $\mathcal{N} = (1/2, 1/2)$  superspace as in [\[12\]](#page-30-1). The non(anti)commutative  $\mathcal{N}=1$  superspace is introduced by imposing nonanticommutativity for Grassmann coordinates  $\theta^{\alpha}$ :

$$
\{\theta^{\alpha}, \theta^{\beta}\}_{*} = C^{\alpha\beta},\tag{1}
$$

whereas the chiral coordinates  $y^{\mu} = x^{\mu} + i\theta \sigma^{\mu} \bar{\theta}$  commute with other coordinates and  $\bar{\theta}^{\dot{\alpha}}$  (anti)commute. The \*-product  $f * g$  is defined by  $f * g = f \exp(P)g$ , where  $P =$  $-\frac{1}{2}$  $\frac{1}{2}\overleftarrow{Q}_{\alpha}C^{\alpha\beta}\overrightarrow{Q}_{\beta}$  defines the Poisson structure on the superspace. Since this Q-deformation preserves chirality, *i.e.* the Poisson structure  $P$  commutes with the supercovariant derivatives  $D_{\alpha} = \frac{\partial}{\partial \theta^{\alpha}} + i(\sigma^{\mu} \bar{\theta})_{\alpha} \partial_{\mu}$  and  $\bar{D}_{\dot{\alpha}} = -\frac{\partial}{\partial \bar{\theta}^{\dot{\alpha}}} - i(\theta \sigma^{\mu})^{\dot{\alpha}} \partial_{\mu}$ , we can define (anti)chiral superfields  $\Phi$  ( $\bar{\Phi}$ ) satisfying  $\bar{D}_{\dot{\alpha}}\Phi = 0$  ( $D_{\alpha}\bar{\Phi} = 0$ ). We can also introduce the vector superfield  $V(y, \theta, \theta)$  for representing gauge fields. If we take the WZ gauge for the vector superfield, we are able to write down the Lagrangian in terms of component fields by replacing the product to the ∗-product. The component fields do not transform canonically under the gauge transformation due to the deformation parameter  $C$ . But in [\[6,](#page-29-1) [7\]](#page-29-2), the field redefinition was found such that the component fields transform canonically under the gauge transformation.

For  $\mathcal{N} = 2 U(1)$  gauge theory, matter fields  $\Phi$ ,  $\bar{\Phi}$  in the adjoint representation (under the  $*$ -product) and the vector superfield V in the WZ gauge are given by

$$
\Phi(y,\theta) = A(y) + \sqrt{2}\theta\psi(y) + \bar{\theta}\bar{\theta}F(y),
$$
  

$$
\bar{\Phi}(\bar{y},\bar{\theta}) = \bar{A}(\bar{y}) + \sqrt{2}\bar{\theta}\bar{\psi}(\bar{y}) + \bar{\theta}\bar{\theta}(\bar{F} + 2iC^{\mu\nu}\partial_{\mu}(\bar{A}v_{\nu}))(\bar{y}),
$$

$$
V(y,\theta,\bar{\theta}) = -\theta\sigma^{\mu}\bar{\theta}v_{\mu}(y) + i\theta\theta\bar{\theta}\bar{\lambda}(y) - i\bar{\theta}\bar{\theta}\theta^{\alpha}\left(\lambda_{\alpha} + \frac{1}{2}\varepsilon_{\alpha\beta}C^{\beta\gamma}(\sigma^{\mu}\bar{\lambda})_{\gamma}v_{\mu}\right)(y) + \frac{1}{2}\theta\theta\bar{\theta}\bar{\theta}(D - i\partial^{\mu}v_{\mu})(y),
$$
\n(2)

where  $\bar{y}^{\mu} = x^{\mu} - i\theta\sigma^{\mu}\bar{\theta}$  is the anti-chiral coordinates. The deformed action is defined by

$$
S = \int d^4x d^2\theta d^2\bar{\theta}\bar{\Phi} * e^V * \Phi * e^{-V} + \frac{1}{16g^2} \left( \int d^2\theta W^\alpha * W_\alpha + \int d^2\bar{\theta}\bar{W}_{\dot{\alpha}} * \bar{W}^{\dot{\alpha}} \right). \tag{3}
$$

Here  $q$  denotes the gauge coupling constant. The chiral and anti-chiral field strength are

<span id="page-4-0"></span>
$$
W_{\alpha} = -\frac{1}{4}\bar{D}\bar{D}e^{-V}D_{\alpha}e^{V}, \quad \bar{W}_{\dot{\alpha}} = \frac{1}{4}DDe^{V}\bar{D}_{\dot{\alpha}}e^{-V}
$$
(4)

where the multiplication of superfields is defined by using the ∗-product. After rescaling V to 2gV and  $C^{\alpha\beta}$  to  $\frac{1}{2g}C^{\alpha\beta}$ , this deformed action is expressed as

$$
S = \int d^4x (\mathcal{L}^{(0)} + \mathcal{L}^{(1)}) \tag{5}
$$

where

$$
\mathcal{L}^{(0)} = -\frac{1}{4}v_{\mu\nu}(v^{\mu\nu} + \tilde{v}^{\mu\nu}) - i\bar{\lambda}\bar{\sigma}^{\mu}\partial_{\mu}\lambda + \frac{1}{2}D^2 - \partial^{\mu}\bar{A}\partial_{\mu}A - i\bar{\psi}\bar{\sigma}^{\mu}\partial_{\mu}\psi + \bar{F}F, \qquad (6)
$$

$$
\mathcal{L}^{(1)} = -\frac{i}{2} C^{\mu\nu} v_{\mu\nu} (\bar{\lambda}\bar{\lambda}) + \sqrt{2} C^{\alpha\beta} \psi_{\alpha} (\sigma^{\mu} \bar{\lambda})_{\beta} \partial_{\mu} \bar{A} + i C^{\mu\nu} v_{\mu\nu} \bar{A} F, \tag{7}
$$

and  $v_{\mu\nu} = \partial_{\mu}v_{\nu} - \partial_{\nu}v_{\mu}$ . This action is invariant under the gauge transformation  $\delta_{\lambda}v_{\mu} =$  $-\partial_\mu \lambda$ . Undeformed theory has  $\mathcal{N} = 2$  extended supersymmetry, where only  $\mathcal{N} = 1$ symmetry generated by  $Q_{\alpha}$  and  $\overline{Q}_{\dot{\alpha}}$  is manifest in  $\mathcal{N}=1$  superspace formalism. In deformed theory, the generator  $\overline{Q}_{\dot{\alpha}}$  does not commute with the Poisson structure P and is not a symmetry of the theory. On the other hand,  $Q_{\alpha}$  generates a symmetry of the theory. Since Q transformation does not preserve the WZ gauge, we need to do gauge transformation to retain the WZ gauge. The deformed supersymmetry transformation is

<span id="page-4-1"></span>
$$
\delta_{\xi}^{*} A = \sqrt{2} \xi \psi, \qquad \delta_{\xi}^{*} \bar{A} = 0,
$$
  
\n
$$
\delta_{\xi}^{*} \psi_{\alpha} = \sqrt{2} \xi_{\alpha} F, \qquad \delta_{\xi}^{*} \bar{\psi}_{\dot{\alpha}} = -i \sqrt{2} (\xi \sigma^{\mu})_{\dot{\alpha}} \partial_{\mu} \bar{A},
$$
  
\n
$$
\delta_{\xi}^{*} F = 0, \qquad \delta_{\xi}^{*} \bar{F} = i \sqrt{2} (\xi \sigma^{\mu} \partial_{\mu} \bar{\psi}) - 2 C^{\alpha \beta} \xi_{\alpha} \sigma^{\mu}_{\beta \dot{\alpha}} \partial_{\mu} (\bar{\lambda}^{\dot{\alpha}} \bar{A}),
$$
  
\n
$$
\delta_{\xi}^{*} v_{\mu} = i \xi \sigma_{\mu} \bar{\lambda},
$$
  
\n
$$
\delta_{\xi}^{*} \lambda_{\alpha} = (\sigma^{\mu \nu} \xi)_{\alpha} \left\{ v_{\mu \nu} + \frac{i}{2} C_{\mu \nu} (\bar{\lambda} \bar{\lambda}) \right\} + i \xi_{\alpha} D, \qquad \delta_{\xi}^{*} \bar{\lambda}_{\dot{\alpha}} = 0,
$$
  
\n
$$
\delta_{\xi}^{*} D = -(\xi \sigma^{\mu} \partial_{\mu} \bar{\lambda}).
$$
\n(8)

Remaining  $\mathcal{N} = 1$  supersymmetry, however, is not manifest in this formalism. In order to obtain this supersymmetry transformation more systematically, it is convenient to introduce non(anti)commutative extended superspace. In the next section, we will discuss  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in non(anti)commutative harmonic superspace.

### 2.2 Non(anti)commutative  $\mathcal{N}=2$  harmonic superspace

Next we review non(anti)commutative deformation of  $\mathcal{N}=2$  harmonic superspace and  $\mathcal{N}=2$  supersymmetric  $U(1)$  gauge theory in this superspace [\[13,](#page-30-2) [14\]](#page-30-3). Let  $(x^{\mu}, \theta_{\alpha}^{i}, \bar{\theta}_{\dot{\alpha}}^{i})$  be the coordinates of  $\mathcal{N} = 2$  (rigid) superspace. The index  $i = 1, 2$  labels the doublet of the  $SU(2)_R$  R-symmetry. The supersymmetry generators  $Q^i_{\alpha}$ ,  $\bar{Q}_{\dot{\alpha}i}$  and the supercovariant derivatives  $D^i_\alpha$ ,  $\bar{D}_{\dot{\alpha}i}$  are defined by

$$
Q_{\alpha}^{i} = \frac{\partial}{\partial \theta_{i}^{\alpha}} - i(\sigma^{\mu})_{\alpha\dot{\alpha}} \bar{\theta}^{\dot{\alpha}i} \frac{\partial}{\partial x^{\mu}}, \quad \bar{Q}_{\dot{\alpha}i} = -\frac{\partial}{\partial \bar{\theta}^{\dot{\alpha}i}} + i\theta_{i}^{\alpha}(\sigma^{\mu})_{\alpha\dot{\alpha}} \frac{\partial}{\partial x^{\mu}},
$$
  
\n
$$
D_{\alpha}^{i} = \frac{\partial}{\partial \theta_{i}^{\alpha}} + i(\sigma^{\mu})_{\alpha\dot{\alpha}} \bar{\theta}^{\dot{\alpha}i} \frac{\partial}{\partial x^{\mu}}, \quad \bar{D}_{\dot{\alpha}i} = -\frac{\partial}{\partial \bar{\theta}^{\dot{\alpha}i}} - i\theta_{i}^{\alpha}(\sigma^{\mu})_{\alpha\dot{\alpha}} \frac{\partial}{\partial x^{\mu}}.
$$
\n(9)

The  $\mathcal{N} = 2$  harmonic superspace [\[18\]](#page-30-5) is introduced by adding the harmonic variables  $u_i^{\pm}$  $\frac{1}{i}$  to the  $\mathcal{N}=2$  superspace coordinates. The variables  $u_i^{\pm}$  $i<sub>i</sub>$  form an  $SU(2)$  matrix and satisfy the conditions  $u^{+i}u_i^{-} = 1$  and  $\overline{u^{+i}} = u_i^{-}$  $\frac{1}{i}$ . The completeness condition for  $u_i^{\pm}$  $\frac{1}{i}$  reads  $u_i^+ u_j^- - u_j^+ u_i^- = \epsilon_{ij}$ . Using  $u_i^{\pm}$  $_i^{\pm}$ , the  $SU(2)_R$  indices can be projected into two parts with  $\pm 1$   $U(1)$ ( $\subset$   $SU(2)_R$ ) charges. For example, we define the supercovariant derivatives  $D^{\pm}_{\alpha}$ and  $\bar{D}_{\alpha}^{\pm}$  by  $D_{\alpha}^{\pm} = u_i^{\pm} D_{\alpha}^i$ ,  $\bar{D}_{\alpha}^{\pm} = u_i^{\pm} \bar{D}_{\alpha}^i$ .  $D_{\alpha}^i$  is solved by  $D_{\alpha}^{\pm}$  such as  $D_{\alpha}^{\pm} = u_i^{\pm} D_{\alpha}^- - u_i^- D_{\alpha}^+$ with the help of the completeness condition. In the harmonic superspace formalism, an important ingredient is an analytic superfield rather than the  $\mathcal{N}=2$  chiral superfield. An analytic superfield  $\Phi(x, \theta, \bar{\theta}, u)$  is defined by  $D^+_{\alpha} \Phi = \bar{D}^+_{\dot{\alpha}} \Phi = 0$ . It is convenient to write this analytic superfield in terms of analytic basis:  $x_A^\mu = x^\mu - i(\theta^i \sigma^\mu \bar{\theta}^j + \theta^j \sigma^\mu \bar{\theta}^i) u_i^+ u_j^$  $j^-, \theta_{\alpha}^{\pm} =$  $u_i^{\pm}$  $\frac{1}{i} \theta_{\alpha}^{i}$  and  $\bar{\theta}_{\dot{\alpha}}^{\pm} = u_{i}^{\pm}$  $\pm \bar{\theta}^i_\alpha$ . In this basis, an analytic superfield  $\Phi$  is functions of  $(x^\mu_A, \theta^+, \bar{\theta}^+, u)$ :  $\Phi = \Phi(x_A^{\mu}, \theta^+, \bar{\theta}^+, u).$ 

We now introduce the nonanticommutativity in the  $\mathcal{N}=2$  harmonic superspace by using the ∗-product:

$$
f * g(\theta) = f(\theta) \exp(P)g(\theta), \quad P = -\frac{1}{2}\overleftarrow{Q}_\alpha^i C_{ij}^{\alpha\beta} \overrightarrow{Q}_\beta^j,
$$
(10)

where  $C_{ij}^{\alpha\beta}$  is some constants. With this \*-product, we have following (anti)commutation relations:

$$
\{\theta_i^{\alpha}, \theta_j^{\beta}\}_* = C_{ij}^{\alpha\beta}, \quad [x_L^{\mu}, x_L^{\nu}]_* = [x_L^{\mu}, \theta_i^{\alpha}]_* = [x_L^{\mu}, \bar{\theta}^{\dot{\alpha}i}]_* = 0, \quad \{\bar{\theta}^{\dot{\alpha}i}, \bar{\theta}^{\dot{\beta}j}\}_* = \{\bar{\theta}^{\dot{\alpha}i}, \theta_j^{\alpha}\}_* = 0,
$$
\n(11)

where  $x_L^{\mu} \equiv x^{\mu} + i \theta_i \sigma^{\mu} \bar{\theta}^i$  are  $\mathcal{N} = 2$  chiral coordinates. The deformation parameter  $C_{ij}^{\alpha\beta}$  is symmetric under the exchange of pairs of indices  $(\alpha i)$ , $(\beta j)$ :  $C_{ij}^{\alpha\beta} = C_{ji}^{\beta\alpha}$ . We decompose the nonanticommutative parameter  $C_{ij}^{\alpha\beta}$  into the symmetric and antisymmetric parts with respect to the  $SU(2)$  indices, such as

$$
C_{ij}^{\alpha\beta} = C_{(ij)}^{\alpha\beta} + \frac{1}{4} \epsilon_{ij} \varepsilon^{\alpha\beta} C_s.
$$
 (12)

Here we denote  $A_{(i_1\cdots i_n)}$  by the symmetrized sum of  $A_{i_1\cdots i_n}$  over indices  $i_1, \cdots, i_n$ .  $C_{ij}^{\alpha\beta}$ ij with zero  $C_{(ii)}^{\alpha\beta}$  $\binom{\alpha\beta}{(ij)}$  corresponds to the singlet deformation [\[12,](#page-30-1) [15,](#page-30-6) [16\]](#page-30-0). Since P commutes with the supercovariant derivatives  $D$ , the chiral structure is preserved by this deformation. Since we will consider the non-singlet deformation in the following, we put  $C_s = 0$ .

We now construct the action of  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in this non(anti)commutative superspace. We introduce an analytic superfield  $V^{++}(\zeta, u)$  with  $\zeta = (x_A^{\mu}, \theta^+, \bar{\theta}^+)$  by covariantizing the harmonic derivative  $D^{++} = u^{+i} \frac{\partial}{\partial u^{-i}} - 2i \theta^+ \sigma^{\mu} \bar{\theta}^+ \frac{\partial}{\partial x_A^{\mu}} +$ A  $\theta^{+\alpha}\frac{\partial}{\partial\theta^{-\alpha}} + \bar{\theta}^{+\dot{\alpha}}\frac{\partial}{\partial\theta^{-\dot{\alpha}}} \to \nabla^{++} = D^{++} + iV^{++}$ . Generalizing the construction in [\[19,](#page-30-7) [20\]](#page-30-8), the action is given by

$$
S_{*} = \frac{1}{2} \sum_{n=2}^{\infty} \int d^{4}x d^{8} \theta du_{1} \dots du_{n} \frac{(-i)^{n}}{n} \frac{V^{++}(1) * \dots * V^{++}(n)}{(u_{1}^{+} u_{2}^{+}) \dots (u_{n}^{+} u_{1}^{+})},
$$
(13)

<span id="page-6-0"></span>where  $V^{++}(i) = V^{++}(\zeta_i, u_i)$ ,  $\zeta_i = (x_A, \theta_i^+, \bar{\theta}_i^+)$  and  $d^8\theta = d^4\theta^+d^4\theta^-$  with  $d^4\theta^{\pm} = d^2\theta^{\pm}d^2\bar{\theta}^{\pm}$ . The harmonic integral is defined by the rules: (i)  $\int du f(u) = 0$  for  $f(u)$  with non-zero  $U(1)$  charge. (ii)  $\int du_1 = 1$ . (iii)  $\int du u_{(i_1}^+ \cdots u_{i_n}^+ u_{j_1}^ \bar{j}_1 \cdots \bar{i}_{j_n} = 0$ ,  $(n \ge 1)$ . The action [\(13\)](#page-6-0) is invariant under the gauge transformation

<span id="page-6-1"></span>
$$
\delta_{\Lambda}^* V^{++} = -D^{++} \Lambda + i[\Lambda, V^{++}]_*,\tag{14}
$$

with an analytic superfield  $\Lambda$ . The generic superfield  $V^{++}(\zeta, u)$  includes infinitely many auxiliary fields. Most of these fields are gauged away except the lowest component fields in the harmonic expansion. One can take the WZ gauge

$$
V_{WZ}^{++}(x_A, \theta^+, \bar{\theta}^+, u) = -i\sqrt{2}(\theta^+)^2 \bar{\phi}(x_A) + i\sqrt{2}(\bar{\theta}^+)^2 \phi(x_A) - 2i\theta^+ \sigma^\mu \bar{\theta}^+ A_\mu(x_A) + 4(\bar{\theta}^+)^2 \theta^+ \psi^i(x_A) u_i^- - 4(\theta^+)^2 \bar{\theta}^+ \bar{\psi}^i(x_A) u_i^- + 3(\theta^+)^2 (\bar{\theta}^+)^2 D^{ij}(x_A) u_i^- u_j^-,
$$
(15)

which is convenient to study the theory in the component formalism.

The component action  $S_*$  in the WZ gauge can be expanded in a power series of the deformation parameter C. In [\[14\]](#page-30-3), we have computed the  $O(C)$  action explicitly. The quadratic part  $S_{*,2}$  in  $S_*$  is the same as the commutative one:

$$
S_{*,2} = \int d^4x \left\{ -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{4} F_{\mu\nu} \tilde{F}^{\mu\nu} + \phi \partial^2 \bar{\phi} - i \psi^i \sigma^\mu \partial_\mu \bar{\psi}_i + \frac{1}{4} D^{ij} D_{ij} \right\}.
$$
 (16)

The cubic part  $S_{*,3}$  in  $S_*$  is of order  $O(C)$  and given by

$$
S_{*,3} = \int d^4x \left[ -\frac{2\sqrt{2}}{3} i C^{\alpha\beta}_{(ij)} \psi^i_{\alpha} (\sigma^\nu \partial_\nu \bar{\psi}^j)_{\beta} \bar{\phi} - 2\sqrt{2} i C^{\alpha\beta}_{(ij)} \psi^i_{\alpha} (\sigma^\nu \bar{\psi}^j)_{\beta} \partial_\nu \bar{\phi} \right. + \frac{2}{3} i C^{\alpha\beta}_{(ij)} A_\mu (\sigma^\mu \bar{\psi}^i)_{\alpha} (\sigma^\nu \partial_\nu \bar{\psi}^j)_{\beta} - i C^{\mu\nu}_{(ij)} \bar{\psi}^i \bar{\psi}^j F_{\mu\nu} + \sqrt{2} C^{\mu\nu}_{(ij)} D^{ij} A_\mu \partial_\nu \bar{\phi} + \frac{1}{\sqrt{2}} C^{\mu\nu}_{(ij)} D^{ij} F_{\mu\nu} \bar{\phi} \right]. \tag{17}
$$

Note that here we have already dropped the  $C_s$  dependent terms. We will refer  $S_{*,2} + S_{*,3}$ as the  $O(C)$  action.

In the commutative case, the gauge parameter  $\Lambda = \chi(x_A)$  preserves the WZ gauge and gives rise to the gauge transformation for component fields. In the non(anti)commutative case, however, the gauge transformation [\(14\)](#page-6-1) with the same gauge parameter does not preserve the WZ gauge because of the C-dependent terms arising from the commutator. In order to preserve the WZ gauge, one must include the C-dependent terms. The gauge parameter is shown to take the form

$$
\lambda_C(\zeta, u) = \chi(x_A) + \theta^+ \sigma^\mu \bar{\theta}^+ \lambda_\mu^{(-2)}(x_A, u; C) + (\bar{\theta}^+)^2 \lambda^{(-2)}(x_A, u; C) \n+ (\bar{\theta}^+)^2 \theta^{+\alpha} \lambda_\alpha^{(-3)}(x_A, u; C) + (\theta^+)^2 (\bar{\theta}^+)^2 \lambda^{(-4)}(x_A, u; C),
$$
\n(18)

which has been determined by solving the WZ gauge preserving conditions expanded in harmonic modes [\[14\]](#page-30-3). The gauge transformation is also fully determined, which reads

<span id="page-7-0"></span>
$$
\delta_{\lambda_C}^* A_\mu = -\partial_\mu \chi + O(C^2),
$$

$$
\delta_{\lambda_C}^* \phi = O(C^2),
$$
\n
$$
\delta_{\lambda_C}^* \psi_{\alpha i} = \frac{2}{3} (\varepsilon C_{(ij)} \sigma^\mu \bar{\psi}^j)_{\alpha} \partial_\mu \chi + O(C^2),
$$
\n
$$
\delta_{\lambda_C}^* D_{ij} = 2\sqrt{2} C_{(ij)}^{\mu\nu} \partial_\mu \chi \partial_\nu \bar{\phi} + O(C^2),
$$
\n
$$
\delta_{\lambda_C}^* \text{(others)} = 0.
$$
\n(19)

The  $O(C)$  action is invariant under the  $O(C)$  gauge transformation [\(19\)](#page-7-0).

<span id="page-8-0"></span>These gauge transformations are not canonical. But if we redefine the component fields such as

$$
\hat{A}_{\mu} = A_{\mu} + O(C^2),
$$
\n
$$
\hat{\phi} = \phi + O(C^2), \quad \hat{\phi} = \bar{\phi},
$$
\n
$$
\hat{\psi}_{\alpha i} = \psi_{\alpha i} + \frac{2}{3} (\varepsilon C_{(ij)} \sigma^{\mu} \bar{\psi}^j)_{\alpha} A_{\mu} + O(C^2), \quad \hat{\bar{\psi}}^{\dot{\alpha}} = \bar{\psi}^{\dot{\alpha}},
$$
\n
$$
\hat{D}_{ij} = D_{ij} + 2\sqrt{2} C_{(ij)}^{\mu\nu} A_{\mu} \partial_{\nu} \bar{\phi} + O(C^2),
$$
\n(20)

<span id="page-8-1"></span>the newly defined fields are shown to transform canonically:  $\delta^*_{\lambda_C} \hat{A}_\mu = -\partial_\mu \chi$ ,  $\delta^*_\lambda$  $\chi^*_{c}$  (others) = 0. In terms of redefined fields, the  $O(C)$  action can be written as

$$
S_{*,2} + S_{*,3} = \int d^4x \left[ -\frac{1}{4} \hat{F}_{\mu\nu} (\hat{F}^{\mu\nu} + \tilde{\hat{F}}^{\mu\nu}) + \hat{\phi} \partial^2 \hat{\phi} - i \hat{\psi}^i \sigma^\mu \partial_\mu \hat{\psi}_i + \frac{1}{4} \hat{D}^{ij} \hat{D}_{ij} \right. \\ - 2\sqrt{2} i C_{(ij)}^{\alpha\beta} \hat{\psi}_\alpha^i (\sigma^\mu \hat{\bar{\psi}}^j)_\beta \partial_\mu \hat{\phi} - \frac{2\sqrt{2}}{3} i C_{(ij)}^{\alpha\beta} \hat{\psi}_\alpha^i (\sigma^\mu \partial_\mu \hat{\bar{\psi}}^j)_\beta \hat{\phi} \\ - i C_{(ij)}^{\mu\nu} \hat{\bar{\psi}}^i \hat{\psi}^j \hat{F}_{\mu\nu} + \frac{1}{\sqrt{2}} C_{(ij)}^{\mu\nu} \hat{D}^{ij} \hat{F}_{\mu\nu} \hat{\phi} + O(C^2) \right], \tag{21}
$$

where  $\hat{F}_{\mu\nu} = \partial_{\mu} \hat{A}_{\nu} - \partial_{\nu} \hat{A}_{\mu}$ .

The  $O(C)$  deformed  $\mathcal{N}=2$  supersymmetry transformation of the component fields in the WZ gauge is determined in a similar way[\[13\]](#page-30-2). After the field redefinition [\(20\)](#page-8-0), the transformation is given by

<span id="page-8-2"></span>
$$
\delta_{\xi}^{*}\hat{\phi} = -\sqrt{2}i\xi^{i}\hat{\psi}_{i} - \frac{8}{3}i(\xi^{j}\varepsilon C_{(jk)}\hat{\psi}^{k})\hat{\phi} + O(C^{2}),
$$
\n
$$
\delta_{\xi}^{*}\hat{\phi} = 0,
$$
\n
$$
\delta_{\xi}^{*}\hat{A}_{\mu} = i\xi^{i}\sigma_{\mu}\hat{\psi}_{i} + 2\sqrt{2}i(\xi^{j}\varepsilon C_{(jk)}\sigma_{\mu}\hat{\psi}^{k})\hat{\phi} + O(C^{2}),
$$
\n
$$
\delta_{\xi}^{*}\hat{\psi}^{\alpha i} = -(\xi^{i}\sigma^{\mu\nu})^{\alpha}\hat{F}_{\mu\nu} - \hat{D}^{ij}\xi_{j}^{\alpha} - i(\xi^{i}\sigma_{\mu\nu})^{\alpha}C_{(jk)}^{\mu\nu}(\hat{\psi}^{j}\hat{\psi}^{k}) + 2\sqrt{2}\hat{D}^{(ij}(\xi^{k})\varepsilon C_{(jk)})^{\alpha}\hat{\phi} - \left\{2\sqrt{2}(\xi^{j}\varepsilon C_{(jk)}\sigma^{\mu\nu})^{\alpha} + \frac{2\sqrt{2}}{3}(\xi^{j}\sigma^{\mu\nu}\varepsilon C_{(jk)})^{\alpha} + \sqrt{2}C_{(jk)}^{\mu\nu}\xi^{\alpha j}\right\}\epsilon^{ki}\hat{\phi}\hat{F}_{\mu\nu} + O(C^{2}),
$$

$$
\delta_{\xi}^{*} \hat{\psi}_{\dot{\alpha}}^{i} = +\sqrt{2} (\xi^{i} \sigma^{\nu})_{\dot{\alpha}} \partial_{\nu} \hat{\phi} + 2 (\xi^{j} \varepsilon C_{(jk)} \sigma^{\nu})_{\dot{\alpha}} \partial_{\nu} (\hat{\phi}^{2}) \epsilon^{ki} + O(C^{2}),
$$
\n
$$
\delta_{\xi}^{*} \hat{D}^{ij} = -2i \xi^{(i} \sigma^{\nu} \partial_{\nu} \hat{\psi}^{j)}
$$
\n
$$
- 6\sqrt{2} i \epsilon^{k(l} \partial_{\nu} \{ (\xi^{i} \varepsilon C_{(kl)} \sigma^{\nu} \hat{\psi}^{j)}) \hat{\phi} \} + 2\sqrt{2} i \epsilon^{il} \epsilon^{jm} (\xi^{k} \varepsilon C_{(lm)} \sigma^{\nu} \hat{\psi}_{k}) \partial_{\nu} \hat{\phi} + O(C^{2}).
$$
\n(22)

### 2.3 Reduction to the deformed  $\mathcal{N}=1$  superspace

Since the deformed  $\mathcal{N} = 2$  action [\(5\)](#page-4-0) in the deformed  $\mathcal{N} = 1$  superspace contains only  $O(C)$  corrections, it is important to compare  $\mathcal{N} = 1$  superspace formalism to non(anti)commutative harmonic superspace formalism. We will examine this correspondence in terms of component fields because it is difficult to find the relationship between  $(\Phi,\bar{\Phi},V)$  and  $V_{WZ}^{++}$  explicitly at the superfield level.

We impose the condition for the deformation parameters  $C_{\alpha\beta}^{ij}$  such as

<span id="page-9-2"></span>
$$
C_{ij}^{\alpha\beta} = C_{11}^{\alpha\beta} \delta_i^1 \delta_j^1. \tag{23}
$$

This reduction of the parameters implies that only the  $\theta_{\alpha}^{i}$  coordinates are nonanticommutative and  $\theta_{\alpha}^2$  are ordinary Grassmann coordinates. Then the action [\(21\)](#page-8-1) with redefined component fields becomes

<span id="page-9-0"></span>
$$
S_{*,2} + S_{*,3} = \int d^4x \left[ -\frac{1}{4} \hat{F}_{\mu\nu} (\hat{F}^{\mu\nu} + \tilde{F}^{\mu\nu}) - i \hat{\psi}^i \sigma^\mu \partial_\mu \hat{\psi}_i + \hat{\phi} \partial^\mu \partial_\mu \hat{\phi} + \frac{1}{4} \hat{D}_{ij} \hat{D}^{ij} \right. \\ - \frac{2\sqrt{2}}{3} i C_{11}^{\alpha\beta} \hat{\psi}_\alpha^1 (\sigma^\nu \partial_\nu \hat{\psi}^1)_{\beta} \hat{\phi} - 2\sqrt{2} i C_{11}^{\alpha\beta} \hat{\psi}_\alpha^1 (\sigma^\nu \hat{\psi}^1)_{\beta} \partial_\nu \hat{\phi} \\ - i C_{11}^{\mu\nu} \hat{\psi}^1 \hat{\psi}^1 \hat{F}_{\mu\nu} + \frac{1}{\sqrt{2}} C_{11}^{\mu\nu} \hat{D}^{11} \hat{F}_{\mu\nu} \hat{\phi} \right]. \tag{24}
$$

<span id="page-9-1"></span>Since both component fields  $(\hat{\phi}, \hat{\phi}, \hat{\psi}^i, \hat{\psi}^i, \hat{D}^{ij})$  and  $(A, \bar{A}, \psi, \bar{\psi}, \lambda, \bar{\lambda}, v_\mu, F, \bar{F}, D)$  transform canonically under the gauge transformation, these fields must be related to each other in gauge invariant way. By comparing [\(5\)](#page-4-0) and [\(24\)](#page-9-0), we find

$$
v_{\mu} = \hat{A}_{\mu}, \quad A = -i\hat{\phi}, \quad \bar{A} = i\hat{\phi},
$$
  
\n
$$
D = i\hat{D}^{12}, \quad F = -\frac{1}{\sqrt{2}}\hat{D}^{11}, \quad \bar{F} = -\frac{1}{\sqrt{2}}(\hat{D}^{22} - \sqrt{2}C_{11}^{\mu\nu}\hat{F}_{\mu\nu}\hat{\phi}),
$$
  
\n
$$
\lambda^{\beta} = \hat{\psi}^{2\beta} - \frac{2\sqrt{2}}{3}C_{11}^{\alpha\beta}\hat{\psi}_{\alpha}^1\hat{\phi}, \quad \psi = \hat{\psi}^1, \quad \bar{\psi} = \hat{\psi}_1, \quad \bar{\lambda} = \hat{\psi}_2.
$$
\n(25)

where the deformation parameter  $C_{11}^{\alpha\beta}$  is related to  $C^{\alpha\beta}$  by

<span id="page-9-3"></span>
$$
C_{11}^{\alpha\beta} = \frac{1}{2}C^{\alpha\beta}.
$$
\n(26)

Note that for  $C = 0$  we get the field redefinition for undeformed theory.

We now consider the deformed supersymmetry transformations. Taking the Grassmann parameters as  $(\xi^1, \xi^2) = (-\eta, \xi)$  and using the supersymmetry transformations [\(22\)](#page-8-2) and identifications [\(25\)](#page-9-1), it is shown that the transformation associated with  $\xi$  becomes [\(8\)](#page-4-1). The  $Q^2$  supersymmetry transformation associated with the parameter  $\eta$  would give a remaining transformation. But it turns out that  $O(C)$  transformation with the same identification does not keep the action invariant. It is necessary to introduce  $O(C^2)$  correction to the  $Q^2$  supersymmetry transformation. We found that the following transformation keep the action invariant:

<span id="page-10-0"></span>
$$
\delta_{\eta}^{*} A = \sqrt{2} \eta \lambda - 2i \eta \varepsilon C \psi \bar{A}, \qquad \delta_{\eta}^{*} \bar{A} = 0,
$$
  
\n
$$
\delta_{\eta}^{*} \psi^{\alpha} = (\eta \sigma_{\mu\nu})^{\alpha} \left\{ v^{\mu\nu} + \frac{i}{2} C^{\mu\nu} (\bar{\lambda} \bar{\lambda} - 2F \bar{A}) \right\} + i \eta^{\alpha} D, \qquad \delta_{\eta}^{*} \bar{\psi}_{\dot{\alpha}} = -(\eta \varepsilon C \sigma^{\nu})_{\dot{\alpha}} \partial_{\nu} (\bar{A}^{2}),
$$
  
\n
$$
\delta_{\eta}^{*} F = \sqrt{2} i \eta \sigma^{\nu} \partial_{\nu} \bar{\lambda}, \qquad \delta_{\eta}^{*} \bar{F} = 2\sqrt{2} i \det C \eta \sigma^{\nu} \partial_{\nu} (\bar{\lambda} \bar{A}^{2}),
$$
  
\n
$$
\delta_{\eta}^{*} v_{\mu} = -i \eta \sigma_{\mu} \bar{\psi} + \sqrt{2} (\eta \varepsilon C \sigma_{\mu} \bar{\lambda}) \bar{A},
$$
  
\n
$$
\delta_{\eta}^{*} \lambda^{\alpha} = \sqrt{2} \eta^{\alpha} \bar{F} + \sqrt{2} (\eta \varepsilon C)^{\alpha} \bar{A} D - \sqrt{2} i (\eta \sigma^{\mu\nu} \varepsilon C)^{\alpha} v_{\mu\nu} \bar{A} + \sqrt{2} \det C \eta^{\alpha} \bar{\lambda} \bar{\lambda} \bar{A},
$$
  
\n
$$
\delta_{\eta}^{*} \bar{\lambda}_{\dot{\alpha}} = -\sqrt{2} i (\eta \sigma^{\nu})_{\dot{\alpha}} \partial_{\nu} \bar{A},
$$
  
\n
$$
\delta_{\eta}^{*} D = -\eta \sigma^{\nu} \partial_{\nu} \bar{\psi} + \sqrt{2} i \eta \varepsilon C \sigma^{\nu} \partial_{\nu} (\bar{\lambda} \bar{A}),
$$
  
\n(27)

where  $(\varepsilon C)_{\alpha}{}^{\beta} \equiv \varepsilon_{\alpha\gamma} C^{\gamma\beta}$ . Hence we have found the  $\mathcal{N} = (1,0)$  supersymmetry of the  $\mathcal{N}=2$  action in the deformed  $\mathcal{N}=1$  superspace.

In  $(27)$ , we add  $O(C^2)$  correction to the transformation by hand. This correction is not unique because the action [\(5\)](#page-4-0) is invariant under the transformation  $\tilde{\delta}_{\eta}$  with

$$
\tilde{\delta}_{\eta} \lambda_{\alpha} = \eta_{\alpha} f_1(\bar{A}) F,
$$
  
\n
$$
\tilde{\delta}_{\eta} \bar{F} = i f_1(\bar{A}) (\eta \sigma^{\mu} \partial_{\mu} \bar{\lambda}),
$$
\n(28)

for arbitrary function  $f_1(\overline{A})$  of  $\overline{A}$ . In the next section, we will show that we are able to construct exact supersymmetry transformation for the reduced deformation parameter [\(23\)](#page-9-2). If we use the field redefinition [\(25\)](#page-9-1), we can fix  $f_1(\overline{A}) = 0$ . The result is shown to be exactly equal to [\(27\)](#page-10-0).

We note also that  $\bar{Q}_2$  supersymmetry transformation commutes with the Poisson structure P. The theory is expected to have  $\mathcal{N} = (1, 1/2)$  extended supersymmetry [\[12\]](#page-30-1). In the next section we will construct the exact  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation in the framework of the harmonic superspace formalism.

## <span id="page-11-1"></span>3  $\mathcal{N} = (1, 1/2)$  Supersymmetry

In this section, we will present the  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation generated by  $Q^1$ ,  $Q^2$  and  $\bar{Q}_2$ , within the harmonic superspace formalism. Under the restriction of the deformation parameter [\(23\)](#page-9-2), we can determine the  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation laws exactly. The main concern is the contribution from the deformed gauge transformation to retain the WZ gauge. Although the calculation is much simpler than the case of the generic deformation parameter, it turns out to be considerably lengthy even under the restriction. On the other hand, one finds that the determination of the exact gauge transformation laws is accomplished in a similar way but is much easier than the supersymmetry. Therefore, in order to illustrate how to determine the exact transformations, first we derive in sect. [3.1](#page-11-0) the exact gauge transformation laws in [\[14\]](#page-30-3), restricting ourselves to the reduced deformation parameter [\(23\)](#page-9-2). The exact  $\mathcal{N} = (1, 1/2)$ supersymmetry transformation is given in sect. [3.2,](#page-14-0) though the details of the actual calculation are presented in appendix [B.](#page-20-0) From the exact gauge transformation, we can find a field redefinition that leads to the canonical component gauge transformation. We also give the  $\mathcal{N} = (1, 1/2)$  transformation after this field redefinition.

#### <span id="page-11-0"></span>3.1 Determination of the exact transformation laws

In this subsection, in order to demonstrate how to determine the exact transformation laws under the restriction [\(23\)](#page-9-2), we will derive the exact gauge transformation laws in [\[14\]](#page-30-3). We will denote the analytic gauge parameter as

<span id="page-11-2"></span>
$$
\Lambda(\zeta, u) = \chi(x_A) + \bar{\theta}_{\dot{\alpha}}^+ \lambda^{(0,1)\dot{\alpha}}(x_A, u) + \theta^{+\alpha} \lambda_{\alpha}^{(1,0)}(x_A, u) + (\bar{\theta}^+)^2 \lambda^{(0,2)}(x_A, u) \n+ (\theta^+)^2 \lambda^{(2,0)}(x_A, u) + \theta^+ \sigma^\mu \bar{\theta}^+ \lambda_{\mu}^{(1,1)}(x_A, u) + (\bar{\theta}^+)^2 \theta^{+\alpha} \lambda_{\alpha}^{(1,2)}(x_A, u) \n+ (\theta^+)^2 \bar{\theta}_{\dot{\alpha}}^+ \lambda^{(2,1)\dot{\alpha}}(x_A, u) + (\theta^+)^2 (\bar{\theta}^+)^2 \lambda^{(2,2)}(x_A, u),
$$
\n(29)

where  $\lambda^{(n,m)}(x_A, u)$  is the  $(\theta^+)^n(\bar{\theta}^+)^m$ -component.

The equations to determine the deformed gauge transformation are given in [\[14\]](#page-30-3):

<span id="page-12-0"></span>
$$
-2i\delta_{\Lambda}^{*}A_{\mu} = 2i\partial_{\mu}\chi + 2\sqrt{2}iC^{(+-)\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\partial_{\nu}\chi\bar{\phi} - \partial^{++}\lambda_{\mu}^{(1,1)} - \sqrt{2}C^{++\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\lambda_{\nu}^{(1,1)}\bar{\phi},
$$
\n(30)

$$
\sqrt{2}i\delta_{\Lambda}^*\phi = -2iC^{(+-)\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\partial_{\nu}\chi A^{\mu} -C^{++\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\lambda_{\nu}^{(1,1)}A^{\mu} - \partial^{++}\lambda^{(0,2)},
$$
\n(31)

$$
3\delta_{\Lambda}^{*}D^{ij}u_{i}^{-}u_{j}^{-} = 4\sqrt{2}C^{-\mu\nu}\partial_{\mu}\chi \partial_{\nu}\bar{\phi}
$$

$$
-i\partial^{\mu}\lambda_{\mu}^{(1,1)} - \sqrt{2}iC^{(+-)\alpha\beta}(\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\partial_{\nu}(\lambda_{\mu}^{(1,1)}\bar{\phi}) - \partial^{++}\lambda^{(2,2)}, \tag{32}
$$

$$
4\delta_{\Lambda}^{*}\psi_{\alpha}^{i}u_{i}^{-} = -4C^{(+-)\beta\gamma}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\beta\gamma}\partial_{\nu}\chi\,(\sigma^{\mu}\bar{\psi}^{i})_{\alpha}u_{i}^{-} - 2iC^{++\beta\gamma}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\beta\gamma}\lambda_{\nu}^{(1,1)}(\sigma^{\mu}\bar{\psi}^{i})_{\alpha}u_{i}^{-} - \partial^{++}\lambda_{\alpha}^{(1,2)} - 2\sqrt{2}(\varepsilon C^{++}\lambda^{(1,2)})_{\alpha}\bar{\phi}.
$$
\n(33)

Here we have already set  $C_s = 0$ . The transformation laws for  $\bar{\phi}$  and  $\bar{\psi}_i$  are not deformed, because their equations are not affected by the ∗-product.

Now we take into account the restriction [\(23\)](#page-9-2). We have  $C^{(+)}^{\alpha\beta} = C_{11}^{\alpha\beta} u^{+1} u^{-1}$  and so on. First we find that eq.[\(30\)](#page-12-0) is solved in terms of power series of  $u^{+1}$  and  $u^{-1}$  by

<span id="page-12-1"></span>
$$
\delta_{\Lambda}^* A_{\mu} = -\partial_{\mu} \chi, \tag{34}
$$

$$
\lambda_{\mu}^{(1,1)} = \sum_{n=1}^{\infty} \lambda_{\mu}^{(n)} (u^{+1})^{n-1} (u^{-1})^{n+1}, \tag{35}
$$

where

$$
\lambda_{\mu}^{(n)} \equiv -i \frac{(-2\sqrt{2})^n}{(n+1)!} \left( \overbrace{C_{11}}^n \overbrace{c_{11}}^n \cdots \overbrace{c_{11}}^n \right)^{\alpha \beta} \left( \sigma_{\mu} \overline{\sigma}^{\nu} \varepsilon \right)_{\alpha \beta} \overline{\phi}^n \partial_{\nu} \chi. \tag{36}
$$

Note that the harmonic expansion of  $(u^{+1})^n(u^{-1})^m$  is

$$
(u^{+1})^{n}(u^{-1})^{m} = u^{+(1} \cdots u^{+1} u^{-1} \cdots u^{-1}) = \delta_{(i_{1}}^{1} \cdots \delta_{i_{n}}^{1} \delta_{j_{1}}^{1} \cdots \delta_{j_{m}}^{1} u^{+(i_{1}} \cdots u^{+i_{n}} u^{-j_{1}} \cdots u^{-j_{m})}.
$$
\n(37)

Eq.[\(34\)](#page-12-1) is directly checked by substituting  $\lambda_{\mu}^{(1,1)}$  into the right hand side of eq.[\(30\)](#page-12-0), noting that  $\lambda_{\mu}^{(n)}$  obeys

$$
C_{11}^{\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\lambda_{\nu}^{(n)}\bar{\phi} = -\frac{n+2}{\sqrt{2}}\lambda_{\mu}^{(n+1)} \qquad (n \ge 1).
$$
 (38)

Substituting eq.[\(35\)](#page-12-1) into [\(31\)](#page-12-0) , we find that all the terms in the right hand side of eq.[\(31\)](#page-12-0) is gauged away with  $\lambda^{(2,0)}$ , because every term has at least one  $u^+$ . This leads to

$$
\delta_{\Lambda}^* \phi = 0. \tag{39}
$$

Although the precise form of  $\lambda^{(0,2)}$  is not needed, it is determined as

$$
\lambda^{(0,2)} = -i C_{11}^{\alpha\beta} (\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta} \partial_{\nu} \chi A_{\mu} (u^{-1})^2 + \sum_{n=2}^{\infty} \frac{-1}{(n+1)} C_{11}^{\alpha\beta} (\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta} \lambda_{\nu}^{(n-1)} A_{\mu} (u^{+1})^{n-1} (u^{-1})^{n+1}.
$$
 (40)

Similarly, from eq.[\(32\)](#page-12-0) we find

$$
\delta_{\Lambda}^{*} D_{ij} = \begin{cases} 2\sqrt{2}C_{11}^{\mu\nu}\partial_{\mu} \chi \partial_{\nu} \bar{\phi}, & (i,j) = (1,1) \\ 0, & (i,j) \neq (1,1). \end{cases}
$$
(41)

This comes from the fact that only the first term in the r.h.s. does not contain  $u^+$  and it contributes to  $\delta_{\Lambda}^* D_{11}$  because it is proportional to  $(u^{-1})^2$ .  $\lambda^{(2,2)}$  is then determined as

$$
\lambda^{(2,2)} = -i \sum_{n=2}^{\infty} \partial^{\mu} \lambda_{\mu}^{(n)} (u^{+1})^{n-2} (u^{-1})^{n+2}.
$$
 (42)

<span id="page-13-0"></span>Eq.[\(33\)](#page-12-0) is solved by

$$
\delta_{\Lambda}^{*} \psi_{\alpha}^{i} = \delta_{2}^{i} \frac{2}{3} (\varepsilon C_{11} \sigma^{\mu} \bar{\psi}^{1})_{\alpha} \partial_{\mu} \chi, \tag{43}
$$
\n
$$
\lambda_{\alpha}^{(1,2)} = \frac{4}{3} (\sigma^{\mu} \bar{\psi}_{i})_{\alpha} C_{11}^{\gamma \delta} (\sigma_{\mu} \bar{\sigma}^{\nu} \varepsilon)_{\gamma \delta} \partial_{\nu} \chi u^{-(i} u^{-1} u^{-1}) + \sum_{n=2}^{\infty} i (\sigma^{\mu} \bar{\psi}_{i})_{\alpha} C_{11}^{\gamma \delta} (\sigma_{\mu} \bar{\sigma}^{\nu} \varepsilon)_{\gamma \delta} \lambda_{\nu}^{(n-1)} + \times \left[ a_{n} u^{+(1} \dots u^{+1} u^{-i} \dots u^{-1}) + b_{n} \epsilon^{i} (u^{+1})^{n-2} (u^{-1})^{n+1} \right], \tag{44}
$$

with  $a_n$  and  $b_n$  being appropriate constants. Indeed, substituting [\(35\)](#page-12-1) and [\(44\)](#page-13-0) in ap-pendix [A](#page-19-0) into eq.[\(33\)](#page-12-0) and using the formulas  $(57)$  and  $(58)$  we can see that only the first term in the r.h.s. of [\(33\)](#page-12-0) gives a contribution to  $\delta_{\Lambda}^* \psi^2$  that cannot be gauged away; the other terms are absorbed into  $\lambda_{\alpha}^{(1,2)}$  (with the appropriate choice of  $a_n$  and  $b_n$ ).

As described above, the exact deformed gauge transformation is determined as

<span id="page-13-1"></span>
$$
\delta_{\Lambda}^{*} A_{\mu} = -\partial_{\mu} \chi,
$$
  
\n
$$
\delta_{\Lambda}^{*} \psi_{\alpha}^{2} = -\frac{2}{3} (\varepsilon C_{11} \sigma^{\mu} \bar{\psi}_{2})_{\alpha} \partial_{\mu} \chi,
$$
  
\n
$$
\delta_{\Lambda}^{*} D_{11} = -2\sqrt{2} C_{11}^{\mu \nu} \partial_{\mu} (\partial_{\nu} \chi \bar{\phi}),
$$
  
\n
$$
\delta_{\Lambda}^{*} (\text{others}) = 0.
$$
\n(45)

Note that this result agrees with the deformed gauge transformation for the generic  $C_{ij}$ in [\[14\]](#page-30-3), after setting  $C_{ij}^{\alpha\beta} = C_{11}^{\alpha\beta} \delta_i^1 \delta_j^1$ . For example,  $\delta_{\Lambda}^* A_{\mu}$  was determined as  $\delta_{\Lambda}^* A_{\mu} =$  $-\{1+f(\bar{\phi})\bar{\phi}\}\partial_\mu\chi$  with  $f(\bar{\phi})$  being a function proportional to  $C_{12}$  or  $C_{21}$  (see [\[14\]](#page-30-3) for the definition of f, so that  $f(\bar{\phi}) = 0$  when  $C_{ij} = C_{11} \delta_i^1 \delta_j^1$ , which leads to eq.[\(34\)](#page-12-1).

The determination of the exact  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation is accomplished in a similar way as the one described above, since the deformation of the transformation laws comes from the associated gauge transformation to preserve the WZ gauge. We will give the result in the next subsection. The details of the derivation are found in appendix [B.](#page-20-0)

### <span id="page-14-0"></span>3.2 Exact  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation

Since the theory is expected to have  $\mathcal{N} = (1, 1/2)$  supersymmetry, we will concentrate on this symmetry in the following. For later convenience, we will split the  $\mathcal{N} = (1, 1/2)$ supersymmetry transformation generated by  $Q^1$ ,  $Q^2$  and  $\bar{Q}_2$  into the  $\mathcal{N} = (1,0)$  transformation generated by  $Q^1$ ,  $Q^2$  and  $\mathcal{N} = (0, 1/2)$  by  $\overline{Q}_2$ . The  $\mathcal{N} = (1, 0)$  transformation is defined by

<span id="page-14-2"></span>
$$
\delta_{\xi}^* V_{WZ}^{++} = \tilde{\delta}_{\xi} V_{WZ}^{++} + \delta_{\Lambda}^* V_{WZ}^{++},\tag{46}
$$

and the  $\mathcal{N} = (0, 1/2)$  transformation is

<span id="page-14-3"></span>
$$
\delta_{\xi^2}^* V_{WZ}^{++} = \tilde{\delta}_{\xi^2} V_{WZ}^{++} + \delta_{\Lambda'}^* V_{WZ}^{++}.
$$
\n(47)

<span id="page-14-1"></span>Here

$$
\tilde{\delta}_{\xi} V_{WZ}^{++} = \xi_i Q^i V_{WZ}^{++} = \left( -\xi^{+\alpha} Q_{\alpha}^- + \xi^{-\alpha} Q_{\alpha}^+ \right) V_{WZ}^{++},\tag{48}
$$

$$
\tilde{\delta}_{\bar{\xi}^2} V_{WZ}^{++} = \bar{\xi}^i \bar{Q}_i V_{WZ}^{++} = \left( \bar{\xi}^+_{\dot{\alpha}} \bar{Q}^{-\dot{\alpha}} - \bar{\xi}^-_{\dot{\alpha}} \bar{Q}^{+\dot{\alpha}} \right) V_{WZ}^{++} \tag{49}
$$

and  $Q_{\alpha}^{+} = \frac{\partial}{\partial \theta^{-\alpha}} - 2i\sigma_{\alpha\dot{\alpha}}^{\mu}\bar{\theta}^{+\dot{\alpha}}\frac{\partial}{\partial x_{A}^{\mu}}, \ Q_{\alpha}^{-} = -\frac{\partial}{\partial \theta^{+\alpha}}, \ \bar{Q}_{\dot{\alpha}}^{+} = \frac{\partial}{\partial \bar{\theta}^{-\dot{\alpha}}} + 2i\theta^{+\alpha}\sigma_{\alpha}^{\mu}$  $\frac{\mu}{\alpha\dot{\alpha}}\frac{\partial}{\partial x^{\mu}_{A}},\ \bar{Q}^{-}_{\dot{\alpha}}\,=\,-\frac{\partial}{\partial\bar{\theta}^{+\dot{\alpha}}}.$ Note that  $\bar{\xi}^1$  is implicitly set to be zero in eq.[\(49\)](#page-14-1), because we are now considering the  $\mathcal{N} = (1, 1/2)$  supersymmetry.  $\delta_{\Lambda}^* V_{WZ}^{++}$  ( $\delta_{\Lambda'}^* V_{WZ}^{++}$ ) is a deformed gauge transformation of  $V_{WZ}^{++}$  with an appropriate analytic gauge parameter  $\Lambda(\zeta, u)$   $(\Lambda'(\zeta, u))$  to retain the WZ gauge.

The exact  $\mathcal{N} = (1,0)$  transformation generated by  $Q^1$  and  $Q^2$  is given below (for the derivation, see appendix [B.1\)](#page-20-1):

$$
\delta_{\xi}^{*}\phi = -\sqrt{2}i\xi^{i}\psi_{i} + \frac{8}{3}i(\xi_{2}\varepsilon C_{11}\psi^{1})\bar{\phi} - \frac{2\sqrt{2}}{3}i(\xi_{2}\varepsilon C_{11}\sigma^{\nu}\bar{\psi}_{2})A_{\nu},
$$
\n
$$
\delta_{\xi}^{*}\bar{\phi} = 0,
$$
\n
$$
\delta_{\xi}^{*}A_{\mu} = i\xi^{i}\sigma_{\mu}\bar{\psi}_{i} + 2\sqrt{2}i(\xi^{1}\varepsilon C_{11}\sigma_{\mu}\bar{\psi}^{1})\bar{\phi},
$$
\n
$$
\delta_{\xi}^{*}\psi^{\alpha i} = -(\xi^{i}\sigma^{\mu\nu})^{\alpha}F_{\mu\nu} - D^{ij}\xi_{j}^{\alpha} + \sqrt{2}C_{11}^{\mu\nu}(\xi^{(i}\sigma_{\mu\nu})^{\alpha}D^{11})\bar{\phi} - iC_{11}^{\mu\nu}(\xi^{(i}\sigma_{\mu\nu})^{\alpha}(\bar{\psi}^{1}\bar{\psi}^{1}))
$$
\n
$$
+ \delta_{2}^{i}\left\{2\sqrt{2}(\xi^{1}\varepsilon C_{11}\sigma^{\mu\nu})^{\alpha} + \frac{2\sqrt{2}}{3}(\xi^{1}\sigma^{\mu\nu}\varepsilon C_{11})^{\alpha} + \sqrt{2}C_{11}^{\mu\nu}\xi^{\alpha 1}\right\}\bar{\phi}F_{\mu\nu}
$$
\n
$$
- \delta_{2}^{i}\left\{\frac{4\sqrt{2}}{3}(\xi^{1}\sigma^{\mu\nu}\varepsilon C_{11})^{\alpha} + 2\sqrt{2}C_{11}^{\mu\nu}\xi^{\alpha 1}\right\}\partial_{\mu}\bar{\phi}A_{\nu} + \delta_{2}^{i}\frac{2\sqrt{2}}{3}(\xi^{1}\varepsilon C_{11})^{\alpha}\partial^{\mu}\bar{\phi}A_{\mu}
$$
\n
$$
+ \delta_{2}^{i}\det C_{11}\left\{\frac{8}{3}\xi^{\alpha 1}D^{11}\bar{\phi}^{2} - \frac{8\sqrt{2}}{3}i\xi^{\alpha 1}(\bar{\psi}^{1}\bar{\psi}^{1})\bar{\phi}\right\},
$$
\n
$$
\delta_{\xi}^{*}\bar{\psi}_{\alpha i} = \sqrt
$$

Note that we have used  $(\varepsilon C_{11} \varepsilon C_{11})_{\alpha}{}^{\beta} = -\delta_{\alpha}^{\beta} \det C_{11}.$ 

The  $\mathcal{N} = (0, 1/2)$  supersymmetry transformation generated by  $\overline{Q}_2$  is as follows (these are read from eqs.[\(101\)](#page-27-0)–[\(106\)](#page-27-0) in appendix [B.2](#page-24-0) by setting  $\bar{\xi}^1 = 0$ ):

$$
\delta_{\xi^2}^* \phi = 0,
$$
\n
$$
\delta_{\xi^2}^* \bar{\phi} = \sqrt{2} i \bar{\xi}^2 \bar{\psi}^1,
$$
\n
$$
\delta_{\xi^2}^* \bar{\phi} = \sqrt{2} i \bar{\xi}^2 \bar{\psi}^1,
$$
\n
$$
\delta_{\xi^2}^* \psi^{\alpha i} = \delta_{2}^i \left\{ -\sqrt{2} (\bar{\xi}^2 \bar{\sigma}^{\mu})^{\alpha} \partial_{\mu} \phi - \frac{2}{3} D^{11} (\bar{\xi}^2 \bar{\sigma}^{\mu} \varepsilon C_{11})^{\alpha} A_{\mu} + \frac{4}{3} i \bar{\xi}^2 \bar{\psi}^1 (\psi^1 \varepsilon C_{11})^{\alpha} \right\},
$$
\n
$$
\delta_{\xi^2}^* \bar{\psi}_{\dot{\alpha}i} = \delta_{i}^1 (\bar{\xi}^2 \bar{\sigma}^{\mu \nu})_{\dot{\alpha}} F_{\mu \nu} - \bar{\xi}_{\dot{\alpha}}^2 D_{i2},
$$
\n
$$
\delta_{\xi^2}^* D^{11} = 0,
$$
\n
$$
\delta_{\xi^2}^* D^{12} = \delta_{\xi^2}^* D^{21} = -i \partial_{\mu} (\bar{\xi}^2 \bar{\sigma}^{\mu} \psi^1),
$$
\n
$$
\delta_{\xi^2}^* D^{22} = -2i \partial_{\mu} (\bar{\xi}^2 \bar{\sigma}^{\mu} \psi^2)
$$
\n
$$
- \frac{2}{3} i \partial_{\mu} \left[ 2 (\bar{\xi}^2 \bar{\sigma}^{\nu} \varepsilon C_{11} \sigma^{\mu} \bar{\psi}^1) A_{\nu} - C_{11}^{\alpha \beta} (\sigma^{\nu} \bar{\sigma}^{\mu} \varepsilon)_{\alpha \beta} A_{\nu} \bar{\xi}^2 \bar{\psi}^1 + \sqrt{2} (\bar{\xi}^2 \bar{\sigma}^{\mu} \varepsilon C_{11} \psi^1) \bar{\phi} \right].
$$
\n(51)

From the exact gauge transformation [\(45\)](#page-13-1), we easily find a field redefinition that leads to the canonical gauge transformation:

$$
\hat{\psi}_{\alpha}^2 \equiv \psi_{\alpha}^2 - \frac{2}{3} (\varepsilon C_{11} \sigma^{\mu} \bar{\psi}_2)_{\alpha} A_{\mu}, \tag{52}
$$

$$
\hat{D}_{11} \equiv D_{11} - 2\sqrt{2}C_{11}^{\mu\nu}A_{\nu}\partial_{\mu}\bar{\phi}, \qquad (53)
$$

and the other newly defined fields with hat are the same as the original fields.

Using this field redefinition, the exact  $\mathcal{N} = (1, 1/2)$  transformation becomes as follows. The  $\mathcal{N}=(1,0)$  supersymmetry transformation is

$$
\delta_{\xi}^{*}\hat{\phi} = -\sqrt{2}i\xi^{i}\hat{\psi}_{i} + \frac{8}{3}i(\xi_{2}\varepsilon C_{11}\hat{\psi}^{1})\hat{\phi},
$$
\n
$$
\delta_{\xi}^{*}\hat{\phi} = 0,
$$
\n
$$
\delta_{\xi}^{*}\hat{A}_{\mu} = i\xi^{i}\sigma_{\mu}\hat{\psi}_{i} + 2\sqrt{2}i(\xi^{1}\varepsilon C_{11}\sigma_{\mu}\hat{\psi}^{1})\hat{\phi},
$$
\n
$$
\delta_{\xi}^{*}\hat{\psi}^{\alpha i} = -(\xi^{i}\sigma^{\mu\nu})^{\alpha}\hat{F}_{\mu\nu} - \hat{D}^{ij}\xi_{j}^{\alpha} + \sqrt{2}C_{11}^{\mu\nu}(\xi^{(i}\sigma_{\mu\nu})^{\alpha}\hat{D}^{11})\hat{\phi} - iC_{11}^{\mu\nu}(\xi^{i}\sigma_{\mu\nu})^{\alpha}(\hat{\psi}^{1}\hat{\psi}^{1})
$$
\n
$$
+ \delta_{2}^{i}\left\{2\sqrt{2}(\xi^{1}\varepsilon C_{11}\sigma^{\mu\nu})^{\alpha} + \frac{2\sqrt{2}}{3}(\xi^{1}\sigma^{\mu\nu}\varepsilon C_{11})^{\alpha} + \sqrt{2}C_{11}^{\mu\nu}\xi^{\alpha 1}\right\}\hat{\phi}\hat{F}_{\mu\nu}
$$
\n
$$
+ \frac{8}{3}\delta_{2}^{i}\det C_{11}\xi^{\alpha 1}\left\{\hat{D}^{11}\hat{\phi}^{2} - 2\sqrt{2}i(\hat{\psi}^{1}\hat{\psi}^{1})\hat{\phi}\right\},
$$
\n
$$
\delta_{\xi}^{*}\hat{\psi}_{\alpha i} = \sqrt{2}(\xi_{i}\sigma^{\nu})_{\alpha}\partial_{\nu}\hat{\phi} + 2\delta_{i}^{1}(\xi_{2}\varepsilon C_{11}\sigma^{\nu})_{\alpha}\partial_{\nu}(\hat{\phi}^{2}),
$$
\n
$$
\delta_{\xi}^{*}\hat{\psi}^{11} = -2i\xi^{1}\sigma^{\nu}\partial_{\nu}\hat{\psi}^{1},
$$
\n
$$
\delta_{\xi}^{*}\hat{D}^{12} = \delta_{\xi}^{*}\hat{D}^{21} = -2i\xi^{(1}\sigma^
$$

<span id="page-16-0"></span>where  $\hat{F}_{\mu\nu} \equiv \partial_{\mu}\hat{A}_{\nu} - \partial_{\nu}\hat{A}_{\mu}$ . The  $\mathcal{N} = (0, 1/2)$  supersymmetry transformation becomes

$$
\delta_{\xi^2}^* \hat{\phi} = 0,
$$
\n
$$
\delta_{\xi^2}^* \hat{\phi} = \sqrt{2} i \bar{\xi}^2 \hat{\psi}^1,
$$
\n
$$
\delta_{\xi^2}^* \hat{A}_{\mu} = -i \bar{\xi}^2 \bar{\sigma}_{\mu} \hat{\psi}^1,
$$
\n
$$
\delta_{\xi^2}^* \hat{\psi}^{\alpha i} = \delta_2^i \left\{ -\sqrt{2} (\bar{\xi}^2 \bar{\sigma}^{\mu})^{\alpha} \partial_{\mu} \hat{\phi} + \frac{8}{3} i \bar{\xi}^2 \hat{\psi}^1 (\hat{\psi}^1 \varepsilon C_{11})^{\alpha} \right\},
$$
\n
$$
\delta_{\xi^2}^* \hat{\psi}_{\dot{\alpha}i} = \delta_i^1 (\bar{\xi}^2 \bar{\sigma}^{\mu \nu})_{\dot{\alpha}} \hat{F}_{\mu \nu} - \bar{\xi}_{\dot{\alpha}}^2 \hat{D}_{i2},
$$
\n
$$
\delta_{\xi^2}^* \hat{D}^{11} = 0,
$$

$$
\delta_{\xi^2}^* \hat{D}^{12} = \delta_{\xi^2}^* \hat{D}^{21} = -i \partial_\mu (\bar{\xi}^2 \bar{\sigma}^\mu \hat{\psi}^1),
$$
\n
$$
\delta_{\xi^2}^* \hat{D}^{22} = -2i \partial_\mu (\bar{\xi}^2 \bar{\sigma}^\mu \hat{\psi}^2) + 2i C_{11}^{\mu\nu} \bar{\xi}^2 \hat{\bar{\psi}}^1 \hat{F}_{\mu\nu}
$$
\n
$$
-2\sqrt{2}i(\bar{\xi}^2 \bar{\sigma}^\mu \varepsilon C_{11} \hat{\psi}^1) \partial_\mu \hat{\bar{\phi}} - \frac{2\sqrt{2}}{3}i \partial_\mu \left[ (\bar{\xi}^2 \bar{\sigma}^\mu \varepsilon C_{11} \hat{\psi}^1) \hat{\bar{\phi}} \right].
$$
\n(55)

In terms of the newly defined fields, the action in the WZ gauge should now be invariant under the ordinary gauge transformation and the above  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation.

Note that the  $O(C)$  action is exactly gauge and  $\mathcal{N} = (1, 1/2)$  supersymmetry invariant by itself (in order to show this we have used  $(\bar{\psi}^1)^3 = 0$  as a result of the  $U(1)$  gauge group). An immediate consequence of this fact is that the terms arising from  $(V^{++})^n$   $(n \ge 4)$  in the action, if there exist, have to be also gauge and  $\mathcal{N} = (1, 1/2)$  supersymmetry invariant by themselves.

<span id="page-17-0"></span>Using the field redefinitions [\(25\)](#page-9-1) and [\(26\)](#page-9-3) and transformation [\(55\)](#page-16-0), we find that the action [\(5\)](#page-4-0) is invariant under the  $\mathcal{N} = (0, 1/2)$  transformation:

$$
\delta_{\bar{\eta}}^* A = 0, \quad \delta_{\bar{\eta}}^* \bar{A} = \sqrt{2} \bar{\eta} \bar{\lambda},
$$
  
\n
$$
\delta_{\bar{\eta}}^* \psi^{\alpha} = 0, \quad \delta_{\bar{\eta}}^* \bar{\psi}_{\dot{\alpha}} = (\bar{\eta} \bar{\sigma}^{\mu \nu})_{\dot{\alpha}} v_{\mu \nu} - i \bar{\eta}_{\dot{\alpha}} D,
$$
  
\n
$$
\delta_{\bar{\eta}}^* F = 0, \quad \delta_{\bar{\eta}}^* \bar{F} = \sqrt{2} i \bar{\eta} \bar{\sigma}^{\mu} \partial_{\mu} \lambda + 2 \bar{\eta} \bar{\sigma}^{\mu} \varepsilon C \partial_{\mu} (\psi \bar{A}),
$$
  
\n
$$
\delta_{\bar{\eta}}^* v_{\mu} = -i \bar{\eta} \bar{\sigma}_{\mu} \psi,
$$
  
\n
$$
\delta_{\bar{\eta}}^* \lambda^{\alpha} = -i \sqrt{2} (\bar{\eta} \bar{\sigma}^{\mu})^{\alpha} \partial_{\mu} A - 2 i \bar{\eta} \bar{\lambda} (\psi \varepsilon C)^{\alpha}, \quad \delta_{\bar{\eta}}^* \bar{\lambda}_{\dot{\alpha}} = \sqrt{2} \bar{\eta}_{\dot{\alpha}} F,
$$
  
\n
$$
\delta_{\bar{\eta}}^* D = \bar{\eta} \bar{\sigma}^{\mu} \partial_{\mu} \psi.
$$
\n(56)

Thus we have obtained deformed  $\mathcal{N} = (1, 1/2)$  supersymmetry transformations [\(8\)](#page-4-1), [\(27\)](#page-10-0) and [\(56\)](#page-17-0) in  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in the deformed  $\mathcal{N} = 1$  superspace.

### 4 Conclusions and Discussion

In this paper, we studied  $\mathcal{N} = 2$  supersymmetric  $U(1)$  gauge theory in non(anti)commutative  $\mathcal{N}=1$  and  $\mathcal{N}=2$  harmonic superspaces. We considered the reduction of the deformation parameters in non(anti)commutative harmonic superspace to the deformed  $\mathcal{N} = 1$ superspace. We found that the  $O(C)$  action in harmonic superspace reduces to the one

in the deformed  $\mathcal{N} = 1$  superspace by the field redefinition. We calculated the gauge and  $\mathcal{N} = (1, 1/2)$  supersymmetry transformations exactly. Using the field redefinition, we confirmed  $\mathcal{N} = (1, 1/2)$  supersymmetry of the action in the deformed  $\mathcal{N} = 1$  superspace.

It is known that even in the  $U(1)$  gauge group there exists higher order C-corrections to the action in the deformed harmonic superspace[\[14,](#page-30-3) [15\]](#page-30-6). It is not clear whether these higher order contributions in the action disappear after the reduction of the deformation parameters[\(23\)](#page-9-2). Even if these terms exist after the reduction, these must be gauge- and  $\mathcal{N} = (1, 1/2)$  supersymmetry invariant by themselves and reduce to the deformed action in the  $\mathcal{N} = 1$  superspace formalism by appropriate field redefinition. Some detailed analysis will be studied elsewhere.

In this paper, the component  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation for the action in the deformed  $\mathcal{N} = 1$  superspace has not fixed completely, because it is unclear whether the field identification [\(25\)](#page-9-1) is exact or not. To fix the component transformation, it is useful to work with the  $\mathcal{N} = 2$  rigid superspace formalism. The deformed action could be written in terms of the  $\mathcal{N} = 2$  chiral superfield that describes the  $\mathcal{N} = 2$ vector multiplet and give directly the action in the deformed  $\mathcal{N} = 1$  superspace. If we can construct the appropriately deformed  $\mathcal{N} = 2$  chiral superfield, the component transformation that is not fixed in this paper will be determined. This formalism would be also useful to study a generalization to non-abelian gauge group.

Another interesting issue would be a central extension of  $\mathcal{N} = (1, 1/2)$  supersymmetry algebra. It is also interesting to study properties of deformed properties of solitons in such as monopoles and instantons in such theories [\[9\]](#page-29-4).

Acknowledgments: T. A. is supported by the Grant-in-Aid for Scientific Research in Priority Areas (No.14046201) from the Ministry of Education, Culture, Sports, Science and Technology. A. O. is supported by a 21st Century COE Program at Tokyo Tech "Nanometer-Scale Quantum Physics" by the Ministry of Education, Culture, Sports, Science and Technology.

## <span id="page-19-0"></span>A Useful formulas

<span id="page-19-1"></span>In this appendix, we describe some useful reduction formulas of harmonic variables, which we have used in sect. [3](#page-11-1) to determine the exact transformation laws in the harmonic superspace formalism. They are

u +(1u +1) n z }| { u +(1 · · · u +1 m z }| { u −i u −j u −k u −1 · · · u −1) = n+2 z }| { u +(1 · · · u +1 m z }| { u <sup>−</sup><sup>i</sup>u <sup>−</sup><sup>j</sup>u <sup>−</sup><sup>k</sup>u −1 · · · u −1) + 2m (n + m + 2)(n + m) × ǫ i1 n+1 z }| { u +(1 · · · u +1 m−1 z }| { u <sup>−</sup><sup>j</sup>u <sup>−</sup><sup>k</sup>u −1 · · · u <sup>−</sup>1) +(cyclic permut. of (i, j, k)) + 2m(m − 1) (n + m + 1)(n + m) 2 (n + m − 1) × ǫ i1 ǫ j1 n z }| { u +(1 · · · u +1 m−2 z }| { u <sup>−</sup><sup>k</sup>u −1 · · · u <sup>−</sup>1) +(cyclic permut. of (i, j, k)) , (57) u −k n z }| { u +(1 · · · u +1 m z }| { u −i u −j u −1 · · · u −1) = n z }| { u +(1 · · · u +1 m+1 z }| { u <sup>−</sup><sup>i</sup>u <sup>−</sup><sup>j</sup>u <sup>−</sup><sup>k</sup>u −1 · · · u −1) + 2n (n + m + 1)(n + m) 1 2 n ǫ ki n−1 z }| { u +(1 · · · u +1 m z }| { u <sup>−</sup><sup>j</sup>u −1 · · · u <sup>−</sup>1) +(<sup>i</sup> <sup>↔</sup> <sup>j</sup>) o + n(n + m − 2) (n + m + 1)(n + m) ǫ k1 n−1 z }| { u +(1 · · · u +1 m z }| { u <sup>−</sup><sup>i</sup>u <sup>−</sup><sup>j</sup>u −1 · · · u −1) , (58) u <sup>−</sup>(<sup>k</sup>u −l) n z }| { u +(<sup>i</sup>u +1 · · · u +1 m z }| { u −1 · · · u −1) = n z }| { u +(1 · · · u +1 m+2 z }| { u <sup>−</sup><sup>i</sup>u <sup>−</sup><sup>k</sup>u <sup>−</sup><sup>l</sup>u −1 · · · u −1) + 2n(n + m − 1) (n + m + 2)(n + m) 1 2 n ǫ k1 n−1 z }| { u +(1 · · · u +1 m+1 z }| { u <sup>−</sup><sup>l</sup>u <sup>−</sup><sup>i</sup>u −1 · · · u <sup>−</sup>1) +(<sup>k</sup> <sup>↔</sup> <sup>l</sup>) o + 2n (n + m + 2)(n + m) 1 2 n ǫ ki n−1 z }| { u +(1 · · · u +1 m+1 z }| { u −l u −1 · · · u <sup>−</sup>1) +(<sup>k</sup> <sup>↔</sup> <sup>l</sup>) o + n(n − 1)(n + m − 2) (n + m + 1)(n + m) 2 ǫ l1 ǫ k1 n−2 z }| { u +(1 · · · u +1 m z }| { u <sup>−</sup><sup>i</sup>u −1 · · · u −1)

$$
+\frac{2n(n-1)}{(n+m+1)(n+m)^2} \frac{1}{2} \Big\{ \epsilon^{l_1} \epsilon^{k_i} u^{+(1} \cdots u^{+1} u^{-1} \cdots u^{-1}) + (k \leftrightarrow l) \Big\}, \qquad (59)
$$
  

$$
u^{+(1}u^{-1}) u^{+(i}u^{+1} \cdots u^{+1} u^{-k} u^{-1} \cdots u^{-1})
$$
  

$$
= u^{+(i}u^{+1} \cdots u^{+1} u^{-k} u^{-1} \cdots u^{-1})
$$
  

$$
-\frac{2(n-m)}{(n+m+2)(n+m)} \frac{1}{2} \Big( \epsilon^{i_1} u^{+(1} \cdots u^{+1} u^{-k} u^{-1} \cdots u^{-1}) + (i \leftrightarrow k) \Big)
$$
  

$$
-\frac{2nm}{(n+m+1)(n+m)^2(n+m-1)} \epsilon^{i_1} \epsilon^{k_1} u^{+(1} \cdots u^{+1} u^{-1} \cdots u^{-1}). \qquad (60)
$$

# <span id="page-20-0"></span>**B**  $\mathcal{N} = (1, 1/2)$  **Supersymmetry**

Here we will present the details of calculation of the exact  $\mathcal{N} = (1, 1/2)$  supersymmetry transformation of the component fields when the deformation parameters are restricted as [\(23\)](#page-9-2). The result has been summarized in sect. [3.2.](#page-14-0)

## <span id="page-20-1"></span>**B.1**  $\mathcal{N} = (1, 0)$  supersymmetry

<span id="page-20-2"></span>The equations to determine the deformed  $\mathcal{N}=(1,0)$  supersymmetry transformation are eqs. $(61)$ – $(68)$  which are obtained from eq. $(46)$  and  $(29)$ :

$$
0 = 2i(\xi^+\sigma^\mu)_{\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}}A_\mu - \partial^{++}\lambda^{(0,1)\dot{\alpha}} - 2\lambda^{(1,0)\alpha}(\varepsilon C^{++}\sigma^\mu)_{\alpha\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}}A_\mu,\tag{61}
$$

$$
0 = -2\sqrt{2}i\xi_{\alpha}^{+}\bar{\phi} - \partial^{++}\lambda_{\alpha}^{(1,0)} - 2\sqrt{2}(\varepsilon C^{++}\lambda^{(1,0)})_{\alpha}\bar{\phi}, \qquad (62)
$$

$$
\sqrt{2}i\delta_{\xi}^{*}\phi = 4\xi^{+}\psi^{i}u_{i}^{-} + 4i\lambda^{(1,0)\alpha}(\varepsilon C^{++}\psi^{i})_{\alpha}u_{i}^{-}
$$

$$
-\partial^{++}\lambda^{(0,2)} - C^{++\alpha\beta}(\sigma^{\nu}\bar{\sigma}^{\mu}\varepsilon)_{\alpha\beta}\lambda_{\mu}^{(1,1)}A_{\nu},
$$
(63)

$$
-\sqrt{2}i\delta_{\xi}^*\bar{\phi} = 0, \qquad (64)
$$

$$
-2i\delta_{\xi}^{*}A_{\mu} = 4\xi^{+}\sigma_{\mu}\bar{\psi}^{i}u_{i}^{-} + 4i\lambda^{(1,0)\alpha}(\varepsilon C^{++}\sigma_{\mu}\bar{\psi}^{i})_{\alpha}u_{i}^{-}
$$

$$
- \partial^{++}\lambda_{\mu}^{(1,1)} - \sqrt{2}C^{++\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\lambda_{\nu}^{(1,1)}\bar{\phi}, \qquad (65)
$$

$$
4\delta_{\xi}^{*}\psi_{\alpha}^{i}u_{i}^{-} = -2(\sigma^{\mu}\bar{\sigma}^{\nu}\xi^{-})_{\alpha}\partial_{\nu}A_{\mu} + 6\xi_{\alpha}^{+}D^{ij}u_{i}^{-}u_{j}^{-}
$$
  

$$
-i(\sigma^{\nu}\partial_{\nu}\lambda^{(0,1)})_{\alpha} - 2\sqrt{2}i(\varepsilon C^{+-}\sigma^{\nu}\partial_{\nu}\lambda^{(0,1)})_{\alpha}\bar{\phi}
$$
  

$$
-6i(\varepsilon C^{++}\lambda^{(1,0)})_{\alpha}D^{ij}u_{i}^{-}u_{j}^{-} + 2i\lambda^{(1,0)\beta}(\varepsilon C^{+-}\sigma^{\nu}\bar{\sigma}^{\mu}\varepsilon)_{\beta\alpha}\partial_{\nu}A_{\mu}
$$

+ 
$$
2i\partial_{\nu}\lambda_{\alpha}^{(1,0)}C^{+-\beta\gamma}(\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\beta\gamma}A_{\mu} - 2i(\sigma^{\mu}\bar{\psi}^{i})_{\alpha}u_{i}^{-}C^{++\gamma\delta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\gamma\delta}\lambda_{\nu}^{(1,1)}
$$
  
\n $2i+1$  (1,2)  $2i\sqrt{2}(\epsilon C^{++}+1)$  (1,2))  $\bar{\phi} = 2(\epsilon C^{++}+\epsilon^{\mu}1)$  (2,1))  $\bar{\phi} = 2\epsilon C^{+}(\epsilon C^{++}+\epsilon^{\mu}1)$ 

$$
-\partial^{++}\lambda_{\alpha}^{(1,2)} - 2\sqrt{2}(\varepsilon C^{++}\lambda^{(1,2)})_{\alpha}\bar{\phi} - 2(\varepsilon C^{++}\sigma^{\mu}\lambda^{(2,1)})_{\alpha}A_{\mu},\tag{66}
$$

$$
-4\delta_{\xi}^{*}\bar{\psi}^{\dot{\alpha}i}u_{i}^{-} = 2\sqrt{2}(\xi^{-}\sigma^{\mu})_{\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}}\partial_{\mu}\bar{\phi} + i\partial_{\mu}\lambda^{(1,0)\alpha}\sigma^{\mu}_{\alpha\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}} + 2\sqrt{2}i\partial_{\nu}\left\{\lambda^{(1,0)\alpha}(\varepsilon C^{+-}\sigma^{\nu})_{\alpha\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}}\bar{\phi}\right\} - \partial^{++}\lambda^{(2,1)\dot{\alpha}},
$$
\n(67)

$$
3\delta_{\xi}^{*}D^{ij}u_{i}^{-}u_{j}^{-} = -4i\xi^{-}\sigma^{\mu}\partial_{\mu}\bar{\psi}^{i}u_{i}^{-} + 4\partial_{\nu}\left\{\lambda^{(1,0)\alpha}(\varepsilon C^{+-}\sigma^{\nu}\bar{\psi}^{i})_{\alpha}u_{i}^{-}\right\} - i\partial^{\mu}\lambda_{\mu}^{(1,1)} - \sqrt{2}iC^{+-\alpha\beta}(\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\partial_{\nu}(\lambda_{\mu}^{(1,1)}\bar{\phi}) - \partial^{++}\lambda^{(2,2)}.
$$
(68)

<span id="page-21-2"></span>Note that under the restriction [\(23\)](#page-9-2), we have  $C^{+-\alpha\beta} = C_{11}^{\alpha\beta} u^{+1} u^{-1}$  and so on.

Eq. $(62)$  is solved by

$$
\lambda^{(1,0)\alpha} = 2\sqrt{2}i\xi_i^{\alpha}\bar{\phi} u^{-i} + \sum_{n=1}^{\infty} 2\sqrt{2}i\xi_i^{(n)\alpha}\bar{\phi}
$$
  
 
$$
\times \left[ \alpha_n^{(1,0)} u^{+(1} \cdots u^{+1} u^{-i} u^{-1} \cdots u^{-1}) + \beta_n^{(1,0)} \epsilon^{i1} (u^{+1})^{n-1} (u^{-1})^n \right], \quad (69)
$$

where

$$
\xi_i^{(n)} \alpha \equiv (\xi_i \overbrace{\varepsilon C_{11} \cdots \varepsilon C_{11}}^n)^{\alpha} (2\sqrt{2}\overline{\phi})^n
$$
\n(70)

<span id="page-21-0"></span>and

$$
(\alpha_1^{(1,0)}, \beta_1^{(1,0)}) = \left(\frac{1}{2}, \frac{2}{3}\right),
$$
  
\n
$$
\alpha_n^{(1,0)} = \frac{1}{n+1} \alpha_{n-1}^{(1,0)}, \quad \beta_n^{(1,0)} = \frac{1}{n} \left( \frac{2n}{(2n+1)(2n-1)} \alpha_{n-1}^{(1,0)} + \beta_{n-1}^{(1,0)} \right).
$$
 (71)

We then obtain  $\alpha_n^{(1,0)} = \frac{1}{(n+1)!}$ ,  $\beta_n^{(1,0)} = \frac{2}{(n-1)!(2n+1)}$ . To check [\(71\)](#page-21-0), we need [\(57\)](#page-19-1), [\(58\)](#page-19-1) and

$$
2\sqrt{2}(\xi_i^{(n)}\varepsilon C_{11})^{\alpha}\overline{\phi} = \xi_i^{(n+1)\alpha}.
$$
\n(72)

From eq.[\(61\)](#page-20-2) we find

<span id="page-21-1"></span>
$$
\lambda_{\alpha}^{(0,1)} = 2i(\xi_i \sigma^{\mu})_{\dot{\alpha}} A_{\mu} u^{-i} + \sum_{n=1}^{\infty} 2i(\xi_i^{(n)} \sigma^{\mu})_{\dot{\alpha}} A_{\mu} \Big[ \alpha_n^{(1,0)} u^{+(1} \cdots u^{+1} u^{-i} u^{-1} \cdots u^{-1}) + \beta_n^{(1,0)} \epsilon^{i1} (u^{+1})^{n-1} (u^{-1})^n \Big].
$$
\n(73)

<span id="page-22-0"></span>In eq.[\(65\)](#page-20-2),  $\lambda_{\mu}^{(1,1)}$  is given by

$$
\lambda_{\mu}^{(1,1)} = 2\xi_i \sigma_{\mu} \bar{\psi}_j u^{-(i} u^{-j)} \n+ \xi_i^{(1)} \sigma_{\mu} \bar{\psi}_j \left[ 2u^{+(1} u^{-i} u^{-j} u^{-1)} + \frac{3}{2} \{ \epsilon^{i1} u^{-(j} u^{-1)} + (i \leftrightarrow j) \} \right] \n+ \sum_{n=2}^{\infty} \xi_i^{(n)} \sigma_{\mu} \bar{\psi}_j \left[ \alpha_n^{(1,1)} u^{+(1} \cdots u^{+1} u^{-i} u^{-j} u^{-1} \cdots u^{-1}) + \beta_n^{(1,1)} \frac{1}{2} \{ \epsilon^{i1} u^{+(1} \cdots u^{+1} u^{-j} u^{-1} \cdots u^{-1}) + (i \leftrightarrow j) \} \right] \n+ \gamma_n^{(1,1)} \epsilon^{i1} \epsilon^{j1} (u^{+1})^{n-2} (u^{-1})^n + \delta_n^{(1,1)} \epsilon^{j i} (u^{+1})^{n-1} (u^{-1})^{n+1} \right], \qquad (74)
$$

where  $(\alpha_2^{(1,1)}$  $\beta_2^{(1,1)}, \beta_2^{(1,1)}, \gamma_2^{(1,1)}$  $\delta_2^{(1,1)}, \delta_2^{(1,1)}$ ) =  $(1, \frac{8}{3})$  $\frac{8}{3}, \frac{8}{5}$  $\frac{8}{5}, -\frac{1}{3}$  $\frac{1}{3}$ ) and

$$
\alpha_n^{(1,1)} = \frac{1}{n+1} \left( 4\alpha_{n-1}^{(1,0)} + \alpha_{n-1}^{(1,1)} \right),
$$
  
\n
$$
\beta_n^{(1,1)} = \frac{1}{n+1} \left( \frac{4n}{2n-1} \alpha_{n-1}^{(1,0)} + 4\beta_{n-1}^{(1,0)} + \frac{1}{n} \alpha_{n-1}^{(1,1)} + \beta_{n-1}^{(1,1)} \right),
$$
  
\n
$$
\gamma_n^{(1,1)} = \frac{1}{n} \left( \frac{4n}{(2n+1)(2n-1)} \alpha_{n-1}^{(1,0)} + 2\beta_{n-1}^{(1,0)} + \frac{n+1}{(2n+1)2n(2n-1)} \alpha_{n-1}^{(1,1)} + \frac{1}{2n-2} \beta_{n-1}^{(1,1)} + \gamma_{n-1}^{(1,1)} \right),
$$
  
\n
$$
\delta_n^{(1,1)} = \frac{1}{n+1} \left( \frac{2(n-1)}{2n-1} \alpha_{n-1}^{(1,0)} - 2\beta_{n-1}^{(1,0)} + \delta_{n-1}^{(1,1)} \right).
$$
\n(75)

Substituting eq.[\(74\)](#page-22-0) to the r.h.s. of eq.[\(65\)](#page-20-2) and using  $(57)$ ,  $(58)$  and

$$
-\sqrt{2}C^{++\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}(\xi_i^{(n)}\sigma_{\nu}\bar{\psi}_j)\bar{\phi}=(\xi_i^{(n+1)}\sigma_{\mu}\bar{\psi}_j)(u^{+1})^2,
$$
\n(76)

we can see that all the  $O(C^2)$  terms are gauged away and the contributions to  $\delta_{\xi}^* A_{\mu}$  come only from the first term and the order  $C$  part of the second term in [\(65\)](#page-20-2).

Eq.[\(63\)](#page-20-2) have the same structure as eq.[\(65\)](#page-20-2) in terms of the harmonic variables. Therefore, it is solved in a way similar to  $(65)$ : The  $O(C<sup>2</sup>)$  terms in the r.h.s. of eq.[\(63\)](#page-20-2) are completely gauged away, so that we need to consider only the first and the second term to determine  $\delta_{\xi}^* \phi$ .

<span id="page-22-1"></span>In eq.[\(67\)](#page-20-2),  $\lambda_{\dot{\alpha}}^{(2,1)}$  $\alpha^{(2,1)}$  is given by

$$
\lambda^{(2,1)}_{\dot{\alpha}} = \sqrt{2} \partial_{\mu} \left\{ (\xi^{(1)}_i \sigma^{\mu})_{\dot{\alpha}} \bar{\phi} \right\} u^{-(i} u^{-1} u^{-1)}
$$

$$
+\sum_{n=2}^{\infty} 2\sqrt{2}\partial_{\mu} \left\{ (\xi_i^{(n)}\sigma^{\mu})_{\dot{\alpha}}\bar{\phi} \right\} \left[ \alpha_n^{(2,1)} \overbrace{u^{+(1}\cdots u^{+1}}^{n-1} \overbrace{u^{-i}u^{-1}\cdots u^{-1}}^{n+2} ) + \beta_n^{(2,1)} \epsilon^{i_1} (u^{+1})^{n-2} (u^{-1})^{n+1} \right], \tag{77}
$$

<span id="page-23-0"></span>where

$$
\alpha_n^{(2,1)} = \frac{1}{n+2} (\alpha_n^{(1,0)} + \alpha_{n-1}^{(1,0)}), \quad \beta_n^{(2,1)} = \frac{1}{n+1} \left( \frac{1}{(2n+1)(2n-1)} \alpha_{n-1}^{(1,0)} + \beta_n^{(1,0)} + \beta_{n-1}^{(1,0)} \right). \tag{78}
$$

We can easily find that in the r.h.s. of [\(67\)](#page-20-2) the terms proportional to  $(C_{11})^n$   $(n \ge 2)$  are completely cancelled by this gauge parameter, so that there are no  $O((C_{11})^2)$  terms in  $\delta_\xi^* \bar{\psi}^i$ .

Substituting  $(73)$ ,  $(69)$ ,  $(74)$  and  $(77)$  to eq. $(66)$ , we can check that the following form of the gauge parameter is sufficient :

<span id="page-23-1"></span>
$$
\lambda_{\alpha}^{(1,2)} = \sum_{n=0}^{\infty} \chi_{\alpha i j k(n)} \overbrace{u^{+(1} \dots u^{+1} u^{-i} u^{-j} u^{-k} u^{-1} \dots u^{-1})}^{n+3} \n+ \sum_{n=1}^{\infty} \chi_{\alpha i j(n)} \overbrace{u^{+(1} \dots u^{+1} u^{-i} u^{-j} u^{-1} \dots u^{-1})}^{n+2} \n+ \sum_{n=1}^{\infty} \chi_{\alpha i(n)} \overbrace{u^{+(1} \dots u^{+1} u^{-i} u^{-1} \dots u^{-1})}^{n+2} + \sum_{n=2}^{\infty} \tilde{\chi}_{\alpha i(n)} \overbrace{u^{+(1} \dots u^{+1} u^{-i} u^{-1} \dots u^{-1})}^{n-2} \n+ \sum_{n=2}^{\infty} \chi_{\alpha(n)} \overbrace{u^{+(1} \dots u^{+1} u^{-1} \dots u^{-1})}^{n+1} + \sum_{n=3}^{\infty} \tilde{\chi}_{\alpha(n)} \overbrace{u^{+(1} \dots u^{+1} u^{-1} \dots u^{-1})}^{n-3}, \quad (79)
$$

where the subscript n denotes the power of  $C_{11}$ -dependence of each quantity. To see this is sufficient, we need  $(57)-(60)$  $(57)-(60)$ . Note that in the r.h.s. of eq.[\(66\)](#page-20-2) the terms proportional to  $(C_{11})^n$   $(n \geq 3)$  are completely gauged away, so that there are no  $O((C_{11})^3)$  terms in  $\delta_{\xi}^* \psi^i$ . In fact, in order to determine  $\delta_{\xi}^* \psi^i$ , only the precise forms of  $\chi_{ijk(0)}^{\alpha}$  and  $\chi_{ij(1)}^{\alpha}$  are needed:

$$
\chi_{ijk(0)}^{\alpha} = 2\xi_{(i}^{\alpha}D_{jk)},
$$
\n
$$
\chi_{\alpha}^{\alpha} = \frac{32}{2} \left[ f_{i(\bar{q}_{i},\bar{q}_{i})} - \sqrt{2} D_{\cdots} \bar{\phi} \right] (\xi, \xi C_{ij})^{\alpha} + \left( \text{cyclic permut of } (ijk) \right) \left[ \xi^{k1} \right] (\hat{\phi}^1)
$$

$$
\chi_{ij}^{\alpha}(1) = \frac{32}{15} \left[ \left\{ i(\bar{\psi}_i \bar{\psi}_j) - \sqrt{2} D_{ij} \bar{\phi} \right\} (\xi_k \varepsilon C_{11})^{\alpha} + \left( \text{cyclic permut. of } (ijk) \right) \right] \epsilon^{k1}.
$$
 (81)

Substituting [\(69\)](#page-21-2) and [\(74\)](#page-22-0) to eq.[\(68\)](#page-20-2) and collecting  $u^+$ -independent terms, we obtain

 $\delta_{\xi}^* D^{ij}$ . We do not need the precise form of  $\lambda^{(2,2)}$ . To determine  $\delta_{\xi}^* D^{ij}$ , we can use [\(60\)](#page-19-1) and [\(58\)](#page-19-1).

## <span id="page-24-0"></span>**B.2**  $\mathcal{N} = (0, 1/2)$  supersymmetry

<span id="page-24-1"></span>The equations to determine the deformed  $\mathcal{N} = (0, 1/2)$  supersymmetry transformation are eqs. $(82)$ – $(89)$  which are obtained from eq. $(47)$  (note that here the expression  $(29)$  is used for the analytic gauge parameter  $\Lambda'$ ):

$$
0 = 2\sqrt{2}i\bar{\xi}^{+\dot{\alpha}}\phi - \partial^{++}\lambda^{(0,1)\dot{\alpha}} - 2\lambda^{(1,0)\alpha}(\varepsilon C^{++}\sigma^{\mu})_{\alpha\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}}A_{\mu}, \tag{82}
$$

$$
0 = -2i(\sigma^{\mu}\bar{\xi}^{+})_{\alpha}A_{\mu} - \partial^{++}\lambda_{\alpha}^{(1,0)} - 2\sqrt{2}(\varepsilon C^{++}\lambda^{(1,0)})_{\alpha}\bar{\phi}, \qquad (83)
$$

<span id="page-24-3"></span>
$$
\sqrt{2}i\delta_{\xi}^{*}\phi = 4i\lambda^{(1,0)\alpha}(\varepsilon C^{++}\psi^{i})_{\alpha}u_{i}^{-} \n+ \frac{\sqrt{2}}{3}\partial_{\mu}\lambda^{(2,0)}\partial_{\nu}\bar{\phi} C_{i_{1}j_{1}}^{\alpha_{1}\beta_{1}}C_{i_{2}j_{2}}^{\alpha_{2}\beta_{2}}C_{i_{3}j_{3}}^{\alpha_{3}\beta_{3}} \n(-u^{-i_{1}}u^{+i_{2}}u^{+i_{3}}\varepsilon_{\alpha_{3}\alpha_{2}}\sigma_{\alpha_{1}\dot{\alpha}}^{\mu} + u^{+i_{1}}u^{-i_{2}}u^{+i_{3}}\varepsilon_{\alpha_{3}\alpha_{1}}\sigma_{\alpha_{2}\dot{\alpha}}^{\mu} - u^{+i_{1}}u^{+i_{2}}u^{-i_{3}}\varepsilon_{\alpha_{1}\alpha_{2}}\sigma_{\alpha_{3}\dot{\alpha}}^{\mu}) \n(-u^{-j_{1}}u^{+j_{2}}u^{+j_{3}}\varepsilon_{\beta_{3}\beta_{2}}\sigma_{\beta_{1}\dot{\beta}}^{\nu} + u^{+j_{1}}u^{-j_{2}}u^{+j_{3}}\varepsilon_{\beta_{3}\beta_{1}}\sigma_{\beta_{2}\dot{\beta}}^{\nu} - u^{+j_{1}}u^{+j_{2}}u^{-j_{3}}\varepsilon_{\beta_{1}\beta_{2}}\sigma_{\beta_{3}\dot{\beta}}^{\nu}) \n- \partial^{++}\lambda^{(0,2)} - C^{++\alpha\beta}(\sigma^{\nu}\bar{\sigma}^{\mu}\varepsilon)_{\alpha\beta}\lambda_{\mu}^{(1,1)}A_{\nu},
$$
\n(84)

<span id="page-24-2"></span>
$$
-\sqrt{2}i\delta_{\xi}^{*}\bar{\phi} = -4\bar{\xi}^{+}\bar{\psi}^{i}u_{i}^{-} - \partial^{++}\lambda^{(2,0)},\tag{85}
$$

$$
-2i\delta_{\xi}^{*}A_{\mu} = -4\psi^{i}\sigma_{\mu}\bar{\xi}^{+}u_{i}^{-} + 4i\lambda^{(1,0)\alpha}(\varepsilon C^{++}\sigma_{\mu}\bar{\psi}^{i})_{\alpha}u_{i}^{-} - 2C^{++\alpha\beta}(\sigma^{\nu}\bar{\sigma}_{\mu}\varepsilon)_{\alpha\beta}A_{\nu}\lambda^{(2,0)} - \partial^{++}\lambda_{\mu}^{(1,1)} - \sqrt{2}C^{++\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\lambda_{\nu}^{(1,1)}\bar{\phi},
$$
\n(86)

$$
4\delta_{\xi}^{*}\psi_{\alpha}^{i}u_{i}^{-} = 2\sqrt{2}(\sigma^{\mu}\bar{\xi}^{-})_{\alpha}\partial_{\mu}\phi
$$
  
\n
$$
-i(\sigma^{\nu}\partial_{\nu}\lambda^{(0,1)})_{\alpha} - 2\sqrt{2}i(\varepsilon C^{+-}\sigma^{\nu}\partial_{\nu}\lambda^{(0,1)})_{\alpha}\bar{\phi}
$$
  
\n
$$
-6i(\varepsilon C^{++}\lambda^{(1,0)})_{\alpha}D^{ij}u_{i}^{-}u_{j}^{-} + 2i\lambda^{(1,0)\beta}(\varepsilon C^{+-}\sigma^{\nu}\bar{\sigma}^{\mu}\varepsilon)_{\beta\alpha}\partial_{\nu}A_{\mu}
$$
  
\n
$$
+ 2i\partial_{\nu}\lambda_{\alpha}^{(1,0)}C^{+-\beta\gamma}(\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\beta\gamma}A_{\mu} + 8i(\varepsilon C^{++}\psi^{-})_{\alpha}\lambda^{(2,0)}
$$
  
\n
$$
- 2i(\sigma^{\mu}\bar{\psi}^{i})_{\alpha}u_{i}^{-}C^{++\gamma\delta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\gamma\delta}\lambda_{\nu}^{(1,1)}
$$
  
\n
$$
- \partial^{++}\lambda_{\alpha}^{(1,2)} - 2\sqrt{2}(\varepsilon C^{++}\lambda^{(1,2)})_{\alpha}\bar{\phi} - 2(\varepsilon C^{++}\sigma^{\mu}\lambda^{(2,1)})_{\alpha}A_{\mu}, \qquad (87)
$$

$$
-4\delta_{\xi}^{*}\bar{\psi}^{\dot{\alpha}i}u_{i}^{-} = 6\bar{\xi}^{+\dot{\alpha}}D^{ij}u_{i}^{-}u_{j}^{-} - 2(\bar{\sigma}^{\nu}\sigma^{\mu}\bar{\xi}^{-})^{\dot{\alpha}}\partial_{\mu}A_{\nu} + i\partial_{\mu}\lambda^{(1,0)\alpha}\sigma_{\alpha\dot{\beta}}^{\mu}\varepsilon^{\dot{\beta}\dot{\alpha}} + 2\sqrt{2}i\partial_{\nu}\left\{\lambda^{(1,0)\alpha}(\varepsilon C^{+-}\sigma^{\nu})_{\alpha\dot{\beta}}\varepsilon^{\dot{\beta}\dot{\alpha}}\bar{\phi}\right\} - \partial^{++}\lambda^{(2,1)\dot{\alpha}},
$$
\n
$$
3\delta_{\xi}^{*}D^{ij}u_{i}^{-}u_{j}^{-} = -4i\bar{\xi}^{-}\bar{\sigma}^{\mu}\partial_{\mu}\psi^{i}u_{i}^{-} + 4\partial_{\nu}\left\{\lambda^{(1,0)\alpha}(\varepsilon C^{+-}\sigma^{\nu}\bar{\psi}^{i})_{\alpha}u_{i}^{-}\right\}
$$
\n
$$
(88)
$$

+ 
$$
2iC^{+-\alpha\beta}(\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\partial_{\nu}(\lambda^{(2,0)}A_{\mu})
$$
  
\n-  $i\partial^{\mu}\lambda^{(1,1)}_{\mu} - \sqrt{2}iC^{+-\alpha\beta}(\sigma^{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\partial_{\nu}(\lambda^{(1,1)}_{\mu}\bar{\phi}) - \partial^{++}\lambda^{(2,2)}.$  (89)

Here we should understand that  $\bar{\xi}^1$  is implicitly set to be zero. Note that under the restriction [\(23\)](#page-9-2), we have  $C^{+-\alpha\beta} = C_{11}^{\alpha\beta} u^{+1} u^{-1}$  and so on.

<span id="page-25-2"></span>Eq.[\(83\)](#page-24-1) is solved by

$$
\lambda^{(1,0)\alpha} = -2i(\bar{\xi}_i \bar{\sigma}^{\mu})^{\alpha} A_{\mu} u^{-i} + \sum_{n=1}^{\infty} (-2i) (\bar{\xi}_i \bar{\sigma}^{\mu} \varepsilon \mathcal{C}^{(n)})^{\alpha} A_{\mu} (2\sqrt{2}\bar{\phi})^n
$$

$$
\times \left[ \alpha_n^{(1,0)} \overbrace{u^{+(1} \cdots u^{+1}}^{n} \overbrace{u^{-i}u^{-1} \cdots u^{-1}}^{n+1} + \beta_n^{(1,0)} \epsilon^{i1} (u^{+1})^{n-1} (u^{-1})^n \right], \quad (90)
$$

where

$$
\mathcal{C}^{(n)\alpha\beta} \equiv \left(\overline{C_{11}\varepsilon C_{11} \cdots \varepsilon C_{11}}\right)^{\alpha\beta} \tag{91}
$$

<span id="page-25-1"></span>and  $\alpha_n^{(1,0)}, \beta_n^{(1,0)}$  are given in [\(71\)](#page-21-0). To check this, we need [\(57\)](#page-19-1) and [\(58\)](#page-19-1).

Then from eq.[\(82\)](#page-24-1) we find

$$
\lambda_{\alpha}^{(0,1)} = -2\sqrt{2}i\bar{\xi}_{\dot{\alpha}i}\phi u^{-i} + \sum_{n=1}^{\infty}(-4i)(\bar{\xi}_{i}\bar{\sigma}^{\mu}\varepsilon\mathcal{C}^{(n)}\sigma^{\nu})_{\dot{\alpha}}A_{\mu}A_{\nu}(2\sqrt{2}\bar{\phi})^{n-1} \times \left[\alpha_{n}^{(1,0)}\overbrace{u^{+(1}\cdots u^{+1}}^{n}\overbrace{u^{-i}u^{-1}\cdots u^{-1}}^{n+1} + \beta_{n}^{(1,0)}\epsilon^{i1}(u^{+1})^{n-1}(u^{-1})^{n}\right].
$$
\n(92)

From eq.[\(85\)](#page-24-2), we find  $\delta_{\bar{\xi}}^* \bar{\phi} = \sqrt{2} i \bar{\xi}_i \bar{\psi}^i$  and

<span id="page-25-3"></span>
$$
\lambda^{(2,0)} = -2\bar{\xi}_k \bar{\psi}_l u^{-(k} u^{-l)}.
$$
\n(93)

In eq.[\(86\)](#page-24-2),  $\lambda_{\mu}^{(1,1)}$  is given by

<span id="page-25-0"></span>
$$
\lambda_{\mu}^{(1,1)} = -2\psi_{i}\sigma_{\mu}\bar{\xi}_{j} u^{-(i}u^{-j)} \n+ \left(\lambda_{\mu}^{(1,1)}(1) + \tilde{\lambda}_{\mu}^{(1,1)}(1)\right) \left[2u^{+(1}u^{-i}u^{-j}u^{-1)} + \frac{3}{2}\left\{\epsilon^{i1}u^{-(j}u^{-1)} + (i \leftrightarrow j)\right\}\right] \n+ \sum_{n=2}^{\infty} \lambda_{\mu}^{(1,1)}(n) \left[\alpha_{n}^{(1,1)}u^{+(1} \cdots u^{+1}u^{-i}u^{-j}u^{-1} \cdots u^{-1}) + \beta_{n}^{(1,1)}\frac{1}{2}\left\{\epsilon^{i1}u^{+(1} \cdots u^{+1}u^{-j}u^{-1} \cdots u^{-1}) + (i \leftrightarrow j)\right\}\right]
$$

$$
+\gamma_n^{(1,1)}\epsilon^{i1}\epsilon^{j1}(u^{+1})^{n-2}(u^{-1})^n + \delta_n^{(1,1)}\epsilon^{ji}(u^{+1})^{n-1}(u^{-1})^{n+1}\n+ \sum_{n=2}^{\infty} \tilde{\lambda}_{\mu\ ij\ (n)}^{(1,1)}\left[\tilde{\alpha}_n^{(1,1)}\overbrace{u^{+(1}\cdots u^{+1}}^{n}\overbrace{u^{-i}u^{-j}u^{-1}\cdots u^{-1}}^{n+2} + \tilde{\beta}_n^{(1,1)}\frac{1}{2}\left\{\epsilon^{i1}\overbrace{u^{+(1}\cdots u^{+1}}^{n}\overbrace{u^{-j}u^{-1}\cdots u^{-1}}^{n+1} + \tilde{\gamma}_n^{(1,1)}\epsilon^{i1}\epsilon^{j1}(u^{+1})^{n-2}(u^{-1})^n\right],
$$
\n(94)

where

$$
\lambda_{\mu\ ij\ (n)}^{(1,1)} \equiv -\frac{4}{3} (\bar{\xi}_{(i}\bar{\sigma}^{\nu}\varepsilon \mathcal{C}^{(n)}\sigma_{\mu}\bar{\psi}_{j)}) A_{\nu} (2\sqrt{2}\bar{\phi})^{n-1},
$$
\n
$$
\tilde{\lambda}_{\mu\ ij\ (n)}^{(1,1)} \equiv \frac{2}{3} \mathcal{C}^{(n)\alpha\beta} (\sigma^{\nu}\bar{\sigma}_{\mu}\varepsilon)_{\alpha\beta} A_{\nu} (2\sqrt{2}\bar{\phi})^{n-1} \bar{\xi}_{(i}\bar{\psi}_{j)} - \frac{1}{3} (\psi_{(i}\varepsilon \mathcal{C}^{(n)}\sigma_{\mu}\bar{\xi}_{j)}) (2\sqrt{2}\bar{\phi})^{n}, \ (95)
$$

and

$$
(\alpha_{2}^{(1,1)}, \beta_{2}^{(1,1)}, \gamma_{2}^{(1,1)}, \delta_{2}^{(1,1)}) = \left(\frac{5}{4}, \frac{10}{3}, 2, -\frac{1}{2}\right), \alpha_{n}^{(1,1)} = \frac{1}{n+1} \left(6\alpha_{n-1}^{(1,0)} + \alpha_{n-1}^{(1,1)}\right),
$$
  
\n
$$
\beta_{n}^{(1,1)} = \frac{1}{n+1} \left(\frac{6n}{2n-1}\alpha_{n-1}^{(1,0)} + 6\beta_{n-1}^{(1,0)} + \frac{1}{n}\alpha_{n-1}^{(1,1)} + \beta_{n-1}^{(1,1)}\right),
$$
  
\n
$$
\gamma_{n}^{(1,1)} = \frac{1}{n} \left(\frac{6n}{(2n+1)(2n-1)}\alpha_{n-1}^{(1,0)} + 3\beta_{n-1}^{(1,0)}
$$
  
\n
$$
+ \frac{n+1}{(2n+1)2n(2n-1)}\alpha_{n-1}^{(1,1)} + \frac{1}{2n-2}\beta_{n-1}^{(1,1)} + \gamma_{n-1}^{(1,1)}\right),
$$
  
\n
$$
\delta_{n}^{(1,1)} = \frac{1}{n+1} \left(\frac{3(n-1)}{2n-1}\alpha_{n-1}^{(1,0)} - 3\beta_{n-1}^{(1,0)} + \delta_{n-1}^{(1,1)}\right),
$$
  
\n
$$
(\tilde{\alpha}_{2}^{(1,1)}, \tilde{\beta}_{2}^{(1,1)}, \tilde{\gamma}_{2}^{(1,1)}) = \left(\frac{1}{2}, \frac{4}{3}, \frac{4}{5}\right),
$$
  
\n
$$
\tilde{\alpha}_{n}^{(1,1)} = \frac{1}{n+2}\tilde{\alpha}_{n-1}^{(1,1)}, \quad \tilde{\beta}_{n}^{(1,1)} = \frac{1}{n+1} \left(\frac{1}{n}\tilde{\alpha}_{n-1}^{(1,1)} + \tilde{\beta}_{n-1}^{(1,1)}\right),
$$
  
\n
$$
\tilde{\gamma}_{n}^{(1,1)} = \frac{1}{n} \left(\frac{n+1}{(2n+1)2n(2n-1)}\tilde{\alpha}_{n-
$$

Substituting eq.[\(94\)](#page-25-0) to the r.h.s. of eq.[\(86\)](#page-24-2) and using [\(57\)](#page-19-1), [\(58\)](#page-19-1) and

$$
-\sqrt{2}C_{11}^{\alpha\beta}(\sigma_{\mu}\bar{\sigma}^{\nu}\varepsilon)_{\alpha\beta}\tilde{\lambda}_{\nu\ ij}^{(1,1)}(n)\bar{\phi}=\tilde{\lambda}_{\mu\ ij}^{(1,1)}(n+1),\tag{98}
$$

we can see that all the  $O((C_{11})^2)$  terms are gauged away and the contributions to  $\delta_{\xi}^* A_{\mu}$ come only from the  $(C_{11})^1$  part of the second term besides the first term.

The  $C^3$  term in [\(84\)](#page-24-3) becomes

$$
+\frac{16\sqrt{2}}{3}\det C_{11}C_{11}^{\mu\nu}\partial_{\mu}\lambda^{(2,0)}\partial_{\nu}\bar{\phi}~(u^{+1})^4(u^{-1})^2,
$$

which will be gauged away (because of the excessive number of  $u^+$ ). As a result, we can regard eq.[\(84\)](#page-24-3) as having the same structure as eq.[\(86\)](#page-24-2) in terms of the harmonic variables. Therefore, it is solved in a way similar to [\(86\)](#page-24-2): The  $O((C_{11})^2)$  terms in the r.h.s. of eq.[\(84\)](#page-24-3) are completely gauged away, and the relevant contributions to  $\delta_{\xi}^* \phi$  are coming from each  $O((C_{11})^0)$  part of  $\lambda_{\alpha}^{(1,0)}$  and  $\lambda_{\mu}^{(1,1)}$ .

In eq.[\(88\)](#page-24-2),  $\lambda_{\dot{\alpha}}^{(2,1)}$  $\alpha^{(2,1)}$  is given by

<span id="page-27-1"></span>
$$
\lambda_{\dot{\alpha}}^{(2,1)} = -2\bar{\xi}_{\dot{\alpha}(i}D_{jk)}u^{-(i}u^{-j}u^{-k)} - (\bar{\xi}_{i}\bar{\sigma}^{\nu}\varepsilon C_{11}\sigma^{\mu})_{\dot{\alpha}}\partial_{\mu}(A_{\nu}\bar{\phi})u^{-(i}u^{-1}u^{-1}) \n- \sum_{n=2}^{\infty} 2(\bar{\xi}_{i}\bar{\sigma}^{\nu}\varepsilon C^{(n)}\sigma^{\mu})_{\dot{\alpha}}\partial_{\mu}\left\{A_{\nu}(2\sqrt{2}\bar{\phi})^{n}\right\}\left[\alpha_{n}^{(2,1)}\overbrace{u^{+(1}\cdots u^{+1}}^{n-1}\overbrace{u^{-i}u^{-1}\cdots u^{-1}}^{n+2}) + \beta_{n}^{(2,1)}\epsilon^{i1}(u^{+1})^{n-2}(u^{-1})^{n+1}\right],
$$
\n(99)

where  $\alpha_n^{(2,1)}, \beta_n^{(2,1)}$  are given in [\(78\)](#page-23-0). We can easily find that in the r.h.s. of [\(88\)](#page-24-2) the terms proportional to  $(C_{11})^n$   $(n \geq 2)$  are completely cancelled by this gauge parameter, so that there are no  $O((C_{11})^2)$  terms in  $\delta_{\xi}^* \bar{\psi}^i$ .

Substituting  $(92)$ ,  $(90)$ ,  $(93)$ ,  $(94)$  and  $(99)$  into eq. $(87)$ , we can check that the gauge parameter  $\lambda_{\alpha}^{(1,2)}$  having the same form as [\(79\)](#page-23-1) is sufficient. To see it is sufficient, we need [\(57\)](#page-19-1)–[\(60\)](#page-19-1). In fact, in order to determine  $\delta_{\xi}^* \psi^i$ , only the precise forms of  $\chi_{\alpha i j k(0)}$  and  $\chi_{\alpha i j(1)}$ are needed as in the  $\mathcal{N} = (1, 0)$  case:

$$
\chi_{\alpha ijk(0)} = 0, \quad \chi_{\alpha i j(1)} = \frac{32}{5} \left[ D_{(ij} (\bar{\xi}_k) \bar{\sigma}^\mu \varepsilon C_{11})^\beta \varepsilon_{\alpha \beta} A_\mu + 2i (\varepsilon C_{11} \psi_{(i)} \alpha \bar{\xi}_j \bar{\psi}_k) \right] \epsilon^{k1}.
$$
 (100)

Substituting  $(90)$ ,  $(93)$  and  $(94)$  to eq. $(89)$  and collecting  $u^+$ -independent terms, we obtain  $\delta_{\xi}^* D^{ij}$ . We do not need the precise form of  $\lambda^{(2,2)}$ . To determine  $\delta_{\xi}^* D^{ij}$ , we can use [\(60\)](#page-19-1) and [\(58\)](#page-19-1).

The  $\mathcal{N} = (0, 1/2)$  supersymmetry transformation can be read from the following result by setting  $\bar{\xi}^1 = 0$ :

<span id="page-27-0"></span>
$$
\delta_{\bar{\xi}}^* \phi = c_1 i(\bar{\xi}^1 \bar{\sigma}^\mu \varepsilon C_{11} \psi^1) A_\mu, \tag{101}
$$

$$
\delta_{\bar{\xi}}^* \bar{\phi} = -\sqrt{2} i \bar{\xi}^i \bar{\psi}_i, \tag{102}
$$

$$
\delta_{\xi}^{*} A_{\mu} = i \bar{\xi}^{i} \bar{\sigma}_{\mu} \psi_{i} + c_{2} i (\bar{\xi}^{1} \bar{\sigma}^{\nu} \varepsilon C_{11} \sigma_{\mu} \bar{\psi}^{1}) A_{\nu} \n+ c_{3} i C_{11}^{\alpha \beta} (\sigma^{\nu} \bar{\sigma}_{\mu} \varepsilon)_{\alpha \beta} \bar{\xi}^{1} \bar{\psi}^{1} A_{\nu} + c_{4} i (\bar{\xi}^{1} \bar{\sigma}_{\mu} \varepsilon C_{11} \psi^{1}) \bar{\phi},
$$
\n(103)

$$
\delta_{\xi}^{*} \psi^{\alpha i} = -\sqrt{2} (\bar{\xi}^{i} \bar{\sigma}^{\mu})^{\alpha} \partial_{\mu} \phi - 2D^{(i)} (\bar{\xi}^{1}) \bar{\sigma}^{\mu} \varepsilon C_{11})^{\alpha} A_{\mu} + 4i \bar{\xi}^{(i} \bar{\psi}^{1} (\psi^{1}) \varepsilon C_{11})^{\alpha} \n+ \delta_{2}^{i} \left[ c_{5} (\bar{\xi}^{1} \bar{\sigma}^{\mu} \varepsilon C_{11})^{\alpha} \partial_{\mu} \phi \bar{\phi} + (\bar{\xi}^{1} \bar{\sigma}^{\mu} \varepsilon C_{11} \sigma^{\nu} \bar{\sigma}^{\rho})^{\alpha} \left\{ c_{6} \partial_{\rho} (A_{\mu} A_{\nu}) + c_{7} A_{\mu} \partial_{\nu} A_{\rho} \right\} \n+ c_{8} (\bar{\xi}^{1} \bar{\sigma}^{\mu})^{\alpha} C_{11}^{\beta \gamma} (\sigma^{\nu} \bar{\sigma}^{\rho} \varepsilon)_{\beta \gamma} \partial_{\rho} A_{\mu} A_{\nu} \right] \n+ \delta_{2}^{i} \det C_{11} \left[ (\bar{\xi}^{1} \bar{\sigma}^{\mu})^{\alpha} \left\{ c_{9} D^{11} A_{\mu} \bar{\phi} + c_{10} i \bar{\psi}^{1} \bar{\psi}^{1} A_{\mu} \right\} + c_{11} i \bar{\xi}^{1} \bar{\psi}^{1} \psi^{\alpha 1} \bar{\phi} \right], \tag{104}
$$

$$
\delta_{\bar{\xi}}^* \bar{\psi}_{\dot{\alpha}i} = (\bar{\xi}_i \bar{\sigma}^{\mu\nu})_{\dot{\alpha}} F_{\mu\nu} - \bar{\xi}_{\dot{\alpha}}^j D_{ij} + \delta_i^1 c_{12} \partial_\mu \left[ (\bar{\xi}^1 \bar{\sigma}^\nu \varepsilon C_{11} \sigma^\mu)_{\dot{\alpha}} A_\nu \bar{\phi} \right], \tag{105}
$$

$$
\delta_{\xi}^{*}D^{11} = c_{13}i\partial_{\mu}(\bar{\xi}^{1}\bar{\sigma}^{\mu}\psi^{1}),
$$
\n
$$
\delta_{\xi}^{*}D^{12} = -2i\partial_{\mu}(\bar{\xi}^{(1}\bar{\sigma}^{\mu}\psi^{2}))
$$
\n
$$
+ i\partial_{\mu} \left[c_{14}(\bar{\xi}^{1}\bar{\sigma}^{\nu}\varepsilon C_{11}\sigma^{\mu}\bar{\psi}^{1})A_{\nu} + c_{15}C_{11}^{\alpha\beta}(\sigma^{\nu}\bar{\sigma}^{\mu}\varepsilon)_{\alpha\beta}A_{\nu}\bar{\xi}^{1}\bar{\psi}^{1} + c_{16}(\bar{\xi}^{1}\bar{\sigma}^{\mu}\varepsilon C_{11}\psi^{1})\bar{\phi}\right],
$$
\n
$$
\delta_{\xi}^{*}D^{22} = -2i\partial_{\mu}(\bar{\xi}^{2}\bar{\sigma}^{\mu}\psi^{2}) - \frac{4}{3}i\partial_{\mu} \left[2(\bar{\xi}^{(1}\bar{\sigma}^{\nu}\varepsilon C_{11}\sigma^{\mu}\bar{\psi}^{2}))A_{\nu} - C_{11}^{\alpha\beta}(\sigma^{\nu}\bar{\sigma}^{\mu}\varepsilon)_{\alpha\beta}A_{\nu}\bar{\xi}^{(1}\bar{\psi}^{2)} + \sqrt{2}(\bar{\xi}^{(1}\bar{\sigma}^{\mu}\varepsilon C_{11}\psi^{2}))\bar{\phi}\right]
$$
\n
$$
+ i \det C_{11}\partial_{\mu} \left[c_{17}(\bar{\xi}^{1}\bar{\sigma}^{\nu}\sigma^{\mu}\bar{\psi}^{1})A_{\nu}\bar{\phi} + c_{18}\bar{\xi}^{1}\bar{\psi}^{1}A^{\mu}\bar{\phi} + c_{19}(\bar{\xi}^{1}\bar{\sigma}^{\mu}\psi^{1})\bar{\phi}^{2}\right], \qquad (106)
$$

where  $c_i$ 's are certain constants.

### <span id="page-28-0"></span>References

- <span id="page-28-1"></span>[1] J. H. Schwarz and P. Van Nieuwenhuizen, Lett. Nuovo Cim. 34, 21 (1982).
- [2] D. Klemm, S. Penati and L. Tamassia, Class. Quant. Grav. 20 (2003) 2905, [hep-th/0104190;](http://arxiv.org/abs/hep-th/0104190) S. Ferrara and M. A. Lledo, JHEP 0005 (2000) 008, [hep-th/0002084;](http://arxiv.org/abs/hep-th/0002084) S. Ferrara, M. A. Lledo and O. Macia, JHEP 0309 (2003) 068, [hep-th/0307039.](http://arxiv.org/abs/hep-th/0307039)
- <span id="page-28-3"></span><span id="page-28-2"></span>[3] H. Ooguri and C. Vafa, Adv. Theor. Math. Phys. 7 (2003) 53, [hep-th/0302109;](http://arxiv.org/abs/hep-th/0302109) Adv. Theor. Math. Phys. 7 (2004) 405, [hep-th/0303063.](http://arxiv.org/abs/hep-th/0303063)
- [4] N. Berkovits and N. Seiberg, JHEP 0307 (2003) 010, [hep-th/0306226.](http://arxiv.org/abs/hep-th/0306226)
- <span id="page-29-1"></span><span id="page-29-0"></span>[5] J. de Boer, P. A. Grassi and P. van Nieuwenhuizen, Phys. Lett. B574 (2003) 98, [hep-th/0302078.](http://arxiv.org/abs/hep-th/0302078)
- <span id="page-29-2"></span>[6] N. Seiberg, JHEP 0306, 010 (2003), [hep-th/0305248.](http://arxiv.org/abs/hep-th/0305248)
- <span id="page-29-3"></span>[7] T. Araki, K. Ito and A. Ohtsuka, Phys. Lett. B573 (2003) 209, [hep-th/0307076.](http://arxiv.org/abs/hep-th/0307076)
- [8] R. Britto, B. Feng and S. J. Rey, JHEP 0307 (2003) 067, [hep-th/0306215;](http://arxiv.org/abs/hep-th/0306215) S. Terashima and J. T. Yee, JHEP 0312 (2003) 053, [hep-th/0306237;](http://arxiv.org/abs/hep-th/0306237) M. Hatsuda, S. Iso and H. Umetsu, Nucl. Phys. B671 (2003) 217, [hep-th/0306251;](http://arxiv.org/abs/hep-th/0306251) R. Britto, B. Feng and S. J. Rey, JHEP 0308 (2003) 001, [hep-th/0307091;](http://arxiv.org/abs/hep-th/0307091) M. T. Grisaru, S. Penati and A. Romagnoni, JHEP 0308 (2003) 003, [hep-th/0307099;](http://arxiv.org/abs/hep-th/0307099) R. Britto and B. Feng, Phys. Rev. Lett. 91 (2003) 201601, [hep-th/0307165;](http://arxiv.org/abs/hep-th/0307165) A. Romagnoni, JHEP 0310 (2003) 016, [hep-th/0307209;](http://arxiv.org/abs/hep-th/0307209) O. Lunin and S. J. Rey, JHEP 0309 (2003) 045, [hep-th/0307275;](http://arxiv.org/abs/hep-th/0307275)
	- D. Berenstein and S. J. Rey, Phys. Rev. D68 (2003) 121701, [hep-th/0308049;](http://arxiv.org/abs/hep-th/0308049)

M. Alishahiha, A. Ghodsi and N. Sadooghi, Nucl. Phys. B 691 (2004) 111, [hep-th/0309037;](http://arxiv.org/abs/hep-th/0309037)

A. Imaanpur and S. Parvizi, JHEP 0407 (2004) 010, [hep-th/0403174;](http://arxiv.org/abs/hep-th/0403174)

A. T. Banin, I. L. Buchbinder and N. G. Pletnev, JHEP 0407 (2004) 011 [hep-th/0405063;](http://arxiv.org/abs/hep-th/0405063)

- A. Sako and T. Suzuki, JHEP 0411 (2004) 010, [hep-th/0408226;](http://arxiv.org/abs/hep-th/0408226)
- I. Jack, D.R.T. Jones and L.A. Worthy, [hep-th/0412041;](http://arxiv.org/abs/hep-th/0412041)
- S. Penati and A. Romagnoni, JHEP 0502 (2005) 064, [hep-th/0412041;](http://arxiv.org/abs/hep-th/0412041)
- O. D. Azorkina, A. T. Banin, I. L. Buchbinder and N. G. Pletnev, [hep-th/0502008;](http://arxiv.org/abs/hep-th/0502008)
- T. Hatanaka, S. V. Ketov, Y. Kobayashi and S. Sasaki, [hep-th/0502026;](http://arxiv.org/abs/hep-th/0502026)

M. Billo, M. Frau, F. Lonegro and A. Lerda, [hep-th/0502084.](http://arxiv.org/abs/hep-th/0502084)

<span id="page-29-4"></span>[9] A. Imaanpur, JHEP 0309 (2003) 077, [hep-th/0308171;](http://arxiv.org/abs/hep-th/0308171) JHEP 0312 (2003) 009, [hep-th/0311137;](http://arxiv.org/abs/hep-th/0311137)

P. A. Grassi, R. Ricci and D. Robles-Llana, JHEP 0407 (2004) 065, [hep-th/0311155;](http://arxiv.org/abs/hep-th/0311155) R. Britto, B. Feng, O. Lunin and S. J. Rey, Phys. Rev. D 69 (2004) 126004, [hep-th/0311275;](http://arxiv.org/abs/hep-th/0311275)

M. Billo, M. Frau, I. Pesando and A. Lerda, JHEP 0405 (2004) 023, [hep-th/0402160.](http://arxiv.org/abs/hep-th/0402160)

- [10] B. Chandrasekhar and A. Kumar, JHEP 0403 (2004) 013, [hep-th/0310137;](http://arxiv.org/abs/hep-th/0310137) T. Inami and H. Nakajima, Prog. Theor. Phys. 111 (2004) 961, [hep-th/0402137;](http://arxiv.org/abs/hep-th/0402137) B. Chandrasekhar, Phys. Rev. D70 (2004) 125003, [hep-th/0408184;](http://arxiv.org/abs/hep-th/0408184) L. Alvarez-Gaume and M. A. Vazquez-Mozo, [hep-th/0503016.](http://arxiv.org/abs/hep-th/0503016)
- [11] S. Ferrara and E. Sokatchev, Phys. Lett. B579 (2004) 226, [hep-th/0308021;](http://arxiv.org/abs/hep-th/0308021) S.V. Ketov and S. Sasaki, Phys. Lett. B 595 (2004) 530, [hep-th/0404119;](http://arxiv.org/abs/hep-th/0404119) Phys. Lett. B 597 (2004) 105, [hep-th/0405278;](http://arxiv.org/abs/hep-th/0405278) [hep-th/0407211.](http://arxiv.org/abs/hep-th/0407211)
- <span id="page-30-2"></span><span id="page-30-1"></span>[12] E. Ivanov, O. Lechtenfeld and B. Zupnik, JHEP 0402 (2004) 012, [hep-th/0308012;](http://arxiv.org/abs/hep-th/0308012) Nucl. Phys. B 707 (2005) 69, [hep-th/0408146.](http://arxiv.org/abs/hep-th/0408146)
- <span id="page-30-3"></span>[13] T. Araki, K. Ito and A. Ohtsuka, Phys. Lett. B606 (2995) 202, [hep-th/0410203.](http://arxiv.org/abs/hep-th/0410203)
- <span id="page-30-6"></span>[14] T. Araki, K. Ito and A. Ohtsuka, JHEP 0401 (2004) 046, [hep-th/0401012.](http://arxiv.org/abs/hep-th/0401012)
- <span id="page-30-0"></span>[15] T. Araki and K. Ito, Phys. Lett. B595 (2004) 513, [hep-th/0404250.](http://arxiv.org/abs/hep-th/0404250)
- <span id="page-30-4"></span>[16] S. Ferrara, E. Ivanov, O. Lechtenfeld, E. Sokatchev and B. Zupnik, Nucl. Phys. B 704 (2005) 154, [hep-th/0405049.](http://arxiv.org/abs/hep-th/0405049)
- <span id="page-30-5"></span>[17] J. Wess and J. Bagger, "Supersymmetry and Supergravity," Princeton University Press, 1992.
- <span id="page-30-7"></span>[18] A. Galperin, E. Ivanov, V. Ogievetsky and E. Sokatchev, "Harmonic Superspace", Cambridge University Press , 2001.
- <span id="page-30-8"></span>[19] B.M. Zupnik, Phys. Lett. B183 (1987) 175.
- [20] I. L. Buchbinder and I. B. Samsonov, Grav.Cosmol.8 (2002) 17, [hep-th/0109130.](http://arxiv.org/abs/hep-th/0109130)