

Six Gluon Open Superstring Disk Amplitude, Multiple Hypergeometric Series and Euler-Zagier Sums

D. Oprisa ^a and S. Stieberger ^b

^a *Max-Planck-Institut für Physik,
Föhringer Ring 6, 80805 München, Germany*

^b *Arnold-Sommerfeld-Center for Theoretical Physics,
Department für Physik, Ludwig-Maximilians-Universität München,
Theresienstraße 37, 80333 München, Germany*

Abstract

The six gluon disk amplitude is calculated in superstring theory. This amplitude probes the gauge interactions with six external legs on Dp -branes, in particular including *e.g.* F^6 -terms. The full string S -matrix can be expressed by six generalized multiple hypergeometric functions (triple hypergeometric functions), which in the effective action play an important rôle in arranging the higher order α' gauge interaction terms with six external legs (like $F^6, D^4F^4, D^2F^5, D^6F^4, D^2F^6, \dots$).

A systematic and efficient method is found to calculate tree-level string amplitudes by equating seemingly different expressions for one and the same string S -matrix: Comparable to Riemann identities appearing in string-loop calculations, we find an intriguing way of using world-sheet supersymmetry to generate a system of non-trivial equations for string tree-level amplitudes. These equations result in algebraic identities between different multiple hypergeometric functions. Their (six-dimensional) solution gives the ingredients of the string S -matrix. We derive material relevant for any open string six-point scattering process: relations between triple hypergeometric functions, their integral representations and their α' - (momentum)-expansions given by (generalized) Euler-Zagier sums or (related) Witten zeta-functions.

Contents

1. Introduction	2
2. Tree-level scattering of open superstrings on Dp -branes	8
2.1. Open superstring gluon S -matrix	8
2.2. Four gluon open superstring S -matrix revisited	11
3. Six gluon open superstring S -matrix	17
3.1. General expression for the amplitude	18
3.2. Specifying the integration region \mathcal{I}_π	27
3.3. System of equations and its six-element solution	34
3.4. Momentum expansion of the string S -matrix	39
4. Multiple hypergeometric functions, generalized Kampé de Fériet function and Euler-Zagier sums	46
4.1. Generalized hypergeometric functions ${}_pF_q$	46
4.2. Generalized Kampé de Fériet function $F_{C:D}^{A:B}$ and triple hypergeometric function $F^{(3)}$	47
4.3. Harmonic Number and Euler sums	50
4.4. Momentum expansion of hypergeometric functions ${}_3F_2$ and ${}_4F_3$	57
4.5. Multiple zeta values, Euler-Zagier sums and momentum expansion of $F^{(3)}$	59
4.6. Momentum expansion of singular triple hypergeometric functions $F^{(3)}$	69
5. Reducible diagrams and contact interactions	72
6. Concluding remarks	79
Appendix A. Five gluon open superstring S -matrix	82
Appendix B. Series involving the Harmonic Number	93
Appendix C. Momentum expansion of ${}_4F_3 \left[\begin{matrix} 1+b, 1+a, 1+c, -j \\ 2+b+e, 2+a+d, 2+c+f \end{matrix} \right]$	95
Appendix D. Some group theoretical facts	96

1. Introduction

Due to the non-locality of string theory or the presence of infinite many massive string states the world-volume effective action describing the dynamics of massless fields of a Dp -brane is organized as an infinite power series in the string tension α' . A famous example for such a series is represented by the tree-level gauge sector of a Dp -brane. For F the gauge field strength and D the gauge covariant derivative¹, this series may be formally written:

$$\mathcal{L}_{\text{effective}}^{Dp} = \text{Tr} \sum'_{\substack{m \geq 2 \\ n \geq 0}} \alpha'^{\frac{1}{2}n+m-2} D^n F^m . \quad (1.1)$$

Of course, this series contains a lot of redundant terms, which may be eventually eliminated through applying Bianchi identities, equations of motions, partial integrations and field redefinitions. Moreover, due to the relation

$$[D_\mu, D_\nu] F_{\rho\sigma} = -i [F_{\mu\nu}, F_{\rho\sigma}] \quad (1.2)$$

certain combinations of derivative terms may be converted into non-derivative terms and vice versa. To this end, the prime at the sum indicates, that the sum (1.1) should be evaluated over only all independent invariants. In the supersymmetric case the latter are supposed to represent supersymmetric invariants. Furthermore, according to the definition of the effective action, each term in the series (1.1) accounts for a gluon contact interaction represented by an irreducible Feynman diagram of the underlying (higher derivative) gauge theory. Whether the series converges or may be even written as an analytic function in F and DF is of burning interest. To determine the expansion (1.1) for a given string background, in that background one calculates gauge boson string tree-level scattering amplitudes (their string S -matrices) and their α' -expansion matches certain terms of $Eq.$ (1.1) (after proper subtraction of reducible diagrams). The fact, that $Eq.$ (1.2) allows us to write terms in the effective action in various different ways, which are all equivalent from the string S -matrix point of view, is related to the fact, that a string S -matrix does not feel those redefinitions (on-shell). The lowest order terms of (1.1), *i.e.* the α'^0 -order, describes just the ordinary Non-Abelian Yang-Mills theory, *i.e.* the interaction term $\text{Tr}F^2$. While the next leading order α' is absent in the superstring, the α'^2 -order has been completely

¹ F and D are in the adjoint representation of the gauge group. Some group theoretical facts are summarized in appendix D.

matched to pieces of a four–gluon open superstring scattering S –matrix. To this end, up to the order α'^2 , the series (1.1) looks as follows [1,2,3]

$$\mathcal{L}_{\text{effective}}^{Dp} = \text{Tr} \left\{ F_{mn}^2 - \frac{1}{3} (2\pi\alpha')^2 \left(F_{ab}F_{bd}F_{ca}F_{dc} + \frac{1}{2} F_{ab}F_{bc}F_{cd}F_{da} - \frac{1}{4} F_{ab}F_{ba}F_{cd}F_{dc} - \frac{1}{8} F_{ab}F_{cd}F_{ba}F_{dc} \right) + \mathcal{O}(\alpha'^3) \right\} . \quad (1.3)$$

Up to this α' –order derivative terms ($n \neq 0$) may be eliminated as a result of field redefinitions [3]. The series (1.1) is different for the bosonic and superstring. In this article we shall only discuss the more interesting superstring case. Furthermore, it drastically simplifies in the case of an Abelian gauge group, *i.e.* a single Dp –brane, due to the absence of a huge set of gluon interactions. Up to the order α'^2 , displayed *Eq.* (1.3), the terms may be derived from the so–called (generalized) Born–Infeld action [4]

$$\mathcal{L} = \text{STr} \sqrt{\det(\delta_{mn} + 2\pi\alpha' F_{mn})} = \text{STr} \sum_{\nu_i \geq 0} c_{\nu_1 \nu_2 \dots \nu_n} (\text{tr} F^2)^{\nu_1} (\text{tr} F^4)^{\nu_2} \dots (\text{tr} F^{2n})^{\nu_n} , \quad (1.4)$$

with the coefficients

$$c_{\nu_1 \nu_2 \dots \nu_n} = (2\pi\alpha')^{2\nu_1 + 4\nu_2 + \dots + 2n\nu_n} \left(-\frac{1}{4} \right)^{\sum_i \nu_i} \frac{1}{\nu_1! \dots \nu_n!} \frac{1}{2^{\nu_2} 3^{\nu_3} \dots n^{\nu_n}} \quad (1.5)$$

after expansion to fourth order in F and applying the rule of the symmetrized gauge trace STr . The latter “averages” over all group structures

$$\text{STr}(A_1 \dots A_n) := \frac{1}{n!} \text{Tr}(A_{(1} \dots A_n)}) , \quad (1.6)$$

with all permutations of gauge fields taken into account.

However, the series (1.1) departs already at the order α'^3 from this nice and closed form (1.4), due to the presence of F^5 and $D^2 F^4$ –terms. Those terms appear with a $\zeta(3)$ factor in the effective action. While the $D^2 F^4$ can be fixed through expanding the four gluon string S –matrix up to order α'^3 [5], the pure F^5 terms can only be fixed by computing a five gluon string S –matrix [6]. In fact, there are several important works, which investigate the structure of the higher order α' gauge interaction terms in the string effective action. However, only F^4 , $D^{2n} F^4$ and F^5 , $D^{2n} F^5$ terms have been thoroughly investigated on the basis of the four– and five–gluon string S –matrix results, respectively [7,8], since only those two amplitudes have been calculated [1,6]. However, to probe the

full α'^4 order of (1.1), one has to compute six-point gluon scattering amplitudes. This amplitude comprises the necessary information to obtain the structure of the F^6 and also $D^{2n}F^6$ -terms.

α'^0	1	F²			
α'^1	0	F^3	D^2F^2		
α'^2	$\zeta(2)$	F⁴	D^2F^3	D^4F^2	
α'^3	$\zeta(3)$	F⁵	D²F⁴	D^6F^2	
α'^4	$\zeta(4)$	F⁶	D⁴F⁴	D²F⁵	
α'^5	$\zeta(5)$	F⁷	D⁶F⁴	D⁴F⁵	D²F⁶
\vdots	

Table 1: Higher order F -terms appearing at a given α' -order in the supersymmetric D -brane action Eq. (1.1).

The expected higher order α' gauge interaction terms are displayed in Table 1 up to α'^5 . The terms in bold are the interactions, which contribute to the series (1.1), while the others may be eliminated (*cf.* the comment below (1.1)). Furthermore, we show the respective zeta-function coefficients of those terms in the effective action.

Computing the six gluon string S -matrix in superstring theory is the main part of the present article. More concretely, we calculate the superstring tree-level S -matrix with six external gauge bosons. The latter are represented by open strings, hence the string world-sheet topology of this scattering process is a disk. Furthermore, we present the relevant material, which is needed for any open string six-point scattering process involving six open strings: the integral representations of the triple hypergeometric function, various identities between multiple hypergeometric functions and Euler-Zagier sums. Calculating tree-level string S -matrices with more than five external states is certainly quite challenging from the technical point of view. In fact, so far only string tree-level amplitudes involving five external states have been elaborated. More precisely², five gluon tree-level amplitudes have been determined in the heterotic string in *Ref.* [11], while in the type I superstring

² From the technical point of view, five open string tree-level amplitudes are related to *e.g.* three open and one closed string disk amplitudes [9]. Note, that in *Ref.* [10] a specific tree-level string amplitude with two open and two closed strings has been calculated. On the disk, this amplitude is related to an open string six-point function of a very special kind, which has made tractable the problem.

in *Refs.* [12,6]. Hence, it is appropriate to state, that this is the first work, where the problem of tree-level string amplitudes involving six external states has been addressed.

We find a very powerful trick, to tackle the problem of calculating N -point tree-level string S -matrices in the superstring. More precisely, in order to guarantee a total ghost charge of -2 on the disk, two vertex operators of the N gluons have to be chosen in the (-1) -ghost picture, while the remaining $N - 2$ vertices are put into the zero-ghost picture. However, we are completely free, which pair of vertex operators we choose to put into the (-1) -ghost picture. There are $\binom{N}{2} = \frac{1}{2}(N - 1)N$ possibilities, which all have to lead to the same result. For a given space-time kinematics each of these choices leads to a seemingly different expression. Quite generically, such an expression is given by some multiple hypergeometric functions, multiplied with some polynomials in the kinematic invariants and some Γ -functions depending on the latter. All these expressions have to be equated. This gives a set of non-trivial relations between different hypergeometric functions, which allows us to express many of them in terms of just a few. These identities represent algebraic equations between different multiple hypergeometric functions. Their solution gives the ingredients of the string S -matrix. Many of the identities between hypergeometric functions we obtain by the above procedure cannot be found in the literature, see *e.g.* [13]. Hence, from the mathematical point of view identities between different hypergeometric functions may be inferred from higher point string tree-level amplitudes as a result of supersymmetry. We shall point out, that the above described symmetry, following from world-sheet supersymmetry should not be confused with the cyclic invariance of a gluon string S -matrix. Applying this symmetry on a given amplitude would also lead to seemingly different expressions for one and the same kinematic structure. Equating the latter would also give a system of equations. However, this system is much smaller, in fact too small, to write down the full string S -matrix in terms of a few functions, which in contrast is possible using the above described trick using world-sheet supersymmetry.

There is a striking relation between multiple Gaussian hypergeometric functions or their integral representations, which are the relevant objects for tree-level string calculations and zeta-functions of various kinds, which play a key role in modern number theory. A simple example for this relation may be given by the following integral ($\alpha's > -1$):

$$\int_0^1 dx \int_0^1 dy (1-xy)^{\alpha's-1} = \frac{H_{\alpha's}}{\alpha's} = \zeta(2) - \alpha's \zeta(3) + (\alpha's)^2 \zeta(4) - (\alpha's)^3 \zeta(5) + \dots \quad (1.7)$$

This integral may calculate the α' -expansion for a certain kinematics of a five-point string amplitude with s being some kinematic invariant. Besides, H_n is the harmonic number (*cf.* section 4 for more details). Similarly, one may derive

$$\int_0^1 dx \int_0^1 dy \int_0^1 dz (1 - xyz)^{\alpha' s - 2} = \zeta(2) + \alpha' s [\zeta(2) - 2 \zeta(3)] + (\alpha' s)^2 [\zeta(2) - 2 \zeta(3) + \frac{5}{4} \zeta(4)] + \dots \quad (1.8)$$

accounting for a certain (simplified) six-point scattering process³. In the above two examples the basic Riemann zeta-functions appear. The latter are given by the following integer sum:

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} \quad , \quad s \geq 2. \quad (1.10)$$

Generalization of them are the multiple zeta values, *e.g.*:

$$\zeta(s_1, s_2, s_3) = \sum_{\substack{m_i=1 \\ m_3 < m_2 < m_1}}^{\infty} \frac{1}{m_1^{s_1} m_2^{s_2} m_3^{s_3}} \quad , \quad s_1 \geq 2 \quad , \quad s_2, s_3 \geq 1 \quad , \quad (1.11)$$

which appear in the integral:

$$\int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{(1 - xy)^{\alpha' s}}{(1 - xyz)^2} = \zeta(2) - \alpha' s \zeta(3) + (s\alpha')^2 \zeta(2, 1, 1) + \dots \quad , \quad (1.12)$$

with $\zeta(2, 1, 1) = \zeta(4)$. A class of integrals, similar to (1.8) and (1.12) may be evaluated by calculating the relevant multiple zeta values or related Euler sums (*cf.* section 4 for more details). However, the α' -expansion of the full disk six-point string S -matrix, which is eventually expressed through some triple hypergeometric functions, generically leads to new types of zeta-functions. The latter are multiple sums over integers, which in most of the cases we have to evaluate case by case. One simplified example is the following type:

$$\int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{(1 - xy)^{\alpha' s} (1 - yz)^{\alpha' t}}{(1 - xyz)^2} = \zeta(2) - \alpha' (s + t) \zeta(3) + \alpha'^2 [s^2 + t^2 + \frac{7}{4} st] \zeta(4) + \dots \quad (1.13)$$

³ Due to the nine kinematic invariants the integrands, which appear in any six-point scattering process on the disk, are combinations of the nine polynomials:

$$x \quad , \quad (1 - x) \quad , \quad y \quad , \quad (1 - y) \quad , \quad z \quad , \quad (1 - z) \quad , \quad (1 - xy) \quad , \quad (1 - yz) \quad , \quad (1 - xyz) \quad . \quad (1.9)$$

Their integrals may be expressed through the triple hypergeometric function $F^{(3)}$ (*cf.* section 4).

Extracting from the above integral the second order in α' requires to evaluate the following triple sum

$$\sum_{m_i=1}^{\infty} \frac{m_3}{m_1 m_2 (m_1 + m_3) (m_2 + m_3) (m_1 + m_2 + m_3)} = \frac{7}{4} \zeta(4) \quad (1.14)$$

among others. The evaluation of those sums, which is the topic of subsection 4.5 is closely related to the subject of Tornheim double series, Witten zeta-functions and multiple Euler/Zagier sums. Those sums generically may not be simply expressed by the Riemann zeta-function. An example of a Witten zeta-function is

$$W(a, b, c) = \sum_{m, n=1}^{\infty} \frac{1}{m^a n^b (m+n)^c}, \quad (1.15)$$

which also occurs after expanding the triple hypergeometric function w.r.t. α' .

Just as special values of the Riemann zeta-functions (1.10) are related to the Euler characteristics of moduli spaces \mathcal{M}_g of Riemann surfaces of genus g , *i.e.* $\chi(\mathcal{M}_g) \sim \zeta(2g)$, $g \geq 2$, special values of the encountered new types of zeta-functions calculate the volume of certain moduli spaces of vector bundles of curves. Hence the α' -expansion of our six gluon disk amplitude may perhaps be related to some geometric setup described by the relation between the triple hypergeometric function and new types of zeta-functions.

While computing string amplitudes with external gluons allows to determine (1.1) directly by means of extracting the relevant parts of the effective action from the string S -matrix amplitude there have been presented some indirect approaches to tackle this problem: By matching the open string spectrum with a spectrum following from a higher order Yang-Mills action one obtains information about the higher order terms in (1.1), see *Refs.* [14,15] and *Refs.* [16] for a review. More recently, an other approach has been suggested [17]: Requiring the existence of certain *BPS* solutions to the equations of motions allowed to propose the α'^4 -part of (1.1). Since all those approaches are indirect, in particular based on field-theory, it is certainly important to check these findings against the results derived from the string amplitudes [18]. Moreover, the various transcendental zeta values $\zeta(3), \zeta(5), \zeta(3)^2, \dots$, which appear in the α' -expansion of the six gluon string S -matrix, are certainly hard to obtain in any field-theory approach.

Our results may be summarized under the following three points:

- An efficient method to calculate supersymmetric N -point tree – level string amplitudes is presented.
- The six gluon open superstring disk amplitude can be expressed through a basis of six triple hypergeometric functions, which encode the full α' – dependence.
- Material to obtain the α' – expansion of these functions is derived : We calculate many multiple Euler – Zagier sums including multiple harmonic series.

2. Tree-level scattering of open superstrings on Dp -branes

2.1. Open superstring gluon S -matrix

In the following we study the gauge dynamics on the Dp -brane world volume with sixteen supercharges. Our discussion holds for both type I or type IIA/IIB superstring theory. The gauge interactions are described by an effective D -brane action of the sort (1.1), which is derived from open string scattering amplitudes. The α' -expansion of the latter reproduces certain interaction terms of the effective action. The massless open string states of a Dp -brane are the gauge vectors A^μ , which are the longitudinal massless excitations of the NS -ground state and the gauge fermions ψ^μ originating from the R -ground state, with $\mu = 0, \dots, p$. The Dp -brane fields ψ^μ and A^μ obey Neumann boundary conditions. In addition, there are the fields with transverse index $\mu = p + 1, \dots, 9$, which obey Dirichlet boundary conditions. Here, we shall not discuss their contribution to the D -brane action. Their effects may be deduced from T -duality [19]. Hence, without loss of generality, we may focus on an unwrapped $D9$ -brane, *i.e.* $p = 9$. The latter carries 16 supercharges. Hence, for concreteness we perform our calculation in an uncompactified $D = 10$ type I superstring background with $SO(32)$ gauge group. This is no restriction on the dimension nor on the gauge group of the Dp -brane. It only means, that no transversal oscillations of the branes are allowed and the fields A^μ, ψ^μ with $\mu = 0, \dots, 9$, live on the $D9$ -brane world-volume. Qualitative different results are only to be expected for $D3$ -branes due to the reduced number of kinematic invariants in four space-time dimensions (*cf.* comments following later).

A string S -matrix \mathcal{A}_N of N external gluons with momenta k_i , polarizations ξ_i and color indices a_i may be formally arranged as the sum:

$$\mathcal{A}_N(k_1, \xi_1, a_1; \dots; k_N, \xi_N, a_N) = \sum_{\substack{\pi \in \overline{(1,2,\dots,N)} \\ (N-1)! \text{ permutations } \pi}} \text{Tr}(\lambda^{\pi(1)} \lambda^{\pi(2)} \dots \lambda^{\pi(N)}) A^\pi. \quad (2.1)$$

The sum runs over all $(N-1)!$ cyclically inequivalent permutations $(\pi(1), \pi(2), \dots, \pi(N))$ of the N gluons, labeled by the N integers i , with $i = 1, \dots, N$, *i.e.* $\pi \in S_N/\mathbf{Z}_N$. The matrices λ^a are the Chan–Paton factors describing the gauge degrees of freedom of the open string end points. Hence, they are in the adjoint representation of the gauge group under consideration. Throughout the whole article Tr denotes the group trace over the Chan–Paton factors λ^a in the adjoint representation. On the other hand, the trace tr accounts for the trace over the Lorentz–indices (μ, ν) of the gauge field strength $F_{\mu\nu}$.

In the string S -matrix the external gauge bosons are asymptotic states incoming and outgoing at $t = \pm\infty$. For any open string scattering the string world–sheet may be conformally mapped to a Riemann surface with boundaries. The external states are created through vertex operators inserted at the boundary. For open string tree–level scattering the world–sheet topology becomes a disk. The latter may be conformally mapped to the upper half plane $\text{Im}z \geq 0$ with the real axis as its boundary. Hence, all vertex positions z_i are located along the real axis, *i.e.* $z_i \in \mathbf{R}$. The gauge boson vertex operator is given by

$$V_{A^a}^{(-1)}(z, k) = \lambda^a \xi_\mu e^{-\phi(z)} \psi^\mu(z) e^{ik_\rho X^\rho(z)} \quad (2.2)$$

in the (-1) -ghost picture, and by

$$V_{A^a}^{(0)}(z, k) = \lambda^a \xi_\mu [\partial X^\mu(z) + i(k\psi) \psi^\mu(z)] e^{ik_\rho X^\rho(z)} \quad (2.3)$$

in the zero–ghost picture. Here, ξ^μ and k^μ are the polarization vector and space–time momentum of the gauge boson, respectively. The latter obey the on–shell constraints $\sum_{\mu=0}^p \xi^\mu k^\mu = 0$ and $k^2 = 0$.

The string S -matrix (2.1) becomes ($N \geq 3$):

$$\begin{aligned} & \mathcal{A}_N(k_1, \xi_1, a_1; \dots; k_N, \xi_N, a_N) \\ &= V_{\text{CKG}}^{-1} \prod_{r=1}^N \int d^2 z_r \langle V_{A^{a_1}}^{(-1)}(z_1) V_{A^{a_2}}^{(-1)}(z_2) V_{A^{a_3}}^{(0)}(z_3) \dots V_{A^{a_N}}^{(0)}(z_N) \rangle. \end{aligned} \quad (2.4)$$

Here, V_{CKG} is the volume of the conformal Killing group $PSL(2, \mathbf{R})$, which leaves the boundary ($\text{Im}(z) = 0$) of the disk fixed. Two vertex operators have to be chosen in the (-1) -ghost picture (2.2) and the remaining $N - 2$ in the zero-ghost picture, in order to guarantee a total ghost charge of -2 on the disk. This requirement is a consequence of the superdiffeomorphism invariance on the string world sheet. On the disk the conformal Killing volume V_{CKG}^{-1} is cancelled by fixing three positions and introducing the respective c -ghost correlator. We have total momentum conservation $\sum_{i=1}^N k_i = 0$ for the N external momenta k_i .

Essentially to calculate the string S -matrix \mathcal{A}_N , the main work consists of determining, for a given group ordering π , the function A^π contributing to (2.1). The latter captures the full space-time kinematics to all orders in α' . In Eq. (2.4) the contribution of a given group contraction $\text{Tr}(\lambda^{\pi(1)} \lambda^{\pi(2)} \dots \lambda^{\pi(N)})$, described by a permutation π of the N indices i , is given by specifying the integration region \mathcal{I}_π

$$\mathcal{I}_\pi = \{ \text{Im}(z_j) = 0 \mid z_{\pi(1)} < z_{\pi(2)} < \dots < z_{\pi(N)} \} \quad (2.5)$$

along the boundary of the disk. Hence we may write:

$$\begin{aligned} & \text{Tr}(\lambda^{\pi(1)} \lambda^{\pi(2)} \dots \lambda^{\pi(N)}) A^\pi \\ &= V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{j=1}^N d^2 z_j \langle V_{A^{a_1}}^{(-1)}(z_1) V_{A^{a_2}}^{(-1)}(z_2) V_{A^{a_3}}^{(0)}(z_3) \dots V_{A^{a_N}}^{(0)}(z_N) \rangle . \end{aligned} \quad (2.6)$$

For the moment, in Eq. (2.4) we have chosen the first and second gauge boson vertex operator in the (-1) -ghost picture, while the others are in the zero-ghost picture. To clearly mark this, in the following we shall write for (2.6):

$$A^\pi \equiv A^\pi(1, 2, 3, \dots, N) . \quad (2.7)$$

Later we shall relax this special choice. Hence, in the following, the argument of $A^\pi(a, b, i_1, i_2, \dots, i_{N-2})$ should indicate, that the pair (a, b) of gauge vertex operators has been chosen to be in the (-1) -ghost picture, while the others i_k are taken in the zero-ghost picture. We should point out, that this ordering is independent of the group structure and we could equally calculate all group contributions A^π by one and the same choice, *e.g.* with the first and second gauge vertex in the (-1) -ghost picture (2.7). However, as we shall see in a moment, relaxing this choice will lead to non-trivial relations between different expressions for one and the same group structure π .

The correlator in (2.4) or (2.6) may be evaluated by performing Wick contractions. In effect the correlator decomposes into products of various two–point functions. The latter are basic on the disk and given by:

$$\begin{aligned}
\langle X^\mu(z_1)X^\nu(z_2) \rangle &= -g^{\mu\nu} \ln(z_1 - z_2) , \\
\langle \psi^\mu(z_1)\psi^\nu(z_2) \rangle &= -\frac{g^{\mu\nu}}{z_1 - z_2} , \\
\langle e^{ik_1 X(z_1)} e^{ik_2 X(z_2)} \rangle &= |z_1 - z_2|^{k_1 k_2} , \\
\langle e^{-\phi(z_1)} e^{-\phi(z_2)} \rangle &= \frac{1}{z_1 - z_2} .
\end{aligned} \tag{2.8}$$

Throughout the whole article we shall work with the convention $\alpha' = \frac{1}{2}$.

2.2. Four gluon open superstring S –matrix revisited

The open string disk scattering of four gauge bosons is summarized in the string S –matrix:

$$\mathcal{A}_4(k_1, \xi_1, a_1; k_2, \xi_2, a_2; k_3, \xi_3, a_3; k_4, \xi_4, a_4) = \sum_{\substack{\pi \in \overline{(1,2,3,4)} \\ 6 \text{ permutations } \pi}} \text{Tr}(\lambda^{\pi(1)} \lambda^{\pi(2)} \lambda^{\pi(3)} \lambda^{\pi(4)}) A^\pi . \tag{2.9}$$

The sum runs over all six cyclically inequivalent permutations π of the four external gluons labeled by 1, 2, 3 and 4. For the group structure $\text{Tr}(\lambda^1 \lambda^2 \lambda^4 \lambda^3)$, *i.e.* $\pi = (1, 2, 4, 3)$, the function A^π is given by [1,2]

$$A^{(1,2,4,3)} = \frac{\Gamma(s) \Gamma(t)}{\Gamma(1 + s + t)} [tu (\xi_1 \xi_2) (\xi_3 \xi_4) + su (\xi_1 \xi_3) (\xi_2 \xi_4) + st (\xi_1 \xi_4) (\xi_2 \xi_3) + \dots] , \tag{2.10}$$

with the three Mandelstam variables $s = k_1 k_2$, $t = k_1 k_3$ and $u = k_1 k_4$, with $s + t + u = 0$. The dots stand for other kinematical contractions of the form $(\xi\xi)(\xi k)(\xi k)$. With (2.10) the full four gluon string S –matrix (2.9) may be obtained by relabeling gluon indices. Furthermore, its field–theory interpretation, which leads to the terms displayed in (1.3), is fully understood [3]. However, in this subsection we shall discuss the four gauge boson string S –matrix (2.10) from a different perspective.

In fact, rather than studying the amplitude (2.9) for a given (fixed) pair of gluon vertices in the (-1) –ghost picture, we allow for any pairings. Hence we shall focus our

attention on the more general expression (*cf.* also Eq. (2.6)):

$$\begin{aligned} \text{Tr}(\lambda^{\pi(1)} \lambda^{\pi(2)} \dots \lambda^{\pi(N)}) A^\pi(a, b, i, j) \\ = V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^4 d^2 z_r \langle V_{A^{a_a}}^{(-1)}(z_a) V_{A^{a_b}}^{(-1)}(z_b) V_{A^{a_i}}^{(0)}(z_i) V_{A^{a_j}}^{(0)}(z_j) \rangle , \end{aligned} \quad (2.11)$$

with some permutation $(a, b, i, j) \in \overline{(1, 2, 3, 4)}$ of vertex operators. According to the definition (2.11), any choice, *e.g.* $(a, b, i, j) = (1, 4, 2, 3)$ would be as good to determine the full S -matrix (2.10).

After performing the Wick contractions in (2.11), with (2.8) we obtain the following expression for (2.11):

$$\begin{aligned} A^\pi(a, b, i, j) &= A_2^\pi(a, b, i, j) (\xi_a \xi_b) (\xi_i \xi_j) \\ &+ A_1^\pi(a, b, i, j) (\xi_a \xi_i) (\xi_b \xi_j) + A_1^\pi(a, b, j, i) (\xi_a \xi_j) (\xi_b \xi_i) \\ &+ B_2^\pi(a, b, i, j) (\xi_a \xi_b) + B_3^\pi(i, j, a, b) (\xi_i \xi_j) + B_1^\pi(a, i, b, j) (\xi_a \xi_i) \\ &+ B_1^\pi(a, j, b, i) (\xi_a \xi_j) + B_1^\pi(b, i, a, j) (\xi_b \xi_i) + B_1^\pi(b, j, a, i) (\xi_b \xi_j) , \end{aligned} \quad (2.12)$$

with the following integrals:

$$\begin{aligned} A_1^\pi(a, b, i, j) &= \int_{\mathcal{I}_\pi} dz_4 \langle c(z_1) c(z_2) c(z_3) \rangle \mathcal{E} \frac{k_i k_j}{z_{ai} z_{jb} z_{ij}} \frac{(-1)}{z_{ab}} , \\ A_2^\pi(a, b, i, j) &= \int_{\mathcal{I}_\pi} dz_4 \langle c(z_1) c(z_2) c(z_3) \rangle \mathcal{E} \frac{1}{z_{ab}^2 z_{ij}^2} (1 - k_i k_j) , \\ B_1^\pi(a, i, b, j) &= \int_{\mathcal{I}_\pi} dz_4 \langle c(z_1) c(z_2) c(z_3) \rangle \mathcal{E} \frac{1}{z_{ab} z_{ij}} \\ &\times \left\{ \frac{(\xi_b k_i)(\xi_j k_a)}{z_{aj} z_{bi}} + \frac{(\xi_b k_i)(\xi_j k_b)}{z_{ai} z_{bj}} - \frac{(\xi_b k_j)(\xi_j k_i)}{z_{ai} z_{bj}} \right\} , \\ B_2^\pi(a, b, i, j) &= \int_{\mathcal{I}_\pi} dz_4 \langle c(z_1) c(z_2) c(z_3) \rangle \mathcal{E} \frac{(-1)}{z_{ab}^2 z_{ij}^2} \left\{ (\xi_i k_b)(\xi_j k_b) - (\xi_i k_b)(\xi_j k_a) \frac{z_{bj} z_{ia}}{z_{aj} z_{ib}} \right. \\ &\left. + (\xi_i k_a)(\xi_j k_a) + (\xi_i k_a)(\xi_j k_b) \frac{z_{aj} z_{ib}}{z_{bj} z_{ia}} - (\xi_i k_j)(\xi_j k_i) \right\} , \\ B_3^\pi(i, j, a, b) &= \int_{\mathcal{I}_\pi} dz_4 \langle c(z_1) c(z_2) c(z_3) \rangle \mathcal{E} \frac{1}{z_{ab} z_{ij}} \left\{ \frac{(\xi_a k_i)(\xi_b k_j)}{z_{ai} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_i)}{z_{aj} z_{bi}} \right\} , \end{aligned} \quad (2.13)$$

with the three positions z_1, z_2 and z_3 fixed due to $PSL(2, \mathbf{R})$ invariance on the disk. In addition, we have introduced $\mathcal{E} = \prod_{r < s} |z_{rs}|^{k_r k_s}$. The region \mathcal{I}_π of the z_4 -integration (2.5) is determined by the group structure π under consideration. From the sum (2.12), the

meaning of the above five functions becomes clear: While the functions A_r^π appear in front of the $(\xi\xi)(\xi\xi)$ contractions

$$A_1^\pi(a, b, i, j) (\xi_a \xi_i)(\xi_b \xi_j) \quad , \quad A_2^\pi(a, b, i, j) (\xi_a \xi_b)(\xi_i \xi_j) \quad , \quad (2.14)$$

the functions B_s^π capture the contractions $(\xi\xi)(\xi k)(\xi k)$:

$$B_1^\pi(a, i, b, j) (\xi_a \xi_i) \quad , \quad B_2^\pi(a, b, i, j) (\xi_a \xi_b) \quad , \quad B_3^\pi(i, j, a, b) (\xi_i \xi_j) \quad . \quad (2.15)$$

The crucial difference between the two functions A_2^π and A_1^π is, whether the polarizations ξ_a, ξ_b of the two vertices a and b in the (-1) -ghost picture are contracted among themselves (*i.e.* leading to the kinematics $(\xi_a \xi_b)$) or contracted with the other polarizations ξ_i, ξ_j , respectively. Similarly, the two functions B_2^π and B_1^π account for the two different cases, whether the polarization ξ_a is contracted with ξ_b or one of ξ_i, ξ_j , respectively. Moreover, the function B_3^π captures the case, when both ξ_a and ξ_b are contracted with momenta.

In each amplitude $A^\pi(a, b, i, j)$, for a given permutation $(a, b, i, j) \in \overline{(1, 2, 3, 4)}$ all possible space-time kinematics show up once. *E.g.* the kinematics $(\xi_1 \xi_2)(\xi_3 \xi_4)$, $(\xi_1 \xi_3)(\xi_2 \xi_4)$ and $(\xi_1 \xi_4)(\xi_2 \xi_3)$ show up once. However, the five functions A_r, B_s in front of those kinematics are different depending on the permutation $(a, b, i, j) \in \overline{(1, 2, 3, 4)}$ under consideration, *i.e.* which vertex pair (a, b) we have put into the (-1) -ghost picture. More concretely, for the given kinematics

$$(\xi_A \xi_B) (\xi_C \xi_D) \quad (2.16)$$

for one and the same group contraction π we obtain the three seemingly different looking expressions

$$A_1^\pi(A, C, B, D) \quad , \quad A_1^\pi(A, D, B, C) \quad , \quad A_2^\pi(A, B, C, D) \quad , \quad (2.17)$$

depending on which pair of vertex operator has been put into the (-1) ghost picture. Recall, $A_1^\pi(A, C, B, D)$ is the function with the gluon vertices A and C in the (-1) -ghost picture, while $A_1^\pi(A, D, B, C)$ with A and D and $A_2^\pi(A, B, C, D)$ with A and B singled out. In other words, the three functions A_r^π are associated to taking the pairs (A, C) , (A, D) or (A, B) of vertex operators in the (-1) -ghost picture, respectively. However, they all refer to the kinematical contraction $(\xi_A \xi_B)(\xi_C \xi_D)$ with the same group structure π . Hence, they all have to be equal as a result of just choosing different pairs of vertex operators in

the (-1) -ghost picture. In total, for the contractions $(\xi\xi)(\xi\xi)$ we obtain the following set of equations (for a given group contraction π):

$$\begin{aligned}
(\xi_1\xi_2)(\xi_3\xi_4) : \quad & A_2^\pi(1, 2, 3, 4) = A_1^\pi(1, 3, 2, 4) = A_1^\pi(1, 4, 2, 3) , \\
(\xi_1\xi_3)(\xi_2\xi_4) : \quad & A_2^\pi(1, 3, 2, 4) = A_1^\pi(1, 2, 3, 4) = A_1^\pi(1, 4, 3, 2) , \\
(\xi_1\xi_4)(\xi_2\xi_3) : \quad & A_2^\pi(1, 4, 2, 3) = A_1^\pi(1, 2, 4, 3) = A_1^\pi(1, 3, 4, 2) .
\end{aligned} \tag{2.18}$$

Let us focus on the group theoretical factor $\text{Tr}(\lambda^1\lambda^2\lambda^4\lambda^3)$, *i.e.* $\pi = (1, 2, 4, 3)$. We fix the first three vertex positions to $z_1 = -z_\infty, z_2 = 0$ and $z_3 = 1$, *i.e.* $\langle c(z_1)c(z_2)c(z_3) \rangle = -z_\infty^2$, with $z_\infty = \infty$. According to (2.5), for \mathcal{I}_π we integrate z_4 from $z_4 = 0$ until $z_4 = 1$. Then, the integrals A_r^π in (2.13) can be written for $\pi = (1, 2, 4, 3)$

$$\begin{aligned}
A_2^\pi(1, 2, 3, 4) &= (s-1) F_4 \quad , \quad A_1^\pi(1, 3, 2, 4) = -t F_3 \quad , \quad A_1^\pi(1, 4, 2, 3) = u F_1 \quad , \\
A_2^\pi(1, 3, 2, 4) &= (t-1) F_5 \quad , \quad A_1^\pi(1, 2, 3, 4) = -s F_3 \quad , \quad A_1^\pi(1, 4, 3, 2) = -u F_2 \quad , \\
A_2^\pi(1, 4, 2, 3) &= (u-1) F_0 \quad , \quad A_1^\pi(1, 2, 4, 3) = s F_1 \quad , \quad A_1^\pi(1, 3, 4, 2) = -t F_2 \quad ,
\end{aligned} \tag{2.19}$$

with the integrals:

$$F_j = \int_0^1 dx P_j x^t (1-x)^s \quad , \tag{2.20}$$

and the polynomials:

$$\begin{aligned}
P_0 &= 1 \quad , \quad P_1 = \frac{1}{x-1} \quad , \quad P_2 = \frac{1}{x} \quad , \\
P_3 &= \frac{1}{x(x-1)} \quad , \quad P_4 = \frac{1}{(x-1)^2} \quad , \quad P_5 = \frac{1}{x^2} \quad .
\end{aligned} \tag{2.21}$$

Equating the functions (2.19) according to (2.18) yields the following system of six equations for the six functions F_j :

$$\begin{aligned}
(s-1) F_4 &= u F_1 \quad , \quad (s-1) F_4 = -t F_3 \\
(t-1) F_5 &= -s F_3 \quad , \quad (t-1) F_5 = -u F_2 \\
(u-1) F_0 &= s F_1 \quad , \quad (u-1) F_0 = -t F_2 \quad .
\end{aligned} \tag{2.22}$$

Their solution may be written:

$$\begin{aligned}
F_1 &= \frac{u-1}{s} F_0 \quad , \quad F_2 = \frac{1-u}{t} F_0 \quad , \\
F_3 &= \frac{u(1-u)}{st} F_0 \quad , \quad F_4 = \frac{u(1-u)}{s(1-s)} F_0 \quad , \quad F_5 = \frac{u(1-u)}{t(1-t)} F_0 \quad .
\end{aligned} \tag{2.23}$$

Hence, all functions F_j may be expressed by one single function F_0

$$\boxed{F_j = \Lambda_j(s, t, u) F_0} \quad (2.24)$$

with some momentum dependent factor Λ_j like it is given in *Eqs.* (2.23). Of course, the integrals (2.20) are easy to determine with *Eq.* (4.1). The relations (2.23) may be directly checked using the basic identity $\Gamma(x + 1) = x \Gamma(x)$. More concretely, we obtain:

$$\begin{aligned} F_0 &= \frac{\Gamma(s + 1) \Gamma(t + 1)}{\Gamma(2 + s + t)} \quad , \quad F_1 = -\frac{\Gamma(s) \Gamma(t + 1)}{\Gamma(1 + s + t)} \quad , \quad F_2 = \frac{\Gamma(s + 1) \Gamma(t)}{\Gamma(1 + s + t)} \quad , \\ F_3 &= -\frac{\Gamma(s) \Gamma(t)}{\Gamma(s + t)} \quad , \quad F_4 = \frac{\Gamma(s - 1) \Gamma(t + 1)}{\Gamma(s + t)} \quad , \quad F_5 = \frac{\Gamma(s + 1) \Gamma(t - 1)}{\Gamma(s + t)} \quad . \end{aligned} \quad (2.25)$$

However, what this exercise shows is, that with changing the ghost–picture of the vertex operators we obtain different integrals (2.17) for the kinematics $(\xi_A \xi_B)(\xi_C \xi_D)$. These expressions have to be equal and we obtain a system of equations, which allows us, to express all integrals A_r^π , given in (2.19), in terms of a *single* integral. In the concrete case (2.23), this integral is F_0 . We have some freedom, which integral we want to choose to be left over. The integral F_3 could also have been taken. In fact, for higher gluon scattering, it proves to be very convenient to choose an integral, which does not lead to poles in the kinematic invariants. For such an integral its power series in the kinematic invariants may be easier derived.

Similar statements hold for the kinematics $(\xi\xi)(\xi k)(\xi k)$ referring to a fixed group structure π . More concretely, consider the kinematics

$$(\xi_A \xi_B) (\xi_C k)(\xi_D \tilde{k}) \quad , \quad (2.26)$$

with k, \tilde{k} accounting for a set of momenta subject to momentum conservation. In other words, the kinematics (2.26) readily stands for four independent space–time kinematics. According to (2.15) the six functions

$$\begin{aligned} B_2^\pi(A, B, C, D) \quad , \quad B_1^\pi(A, B, C, D) \quad , \quad B_1^\pi(A, B, D, C) \quad , \\ B_1^\pi(B, A, C, D) \quad , \quad B_1^\pi(B, A, D, C) \quad , \quad B_3^\pi(A, B, C, D) \end{aligned} \quad (2.27)$$

appear in (2.12) and are relevant to the kinematics (2.26). They refer to the six cases, in which one of the pairs $(A, B), (A, C), (A, D), (B, C), (B, D)$ or (C, D) of vertex operators

is put into the (-1) -ghost picture, respectively. Similar, as in *Eq.* (2.18) all these six functions (2.27) have to be equated. Since, after definition, the latter still contain some ξk kinematical contractions, what has to be equated, is the relevant piece of those functions referring to the same scalar product $(\xi k)(\xi k)$, after applying momentum conservation. Hence, in what follows⁴ one obtains actually twenty equations, which give rise to relations between different integrals of the type (2.20). After considering all possible kinematics, in addition to the polynomials (2.21), the following set

$$\begin{aligned} P_6 &= \frac{1}{x(x-1)^2} \quad , \quad P_7 = \frac{x}{(x-1)^2} \quad , \quad P_8 = \frac{1}{x^2(x-1)} \\ P_9 &= \frac{x-1}{x} \quad , \quad P_{10} = \frac{x-1}{x^2} \quad , \quad P_{11} = \frac{x}{x-1} \end{aligned} \quad (2.28)$$

appears. The equations following from equating (2.27) for all possible kinematics provide the relations between the integrals (2.20) associated to (2.28) and F_0 :

$$\begin{aligned} F_6 &= F_4 - F_3 \quad , \quad F_7 = F_1 + F_4 \quad , \quad F_8 = F_3 - F_5 \quad , \\ F_9 &= F_0 - F_2 \quad , \quad F_{10} = F_2 - F_5 \quad , \quad F_{11} = F_0 + F_2 + F_3 \quad . \end{aligned} \quad (2.29)$$

In addition, the identity $F_1 - F_2 = F_3$ appears. Of course, the latter relation, which is satisfied by the solutions (2.25) is just a consequence of $P_1 - P_2 = P_3$, *i.e.* $\frac{1}{x-1} - \frac{1}{x} = \frac{1}{x(x-1)}$.

Though, the identities (2.23) are easy to prove in the case at hand, the strategy, outlined above for the four-gluon scattering, yields non-trivial identities in the case of gluon amplitudes involving more than four gluons. From *Eq.* (2.23) we deduce, that those identities involve rational polynomials in the Mandelstam variables s, t and u . These polynomials account for the pole structure of the given kinematics, while the function through which all other functions are expressed may be chosen to be simple in the sense, that it does not contain poles. *E.g.* in the case above, the function F_0 has the power series w.r.t. to the Mandelstam variables s, t, u

$$F_0 = 1 - s - t + s^2 + 2 s t - \zeta(2) s t + t^2 + \dots \quad , \quad (2.30)$$

i.e. it is finite w.r.t. to all invariants. Hence all pole structure of the amplitude is encountered in the rational polynomials in front of the function F_0 , *i.e.* they follow from solving the algebraic equations (2.22) through expressing all functions $F_j, j \geq 1$ through F_0 .

⁴ More concretely, *e.g.* in the case $A = 1, B = 2, C = 3$ and $D = 4$ one reduces the functions to the kinematics $(\xi_3 k_1) (\xi_4 k_2)$ and $(\xi_3 k_2) (\xi_4 k_1)$ [and $(\xi_3 k_1) (\xi_4 k_1)$ and $(\xi_3 k_2) (\xi_4 k_2)$] by eliminating the combinations $\xi_3 k_4$ and $\xi_4 k_3$ as a result of momentum conservation.

In the case of more than four gluons, those identities allow to express each integral accounting for a specific kinematics through a linear combination of integrals without poles, such that the full pole structure originates algebraically. In fact, the case with six gluons shall be discussed in the next section, while the five gluon case is revisited in section 6.

3. Six gluon open superstring S -matrix

In the following we study the string S -matrix describing the string tree-level scattering of six gauge bosons on the disk ($N = 6$ in (2.4)):

$$\begin{aligned} \mathcal{A}_6(k_1, \xi_1, a_1; k_2, \xi_2, a_2; k_3, \xi_3, a_3; k_4, \xi_4, a_4; k_5, \xi_5, a_5; k_6, \xi_6, a_6) \\ = V_{\text{CKG}}^{-1} \prod_{r=1}^6 \int d^2 z_r \langle V_{A^{a_1}}^{(-1)}(z_1) V_{A^{a_2}}^{(-1)}(z_2) V_{A^{a_3}}^{(0)}(z_3) V_{A^{a_4}}^{(0)}(z_4) V_{A^{a_5}}^{(0)}(z_5) V_{A^{a_6}}^{(0)}(z_6) \rangle . \end{aligned} \quad (3.1)$$

We have total momentum conservation:

$$\sum_{i=1}^6 k_i = 0 \quad (3.2)$$

for the six external gluon momenta k_i . Furthermore, we have chosen two gauge vertex operators in the (-1) -ghost picture in order to guarantee a total ghost charge of -2 on the disk. The basic ingredients to start with have been presented in subsection 2.1. However, one has to keep in mind, that the case $p = 3$ is special, since in that case the six order Lorentz invariant $\text{tr}F^6$ may be expressed by the other two $(\text{tr}F^2)^3$ and $\text{tr}F^2\text{tr}F^4$ invariants. In the scattering process of six gluons this difference manifests in a reduced number of kinematical (space-time) invariants. Later we shall comment on this issue.

Recall, that the four-gluon string S -matrix \mathcal{A}_4 , discussed in the previous subsection, gives rise to the following two kinematical structures $(\xi\xi)(\xi\xi)$ and $(\xi\xi)(\xi k)(\xi k)$ entering (2.10). At order $\mathcal{O}(k^4)$ in the external momenta, both kinematics comprise parts of the same higher order gauge interaction terms F^4 in the effective action as a result of gauge invariance. This means, that these interaction terms may be completely inferred from the first contractions $(\xi\xi)(\xi\xi)$ [3], while the string S -matrix result for the second kinematics $(\xi\xi)(\xi k)(\xi k)$ may be related to the first one. The explicit evaluation of the correlator in Eq. (3.1) leads to a sum of integrals with various different (space-time) kinematic contractions as coefficients. More precisely, we encounter the three kinematical structures $(\xi\xi)(\xi\xi)(\xi\xi)$, $(\xi\xi)(\xi\xi)(\xi k)(\xi k)$ and $(\xi\xi)(\xi k)(\xi k)(\xi k)(\xi k)$, which in the following will be denoted by A , B and C -kinematics, respectively. Their contributions in the effective action are all related to each other through gauge-invariance. In other words, the effective action at a given order in the external momenta, may be reproduced from a particular class of those kinematics.

3.1. General expression for the amplitude

In Eq. (3.1) we have chosen the first two gauge vertex operators in the (-1) -ghost picture. We have already stressed in the previous subsection, that we are free to choose which pair (a, b) of the six gauge vertex operators we put into the (-1) -ghost picture. Hence, according to the definition (2.7), for a given group structure described by the permutation π , instead of the particular choice in (3.1), we may generically consider the expression

$$\begin{aligned} \text{Tr}(\lambda^{\pi(1)} \lambda^{\pi(2)} \lambda^{\pi(3)} \lambda^{\pi(4)} \lambda^{\pi(5)} \lambda^{\pi(6)}) A^\pi(a, b, i, j, k, l) &= V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^2 z_r \\ &\langle V_{A^{a_a}}^{(-1)}(z_a) V_{A^{a_b}}^{(-1)}(z_b) V_{A^{a_i}}^{(0)}(z_i) V_{A^{a_j}}^{(0)}(z_j) V_{A^{a_k}}^{(0)}(z_k) V_{A^{a_l}}^{(0)}(z_l) \rangle, \end{aligned} \quad (3.3)$$

with some permutation $(a, b, i, j, k, l) \in \overline{(1, 2, 3, 4, 5, 6)}$ of vertex operators. In Eq. (2.1), we have explained, how this term contributes to the full string S -matrix (3.1).

After performing the Wick contractions the correlator in (3.3) decomposes into products of various two-point functions, given in Eq. (2.8):

$$\begin{aligned} A^\pi(a, b, i, j, k, l) &= V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^2 z_r \sum_{\substack{(i_1, i_2, i_3, i_4) \\ \in (i, j, k, l)}} \left\{ \frac{1}{2} \mathcal{A}_1(a, b, i_1, i_2, i_3, i_4) (\xi_a \xi_{i_1}) (\xi_b \xi_{i_4}) (\xi_{i_2} \xi_{i_3}) \right. \\ &+ \frac{1}{8} \mathcal{A}_2(a, b, i_1, i_2, i_3, i_4) (\xi_a \xi_b) (\xi_{i_1} \xi_{i_2}) (\xi_{i_3} \xi_{i_4}) + \frac{1}{4} \mathcal{B}_2(a, b, i_1, i_2, i_3, i_4) (\xi_a \xi_b) (\xi_{i_1} \xi_{i_2}) \\ &+ \frac{1}{2} \mathcal{B}_1(a, b, i_1, i_2, i_3, i_4) (\xi_a \xi_{i_1}) (\xi_b \xi_{i_2}) + \frac{1}{8} \mathcal{B}_4(i_1, i_2, i_3, i_4, a, b) (\xi_{i_1} \xi_{i_2}) (\xi_{i_3} \xi_{i_4}) \\ &+ \frac{1}{2} \mathcal{B}_3(a, i_1, i_2, i_3, b, i_4) (\xi_a \xi_{i_1}) (\xi_{i_2} \xi_{i_3}) + \frac{1}{2} \mathcal{B}_3(b, i_1, i_2, i_3, a, i_4) (\xi_b \xi_{i_1}) (\xi_{i_2} \xi_{i_3}) \\ &+ \frac{1}{24} \mathcal{C}_1(a, b, i_1, i_2, i_3, i_4) (\xi_a \xi_b) + \frac{1}{4} \mathcal{C}_3(i_1, i_2, a, b, i_3, i_4) (\xi_{i_1} \xi_{i_2}) \\ &\left. + \frac{1}{6} \mathcal{C}_2(a, i_1, b, i_2, i_3, i_4) (\xi_a \xi_{i_1}) + \frac{1}{6} \mathcal{C}_2(b, i_1, a, i_2, i_3, i_4) (\xi_b \xi_{i_1}) \right\}. \end{aligned} \quad (3.4)$$

In Eq. (3.4) in front of the contractions $(\xi_a \xi_i) (\xi_j \xi_k) (\xi_l \xi_b)$ and $(\xi_a \xi_b) (\xi_i \xi_j) (\xi_k \xi_l)$ there appear the two basic functions $\mathcal{A}_1(a, b, i, j, k, l)$ and $\mathcal{A}_2(a, b, i, j, k, l)$, given by

$$\begin{aligned} \mathcal{A}_1(a, b, i, j, k, l) &= -\frac{\mathcal{E}}{z_{ab} z_{ai}} \left\{ \frac{(k_i k_j) (k_k k_l)}{z_{lb} z_{jk} z_{ij} z_{kl}} - \frac{(k_i k_k) (k_j k_l)}{z_{lb} z_{jk} z_{ik} z_{jl}} - \frac{(k_i k_l) (1 - k_j k_k)}{z_{lb} z_{il} z_{jk}^2} \right\}, \\ \mathcal{A}_2(a, b, i, j, k, l) &= -\frac{\mathcal{E}}{z_{ab}^2} \left\{ \frac{(1 - k_i k_j) (1 - k_k k_l)}{z_{ij}^2 z_{kl}^2} - \frac{(k_i k_k) (k_j k_l)}{z_{ij} z_{kl} z_{ik} z_{jl}} + \frac{(k_i k_l) (k_j k_k)}{z_{ij} z_{kl} z_{il} z_{jk}} \right\}, \end{aligned} \quad (3.5)$$

respectively, with:

$$\mathcal{E} = \prod_{r < s}^6 |z_{rs}|^{k_r k_s} . \quad (3.6)$$

In the following, we in particular shall need the integrated expressions

$$A_i^\pi = V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^6 z_r \mathcal{A}_i , \quad B_i^\pi = V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^6 z_r \mathcal{B}_i , \quad C_i^\pi = V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^6 z_r \mathcal{C}_i , \quad (3.7)$$

which enter the expression (3.4). Furthermore we have:

$$\begin{aligned} \mathcal{B}_1(a, b, i, j, k, l) = & -\mathcal{E} \left\{ \frac{z_{ai}(\xi_k k_i)(\xi_l k_i)}{z_{ab}^2 z_{ij} z_{ik} z_{il}} \left(\frac{z_{ai}}{z_{ak} z_{al} z_{ij}} - \frac{z_{ai} k_i k_j}{z_{ak} z_{al} z_{ij}} + \frac{k_j k_k}{z_{al} z_{jk}} + \frac{k_j k_l}{z_{ak} z_{jl}} \right) \right. \\ & - \frac{(\xi_k k_l)(\xi_l k_j)}{z_{ab}^2 z_{ij} z_{jl} z_{kl}} \left(\frac{z_{aj}(1 - k_i k_j)}{z_{ak} z_{ij}} - \frac{k_i k_k}{z_{ik}} - \frac{z_{al} k_i k_l}{z_{ak} z_{il}} \right) + \frac{(\xi_k k_i)(\xi_l k_b)}{z_{ab} z_{al} z_{bl} z_{ij} z_{ik}} \left(\frac{z_{ai}(1 - k_i k_j)}{z_{ak} z_{ij}} + \frac{k_j k_k}{z_{jk}} \right) \\ & + \frac{(\xi_k k_i)(\xi_l k_k)}{z_{ab}^2 z_{ij} z_{ik} z_{kl}} \left(\frac{z_{ai}(1 - k_i k_j)}{z_{al} z_{ij}} + \frac{z_{ak} k_j k_k}{z_{al} z_{jk}} + \frac{k_j k_l}{z_{jl}} \right) + \frac{(\xi_k k_b)(\xi_l k_j)}{z_{ab} z_{ak} z_{bk} z_{ij} z_{jl}} \left(\frac{z_{aj}(1 - k_i k_j)}{z_{al} z_{ij}} - \frac{k_i k_l}{z_{il}} \right) \\ & + \frac{(\xi_k k_j)(\xi_l k_i)}{z_{ab}^2 z_{ij} z_{il} z_{jk}} \left(\frac{z_{ai} z_{aj}(1 - k_i k_j)}{z_{ak} z_{al} z_{ij}} - \frac{z_{ai} k_i k_k}{z_{al} z_{ik}} + \frac{z_{aj} k_j k_l}{z_{ak} z_{jk}} + \frac{k_k k_l}{z_{kl}} \right) + \frac{(\xi_k k_b)(\xi_l k_k)(1 - k_i k_j)}{z_{ab} z_{al} z_{bk} z_{ij}^2 z_{kl}} \\ & + \frac{(\xi_k k_j)(\xi_l k_b)}{z_{ab} z_{al} z_{bl} z_{ij} z_{jk}} \left(\frac{z_{aj}(1 - k_i k_j)}{z_{ak} z_{ij}} - \frac{k_i k_k}{z_{ik}} \right) + \frac{(\xi_k k_b)(\xi_l k_i)}{z_{ab} z_{ak} z_{bk} z_{ij} z_{il}} \left(\frac{z_{ai}(1 - k_i k_j)}{z_{al} z_{ij}} + \frac{k_j k_l}{z_{jl}} \right) \\ & + \frac{z_{aj}(\xi_k k_j)(\xi_l k_j)}{z_{ab}^2 z_{ij} z_{jk} z_{jl}} \left(\frac{z_{aj}(1 - k_i k_j)}{z_{ak} z_{al} z_{ij}} - \frac{k_i k_k}{z_{al} z_{ik}} + \frac{k_i k_l}{z_{ak} z_{il}} \right) + \frac{(\xi_k k_b)(\xi_l k_b)(1 - k_i k_j)}{z_{ak} z_{al} z_{bk} z_{bl} z_{ij}^2} \\ & + \frac{(\xi_k k_j)(\xi_l k_k)}{z_{ab}^2 z_{ij} z_{jk} z_{kl}} \left(\frac{z_{aj}(1 - k_i k_j)}{z_{al} z_{ij}} - \frac{z_{ak} k_i k_k}{z_{al} z_{ik}} - \frac{k_i k_l}{z_{il}} \right) - \frac{(\xi_k k_l)(\xi_l k_b)(1 - k_i k_j)}{z_{ab} z_{ak} z_{bl} z_{ij}^2 z_{kl}} \\ & + \frac{(\xi_k k_i)(\xi_l k_j)}{z_{ab}^2 z_{ij} z_{ik} z_{jl}} \left(\frac{z_{ai} z_{aj}(1 - k_i k_j)}{z_{ak} z_{al} z_{ij}} - \frac{z_{ai} k_i k_l}{z_{ak} z_{il}} + \frac{z_{aj} k_j k_k}{z_{al} z_{jk}} - \frac{k_k k_l}{z_{kl}} \right) \\ & \left. - \frac{(\xi_k k_l)(\xi_l k_i)}{z_{ab}^2 z_{ij} z_{il} z_{kl}} \left(\frac{z_{ai}(1 - k_i k_j)}{z_{ak} z_{ij}} + \frac{k_j k_k}{z_{jk}} + \frac{z_{al} k_j k_l}{z_{ak} z_{jl}} \right) \right\} , \\ \mathcal{B}_2(a, b, i, j, k, l) = & -\mathcal{E} \left\{ \frac{(\xi_k k_j)(\xi_l k_i)}{z_{ab} z_{bj} z_{il} z_{jk}} \left(\frac{z_{aj} k_i k_j}{z_{ak} z_{al} z_{ij}} + \frac{k_i k_k}{z_{al} z_{ik}} - \frac{z_{aj} k_j k_l}{z_{ai} z_{ak} z_{jl}} - \frac{k_k k_l}{z_{ai} z_{kl}} \right) \right. \\ & + \frac{(\xi_k k_j)(\xi_l k_b)}{z_{ai} z_{al} z_{bj} z_{bl} z_{jk}} \left(\frac{z_{aj} k_i k_j}{z_{ak} z_{ij}} + \frac{k_i k_k}{z_{ik}} \right) + \frac{(\xi_k k_l)(\xi_l k_i)}{z_{ab} z_{bj} z_{il} z_{kl}} \left(\frac{z_{al} k_j k_l}{z_{ai} z_{ak} z_{jl}} + \frac{k_j k_k}{z_{ai} z_{jk}} - \frac{k_i k_j}{z_{ak} z_{ij}} \right) \\ & + \frac{(\xi_k k_b)(\xi_l k_j)}{z_{ai} z_{ak} z_{bj} z_{bk} z_{jl}} \left(\frac{z_{aj} k_i k_j}{z_{al} z_{ij}} - \frac{k_i k_l}{z_{il}} \right) + \frac{(\xi_k k_i)(\xi_l k_i)}{z_{ab} z_{bj} z_{ik} z_{il}} \left(\frac{z_{ai} k_i k_j}{z_{ak} z_{al} z_{ij}} - \frac{k_j k_k}{z_{al} z_{jk}} - \frac{k_j k_l}{z_{ak} z_{jl}} \right) \\ & \left. + \frac{(\xi_k k_b)(\xi_l k_i)}{z_{ak} z_{bj} z_{bk} z_{il}} \left(\frac{k_i k_j}{z_{al} z_{ij}} - \frac{k_j k_l}{z_{ai} z_{jl}} \right) + \frac{(\xi_k k_j)(\xi_l k_j)}{z_{ab} z_{ai} z_{bj} z_{jk} z_{jl}} \left(\frac{z_{aj}^2 k_i k_j}{z_{ak} z_{al} z_{ij}} + \frac{z_{aj} k_i k_k}{z_{al} z_{ik}} + \frac{z_{aj} k_i k_l}{z_{ak} z_{il}} \right) \right\} \end{aligned}$$

$$\begin{aligned}
& -\frac{(\xi_k k_l)(\xi_l k_j)}{z_{ab} z_{ai} z_{bj} z_{jl} z_{kl}} \left(\frac{z_{aj} k_i k_j}{z_{ak} z_{ij}} + \frac{k_i k_k}{z_{ik}} + \frac{z_{al} k_i k_l}{z_{ak} z_{il}} \right) + \frac{(\xi_k k_i)(\xi_l k_b)}{z_{al} z_{bj} z_{bl} z_{ik}} \left(\frac{k_i k_j}{z_{ak} z_{ij}} - \frac{k_j k_k}{z_{ai} z_{jk}} \right) \\
& + \frac{(\xi_k k_i)(\xi_l k_j)}{z_{ab} z_{bj} z_{ik} z_{jl}} \left(\frac{z_{aj} k_i k_j}{z_{ak} z_{al} z_{ij}} + \frac{k_i k_l}{z_{ak} z_{il}} - \frac{z_{aj} k_j k_k}{z_{ai} z_{al} z_{jk}} + \frac{k_k k_l}{z_{ai} z_{kl}} \right) \frac{(\xi_k k_b)(\xi_l k_k)}{z_{ai} z_{al} z_{bj}} \frac{k_i k_j}{z_{bk} z_{ij} z_{kl}} \\
& + \frac{(\xi_k k_i)(\xi_l k_k)}{z_{ab} z_{bj} z_{ik} z_{kl}} \left(\frac{k_i k_j}{z_{al} z_{ij}} - \frac{z_{ak} k_j k_k}{z_{ai} z_{al} z_{jk}} - \frac{k_j k_l}{z_{ai} z_{jl}} \right) + \frac{(\xi_k k_b)(\xi_l k_b)}{z_{ai} z_{ak} z_{al}} \frac{z_{ab} k_i k_j}{z_{bj} z_{bk} z_{bl} z_{ij}} \\
& + \frac{(\xi_k k_j)(\xi_l k_k)}{z_{ab} z_{ai} z_{bj} z_{jk} z_{kl}} \left(\frac{z_{aj} k_i k_j}{z_{al} z_{ij}} + \frac{z_{ak} k_i k_k}{z_{al} z_{ik}} + \frac{k_i k_l}{z_{il}} \right) + \frac{(\xi_k k_l)(\xi_l k_b)}{z_{ai} z_{ak} z_{bj}} \frac{k_i k_j}{z_{bl} z_{ij} z_{kl}} \Big\} , \\
\mathcal{B}_3(a, i, j, k, b, l) = & -\mathcal{E} \left\{ \frac{(k_j k_k - 1)}{z_{jk}^2} \left(\frac{(\xi_b k_i)(\xi_l k_b)}{z_{ai} z_{al} z_{bi} z_{bl}} + \frac{(\xi_b k_i)(\xi_l k_i)}{z_{ab} z_{al} z_{bi} z_{il}} + \frac{(\xi_b k_l)(\xi_l k_i)}{z_{ab} z_{ai} z_{bl} z_{il}} \right) \right. \\
& + \frac{(\xi_b k_i)(\xi_l k_j)}{z_{ab} z_{ai} z_{jk} z_{jl}} \left(\frac{(z_{aj})(k_j k_k - 1)}{z_{al} z_{bi} z_{jk}} + \frac{k_k k_l}{z_{bi} z_{kl}} \right) + \frac{(\xi_b k_k)(\xi_l k_i)}{z_{ab} z_{bk} z_{il} z_{jk}} \left(\frac{k_i k_j}{z_{al} z_{ij}} - \frac{k_j k_l}{z_{ai} z_{jl}} \right) \\
& + \frac{(\xi_b k_j)(\xi_l k_i)}{z_{ab} z_{bj} z_{il} z_{jk}} \left(\frac{k_k k_l}{z_{ai} z_{kl}} - \frac{k_i k_k}{z_{al} z_{ik}} \right) + \frac{(\xi_b k_i)(\xi_l k_k)}{z_{ab} z_{ai} z_{bi} z_{kl}} \left(\frac{(z_{ak})(k_j k_k - 1)}{z_{al} z_{jk}} + \frac{k_j k_l}{z_{jl}} \right) \\
& + \frac{k_i k_j}{z_{ai} z_{ij} z_{jk}} \left(\frac{(\xi_b k_k)(\xi_l k_b)}{z_{al} z_{bk} z_{bl}} + \frac{z_{ak}(\xi_b k_k)(\xi_l k_k)}{z_{ab} z_{al} z_{bk} z_{kl}} + \frac{(\xi_b k_l)(\xi_l k_k)}{z_{ab} z_{bl} z_{kl}} \right) \\
& - \frac{k_i k_k}{z_{ai} z_{ik} z_{jk}} \left(\frac{(\xi_b k_j)(\xi_l k_b)}{z_{al} z_{bj} z_{bl}} + \frac{z_{aj}(\xi_b k_j)(\xi_l k_j)}{z_{ab} z_{al} z_{bj} z_{jl}} + \frac{(\xi_b k_l)(\xi_l k_j)}{z_{ab} z_{bl} z_{jl}} \right) \\
& \left. + \frac{(\xi_b k_k)(\xi_l k_j)}{z_{ab} z_{ai} z_{bk} z_{jk} z_{jl}} \left(\frac{z_{aj} k_i k_j}{z_{al} z_{ij}} + \frac{k_i k_l}{z_{il}} \right) - \frac{(\xi_b k_j)(\xi_l k_k)}{z_{ab} z_{ai} z_{bj} z_{jk} z_{kl}} \left(\frac{z_{ak} k_i k_k}{z_{al} z_{ik}} + \frac{k_i k_l}{z_{il}} \right) \right\} , \\
\mathcal{B}_4(i, j, k, l, a, b) = & -\frac{\mathcal{E}}{z_{ab} z_{ij} z_{kl}} \left\{ \frac{(1 - k_k k_l)}{z_{kl}} \left(\frac{(\xi_a k_i)(\xi_b k_j)}{z_{ai} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_i)}{z_{aj} z_{bi}} \right) \right. \\
& + \frac{k_i k_k}{z_{ik}} \left(\frac{(\xi_a k_j)(\xi_b k_l)}{z_{aj} z_{bl}} - \frac{(\xi_a k_l)(\xi_b k_j)}{z_{al} z_{bj}} \right) + \frac{k_i k_l}{z_{il}} \left(\frac{(\xi_a k_k)(\xi_b k_j)}{z_{ak} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_k)}{z_{aj} z_{bk}} \right) \\
& + \frac{k_j k_k}{z_{jk}} \left(\frac{(\xi_a k_l)(\xi_b k_i)}{z_{al} z_{bi}} - \frac{(\xi_a k_i)(\xi_b k_l)}{z_{ai} z_{bl}} \right) + \frac{k_j k_l}{z_{jl}} \left(\frac{(\xi_a k_i)(\xi_b k_k)}{z_{ai} z_{bk}} - \frac{(\xi_a k_k)(\xi_b k_i)}{z_{ak} z_{bi}} \right) \\
& \left. + \frac{(1 - k_i k_j)}{z_{ij}} \left(\frac{(\xi_a k_k)(\xi_b k_l)}{z_{ak} z_{bl}} - \frac{(\xi_a k_l)(\xi_b k_l)}{z_{al} z_{bk}} \right) \right\} . \tag{3.8}
\end{aligned}$$

Besides, the three functions:

$$\begin{aligned}
\mathcal{C}_1(a, b, i, j, k, l) = & -\mathcal{E} \frac{(\xi_j k_b)(\xi_k k_b)(\xi_l k_b) z_{ab}}{z_{ai} z_{bj} z_{bk} z_{bl}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{ak} z_{al} z_{bi}} - \frac{(\xi_i k_j)}{z_{ak} z_{al} z_{ij}} - \frac{(\xi_j k_k)}{z_{aj} z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{aj} z_{ak} z_{il}} \right) \\
& + \frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_b)}{z_{aj} z_{bk} z_{bl} z_{ij}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{ak} z_{al} z_{bi}} - \frac{(\xi_i k_k)}{z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{ak} z_{il}} \right) + \frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_b)}{z_{ak} z_{bj} z_{bl} z_{ik}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{al} z_{bi}} - \frac{(\xi_i k_j)}{z_{al} z_{ij}} - \frac{(\xi_i k_l)}{z_{aj} z_{il}} \right) \\
& + \frac{(\xi_j k_k)(\xi_k k_b)(\xi_l k_b)}{z_{ai} z_{bk} z_{bl} z_{jk}} \left(\frac{(\xi_i k_j)}{z_{al} z_{ij}} - \frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{al} z_{bi}} + \frac{(\xi_i k_k) z_{ak}}{z_{aj} z_{al} z_{ik}} + \frac{(\xi_i k_l)}{z_{aj} z_{il}} \right) + \frac{(\xi_j k_i)(\xi_k k_i)(\xi_l k_b) z_{ai}}{z_{aj} z_{ak} z_{bl} z_{ij} z_{ik}} \left(\frac{(\xi_i k_b)}{z_{al} z_{bi}} - \frac{(\xi_i k_l)}{z_{ab} z_{il}} \right) \\
& + \frac{(\xi_j k_l)(\xi_k k_b)(\xi_l k_b)}{z_{ai} z_{bk} z_{bl} z_{jl}} \left(\frac{(\xi_i k_j)}{z_{ak} z_{ij}} - \frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{ak} z_{bi}} + \frac{(\xi_i k_k)}{z_{aj} z_{ik}} + \frac{(\xi_i k_l) z_{al}}{z_{aj} z_{ak}} \right) + \frac{(\xi_j k_k)(\xi_k k_i)(\xi_l k_b)}{z_{aj} z_{bl} z_{ik} z_{jk}} \left(\frac{(\xi_i k_l)}{z_{ab} z_{il}} - \frac{(\xi_i k_b)}{z_{al} z_{bi}} \right) \\
& + \frac{(\xi_j k_l)(\xi_k k_i)(\xi_l k_b)}{z_{ak} z_{bl} z_{ik} z_{jl}} \left(\frac{(\xi_i k_j)}{z_{ab} z_{ij}} - \frac{(\xi_i k_b)}{z_{aj} z_{bi}} + \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{aj} z_{il}} \right) + \frac{(\xi_j k_b)(\xi_k k_j)(\xi_l k_b)}{z_{ai} z_{bj} z_{bl} z_{jk}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{ak} z_{al} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ak} z_{al} z_{ij}} - \frac{(\xi_i k_k)}{z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{ak} z_{il}} \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{(\xi_j k_l)(\xi_k k_j)(\xi_l k_b)}{z_{ai} z_{bl} z_{jk} z_{jl}} \left(\frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{ak} z_{ij}} - \frac{(\xi_i k_b)}{z_{ak} z_{bi}} + \frac{(\xi_i k_k)}{z_{ab} z_{ik}} + \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{ak} z_{il}} \right) + \frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_i) z_{ai}}{z_{aj} z_{al} z_{bk} z_{ij} z_{il}} \left(\frac{(\xi_i k_b)}{z_{ak} z_{bi}} - \frac{(\xi_i k_k)}{z_{ab} z_{ik}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_l)(\xi_l k_b)}{z_{ai} z_{bj} z_{bl} z_{kl}} \left(\frac{(\xi_i k_j)}{z_{ak} z_{ij}} - \frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{ak} z_{bi}} + \frac{(\xi_i k_k)}{z_{aj} z_{ik}} + \frac{(\xi_i k_l) z_{al}}{z_{aj} z_{ak} z_{il}} \right) + \frac{(\xi_j k_i)(\xi_k k_l)(\xi_l k_b)}{z_{aj} z_{bl} z_{ij} z_{kl}} \left(\frac{(\xi_i k_k)}{z_{ab} z_{ik}} - \frac{(\xi_i k_b)}{z_{ak} z_{bi}} + \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{ak} z_{il}} \right) \\
& + \frac{(\xi_j k_k)(\xi_k k_l)(\xi_l k_b)}{z_{ai} z_{bl} z_{jk} z_{kl}} \left(\frac{(\xi_i k_b)}{z_{aj} z_{bi}} - \frac{(\xi_i k_j)}{z_{ab} z_{ij}} - \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{aj} z_{ik}} - \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{aj} z_{il}} \right) + \frac{(\xi_j k_b)(\xi_k k_b)(\xi_l k_i)}{z_{al} z_{bj} z_{bk} z_{il}} \left(\frac{(\xi_i k_b)}{z_{aj} z_{ak} z_{bi}} - \frac{(\xi_i k_j)}{z_{ak} z_{ij}} - \frac{(\xi_i k_k)}{z_{aj} z_{ik}} \right) \\
& + \frac{(\xi_j k_l)(\xi_k k_l)(\xi_l k_b) z_{al}}{z_{aj} z_{bl} z_{jl} z_{kl}} \left(\frac{(\xi_i k_b)}{z_{aj} z_{ak} z_{bi}} - \frac{(\xi_i k_j)}{z_{ab} z_{ak} z_{ij}} - \frac{(\xi_i k_k)}{z_{ab} z_{aj} z_{ik}} - \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{aj} z_{ak} z_{il}} \right) + \frac{(\xi_j k_i)(\xi_k k_j)(\xi_l k_b)}{z_{ak} z_{bl} z_{ij} z_{jk}} \left(\frac{(\xi_i k_b)}{z_{al} z_{bi}} - \frac{(\xi_i k_l)}{z_{ab} z_{il}} \right) \\
& + \frac{(\xi_j k_k)(\xi_k k_b)(\xi_l k_i)}{z_{al} z_{bk} z_{il} z_{jk}} \left(\frac{(\xi_i k_j)}{z_{ab} z_{ij}} - \frac{(\xi_i k_b)}{z_{aj} z_{bi}} + \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{aj} z_{ik}} \right) + \frac{(\xi_j k_l)(\xi_k k_b)(\xi_l k_i)}{z_{aj} z_{bk} z_{il} z_{jl}} \left(\frac{(\xi_i k_k)}{z_{ab} z_{ik}} - \frac{(\xi_i k_b)}{z_{ak} z_{bi}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_i) z_{ai}}{z_{ak} z_{al} z_{bj} z_{ik} z_{il}} \left(\frac{(\xi_i k_b)}{z_{aj} z_{bi}} - \frac{(\xi_i k_j)}{z_{ab} z_{ij}} \right) + \frac{(\xi_i k_b)(\xi_j k_i)(\xi_k k_i)(\xi_l k_i) z_{ai}^2}{z_{ab} z_{aj} z_{ak} z_{al} z_{bi} z_{ij} z_{ik} z_{il}} - \frac{(\xi_k k_i)(\xi_l k_i) z_{ai}}{z_{ab} z_{aj} z_{bi} z_{ik} z_{il}} \left(\frac{(\xi_i k_b)(\xi_j k_k)}{z_{al} z_{jk}} + \frac{(\xi_i k_b)(\xi_j k_l)}{z_{ak} z_{jl}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_j)(\xi_l k_i)}{z_{al} z_{bj} z_{il} z_{jk}} \left(\frac{(\xi_i k_b)}{z_{al} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{ak} z_{ij}} - \frac{(\xi_i k_k)}{z_{ab} z_{ik}} \right) + \frac{(\xi_k k_j)(\xi_l k_i)}{z_{ab} z_{ak} z_{bi} z_{il} z_{jk}} \left(\frac{(\xi_i k_b)(\xi_j k_i) z_{ai}}{z_{al} z_{ij}} - \frac{(\xi_i k_b)(\xi_j k_l)}{z_{jl}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_l)(\xi_l k_i)}{z_{ak} z_{bj} z_{il} z_{kl}} \left(\frac{(\xi_i k_j)}{z_{ab} z_{ij}} - \frac{(\xi_i k_b)}{z_{aj} z_{bi}} \right) + \frac{(\xi_k k_l)(\xi_l k_i)}{z_{ab} z_{aj} z_{bi} z_{il} z_{kl}} \left(\frac{(\xi_i k_b)(\xi_j k_k)}{z_{jk}} - \frac{(\xi_i k_b)(\xi_j k_i) z_{ai}}{z_{ak} z_{ij}} + \frac{(\xi_i k_b)(\xi_j k_l) z_{al}}{z_{ak} z_{jl}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_b)(\xi_l k_j)}{z_{aj} z_{bj} z_{bk} z_{jl}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{ak} z_{al} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ak} z_{al}} - \frac{(\xi_i k_k)}{z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{ak} z_{il}} \right) + \frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_j)}{z_{al} z_{bk} z_{ij} z_{jl}} \left(\frac{(\xi_i k_b)}{z_{ak} z_{bi}} - \frac{(\xi_i k_k)}{z_{ab} z_{ik}} \right) \\
& + \frac{(\xi_j k_k)(\xi_k k_b)(\xi_l k_j)}{z_{ai} z_{bk} z_{jk} z_{jl}} \left(\frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{al} z_{ij}} - \frac{(\xi_i k_b)}{z_{al} z_{bi}} + \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{al} z_{ik}} + \frac{(\xi_i k_l)}{z_{ab} z_{il}} \right) + \frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_j)}{z_{ak} z_{bj} z_{ik} z_{jl}} \left(\frac{(\xi_i k_b)}{z_{al} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{al} z_{ij}} - \frac{(\xi_i k_l)}{z_{ab} z_{il}} \right) \\
& + \frac{(\xi_k k_i)(\xi_l k_j)}{z_{ab} z_{al} z_{bi} z_{ik} z_{jl}} \left(\frac{(\xi_i k_b)(\xi_j k_i) z_{ai}}{z_{ak} z_{ij}} - \frac{(\xi_i k_b)(\xi_j k_k)}{z_{jk}} \right) + \frac{(\xi_j k_b)(\xi_k k_l)(\xi_l k_j)}{z_{ai} z_{bj} z_{jl} z_{kl}} \left(\frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{ak} z_{ij}} - \frac{(\xi_i k_b)}{z_{ak} z_{bi}} + \frac{(\xi_i k_k)}{z_{ab} z_{ik}} + \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{ak} z_{il}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_j)(\xi_l k_j) z_{aj}}{z_{ai} z_{bj} z_{jk} z_{jl}} \left(\frac{(\xi_i k_b)}{z_{ak} z_{al} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{ak} z_{al} z_{ij}} - \frac{(\xi_i k_k)}{z_{ab} z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{ab} z_{ak} z_{il}} \right) + \frac{(\xi_i k_b)(\xi_j k_i)(\xi_k k_j)(\xi_l k_j) z_{aj}}{z_{ab} z_{ak} z_{al} z_{bi} z_{ij} z_{jk} z_{jl}} \\
& - \frac{(\xi_i k_b)(\xi_j k_i)(\xi_k k_l)(\xi_l k_j)}{z_{ab} z_{ak} z_{bi} z_{ij} z_{jl} z_{kl}} + \frac{(\xi_j k_b)(\xi_k k_b)(\xi_l k_k)}{z_{ai} z_{bj} z_{bk} z_{kl}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{al} z_{bi}} - \frac{(\xi_i k_j)}{z_{al} z_{ij}} - \frac{(\xi_i k_k) z_{ak}}{z_{aj} z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{aj} z_{il}} \right) \\
& + \frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_k)}{z_{aj} z_{bk} z_{ij} z_{kl}} \left(\frac{(\xi_i k_b)}{z_{al} z_{bi}} - \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{ab} z_{il}} \right) + \frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_k)}{z_{al} z_{bj} z_{ik} z_{kl}} \left(\frac{(\xi_i k_b)}{z_{aj} z_{bi}} - \frac{(\xi_i k_j)}{z_{ab} z_{ij}} \right) \\
& + \frac{(\xi_j k_k)(\xi_k k_b)(\xi_l k_k) z_{ak}}{z_{ai} z_{bk} z_{ik} z_{kl}} \left(\frac{(\xi_i k_j)}{z_{ab} z_{al} z_{ij}} - \frac{(\xi_i k_b)}{z_{aj} z_{al} z_{bi}} + \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{aj} z_{al} z_{ik}} + \frac{(\xi_i k_l)}{z_{ab} z_{aj} z_{il}} \right) - \frac{(\xi_i k_b)(\xi_j k_l)(\xi_k k_i)(\xi_l k_j)}{z_{ab} z_{aj} z_{bi} z_{ik} z_{jl} z_{kl}} \\
& + \frac{(\xi_j k_l)(\xi_k k_b)(\xi_l k_k)}{z_{ai} z_{bk} z_{jl} z_{kl}} \left(\frac{(\xi_i k_j)}{z_{ab} z_{ij}} - \frac{(\xi_i k_b)}{z_{aj} z_{bi}} + \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{aj} z_{ik}} + \frac{(\xi_i k_l) z_{al}}{z_{ab} z_{aj} z_{il}} \right) + \frac{(\xi_i k_b)(\xi_k k_i)(\xi_l k_k)}{z_{ab} z_{aj} z_{al} z_{bi} z_{ik} z_{kl}} \left(\frac{(\xi_j k_i) z_{ai}}{z_{ij}} - \frac{(\xi_j k_k) z_{ak}}{z_{jk}} \right) \\
& + \frac{(\xi_j k_b)(\xi_k k_j)(\xi_l k_k)}{z_{ai} z_{bj} z_{jk} z_{kl}} \left(\frac{(\xi_i k_b)}{z_{al} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{al} z_{ij}} - \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{al} z_{ik}} - \frac{(\xi_i k_l)}{z_{ab} z_{il}} \right) + \frac{(\xi_i k_b)(\xi_j k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab} z_{al} z_{bi} z_{ij} z_{jk} z_{kl}} ,
\end{aligned}$$

$$\begin{aligned}
C_2(a, i, b, j, k, l) = & -\mathcal{E} \left\{ \frac{(\xi_k k_b)(\xi_l k_b) z_{ab}}{z_{ai} z_{aj} z_{bi} z_{bk} z_{bl}} \left(\frac{(\xi_b k_i)(\xi_j k_l)}{z_{ak} z_{jl}} + \frac{(\xi_b k_i)(\xi_j k_k)}{z_{al} z_{jk}} - \frac{(\xi_b k_i)(\xi_j k_b) z_{ab}}{z_{ak} z_{al} z_{bj}} \right) \right. \\
& + \frac{(\xi_j k_l)(\xi_k k_i)(\xi_l k_b)}{z_{aj} z_{bl} z_{ik} z_{jl}} \left(\frac{(\xi_b k_i)}{z_{ak} z_{bi}} + \frac{(\xi_b k_k)}{z_{ai} z_{bk}} \right) + \frac{(\xi_b k_i)(\xi_k k_l)(\xi_l k_b)}{z_{ai} z_{aj} z_{bi} z_{bl} z_{kl}} \left(\frac{(\xi_j k_b) z_{ab}}{z_{ak} z_{bj}} - \frac{(\xi_j k_k)}{z_{jk}} - \frac{(\xi_j k_l)}{z_{ak} z_{jl}} \right) \\
& - \frac{(\xi_k k_i)(\xi_l k_b)}{z_{al} z_{bl} z_{ik}} \left(\frac{(\xi_b k_i)(\xi_j k_b) z_{ab}}{z_{aj} z_{ak} z_{bi} z_{bj}} + \frac{(\xi_b k_k)(\xi_j k_b) z_{ab}}{z_{ai} z_{aj} z_{bj} z_{bk}} + \frac{(\xi_b k_i)(\xi_j k_i) z_{ai}}{z_{aj} z_{ak} z_{bi} z_{ij}} + \frac{(\xi_b k_j)(\xi_j k_i)}{z_{bj} z_{ij}} + \frac{(\xi_b k_k)(\xi_j k_i)}{z_{aj} z_{bk} z_{ij}} \right) \\
& + \frac{(\xi_j k_k)(\xi_k k_i)(\xi_l k_b)}{z_{bl} z_{ik} z_{jk}} \left(\frac{(\xi_b k_i)}{z_{aj} z_{al} z_{bi}} + \frac{(\xi_b k_j)}{z_{ai} z_{al} z_{bj}} + \frac{(\xi_b k_k) z_{ak}}{z_{ai} z_{aj} z_{al} z_{bk}} \right) + \frac{(\xi_j k_i)(\xi_k k_l)(\xi_l k_b)}{z_{ak} z_{bl} z_{ij} z_{kl}} \left(\frac{(\xi_b k_i)}{z_{aj} z_{bi}} + \frac{(\xi_b k_j)}{z_{ai} z_{bj}} \right) \\
& - \frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_b) z_{ab}}{z_{ak} z_{al} z_{bk} z_{bl} z_{ij}} \left(\frac{(\xi_b k_i)}{z_{aj} z_{bi}} + \frac{(\xi_b k_j)}{z_{ai} z_{bj}} \right) - \frac{(\xi_j k_i)(\xi_k k_j)(\xi_l k_b)}{z_{al} z_{bl} z_{ij} z_{jk}} \left(\frac{(\xi_b k_i)}{z_{ak} z_{bi}} + \frac{(\xi_b k_j) z_{aj}}{z_{ai} z_{ak} z_{bj}} + \frac{(\xi_b k_k)}{z_{ai} z_{bk}} \right) \\
& - \frac{(\xi_j k_b)(\xi_k k_b)(\xi_l k_i) z_{ab}}{z_{aj} z_{ak} z_{bj} z_{bk} z_{il}} \left(\frac{(\xi_b k_i)}{z_{al} z_{bi}} + \frac{(\xi_b k_l)}{z_{ai} z_{bl}} \right) + \frac{(\xi_j k_b)(\xi_k k_l)(\xi_l k_i)}{z_{aj} z_{bj} z_{il} z_{kl}} \left(\frac{(\xi_b k_i)}{z_{ak} z_{bi}} + \frac{(\xi_b k_k)}{z_{ai} z_{bk}} + \frac{(\xi_b k_l) z_{al}}{z_{ai} z_{ak} z_{bl}} \right) \\
& - \frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_i)}{z_{ak} z_{bk} z_{ij} z_{il}} \left(\frac{(\xi_b k_i) z_{ai}}{z_{aj} z_{al} z_{bi}} + \frac{(\xi_b k_j)}{z_{al} z_{bj} z_{il}} + \frac{(\xi_b k_l)}{z_{aj} z_{bl} z_{il}} \right) + \frac{(\xi_j k_k)(\xi_k k_b)(\xi_l k_i)}{z_{aj} z_{bk} z_{il} z_{jk}} \left(\frac{(\xi_b k_i)}{z_{al} z_{bi}} + \frac{(\xi_b k_l)}{z_{ai} z_{bl}} \right) \\
& + \frac{(\xi_j k_l)(\xi_k k_b)(\xi_l k_i)}{z_{ak} z_{bk} z_{il} z_{jl}} \left(\frac{(\xi_b k_i)}{z_{aj} z_{bi}} + \frac{(\xi_b k_j)}{z_{ai} z_{bj}} + \frac{(\xi_b k_l) z_{al}}{z_{ai} z_{aj} z_{bl}} \right) - \frac{(\xi_j k_b)(\xi_k k_j)(\xi_l k_i)}{z_{ak} z_{bj} z_{il} z_{jk}} \left(\frac{(\xi_b k_i)}{z_{al} z_{bi}} + \frac{(\xi_b k_l)}{z_{ai} z_{bl}} \right)
\end{aligned}$$

$$\begin{aligned}
& -\frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_i)}{z_{aj}z_{bj}z_{ik}z_{il}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{ak}z_{al}z_{bi}} + \frac{(\xi_b k_k)}{z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ak}z_{bl}} \right) + \frac{(\xi_b k_i)(\xi_k k_j)(\xi_l k_b)}{z_{ai}z_{ak}z_{bi}z_{bl}z_{jk}} \left(\frac{(\xi_j k_l)}{z_{jl}} - \frac{(\xi_j k_b)z_{ab}}{z_{al}z_{bj}} \right) \\
& -\frac{(\xi_j k_i)(\xi_k k_i)(\xi_l k_i)z_{ai}}{z_{ab}z_{ij}z_{ik}z_{il}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{aj}z_{ak}z_{al}z_{bi}} + \frac{(\xi_b k_j)}{z_{ak}z_{al}z_{bj}} + \frac{(\xi_b k_k)}{z_{aj}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{aj}z_{ak}z_{bl}} \right) \\
& +\frac{(\xi_j k_k)(\xi_k k_i)(\xi_l k_i)}{z_{ab}z_{ik}z_{il}z_{jk}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{aj}z_{al}z_{bi}} + \frac{(\xi_b k_j)}{z_{al}z_{bj}} + \frac{(\xi_b k_k)z_{ak}}{z_{aj}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{aj}z_{bl}} \right) + \frac{(\xi_b k_i)(\xi_j k_k)(\xi_k k_b)(\xi_l k_j)}{z_{ai}z_{al}z_{bi}z_{bk}z_{jk}z_{jl}} \\
& +\frac{(\xi_j k_l)(\xi_k k_i)(\xi_l k_i)}{z_{aj}z_{ik}z_{il}z_{jl}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{aj}z_{ak}z_{bi}} + \frac{(\xi_b k_j)}{z_{ak}z_{bj}} + \frac{(\xi_b k_k)}{z_{aj}z_{bk}} + \frac{(\xi_b k_l)z_{al}}{z_{aj}z_{ak}z_{bl}} \right) - \frac{(\xi_b k_i)(\xi_j k_b)(\xi_k k_b)(\xi_l k_j)z_{ab}}{z_{ai}z_{ak}z_{al}z_{bi}z_{bj}z_{bk}z_{jl}} \\
& -\frac{(\xi_j k_i)(\xi_k k_j)(\xi_l k_i)}{z_{ab}z_{ij}z_{il}z_{jk}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{ak}z_{al}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ak}z_{al}z_{bj}} + \frac{(\xi_b k_k)}{z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ak}z_{bl}} \right) - \frac{(\xi_b k_i)(\xi_j k_b)(\xi_k k_j)(\xi_l k_j)z_{aj}}{z_{ai}z_{ak}z_{al}z_{bi}z_{bj}z_{jk}z_{jl}} \\
& +\frac{(\xi_j k_l)(\xi_k k_j)(\xi_l k_i)}{z_{ab}z_{il}z_{jk}z_{jl}} \left(\frac{(\xi_b k_i)}{z_{ak}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ai}z_{ak}z_{bj}} + \frac{(\xi_b k_k)}{z_{ai}z_{bk}} + \frac{(\xi_b k_l)z_{al}}{z_{ai}z_{ak}z_{bl}} \right) + \frac{(\xi_b k_i)(\xi_j k_b)(\xi_k k_l)(\xi_l k_j)}{z_{ai}z_{ak}z_{bi}z_{bj}z_{jl}z_{kl}} \\
& +\frac{(\xi_j k_i)(\xi_k k_l)(\xi_l k_i)}{z_{ab}z_{ij}z_{il}z_{kl}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{aj}z_{ak}z_{bi}} + \frac{(\xi_b k_j)}{z_{ak}z_{bj}} + \frac{(\xi_b k_k)}{z_{aj}z_{bk}} + \frac{(\xi_b k_l)z_{al}}{z_{aj}z_{ak}z_{bl}} \right) - \frac{(\xi_b k_i)(\xi_j k_b)(\xi_k k_b)(\xi_l k_k)z_{ab}}{z_{ai}z_{aj}z_{al}z_{bi}z_{bj}z_{bk}z_{kl}} \\
& -\frac{(\xi_j k_k)(\xi_k k_l)(\xi_l k_i)}{z_{ab}z_{bi}z_{il}z_{kl}} \left(\frac{(\xi_b k_i)}{z_{aj}z_{bi}} + \frac{(\xi_b k_j)}{z_{ai}z_{bj}} + \frac{(\xi_b k_k)z_{ak}}{z_{ai}z_{aj}z_{bk}} + \frac{(\xi_b k_l)z_{al}}{z_{ai}z_{aj}z_{bl}} \right) + \frac{(\xi_b k_i)(\xi_j k_k)(\xi_k k_b)(\xi_l k_k)z_{ak}}{z_{ai}z_{aj}z_{al}z_{bi}z_{bk}z_{jk}z_{kl}} \\
& -\frac{(\xi_j k_l)(\xi_k k_l)(\xi_l k_i)z_{al}}{z_{ab}z_{il}z_{jl}z_{kl}} \left(\frac{(\xi_b k_i)}{z_{aj}z_{ak}z_{bi}} + \frac{(\xi_b k_j)}{z_{ai}z_{ak}z_{bj}} + \frac{(\xi_b k_k)}{z_{ai}z_{aj}z_{bk}} + \frac{(\xi_b k_l)z_{al}}{z_{ai}z_{aj}z_{ak}z_{bl}} \right) \\
& -\frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_j)}{z_{ak}z_{bk}z_{ij}z_{jl}} \left(\frac{(\xi_b k_i)}{z_{al}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ai}z_{al}z_{bj}} + \frac{(\xi_b k_l)}{z_{ai}z_{bl}} \right) - \frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_j)}{z_{al}z_{bj}z_{ik}z_{jl}} \left(\frac{(\xi_b k_i)}{z_{ak}z_{bi}} + \frac{(\xi_b k_k)}{z_{ai}z_{bk}} \right) \\
& -\frac{(\xi_j k_i)(\xi_k k_i)(\xi_l k_j)}{z_{ab}z_{ij}z_{ik}z_{jl}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{ak}z_{al}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ak}z_{al}z_{bj}} + \frac{(\xi_b k_k)}{z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ak}z_{bl}} \right) + \frac{(\xi_b k_i)(\xi_j k_l)(\xi_k k_b)(\xi_l k_k)}{z_{ai}z_{aj}z_{bi}z_{bk}z_{jl}z_{kl}} \\
& +\frac{(\xi_j k_k)(\xi_k k_i)(\xi_l k_j)}{z_{ab}z_{ik}z_{jk}z_{jl}} \left(\frac{(\xi_b k_i)}{z_{al}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ai}z_{al}z_{bj}} + \frac{(\xi_b k_k)z_{ak}}{z_{ai}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ai}z_{bl}} \right) - \frac{(\xi_b k_i)(\xi_j k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab}z_{al}z_{bi}z_{ij}z_{jk}z_{kl}} \\
& -\frac{(\xi_j k_i)(\xi_k k_j)(\xi_l k_j)z_{aj}}{z_{ab}z_{ij}z_{jk}z_{jl}} \left(\frac{(\xi_b k_i)}{z_{ak}z_{al}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ai}z_{ak}z_{al}z_{bj}} + \frac{(\xi_b k_k)}{z_{ai}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ai}z_{ak}z_{bl}} \right) \\
& +\frac{(\xi_j k_l)(\xi_k k_l)(\xi_l k_j)}{z_{ab}z_{ij}z_{jl}z_{kl}} \left(\frac{(\xi_b k_i)}{z_{ak}z_{bi}} + \frac{(\xi_b k_j)z_{aj}}{z_{ai}z_{ak}z_{bj}} + \frac{(\xi_b k_k)}{z_{ai}z_{bk}} + \frac{(\xi_b k_l)}{z_{ai}z_{ak}z_{bl}} \right) - \frac{(\xi_b k_j)(\xi_j k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab}z_{ai}z_{al}z_{bj}z_{ij}z_{jk}z_{kl}} \\
& -\frac{(\xi_j k_i)(\xi_k k_b)(\xi_l k_k)}{z_{al}z_{bk}z_{ij}z_{kl}} \left(\frac{(\xi_b k_i)}{z_{aj}z_{bi}} + \frac{(\xi_b k_j)}{z_{ai}z_{bj}} \right) - \frac{(\xi_b k_i)(\xi_j k_b)(\xi_k k_j)(\xi_l k_k)}{z_{ai}z_{al}z_{bi}z_{bj}z_{jk}z_{kl}} \\
& -\frac{(\xi_j k_b)(\xi_k k_i)(\xi_l k_k)}{z_{aj}z_{bj}z_{ik}z_{kl}} \left(\frac{(\xi_b k_i)}{z_{al}z_{bi}} + \frac{(\xi_b k_k)z_{ak}}{z_{ai}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ai}z_{bl}} \right) - \frac{(\xi_b k_k)(\xi_j k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab}z_{ai}z_{al}z_{bk}z_{ij}z_{jk}z_{kl}} \\
& -\frac{(\xi_j k_i)(\xi_k k_i)(\xi_l k_k)}{z_{ab}z_{ij}z_{ik}z_{kl}} \left(\frac{(\xi_b k_i)z_{ai}}{z_{aj}z_{al}z_{bi}} + \frac{(\xi_b k_j)}{z_{al}z_{bj}} + \frac{(\xi_b k_k)z_{ak}}{z_{aj}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{aj}z_{bl}} \right) - \frac{(\xi_b k_l)(\xi_j k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab}z_{ai}z_{bl}z_{ij}z_{jk}z_{kl}} \\
& +\frac{(\xi_j k_k)(\xi_k k_i)(\xi_l k_k)z_{ak}}{z_{ab}z_{ik}z_{jk}z_{kl}} \left(\frac{(\xi_b k_i)z_{aj}}{z_{al}z_{bi}} + \frac{(\xi_b k_j)}{z_{ai}z_{al}z_{bj}} + \frac{(\xi_b k_k)z_{ak}}{z_{ai}z_{aj}z_{al}z_{bk}} + \frac{(\xi_b k_l)}{z_{ai}z_{aj}z_{bl}} \right) \\
& +\frac{(\xi_j k_l)(\xi_k k_i)(\xi_l k_k)}{z_{ab}z_{ik}z_{jl}z_{kl}} \left(\frac{(\xi_b k_i)}{z_{aj}z_{bi}} + \frac{(\xi_b k_j)}{z_{ai}z_{bj}} + \frac{(\xi_b k_k)z_{ak}}{z_{ai}z_{aj}z_{bk}} + \frac{(\xi_b k_l)z_{al}}{z_{ai}z_{aj}z_{bl}} \right),
\end{aligned}$$

$$\begin{aligned}
C_3(i, j, a, b, k, l) = & -\frac{\mathcal{E}}{z_{ij}} \left\{ \frac{(\xi_a k_i)(\xi_b k_j)(\xi_k k_b)(\xi_l k_b)z_{ab}}{z_{ai}z_{ak}z_{al}z_{bj}z_{bk}z_{bl}} + \frac{(\xi_a k_j)(\xi_b k_i)(\xi_k k_b)(\xi_l k_b)z_{ab}}{z_{aj}z_{ak}z_{al}z_{bi}z_{bk}z_{bl}} \right. \\
& -\frac{(\xi_a k_j)(\xi_b k_i)(\xi_k k_i)(\xi_l k_b)z_{ai}}{z_{aj}z_{ak}z_{al}z_{bi}z_{bl}z_{ik}} - \frac{(\xi_a k_j)(\xi_b k_k)(\xi_k k_i)(\xi_l k_b)}{z_{aj}z_{al}z_{bk}z_{bl}z_{ik}} + \frac{(\xi_a k_i + \xi_a k_k)(\xi_b k_j)(\xi_k k_i)(\xi_l k_b)}{z_{al}z_{bj}z_{bl}z_{ik}} \\
& -\frac{(\xi_a k_j + \xi_a k_k)(\xi_b k_i)(\xi_k k_j)(\xi_l k_b)}{z_{ak}z_{al}z_{bi}z_{bl}z_{jk}} + \frac{(\xi_a k_i)(\xi_b k_j)(\xi_k k_j)(\xi_l k_b)z_{aj}}{z_{ai}z_{ak}z_{al}z_{bj}z_{bl}z_{jk}} + \frac{(\xi_a k_i)(\xi_b k_k)(\xi_k k_j)(\xi_l k_b)}{z_{ai}z_{al}z_{bk}z_{bl}z_{jk}} \\
& +\frac{(\xi_a k_j)(\xi_b k_i)(\xi_k k_l)(\xi_l k_b)}{z_{aj}z_{ak}z_{bi}z_{bl}z_{kl}} + \frac{(\xi_a k_i)(\xi_b k_j)(\xi_k k_l)(\xi_l k_b)}{z_{ai}z_{ak}z_{bj}z_{bl}z_{kl}} + \frac{(\xi_a k_i + \xi_a k_k)(\xi_b k_j)(\xi_k k_b)(\xi_l k_i)}{z_{ak}z_{al}z_{bj}z_{bk}z_{il}} \\
& -\frac{(\xi_a k_j)(\xi_b k_l)(\xi_k k_b)(\xi_l k_i)}{z_{aj}z_{ak}z_{bk}z_{bl}z_{il}} + \frac{(\xi_k k_i)(\xi_l k_i)z_{ai}}{z_{ab}z_{ak}z_{al}z_{ik}z_{il}} \left(\frac{1}{z_{bj}} - \frac{z_{ai}}{z_{aj}z_{bi}} \right) - \frac{(\xi_a k_j)(\xi_b k_i)(\xi_k k_b)(\xi_l k_i)z_{ai}}{z_{aj}z_{ak}z_{al}z_{bi}z_{bk}z_{il}} \\
& +\frac{(\xi_a k_k + \xi_a k_l)(\xi_b k_j)(\xi_k k_i)(\xi_l k_i)z_{ai}}{z_{ab}z_{ak}z_{al}z_{bj}z_{ik}z_{il}} - \frac{(\xi_a k_j)(\xi_k k_i)(\xi_l k_i)z_{ai}}{z_{ab}z_{aj}z_{ik}z_{il}} \left(\frac{\xi_b k_k}{z_{al}z_{bk}} + \frac{\xi_b k_l}{z_{ak}z_{bl}} \right) \\
& \left. -\frac{(\xi_a k_j + \xi_a k_k)(\xi_b k_i)(\xi_k k_j)(\xi_l k_i)z_{ai}}{z_{ab}z_{ak}z_{al}z_{bi}z_{il}z_{jk}} + \frac{(\xi_a k_i + \xi_a k_l)(\xi_b k_j)(\xi_k k_j)(\xi_l k_i)z_{aj}}{z_{ab}z_{ak}z_{al}z_{bj}z_{il}z_{jk}} \right\}
\end{aligned}$$

$$\begin{aligned}
& + \frac{(\xi_a k_i + \xi_a k_l)(\xi_b k_k)(\xi_k k_j)(\xi_l k_i)}{z_{ab} z_{al} z_{bk} z_{il} z_{jk}} - \frac{(\xi_a k_j + \xi_a k_k)(\xi_b k_l)(\xi_k k_j)(\xi_l k_i)}{z_{ab} z_{ak} z_{bl} z_{il} z_{jk}} - \frac{(\xi_a k_j)(\xi_b k_i)(\xi_k k_i)(\xi_l k_k) z_{ai}}{z_{ab} z_{aj} z_{al} z_{bi} z_{ik} z_{kl}} \\
& + \frac{(\xi_a k_j)(\xi_b k_i)(\xi_k k_l)(\xi_l k_i) z_{ai}}{z_{ab} z_{aj} z_{ak} z_{bi} z_{il} z_{kl}} - \frac{(\xi_a k_i + \xi_a k_k + \xi_a k_l)(\xi_b k_j)(\xi_k k_l)(\xi_l k_i)}{z_{ab} z_{ak} z_{bj} z_{il} z_{kl}} \\
& + \frac{(\xi_a k_j)(\xi_k k_l)(\xi_l k_i)}{z_{ab} z_{aj} z_{il} z_{kl}} \left(\frac{\xi_b k_k}{z_{bk}} + \frac{\xi_b k_l}{z_{ak} z_{bl}} \right) + \frac{(\xi_k k_b)(\xi_l k_j)}{z_{ai} z_{ak} z_{bk} z_{jl}} \left(\frac{(\xi_a k_i)(\xi_b k_j) z_{aj}}{z_{al} z_{bj}} + \frac{(\xi_a k_i)(\xi_b k_l)}{z_{bl}} \right) \\
& - \frac{(\xi_a k_j + \xi_a k_l)(\xi_b k_i)(\xi_k k_b)(\xi_l k_j)}{z_{ak} z_{al} z_{bi} z_{bk} z_{jl}} + \frac{(\xi_a k_i)(\xi_l k_j) z_{aj}}{z_{ab} z_{ai} z_{ak} z_{jl}} \left(\frac{(\xi_b k_l)(\xi_k k_j) z_{bl}}{z_{jk}} - \frac{(\xi_b k_j)(\xi_k k_l)}{z_{bj} z_{kl}} \right) \\
& + \frac{(\xi_k k_i)(\xi_l k_j)}{z_{ab} z_{ak} z_{al} z_{ik} z_{jl}} \left(\frac{(\xi_a k_i)(\xi_b k_j) z_{aj}}{z_{bj}} - \frac{(\xi_a k_l)(\xi_b k_i) z_{ai}}{z_{bi}} - \frac{(\xi_a k_j)(\xi_b k_i) z_{ai}}{z_{bi}} \right), \\
& + \frac{(\xi_k k_i)(\xi_l k_j)}{z_{ab} z_{ik} z_{jl}} \left(\frac{(\xi_a k_k)(\xi_b k_j) z_{aj}}{z_{ak} z_{al} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_k)}{z_{al} z_{bk}} - \frac{(\xi_a k_l)(\xi_b k_k)}{z_{al} z_{bk}} + \frac{(\xi_a k_i)(\xi_b k_l)}{z_{ak} z_{bl}} + \frac{(\xi_a k_k)(\xi_b k_l)}{z_{ak} z_{bl}} \right) \quad (3.9) \\
& + \frac{(\xi_b k_i)(\xi_k k_l)(\xi_l k_j)}{z_{ab} z_{ak} z_{bi} z_{jl} z_{kl}} ((\xi_a k_j) + (\xi_a k_k) + (\xi_a k_l)) - \frac{(\xi_a k_i)(\xi_k k_l)(\xi_l k_j)}{z_{ab} z_{ai} z_{jl} z_{kl}} \left(\frac{\xi_b k_k}{z_{bk}} + \frac{(\xi_b k_l) z_{al}}{z_{ak} z_{bl}} \right) \\
& + \frac{(\xi_k k_j)(\xi_l k_j) z_{aj}}{z_{ab} z_{al} z_{jk} z_{jl}} \left(\frac{(\xi_a k_i)(\xi_b k_k)}{z_{bk}} + \frac{(\xi_a k_i)(\xi_b k_j) z_{aj}}{z_{ai} z_{ak} z_{bj}} - \frac{(\xi_a k_l)(\xi_b k_i)}{z_{ak} z_{bi}} - \frac{(\xi_a k_k)(\xi_b k_i)}{z_{ak} z_{bi}} - \frac{(\xi_a k_j)(\xi_b k_i)}{z_{ak} z_{bi}} \right) \\
& + \frac{(\xi_k k_b)(\xi_l k_k)}{z_{al} z_{bk} z_{kl}} \left(\frac{(\xi_a k_i)(\xi_b k_j)}{z_{ai} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_i)}{z_{aj} z_{bi}} \right) + \frac{(\xi_b k_j)(\xi_k k_i)(\xi_l k_k)}{z_{ab} z_{al} z_{bj} z_{ik} z_{kl}} ((\xi_a k_i) + (\xi_a k_k) + (\xi_a k_l)) \\
& - \frac{(\xi_a k_j)(\xi_k k_i)(\xi_l k_k)}{z_{ab} z_{aj} z_{ik} z_{kl}} \left(\frac{(\xi_b k_k) z_{ak}}{z_{al} z_{bk}} + \frac{(\xi_b k_l)}{z_{bl}} \right) - \frac{(\xi_b k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab} z_{al} z_{bi} z_{jk} z_{kl}} ((\xi_a k_j) + (\xi_a k_k) + (\xi_a k_l)) \\
& + \frac{(\xi_a k_i)(\xi_k k_j)(\xi_l k_k)}{z_{ab} z_{ai} z_{jk} z_{kl}} \left(\frac{(\xi_b k_j) z_{aj}}{z_{al} z_{bj}} + \frac{(\xi_b k_k) z_{ak}}{z_{al} z_{bk}} + \frac{(\xi_b k_l)}{z_{bl}} \right) \}.
\end{aligned}$$

From *Eqs.* (3.4) we now extract the various kinematics, classified according to their amount of self-contractions $\xi_i \xi_j$ of polarization vectors ξ_k .

Kinematics ($\xi\xi$) ($\xi\xi$) ($\xi\xi$)

The total number of the kinematics ($\xi\xi$)($\xi\xi$)($\xi\xi$) is 15. They are given as follows:

$$\begin{aligned}
\Xi_1 & := (\xi_1 \xi_2) (\xi_3 \xi_4) (\xi_5 \xi_6) \quad , \quad \Xi_2 := (\xi_1 \xi_2) (\xi_3 \xi_5) (\xi_4 \xi_6) \quad , \quad \Xi_3 := (\xi_1 \xi_2) (\xi_3 \xi_6) (\xi_4 \xi_5), \\
\Xi_4 & := (\xi_1 \xi_3) (\xi_2 \xi_4) (\xi_5 \xi_6) \quad , \quad \Xi_5 := (\xi_1 \xi_3) (\xi_2 \xi_5) (\xi_4 \xi_6) \quad , \quad \Xi_6 := (\xi_1 \xi_3) (\xi_2 \xi_6) (\xi_4 \xi_5), \\
\Xi_7 & := (\xi_1 \xi_4) (\xi_2 \xi_3) (\xi_5 \xi_6) \quad , \quad \Xi_8 := (\xi_1 \xi_4) (\xi_2 \xi_5) (\xi_3 \xi_6) \quad , \quad \Xi_9 := (\xi_1 \xi_4) (\xi_2 \xi_6) (\xi_3 \xi_5), \\
\Xi_{10} & := (\xi_1 \xi_5) (\xi_2 \xi_3) (\xi_4 \xi_6) \quad , \quad \Xi_{11} := (\xi_1 \xi_5) (\xi_2 \xi_4) (\xi_3 \xi_6) \quad , \quad \Xi_{12} := (\xi_1 \xi_5) (\xi_2 \xi_6) (\xi_3 \xi_4), \\
\Xi_{13} & := (\xi_1 \xi_6) (\xi_2 \xi_3) (\xi_4 \xi_5) \quad , \quad \Xi_{14} := (\xi_1 \xi_6) (\xi_2 \xi_4) (\xi_3 \xi_5) \quad , \quad \Xi_{15} := (\xi_1 \xi_6) (\xi_2 \xi_5) (\xi_3 \xi_4). \quad (3.10)
\end{aligned}$$

The two integrands (3.5), which appear in (3.4) in the combinations $\mathcal{A}_1(a, b, i, j, k, l)$ ($\xi_a \xi_i$) ($\xi_b \xi_l$) ($\xi_j \xi_k$) and $\mathcal{A}_2(a, b, i, j, k, l)$ ($\xi_a \xi_b$) ($\xi_i \xi_j$) ($\xi_k \xi_l$), account for the two cases, whether the two vertices a, b in the (-1) -ghost picture are contracted among themselves or with other vertices. The function $\mathcal{A}_1(a, b, i_1, i_2, i_3, i_4)$ is invariant under permutations of i_2, i_3 , while the function $\mathcal{A}_2(a, b, i_1, i_2, i_3, i_4)$ shares permutation symmetry within the pairs of indices i_1, i_2 and i_3, i_4 or mutually exchange of the latter pair. Moreover, the function $\mathcal{A}_2(a, b, i_1, i_2, i_3, i_4)$ is invariant under permutation of a and b . On the other hand,

the function $\mathcal{A}_1(a, b, i_1, i_2, i_3, i_4)$ stays invariant under the latter permutation, if in addition the indices i_1 and i_4 are exchanged. It is easy to convince oneself, that under the above symmetries the kinematics $(\xi_a \xi_{i_1})(\xi_b \xi_{i_4})(\xi_{i_2} \xi_{i_3})$ and $(\xi_a \xi_b)(\xi_{i_1} \xi_{i_2})(\xi_{i_3} \xi_{i_4})$, to which the functions $\mathcal{A}_1, \mathcal{A}_2$ refer, respectively, do not change.

Due to those symmetries it is obvious, that within the sum $\{\}$ of $A^\pi(a, b, i, j, k, l)$ any permutation of the four indices i, j, k, l , which label the four vertices in the zero-ghost picture, yields the same expression (3.4) as a result of Bose-symmetry of the NS -operators. Moreover, the symmetry under the permutation of a and b is evident. In each amplitude $A^\pi(a, b, i, j, k, l)$, as given in Eq. (3.4), all 15 space-time kinematical contractions (3.10) show up once. Exchanging indices from the two sets $\{a, b\}$ and $\{i, j, k, l\}$, *i.e.* putting instead of the operators a, b other operators in the (-1) -ghost picture, leads to seemingly different expressions. In other words, putting different vertex operators in the (-1) -ghost picture, provides a different function A_r^π in front of a given kinematics Ξ . The six-point amplitude allows for 15 possibilities to choose which vertex pair (a, b) to put into the (-1) -ghost picture. Therefore, for the given kinematics

$$(\xi_A \xi_B) (\xi_C \xi_D) (\xi_E \xi_F) \quad (3.11)$$

we obtain 15 *a priori*- different expressions A_r , when we examine (3.4) for the choices

$$(a, b) \in \{ (A, B), (A, C), (A, D), (A, E), (A, F), (B, C), (B, D), (B, E), \\ (B, F), (C, D), (C, E), (C, F), (D, E), (D, F), (E, F) \} . \quad (3.12)$$

The latter may be read off from (3.4):

$$\begin{aligned} A_1^\pi(A, C, B, E, F, D) & , \quad A_1^\pi(A, D, B, E, F, C) & , \quad A_1^\pi(A, E, B, C, D, F) , \\ A_1^\pi(A, F, B, C, D, E) & , \quad A_1^\pi(B, C, A, E, F, D) & , \quad A_1^\pi(B, D, A, E, F, C) , \\ A_1^\pi(B, E, A, C, D, F) & , \quad A_1^\pi(B, F, A, C, D, E) & , \quad A_1^\pi(C, E, D, A, B, F) , \\ A_1^\pi(C, F, D, A, B, E) & , \quad A_1^\pi(D, E, C, A, B, F) & , \quad A_1^\pi(D, F, C, A, B, E) , \\ A_2^\pi(A, B, C, D, E, F) & , \quad A_2^\pi(C, D, A, B, E, F) & , \quad A_2^\pi(E, F, A, B, C, D) . \end{aligned} \quad (3.13)$$

They all have to be equated, which gives rise to 14 equations for each kinematics (3.11) under consideration. *E.g.* the kinematics Ξ_1 , *i.e.* $A = 1, B = 2, C = 3, D = 4, E = 5$ and $F = 6$ yields the system of equations:

$$\begin{aligned} A_2^\pi(1, 2, 3, 4, 5, 6) & = A_2^\pi(3, 4, 1, 2, 5, 6) = A_2^\pi(5, 6, 1, 2, 3, 4) \\ & = A_1^\pi(1, 3, 2, 5, 6, 4) = A_1^\pi(1, 4, 2, 5, 6, 3) = A_1^\pi(1, 5, 2, 3, 4, 6) = A_1^\pi(1, 6, 2, 3, 4, 5) \\ & = A_1^\pi(2, 3, 1, 5, 6, 4) = A_1^\pi(2, 4, 1, 5, 6, 3) = A_1^\pi(2, 5, 1, 3, 4, 6) = A_1^\pi(2, 6, 1, 3, 4, 5) \\ & = A_1^\pi(3, 5, 4, 1, 2, 6) = A_1^\pi(3, 6, 4, 1, 2, 5) = A_1^\pi(4, 5, 3, 1, 2, 6) = A_1^\pi(4, 6, 3, 1, 2, 5) . \end{aligned} \quad (3.14)$$

This system of equations involves 15 different functions \tilde{F} . Each of them appears (linearly) three times. Not all of the equations are independent.

Kinematics $(\xi\xi)$ $(\xi\xi)$ (ξk) (ξk)

The kinematics $(\xi\xi)(\xi\xi)(\xi k)(\xi k)$, of which there exist the total number of $\frac{1}{2} \binom{6}{4} \binom{4}{2} \times 4^2 = 45 \times 16$ on-shell, appears in (3.4) with the four different functions B_1, \dots, B_4 , given in Eqs. (3.7) and (3.8). The latter distinguish the way, how the polarizations of the two vertex operators a and b , which are put in the (-1) -ghost picture, are contracted with the other vectors labeled by i, j, k and l . More precisely, they account for the kinematics:

$$\begin{aligned}
B_2^\pi(a, b, i, j, k, l) (\xi_a \xi_b) (\xi_i \xi_j) &\rightarrow (\xi_a \xi_b) (\xi_i \xi_j) (\xi_k k_r) (\xi_l k_s) , \\
B_1^\pi(a, b, i, j, k, l) (\xi_a \xi_i) (\xi_b \xi_j) &\rightarrow (\xi_a \xi_i) (\xi_b \xi_j) (\xi_k k_r) (\xi_l k_s) , \\
B_3^\pi(a, i, j, k, b, l) (\xi_a \xi_i) (\xi_j \xi_k) &\rightarrow (\xi_a \xi_i) (\xi_j \xi_k) (\xi_b k_r) (\xi_l k_s) , \\
B_4^\pi(i, j, k, l, a, b) (\xi_i \xi_j) (\xi_k \xi_l) &\rightarrow (\xi_i \xi_j) (\xi_k \xi_l) (\xi_a k_r) (\xi_b k_s) .
\end{aligned} \tag{3.15}$$

As before, we may obtain 15 different expressions for one given kinematics

$$(\xi_A \xi_B) (\xi_C \xi_D) (\xi_E k) (\xi_F k) \tag{3.16}$$

depending on which pair (a, b) (running over the set (3.12)) of vertex operators we put into the (-1) ghost picture. Hence, the given kinematics (3.16) may be calculated from the following 15 functions:

$$\begin{aligned}
B_2^\pi(A, B, C, D, E, F) &, \quad B_2^\pi(C, D, A, B, E, F) &, \quad B_1^\pi(A, C, B, D, E, F) , \\
B_1^\pi(A, D, B, C, E, F) &, \quad B_1^\pi(B, C, A, D, E, F) &, \quad B_1^\pi(B, D, A, C, E, F) , \\
B_3^\pi(A, B, C, D, E, F) &, \quad B_3^\pi(B, A, C, D, E, F) &, \quad B_3^\pi(C, D, A, B, E, F) , \\
B_3^\pi(D, C, A, B, E, F) &, \quad B_3^\pi(A, B, C, D, F, E) &, \quad B_3^\pi(B, A, C, D, F, E) , \\
B_3^\pi(C, D, A, B, F, E) &, \quad B_3^\pi(D, C, A, B, F, E) &, \quad B_4^\pi(A, B, C, D, E, F) .
\end{aligned} \tag{3.17}$$

They all have to be equated, which gives rise to 14 equations for each kinematics (3.11) under consideration. More precisely, since the functions B_i^π contain the contractions $\xi_E k$ and $\xi_F k$, of which we have $4 \times 4 = 16$ on-shell, we obtain $14 \times 16 = 224$ equations for the kinematics (3.16). Hence in total, after taking into account all B -kinematics we obtain $45 \times 224 = 10,080$ non-trivial relations, of which many are the same. In fact, the length of the set boils down to 5,464.

Kinematics $(\xi\xi)$ (ξk) (ξk) (ξk) (ξk)

The kinematics $(\xi\xi)(\xi k)(\xi k)(\xi k)(\xi k)$, of which there exist the total number $\binom{6}{2} \times 4^4 = 15 \times 256$ on-shell, appears in (3.4) with the three different functions $C_1^\pi, C_2^\pi, C_3^\pi$, given in Eqs. (3.9) and (3.7). The latter distinguish the way, how the polarizations of the two vertex operators a and b , which are put in the (-1) -ghost picture, are contracted with the other vectors labeled by i, j, k and l :

$$\begin{aligned} C_1^\pi(a, b, i, j, k, l) (\xi_a \xi_b) &\rightarrow (\xi_a \xi_b) (\xi_i k) (\xi_j k) (\xi_k k) (\xi_l k) , \\ C_2^\pi(a, i, b, j, k, l) (\xi_a \xi_i) &\rightarrow (\xi_a \xi_i) (\xi_b k) (\xi_j k) (\xi_k k) (\xi_l k) , \\ C_3^\pi(i, j, a, b, k, l) (\xi_i \xi_j) &\rightarrow (\xi_i \xi_j) (\xi_a k) (\xi_b k) (\xi_k k) (\xi_l k) . \end{aligned} \quad (3.18)$$

As before, we may obtain 15 different expressions for one given kinematics

$$(\xi_A \xi_B) (\xi_C k_m) (\xi_D k_n) (\xi_E k_p) (\xi_F k_q) \quad (3.19)$$

depending on which pair (a, b) (running over the set (3.12)) of vertex operators is put into the (-1) ghost picture. Hence, the given kinematics (3.16) may be calculated from the following 15 functions:

$$\begin{aligned} C_1^\pi(A, B, C, D, E, F) &, \quad C_2^\pi(A, B, C, D, E, F) \quad , \quad C_2^\pi(A, B, D, C, E, F) , \\ C_2^\pi(A, B, E, C, D, F) &, \quad C_2^\pi(A, B, F, C, D, E) \quad , \quad C_2^\pi(B, A, C, D, E, F) , \\ C_2^\pi(B, A, D, C, E, F) &, \quad C_2^\pi(B, A, E, C, D, F) \quad , \quad C_2^\pi(B, A, F, C, D, E) , \\ C_3^\pi(A, B, C, D, E, F) &, \quad C_3^\pi(A, B, C, E, , D, F) \quad , \quad C_3^\pi(A, B, C, F, D, E) , \\ C_3^\pi(A, B, D, E, C, F) &, \quad C_3^\pi(A, B, D, F, C, E) \quad , \quad C_3^\pi(A, B, E, F, C, D) . \end{aligned} \quad (3.20)$$

They all have to be equated, which gives rise to 14 equations for each kinematics (3.11) under consideration. More precisely, since the functions C_i^π contain the contractions $\xi_C k$, $\xi_D k$, $\xi_E k$ and $\xi_F k$, of which we have $4^4 = 256$ on-shell, we obtain $14 \times 256 = 3,584$ equations for the kinematics (3.16). Hence in total, after taking into account all C -kinematics we obtain $15 \times 3,584 = 53,760$ non-trivial relations, of which many are the same. In fact, the length of the set boils down to 6,727. From the structure (3.9) of the functions C_i , namely they do not involve self-contracted momenta, we deduce, that all those relations lead to polynomial identities.

3.2. Specifying the integration region \mathcal{I}_π

Due to the $PSL(2, \mathbf{R})$ -invariance on the disk, we may fix three positions of the vertex operators. A convenient choice in (3.1) or in (3.5) is

$$z_1 = -z_\infty \quad , \quad z_2 = 0 \quad , \quad z_3 = 1 \quad , \quad (3.21)$$

which implies the ghost factor $\langle c(z_1)c(z_2)c(z_3) \rangle = -z_\infty^2$. A six-point amplitude involving massless external particles, *i.e.* $k_i^2 = 0$, allows for nine⁵ independent kinematic invariants s_i . We may choose⁶ the scalar products:

$$\begin{aligned} s_1 &= k_2 k_4 \quad , \quad s_2 = k_2 k_5 \quad , \quad s_3 = k_2 k_6 \quad , \\ s_4 &= k_3 k_4 \quad , \quad s_5 = k_3 k_5 \quad , \quad s_6 = k_3 k_6 \quad , \\ s_7 &= k_4 k_5 \quad , \quad s_8 = k_4 k_6 \quad , \\ s_9 &= k_5 k_6 \quad . \end{aligned} \quad (3.23)$$

Note, that this particular choice of invariants corresponds to our choice of vertex operator positions (3.21), *i.e.* it results from fixing translation invariance on the disk. In other words, our amplitude (3.1) will only depend on these nine invariants s_i .

The general structure of all the integrals (3.7) is then given by:

$$\begin{aligned} \tilde{F} \left[\begin{array}{c} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{array} \right] &:= \int_{\mathcal{I}_\pi} dz_4 dz_5 dz_6 \quad |z_4|^{\alpha_{24}} \quad |z_5|^{\alpha_{25}} \quad |z_6|^{\alpha_{26}} \\ &\times |1 - z_4|^{\alpha_{34}} \quad |1 - z_5|^{\alpha_{35}} \quad |1 - z_6|^{\alpha_{36}} \quad |z_4 - z_5|^{\alpha_{45}} \quad |z_4 - z_6|^{\alpha_{46}} \quad |z_5 - z_6|^{\alpha_{56}} \quad . \end{aligned} \quad (3.24)$$

⁵ An N -point amplitude with N external particles has $\frac{1}{2}N(N-3)$ independent kinematic invariants (quadratic in the momenta). However, in $D=4$ space-time dimensions this number is reduced to $3N-10$ [20]. This leads to qualitative different results for $D3$ -branes.

⁶ The remaining scalar products of momenta are expressed by these nine invariants:

$$\begin{aligned} k_1 k_2 &= s_4 + s_5 + s_6 + s_7 + s_8 + s_9 \quad , \\ k_1 k_3 &= s_1 + s_2 + s_3 + s_7 + s_8 + s_9 \quad , \\ k_1 k_4 &= -s_1 - s_4 - s_7 - s_8 \quad , \\ k_1 k_5 &= -s_2 - s_5 - s_7 - s_9 \quad , \\ k_1 k_6 &= -s_3 - s_6 - s_8 - s_9 \quad , \\ k_2 k_3 &= -s_1 - s_2 - s_3 - s_4 - s_5 - s_6 - s_7 - s_8 - s_9 \quad . \end{aligned} \quad (3.22)$$

Here, the powers α_{ij} are real numbers. More precisely they are given by

$$\begin{aligned}
\alpha_{24} &= s_1 + n_{24} \quad , \quad \alpha_{25} = s_2 + n_{25} \quad , \quad \alpha_{26} = s_3 + n_{26} \quad , \\
\alpha_{34} &= s_4 + n_{34} \quad , \quad \alpha_{35} = s_5 + n_{35} \quad , \quad \alpha_{36} = s_6 + n_{36} \quad , \\
\alpha_{45} &= s_7 + n_{45} \quad , \quad \alpha_{46} = s_8 + n_{46} \quad , \\
\alpha_{56} &= s_9 + n_{56} \quad ,
\end{aligned}
\tag{3.25}$$

with some negative or zero integers n_{ij} . The latter are elements $n_{ij} \in \{\pm 2, \pm 1, 0\}$ depending on how many fermion or/and bosonic two-point functions are involved. As we shall see in the next section, the integral (3.24) represents a multiple hypergeometric function \tilde{F} .

The vertex positions z_4, z_5 and z_6 take arbitrary values on the real axis, *i.e.* $z_4, z_5, z_6 \in \mathbf{R}$. It will prove to be convenient to take the following choice

$$z_4 = x \quad , \quad z_5 = xy \quad , \quad z_6 = xyz \quad , \tag{3.26}$$

which has the Jacobian $\det\left(\frac{\partial(z_4, z_5, z_6)}{\partial(x, y, z)}\right) = x^2 y$. Then the string S -matrix (3.1) with six external gluons becomes:

$$\begin{aligned}
&\mathcal{A}_6(k_1, \xi_1, a_1; k_2, \xi_2, a_2; k_3, \xi_3, a_3; k_4, \xi_4, a_4; k_5, \xi_5, a_5; k_6, \xi_6, a_6) \\
&= \int_{\mathbf{R}} dx \int_{\mathbf{R}} dy \int_{\mathbf{R}} dz x^2 y \langle V_{A^{a_1}}^{(-1)}(-\infty) V_{A^{a_2}}^{(-1)}(0) V_{A^{a_3}}^{(0)}(1) V_{A^{a_4}}^{(0)}(x) V_{A^{a_5}}^{(0)}(xy) V_{A^{a_6}}^{(0)}(xyz) \rangle .
\end{aligned}
\tag{3.27}$$

According to (2.5), the arrangement of the positions z_4, z_5, z_6 , *i.e.* x, y, z , along the real axis implies an ordering π of the group factors λ^{a_i} . This way the full integral in (3.27) splits into 60 different regions \mathcal{I}_π . Each of them gives rise to a different ordering π of the Chan-Paton factors λ^{a_i} in the group trace Tr according to the general expression (2.1). In fact, there are 60 different possibilities⁷ how the Chan-Paton factors λ^a may be contracted in a closed loop. In the following, let us focus on the group contraction $\text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5 \lambda^6)$, *i.e.* the permutation $\pi = (1, 2, 3, 4, 5, 6)$. According to the choice (3.21) and (2.5), this ordering corresponds to taking the region

$$\mathcal{I}_\pi = \{ z_4, z_5, z_6 \in \mathbf{R} \mid 1 < z_4 < z_5 < z_6 < \infty \} \quad , \tag{3.28}$$

⁷ For a given hexagonal diagram, which contracts the Chan-Paton factors λ^{a_i} in the order (i, j, k, l, m, n) , there exist 5 equivalent diagrams: $(j, k, l, m, n, i), \dots, (n, i, j, k, l, m)$, corresponding to the cyclic permutations. Furthermore, changing their orientation results in twelve equivalent diagrams. Thus, from the $6!$ possible permutations we only take into account $720/12 = 60$ hexagonal diagrams.

i.e. $1 < x, y, z < \infty$ along the real axis. After subsection 2.1 and Eq. (3.3) we denote that particular part of the full string S -matrix \mathcal{A}_6 by $A^\pi \equiv A^{(1,2,3,4,5,6)}$. All other 59 possible group theoretical contractions π may be obtained from that particular expression $A^{(1,2,3,4,5,6)}$ by relabeling the quantum numbers of the vertex operators. Therefore, in the following we concentrate on that particular group structure π and omit the index π in the following. For that case, the amplitude (3.3) becomes:

$$\begin{aligned} & \text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5 \lambda^6) A^{(1,2,3,4,5,6)} \equiv \text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5 \lambda^6) A^{(1,2,3,4,5,6)}(1, 2, 3, 4, 5, 6) \\ & = \int_1^\infty dx \int_1^\infty dy \int_1^\infty dz x^2 y \langle V_{A^{a_1}}^{(-1)}(-\infty) V_{A^{a_2}}^{(-1)}(0) V_{A^{a_3}}^{(0)}(1) V_{A^{a_4}}^{(0)}(x) V_{A^{a_5}}^{(0)}(xy) V_{A^{a_6}}^{(0)}(xyz) \rangle . \end{aligned} \quad (3.29)$$

With the choices (3.26) and (3.28) the integral (3.24) simplifies drastically. In fact, after performing the change of variables $x \rightarrow 1/x, y \rightarrow 1/y, z \rightarrow 1/z$ the integral (3.24) assumes the generic form

$$\tilde{F} \left[\begin{matrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{matrix} \right] \longrightarrow F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] , \quad (3.30)$$

with

$$\begin{aligned} F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] & := \int_0^1 dx \int_0^1 dy \int_0^1 dz \\ & \times x^a y^b z^c (1-x)^d (1-y)^e (1-z)^f (1-xy)^g (1-yz)^h (1-xyz)^j , \end{aligned} \quad (3.31)$$

and the new powers

$$\begin{aligned} a & = -4 - \alpha_{24} - \alpha_{25} - \alpha_{26} - \alpha_{34} - \alpha_{35} - \alpha_{36} - \alpha_{45} - \alpha_{46} - \alpha_{56} , \\ b & = -3 - \alpha_{25} - \alpha_{26} - \alpha_{35} - \alpha_{36} - \alpha_{45} - \alpha_{46} - \alpha_{56} , \\ c & = -2 - \alpha_{26} - \alpha_{36} - \alpha_{46} - \alpha_{56} , \\ d & = \alpha_{34} \quad , \quad e = \alpha_{45} \quad , \quad f = \alpha_{56} \\ g & = \alpha_{35} \quad , \quad h = \alpha_{46} \quad , \quad j = \alpha_{36} , \end{aligned} \quad (3.32)$$

given in terms of the integers n_{ij} and invariants s_i (3.25). To this end, the integrands (3.5), (3.8) and (3.9) become certain products of powers of the nine polynomials (1.9). Of course, the number of these polynomials just reflects the number of kinematic invariants

s_i . *E.g.* for the kinematics $(\xi\xi)(\xi\xi)(\xi\xi)$, with (3.5) the expressions (3.7) may be casted into the form

$$\begin{aligned}
A_1^\pi(a, b, i, j, k, l) &= \sigma_{lb}\sigma_{ai}\sigma_{ab} \left\{ \sigma_{jk}\sigma_{ij}\sigma_{kl} (k_i k_j) (k_k k_l) \tilde{F} \left[\begin{array}{l} \bar{n}_{lb}, \bar{n}_{jk} = -1, \bar{n}_{ai} = -1 \\ \bar{n}_{ij}, \bar{n}_{kl} = -1, \bar{n}_{ab} = -1 \end{array} \right] \right. \\
&\quad - \sigma_{jk}\sigma_{ik}\sigma_{jl} (k_i k_k) (k_j k_l) \tilde{F} \left[\begin{array}{l} \bar{n}_{lb}, \bar{n}_{jk} = -1, \bar{n}_{ai} = -1 \\ \bar{n}_{ik}, \bar{n}_{jl} = -1, \bar{n}_{ab} = -1 \end{array} \right] \\
&\quad \left. - \sigma_{il} (k_i k_l) (1 - k_j k_k) \tilde{F} \left[\begin{array}{l} \bar{n}_{lb}, \bar{n}_{il} = -1, \bar{n}_{ai} = -1 \\ \bar{n}_{jk} = -2, \bar{n}_{ab} = -1 \end{array} \right] \right\}, \\
A_2^\pi(a, b, i, j, k, l) &= (1 - k_i k_j) (1 - k_k k_l) \tilde{F} \left[\begin{array}{l} \bar{n}_{ij} = -2 \\ \bar{n}_{kl} = -2, \bar{n}_{ab} = -2 \end{array} \right] \\
&\quad - \sigma_{ij}\sigma_{kl}\sigma_{ik}\sigma_{jl} (k_i k_k) (k_j k_l) \tilde{F} \left[\begin{array}{l} \bar{n}_{ij}, \bar{n}_{kl} = -1 \\ \bar{n}_{ik}, \bar{n}_{jl} = -1, \bar{n}_{ab} = -2 \end{array} \right] \\
&\quad + \sigma_{ij}\sigma_{kl}\sigma_{il}\sigma_{jk} (k_i k_l) (k_j k_k) \tilde{F} \left[\begin{array}{l} \bar{n}_{ij}, \bar{n}_{kl} = -1 \\ \bar{n}_{il}, \bar{n}_{jk} = -1, \bar{n}_{ab} = -2 \end{array} \right],
\end{aligned} \tag{3.33}$$

with the function $\tilde{F} \left[\begin{array}{l} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{array} \right]$ introduced in *Eq.* (3.24) and rewritten in (3.30). Furthermore, we have defined $\sigma_{ij} = \text{sign}(i - j)$, and $\bar{n}_{ij} := \begin{cases} n_{ij}, & i < j \\ n_{ji}, & i > j \end{cases}$. The entries \bar{n}_{ij} in the functions \tilde{F} are to be understood such, that they are only of relevance, if they contribute in (3.30) or (3.25). Otherwise they are meaningless as a matter of the choice (3.21). Similar expressions follow for the other kinematics B and C , given in *Eqs.* (3.8) and (3.9). The same is true for the other group contractions.

The full string S -matrix is invariant under cyclic permutation of the numbering of the six gauge bosons. There are five possible cyclic symmetries, which do not affect the group factor $\text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5 \lambda^6)$. The latter are

$$(i) : 1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 4, 4 \rightarrow 5, 5 \rightarrow 6, 6 \rightarrow 1, \quad \left\{ \begin{array}{l} s_1 \rightarrow s_5 \\ s_2 \rightarrow s_6 \\ s_3 \rightarrow s_1 + s_2 + s_3 + s_7 + s_8 + s_9 \\ s_4 \rightarrow s_7 \\ s_5 \rightarrow s_8 \\ s_6 \rightarrow -s_1 - s_4 - s_7 - s_8 \\ s_7 \rightarrow s_9 \\ s_8 \rightarrow -s_2 - s_5 - s_7 - s_9 \\ s_9 \rightarrow -s_3 - s_6 - s_8 - s_9 \end{array} \right., \tag{3.34}$$

and

$$\begin{aligned}
(ii) &: 1 \rightarrow 3, 2 \rightarrow 4, 3 \rightarrow 5, 4 \rightarrow 6, 5 \rightarrow 1, 6 \rightarrow 2, \\
(iii) &: 1 \rightarrow 4, 2 \rightarrow 5, 3 \rightarrow 6, 4 \rightarrow 1, 5 \rightarrow 2, 6 \rightarrow 3, \\
(iv) &: 1 \rightarrow 5, 2 \rightarrow 6, 3 \rightarrow 1, 4 \rightarrow 2, 5 \rightarrow 3, 6 \rightarrow 4, \\
(v) &: 1 \rightarrow 6, 2 \rightarrow 1, 3 \rightarrow 2, 4 \rightarrow 3, 5 \rightarrow 4, 6 \rightarrow 5,
\end{aligned}$$

with some linear transformation on the nine invariants s_i . The choice of positions (3.21) puts the functions \tilde{F} in a certain form (3.31), which no longer reflects the cyclic invariance of the amplitude (3.1). In other words, the maps (3.34) have a non-trivial impact on the appearance of the string S -matrix (3.29). However, cyclic symmetry is still present. It only requires some non-trivial change of the integration variables x, y and z in (3.31). Generically, the integrand of (3.31) looks like

$$\begin{aligned}
&x^{k_2 k_3 + \dots} y^{k_2 k_3 + k_2 k_4 + k_3 k_4 + \dots} z^{k_1 k_6 + \dots} (1-x)^{k_3 k_4 + \dots} (1-y)^{k_4 k_5 + \dots} (1-z)^{k_5 k_6 + \dots} \\
&(1-xy)^{k_3 k_5 + \dots} (1-yz)^{k_4 k_6 + \dots} (1-xyz)^{k_3 k_6 + \dots},
\end{aligned}$$

with the dots accounting for combinations of integers n_{ij} as given in (3.32). The transformations (3.34) change the form of the integrand. However, for each case (i) – (v) there exists a change of variables x, y, z such, that the integrand of (3.31), may be brought back into its original form (3.31)

$$\begin{aligned}
&\tilde{x}^{k_2 k_3 + \dots} \tilde{y}^{k_2 k_3 + k_2 k_4 + k_3 k_4 + \dots} \tilde{z}^{k_1 k_6 + \dots} (1-\tilde{x})^{k_3 k_4 + \dots} (1-\tilde{y})^{k_4 k_5 + \dots} (1-\tilde{z})^{k_5 k_6 + \dots} \\
&\times (1-\tilde{x}\tilde{y})^{k_3 k_5 + \dots} (1-\tilde{y}\tilde{z})^{k_4 k_6 + \dots} (1-\tilde{x}\tilde{y}\tilde{z})^{k_3 k_6 + \dots},
\end{aligned}$$

with some new variables $\tilde{x}, \tilde{y}, \tilde{z}$ and new combinations of integers m_{ij} . In fact, these coordinate transformations may be found:

$$\begin{aligned}
(i) \quad &\tilde{x} = 1 - xyz, \quad \tilde{y} = \frac{1 - yz}{1 - xyz}, \quad \tilde{z} = \frac{1 - z}{1 - yz}, \\
(ii) \quad &\tilde{x} = z, \quad \tilde{y} = \frac{1 - xy}{1 - xyz}, \quad \tilde{z} = \frac{1 - y}{1 - yz} \frac{1 - xyz}{1 - xy}, \\
(iii) \quad &\tilde{x} = \frac{1 - z}{1 - yz}, \quad \tilde{y} = y, \quad \tilde{z} = \frac{1 - x}{1 - xy}, \\
(iv) \quad &\tilde{x} = \frac{1 - y}{1 - xy} \frac{1 - xyz}{1 - yz}, \quad \tilde{y} = \frac{1 - yz}{1 - xyz}, \quad \tilde{z} = x, \\
(v) \quad &\tilde{x} = \frac{1 - x}{1 - xy}, \quad \tilde{y} = \frac{1 - xy}{1 - xyz}, \quad \tilde{z} = 1 - xyz.
\end{aligned} \tag{3.35}$$

After taking into account their Jacobians for a given function $\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right]$, with its integral representation given in *Eq.* (3.31) and its parameter detailed in (3.32), we find the following five maps under the actions (3.34):

$$\tilde{F}\left[\begin{array}{c} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{array}\right] \longrightarrow \tilde{F}\left[\begin{array}{c} m_{24}, m_{25}, m_{26}, m_{34}, m_{35} \\ m_{36}, m_{45}, m_{46}, m_{56} \end{array}\right], \quad (3.36)$$

with:

$$\begin{aligned} (i) \quad & m_{24} = 2 + n_{24} + n_{25} + n_{26} + n_{45} + n_{46} + n_{56}, \quad m_{25} = -2 - n_{24} - n_{34} - n_{45} - n_{46}, \\ & m_{26} = -2 - n_{25} - n_{35} - n_{45} - n_{56}, \\ & m_{34} = -4 - n_{24} - n_{25} - n_{26} - n_{34} - n_{35} - n_{36} - n_{45} - n_{46} - n_{56}, \\ & m_{35} = n_{24}, \quad m_{36} = n_{25}, \quad m_{45} = n_{34}, \quad m_{46} = n_{35}, \quad m_{56} = n_{45}, \\ (ii) \quad & m_{24} = n_{26}, \quad m_{25} = n_{36}, \quad m_{26} = n_{46}, \quad m_{34} = 2 + n_{34} + n_{35} + n_{36} + n_{45} + n_{46} + n_{56}, \\ & m_{35} = 2 + n_{24} + n_{25} + n_{26} + n_{45} + n_{46} + n_{56}, \quad m_{36} = -2 - n_{24} - n_{34} - n_{45} - n_{46}, \\ & m_{45} = -4 - n_{24} - n_{25} - n_{26} - n_{34} - n_{35} - n_{36} - n_{45} - n_{46} - n_{56}, \\ & m_{46} = n_{24}, \quad m_{56} = n_{34}, \\ (iii) \quad & m_{24} = -2 - n_{25} - n_{35} - n_{45} - n_{56}, \quad m_{25} = n_{25}, \quad m_{26} = n_{35}, \\ & m_{34} = -2 - n_{26} - n_{36} - n_{46} - n_{56}, \quad m_{35} = n_{26}, \quad m_{36} = n_{36}, \\ & m_{45} = 2 + n_{34} + n_{35} + n_{36} + n_{45} + n_{46} + n_{56}, \\ & m_{46} = 2 + n_{24} + n_{25} + n_{26} + n_{45} + n_{46} + n_{56}, \\ & m_{56} = -4 - n_{24} - n_{25} - n_{26} - n_{34} - n_{35} - n_{36} - n_{45} - n_{46} - n_{56}, \\ (iv) \quad & m_{24} = n_{46}, \quad m_{25} = -2 - n_{24} - n_{34} - n_{45} - n_{46}, \quad m_{26} = n_{24}, \quad m_{34} = n_{56}, \\ & m_{35} = -2 - n_{25} - n_{35} - n_{45} - n_{56}, \quad m_{36} = n_{25}, \quad m_{45} = -2 - n_{26} - n_{36} - n_{46} - n_{56}, \\ & m_{46} = n_{26}, \quad m_{56} = 2 + n_{34} + n_{35} + n_{36} + n_{45} + n_{46} + n_{56}, \\ (v) \quad & m_{24} = n_{35}, \quad m_{25} = n_{36}, \quad m_{26} = 2 + n_{24} + n_{25} + n_{26} + n_{45} + n_{46} + n_{56}, \\ & m_{34} = n_{45}, \quad m_{35} = n_{46}, \quad m_{36} = -2 - n_{24} - n_{34} - n_{45} - n_{46}, \quad m_{45} = n_{56}, \\ & m_{46} = -2 - n_{25} - n_{35} - n_{45} - n_{56}, \quad m_{56} = -2 - n_{26} - n_{36} - n_{46} - n_{56}. \end{aligned} \quad (3.37)$$

From these equations we find the two parameter class of cyclically invariant functions:

$$\tilde{F}\left[\begin{array}{c} n_{46}, -2 - 2 n_{46} - 2 n_{56}, n_{46}, n_{56}, n_{46} \\ -2 - 2 n_{46} - 2 n_{56}, n_{56}, n_{46}, n_{56} \end{array}\right], \quad n_{46}, n_{56} \in \mathbf{Z}. \quad (3.38)$$

After having fixed three vertex positions as *e.g.* in (3.21), the meaning of the cyclic symmetries (3.34) on the amplitude (3.29) is, that we essentially may fix three other vertex positions z_i while keeping the ordering π . This is why the choice (3.21) does not represent any restriction, since we may easily move to an other choice, *e.g.* $z_2 = -\infty$, $z_3 = 0$, $z_4 = 1$ through applying the cyclic symmetry (*i*). Of course, in total the string S -matrix result for a given group structure does not change as long as we keep the ordering π of vertex positions, which corresponds to a certain group structure or integration region \mathcal{L}_π .

From the amplitude $A^\pi(1, 2, 4, 3, 6, 5)$, given in (3.4) or from Eq. (3.33), the contribution to the five kinematics $\Xi_1, \Xi_2, \Xi_5, \Xi_7$ and Ξ_8 may be read off:

$$\begin{aligned}
& - \left\{ (1 - s_4)(1 - s_9) \tilde{F} \begin{bmatrix} 0, 0, 0, -2, 0 \\ 0, 0, 0, -2 \end{bmatrix} + s_6 s_7 \tilde{F} \begin{bmatrix} 0, 0, 0, -1, 0 \\ -1, -1, 0, -1 \end{bmatrix} - s_5 s_8 \tilde{F} \begin{bmatrix} 0, 0, 0, -1, -1 \\ 0, 0, -1, -1 \end{bmatrix} \right\} \Xi_1 \\
& - \left\{ (1 - s_5)(1 - s_8) \tilde{F} \begin{bmatrix} 0, 0, 0, 0, -2 \\ 0, 0, -2, 0 \end{bmatrix} - s_6 s_7 \tilde{F} \begin{bmatrix} 0, 0, 0, 0, -1 \\ -1, -1, -1, 0 \end{bmatrix} - s_4 s_9 \tilde{F} \begin{bmatrix} 0, 0, 0, -1, -1 \\ 0, 0, -1, -1 \end{bmatrix} \right\} \Xi_2 \\
& - \left\{ s_5(1 - s_8) \tilde{F} \begin{bmatrix} 0, -1, 0, 0, -1 \\ 0, 0, -2, 0 \end{bmatrix} + s_6 s_7 \tilde{F} \begin{bmatrix} 0, -1, 0, 0, 0 \\ -1, -1, -1, 0 \end{bmatrix} + s_4 s_9 \tilde{F} \begin{bmatrix} 0, -1, 0, -1, 0 \\ 0, 0, -1, -1 \end{bmatrix} \right\} \Xi_5 \\
& - \left\{ -s_4(1 - s_9) \tilde{F} \begin{bmatrix} 0, 0, 0, -1, 0 \\ 0, 0, 0, -2 \end{bmatrix} + s_6 s_7 \tilde{F} \begin{bmatrix} 0, 0, 0, 0, 0 \\ -1, -1, 0, -1 \end{bmatrix} - s_5 s_8 \tilde{F} \begin{bmatrix} 0, 0, 0, 0, -1 \\ 0, 0, -1, -1 \end{bmatrix} \right\} \Xi_7 \\
& + \left\{ -s_7(1 - s_6) \tilde{F} \begin{bmatrix} 0, -1, 0, 0, 0 \\ -2, -1, 0, 0 \end{bmatrix} - s_5 s_8 \tilde{F} \begin{bmatrix} 0, -1, 0, 0, -1 \\ -1, 0, -1, 0 \end{bmatrix} + s_4 s_9 \tilde{F} \begin{bmatrix} 0, -1, 0, -1, 0 \\ -1, 0, 0, -1 \end{bmatrix} \right\} \Xi_8 .
\end{aligned} \tag{3.39}$$

The tensor Ξ_8 stays invariant under all five cyclic transformations (3.34). Hence, this symmetry has also to be respected by the last line of (3.39) and the kinematics Ξ_8 represents a kind of one-dimensional irreducible representation of the cyclic symmetries (3.34). This may be verified using (3.34) and (3.37). On the other hand, through applying the cyclic symmetries (3.34) on the four kinematics $\Xi_1, \Xi_2, \Xi_5, \Xi_7$ we may generate all remaining 10 kinematics (3.10) from those. Hence, (3.39) represents the full content for the A -kinematics (3.10). Similar symmetries hold for the other two classes of kinematics B and C .

According to (3.4) the expression (3.39) refers to the case, where the first and the second gauge boson vertices are chosen to be in the (-1) -ghost picture, *i.e.* $a = 1$ and $b = 2$. Any other choice would lead to a seemingly different expression than (3.39). Only after a detailed knowledge about the functions \tilde{F} and their various relations (*cf.* section 4) we would be able to see, that the expressions for the other choices

$$\begin{aligned}
(a, b) \in \{ & (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 3), (2, 4), (2, 5), (2, 6), \\ & (3, 4), (3, 5), (3, 6), (4, 5), (4, 6), (5, 6) \}
\end{aligned} \tag{3.40}$$

are indeed the same. There are $\binom{6}{2} = 15$ such choices. However, it is this lack of knowledge about the other choices (3.40) we precisely want to make use of. We have already explained in subsection 3.1, that for the kinematics (3.39) we shall now derive fourteen other expressions with different vertex pairs (a, b) in the (-1) -ghost picture. We shall equate all these expressions with (3.39) and obtain striking identities between all the functions \tilde{F} , similar as described in subsection 2.2 for the four gluon scattering case (*cf. Eq. (2.22)*). This information will give us sufficient knowledge about all functions \tilde{F} , which are involved in *Eq. (3.39)*. Without that information we would have to really calculate the integrals (3.31) for all the functions \tilde{F} , which appear in (3.39). These would be 14 functions. In addition, there appear a lot more from the kinematics B and C . This program shall be outlined in the next subsection.

3.3. System of equations and its six-element solution

From equating all expressions (3.20) for the C -kinematics we obtain 1,041 linear relations between different hypergeometric functions (3.30). One example for such a relation is:

$$\tilde{F} \begin{bmatrix} 0, -1, 0, 0, -1 \\ -1, -1, -1, 1 \end{bmatrix} = \tilde{F} \begin{bmatrix} 0, 0, -1, 0, -1 \\ -1, -1, -1, 1 \end{bmatrix} - \tilde{F} \begin{bmatrix} 0, -1, -1, 0, -1 \\ -1, -1, -1, 2 \end{bmatrix}. \quad (3.41)$$

After looking closer to the structure (3.31) of the functions involved, we realize, that this identity is just a manifestation of the simple relation

$$\begin{aligned} & \frac{y(1-z)}{(1-y)(1-xy)z(1-yz)(1-xyz)} \\ &= \frac{y(1-z)}{(1-y)(1-xy)(1-yz)(1-xyz)} + \frac{y(1-z)^2}{(1-y)(1-xy)z(1-yz)(1-xyz)}. \end{aligned} \quad (3.42)$$

between the three polynomials appearing in the integrands of those three functions. An other more involved example is:

$$\begin{aligned} \tilde{F} \begin{bmatrix} -1, -1, -1, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} &= \tilde{F} \begin{bmatrix} -1, 0, -1, 0, -1 \\ -1, -1, -1, 0 \end{bmatrix} - \tilde{F} \begin{bmatrix} -1, 0, -1, 0, -1 \\ -1, -1, 0, 1 \end{bmatrix} - \tilde{F} \begin{bmatrix} -1, 0, 0, 2, -1 \\ -1, -1, -1, 0 \end{bmatrix} \\ &- \tilde{F} \begin{bmatrix} 1, -1, -1, -1, 0 \\ 0, -1, -1, 0 \end{bmatrix} + \tilde{F} \begin{bmatrix} 2, -1, -1, -1, 0 \\ 0, -1, -1, 0 \end{bmatrix} - 2 \tilde{F} \begin{bmatrix} 0, -1, -1, -1, 1 \\ -1, -1, 0, 0 \end{bmatrix} \\ &+ 2 \tilde{F} \begin{bmatrix} 0, -1, -1, -1, 0 \\ -1, -1, 1, 0 \end{bmatrix} - 2 \tilde{F} \begin{bmatrix} 0, -1, -1, -1, 0 \\ -1, -1, 0, 1 \end{bmatrix} - \tilde{F} \begin{bmatrix} 0, -1, -1, 0, -1 \\ -1, 0, -1, 1 \end{bmatrix} \\ &+ \tilde{F} \begin{bmatrix} 0, -1, -1, 0, 0 \\ -1, -1, -1, 1 \end{bmatrix}. \end{aligned} \quad (3.43)$$

Again, this relation originates from a simple polynomial relation:

$$\begin{aligned}
\frac{1}{1-xyz} &= -\frac{y(1-x)^2}{x(1-y)(1-xy)(1-yz)(1-xyz)} + \frac{1-yz}{(1-y)(1-xy)z(1-xyz)} - \frac{1-z}{(1-y)(1-xy)z(1-xyz)} \\
&+ \frac{y(1-z)}{(1-y)(1-yz)(1-xyz)} - \frac{y(1-z)}{(1-xy)(1-yz)(1-xyz)} + \frac{y}{x(1-y)(1-yz)(1-x)} - \frac{y}{(1-y)(1-yz)(1-x)} \\
&- 2 \frac{1-xy}{(1-y)(1-xyz)(1-x)} + 2 \frac{1-yz}{(1-y)z(1-xyz)(1-x)} - 2 \frac{1-z}{(1-y)z(1-xyz)(1-x)} .
\end{aligned} \tag{3.44}$$

All relations, which we find from solving the system of 6,727 equations for the C -kinematics, have their origin in polynomial relations. These identities are to be compared with a similar type of identities (2.29), which appear in the case of four-gluon scattering.

Let us now move on to the equations following from the A and B -kinematics, *i.e.* following from equating (3.14) and (3.20) for each kinematics. In *Eq.* (3.14) we have listed all expressions, which calculate the kinematics Ξ_1 . *E.g.* we have:

$$\begin{aligned}
A_2(1, 2, 3, 4, 5, 6) &= -(1-s_4)(1-s_9) \tilde{F} \begin{bmatrix} 0, 0, 0, -2, 0 \\ 0, 0, 0, -2 \end{bmatrix} - s_6 s_7 \tilde{F} \begin{bmatrix} 0, 0, 0, -1, 0 \\ -1, -1, 0, -1 \end{bmatrix} \\
&\quad + s_5 s_8 \tilde{F} \begin{bmatrix} 0, 0, 0, -1, -1 \\ 0, 0, -1, -1 \end{bmatrix} , \\
A_1(1, 3, 2, 5, 6, 4) &= -s_1 (1-s_9) \tilde{F} \begin{bmatrix} -1, 0, 0, -1, 0 \\ 0, 0, 0, -2 \end{bmatrix} - s_2 s_8 \tilde{F} \begin{bmatrix} 0, -1, 0, -1, 0 \\ 0, 0, -1, -1 \end{bmatrix} \\
&\quad + s_3 s_7 \tilde{F} \begin{bmatrix} 0, 0, -1, -1, 0 \\ 0, -1, 0, -1 \end{bmatrix} , \\
A_1(3, 5, 4, 1, 2, 6) &= -s_8 (1-s_4-s_5-s_6-s_7-s_8-s_9) \tilde{F} \begin{bmatrix} 0, 0, 0, -1, -1 \\ 0, 0, -1, -1 \end{bmatrix} \\
&\quad - s_1 (s_3+s_6+s_8+s_9) \tilde{F} \begin{bmatrix} -1, 0, 0, -1, -1 \\ 0, 0, 0, -1 \end{bmatrix} \\
&\quad + s_3 (s_1+s_4+s_7+s_8) \tilde{F} \begin{bmatrix} 0, 0, -1, -1, -1 \\ 0, 0, 0, -1 \end{bmatrix} , \\
&\quad \dots .
\end{aligned} \tag{3.45}$$

All these equations have to be identical, because each of them determines the kinematics Ξ_1 . Hence, after equating them we obtain a set of linear equations with coefficients depending on the kinematic invariants s_i :

$$A_2(1, 2, 3, 4, 5, 6) = A_1(1, 3, 2, 5, 6, 4) = A_1(3, 5, 4, 1, 2, 6) = \dots . \tag{3.46}$$

Above we have only depicted three expressions for the kinematics Ξ_1 . Writing down all of them for all A - and B -kinematics we obtain a huge set of equations, which qualitatively

looks like (3.46). This system gives rise to various non-trivial relations among different hypergeometric functions. *E.g.* we obtain:

$$\begin{aligned}
& (1 - s_6) \left(\tilde{F} \begin{bmatrix} -1, 0, 0, 0, 0 \\ -2, -1, 0, 0 \end{bmatrix} - \tilde{F} \begin{bmatrix} 0, -1, 0, 0, 0 \\ -2, -1, 0, 0 \end{bmatrix} \right) - s_8 \tilde{F} \begin{bmatrix} -1, -1, 0, 0, 0 \\ -1, 0, -1, 0 \end{bmatrix} \\
& = s_9 \tilde{F} \begin{bmatrix} -1, -1, 0, 0, 0 \\ -1, 0, 0, -1 \end{bmatrix} + s_3 \left(\tilde{F} \begin{bmatrix} -1, 0, -1, 0, 0 \\ -1, -1, 0, 0 \end{bmatrix} - \tilde{F} \begin{bmatrix} 0, -1, -1, 0, 0 \\ -1, -1, 0, 0 \end{bmatrix} \right), \tag{3.47} \\
& (1 + s_4) \tilde{F} \begin{bmatrix} 0, -1, -1, -2, 0 \\ 0, 0, 0, 0 \end{bmatrix} = +s_1 \left(\tilde{F} \begin{bmatrix} -1, -1, 0, -1, 0 \\ 0, 0, 0, -1 \end{bmatrix} - \tilde{F} \begin{bmatrix} -1, 0, -1, -1, 0 \\ 0, 0, 0, -1 \end{bmatrix} \right) \\
& \quad - s_7 \tilde{F} \begin{bmatrix} 0, -1, -1, -1, 0 \\ 0, -1, 0, 0 \end{bmatrix} - s_8 \tilde{F} \begin{bmatrix} 0, -1, -1, -1, 0 \\ 0, 0, -1, 0 \end{bmatrix}, \\
& s_2 \tilde{F} \begin{bmatrix} 0, -1, 0, -1, 0 \\ 0, 0, -1, 0 \end{bmatrix} = -(s_4 + s_5 + s_6 + s_7 + s_8 + s_9) \left(\tilde{F} \begin{bmatrix} 0, 0, 0, -1, 0 \\ 0, -1, 0, -1 \end{bmatrix} - \tilde{F} \begin{bmatrix} 0, 0, 0, -1, 0 \\ 0, 0, -1, -1 \end{bmatrix} \right) \\
& - s_3 \tilde{F} \begin{bmatrix} 0, 0, -1, -1, 0 \\ 0, -1, 0, 0 \end{bmatrix} - (1 + s_1 + s_2 + s_3 + s_4 + s_5 + s_6 + s_7 + s_8 + s_9) \tilde{F} \begin{bmatrix} 0, 0, 0, 0, 0 \\ 0, -1, -1, 0 \end{bmatrix}, \\
& s_2 \tilde{F} \begin{bmatrix} 0, 0, -1, 0, -1 \\ 0, -1, -1, 1 \end{bmatrix} = s_3 \tilde{F} \begin{bmatrix} 0, -1, -1, -1, 0 \\ -1, -1, 0, 1 \end{bmatrix} + (s_1 + s_2 - s_7) \tilde{F} \begin{bmatrix} 0, -1, -1, 0, 0 \\ -1, -1, -1, 1 \end{bmatrix} \\
& \quad + (s_2 - s_3 - s_4 + s_9) \tilde{F} \begin{bmatrix} 0, -1, -1, 0, -1 \\ -1, -1, -1, 2 \end{bmatrix}.
\end{aligned}$$

After inserting the definitions (3.24) and (3.31) into the above equations we see, that these relations may be proven through performing partial integration. We have found a huge amount of identities of that type.

Moreover, after inserting the 1,041 solutions from the C -kinematics into the whole system, which we obtain from the A and B -kinematics (3.17) and (3.20), *i.e.* after eliminating 1,041 functions \tilde{F} from the A and B -system, we are left over with $366 + 210$ equations. The latter only contain $1270 - 1041 = 229$ functions \tilde{F} . This system of equations is undetermined. We realize that all equations may be solved in terms of a basis of six functions. Hence, any of our 1,270 hypergeometric function $\tilde{F} \left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix} \right]$ may be represented as a linear combination of six (multiple) hypergeometric functions. This is in complete analogy with the four-gluon string S -matrix (*cf.* Eq. (2.24)), where the basis was just one-dimensional and with the five-gluon string S -matrix with a two-dimensional basis (*cf.* appendix A). Recall, that in the four-gluon case (and five-gluon case), we were free, which functions F_j we choose for our basis. In that case we had chosen F_0 from the set (2.20). The same was true in the five-gluon case. There we had singled out Φ_1 and Φ_2 . In both cases, F_0 and Φ_1, Φ_2 share some special properties in view of their momentum

expansions: *cf.* Eqs. (2.30) and (A.27). Here again, for our six–dimensional basis we may choose a convenient basis of six functions. Following the discussion after Eq. (2.29) one simple choice for these six functions are integrals, which do not involve poles in the Mandelstamm variables s_i . As basis we may choose the following six functions $\tilde{F}[-1, -1, -2, 0, 0]$, $\tilde{F}[-1, -1, 0, 0, 0]$, $\tilde{F}[-1, 0, -2, 0, -1]$, $\tilde{F}[-1, 0, -2, 0, 0]$, $\tilde{F}[0, -2, -1, 0, 0]$ and $\tilde{F}[-2, -1, -1, 0, 0]$ and express all remaining 1,264 multiple hypergeometric functions $\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right]$, which show up in the string S –matrix, as linear combination of those six:

$$\begin{aligned}
\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right] &= \Lambda_{\{n_{ij}\}}^1(s_i) \tilde{F}\left[\begin{smallmatrix} -1, -1, -2, 0, 0 \\ 0, 0, 0, 0 \end{smallmatrix}\right] + \Lambda_{\{n_{ij}\}}^2(s_i) \tilde{F}\left[\begin{smallmatrix} -1, -1, 0, 0, 0 \\ -2, 0, 0, 0 \end{smallmatrix}\right] \\
&+ \Lambda_{\{n_{ij}\}}^3(s_i) \tilde{F}\left[\begin{smallmatrix} -1, 0, -2, 0, -1 \\ 0, 0, 0, 0 \end{smallmatrix}\right] + \Lambda_{\{n_{ij}\}}^4(s_i) \tilde{F}\left[\begin{smallmatrix} -1, 0, -2, 0, 0 \\ -1, 0, 0, 0 \end{smallmatrix}\right] \\
&+ \Lambda_{\{n_{ij}\}}^5(s_i) \tilde{F}\left[\begin{smallmatrix} 0, -2, -1, 0, 0 \\ -1, 0, 0, 0 \end{smallmatrix}\right] + \Lambda_{\{n_{ij}\}}^6(s_i) \tilde{F}\left[\begin{smallmatrix} -2, -1, -1, 0, 0 \\ -1, 0, 0, 0 \end{smallmatrix}\right].
\end{aligned}
\tag{3.48}$$

In other words, our system of equations may be solved through this six–dimensional ansatz (3.48). Each function $\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right]$ has its six coefficients $\Lambda_{\{n_{ij}\}}^l(s_i)$, $l = 1, \dots, 6$ depending non–trivially on the nine kinematic invariants s_i . Note, that the result (3.48) states quite non–trivial mathematical relations between the functions $\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right]$, introduced in (3.24) and (3.31). The identities like (3.41), (3.43) and (3.47), which are unified in the striking solution (3.48), are interesting by there own, as they give non–trivial relations between different multiple hypergeometric functions $\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right]$. These relations are to be compared with the identities for ordinary hypergeometric functions ${}_pF_q$, as *e.g.* in Ref. [13].

Let us now turn to our preferred⁸ basis of six functions. According to (3.24) and (3.31)

⁸ An alternative choice would replace one of the six functions by:

$$\tilde{F}\left[\begin{smallmatrix} -1, -1, -1, 0, 0 \\ -1, 0, 0, 0 \end{smallmatrix}\right] = \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{\mathcal{P}(x, y, z)}{1 - xyz}. \tag{3.49}$$

the six functions, which serve as our basis in (3.48), are given by the following integrals:

$$\begin{aligned}
\tilde{F} \begin{bmatrix} -1, -1, -2, 0, 0 \\ 0, 0, 0, 0 \end{bmatrix} &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \mathcal{P}(x, y, z) , \\
\tilde{F} \begin{bmatrix} -1, -1, 0, 0, 0 \\ -2, 0, 0, 0 \end{bmatrix} &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{\mathcal{P}(x, y, z)}{(1 - xyz)^2} , \\
\tilde{F} \begin{bmatrix} -1, 0, -2, 0, -1 \\ 0, 0, 0, 0 \end{bmatrix} &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{\mathcal{P}(x, y, z)}{1 - xy} , \\
\tilde{F} \begin{bmatrix} -1, 0, -2, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{z \mathcal{P}(x, y, z)}{1 - xyz} , \\
\tilde{F} \begin{bmatrix} 0, -2, -1, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{y \mathcal{P}(x, y, z)}{1 - xyz} , \\
\tilde{F} \begin{bmatrix} -2, -1, -1, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} &= \int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{x \mathcal{P}(x, y, z)}{1 - xyz} .
\end{aligned} \tag{3.50}$$

with

$$\begin{aligned}
\mathcal{P}(x, y, z) &= x^{-s_1 - s_2 - s_3 - s_4 - s_5 - s_6 - s_7 - s_8 - s_9} y^{-s_2 - s_3 - s_5 - s_6 - s_7 - s_8 - s_9} z^{-s_3 - s_6 - s_8 - s_9} \\
&\times (1 - x)^{s_4} (1 - y)^{s_7} (1 - z)^{s_9} (1 - xy)^{s_5} (1 - yz)^{s_8} (1 - xyz)^{s_6} \\
&= x^{k_2 k_3} y^{k_2 k_3 + k_2 k_4 + k_3 k_4} z^{k_1 k_6} (1 - x)^{k_3 k_4} (1 - y)^{k_4 k_5} (1 - z)^{k_5 k_6} \\
&\times (1 - xy)^{k_3 k_5} (1 - yz)^{k_4 k_6} (1 - xyz)^{k_3 k_6} ,
\end{aligned} \tag{3.51}$$

stemming from the contractions \mathcal{E} of the exponentials (3.6).

The result (3.48) is the key to express the whole string six-gluon S -matrix in terms of the six functions (3.50). By knowing the momentum (α' -) expansion of those six functions, we immediately obtain the momentum expansions of the remaining functions through the identity (3.48). In practice, after having obtained the relations (3.48) for all of the 1,264 functions, for each kinematics one inserts those relations into the corresponding expressions A^π (*cf.* Eqs. (3.33) or (3.39)) and similarly for B^π and C^π . Hence, by calculating the momentum expansion of (3.50) we are able to write down the α' -expansion for the full string S -matrix (*cf.* next subsection).

Let us point out, that we are free in choosing the six-dimensional basis, and with Eq. (3.48) we may easily switch to different basis representations. The advantage of our basis (3.50) is, that we have only to deal with six functions, whose momentum (α' -) expansions do not have poles, just like F_0 in (2.30) for the four-gluon amplitude. This drastically simplifies the way, how we obtain those expansions (*cf.* the next section). Eventually, one may wish to choose a basis, which encodes the structure of the irreducible gluon interactions to all orders in α' . This may be achieved with finding a cyclically invariant basis (3.38). For further details see *Ref.* [21].

In addition, we remark, that the relations (3.48) are general, *i.e.* independent on the gauge structure π and the integral region \mathcal{I}_π . Though we have worked for a given group structure, resulting in the representation (3.31) for the functions, an other group structure π would only result in a different integral representation (3.31), specified by the integral region \mathcal{I}_π and its parameterization. However, the relations (3.48) stay the same as a matter of changing coordinates of the integrands. Hence, the relations (3.48) are completely general and hold also for other integral representations. This is just like in the case of relations for ordinary hypergeometric functions, which also hold for different integral representations.

We have already pointed out in the introduction, that the equations we impose follow from world-sheet supersymmetry and not (directly) from the cyclic invariance of a gluon amplitude. In fact, the equations, which derive from the cyclic symmetry are automatically contained in the system of equations we have studied. Nevertheless this means, that qualitative different results are to be expected for writing the final expression for the string S -matrix of the bosonic string compared to the superstring. By that we mean, that one should expect a smaller basis of functions than (3.48) for the much fewer subset of functions, which appear in a bosonic amplitude. On the other hand, the various relations for the triple hypergeometric functions (3.48) following from the superstring amplitude can be used to also simplify the corresponding bosonic string S -matrix. The latter clearly involves much less triple hypergeometric functions from the beginning.

3.4. Momentum expansion of the string S -matrix

In the following, to display our result we shall adapt to a different basis (3.23) of

kinematical invariants. We choose⁹ the more physical basis [22]:

$$\begin{aligned}
s_1 &= k_1 k_2 \quad , \quad s_2 = k_2 k_3 \quad , \quad s_3 = k_3 k_4 \quad , \\
s_4 &= k_4 k_5 \quad , \quad s_5 = k_5 k_6 \quad , \quad s_6 = k_6 k_1 \quad , \\
s_7 &= \frac{1}{2} (k_1 + k_2 + k_3)^2 \quad , \quad s_8 = \frac{1}{2} (k_2 + k_3 + k_4)^2 \quad , \quad s_9 = \frac{1}{2} (k_3 + k_4 + k_5)^2 \quad .
\end{aligned} \tag{3.52}$$

In this basis *e.g.* the cyclic symmetries (3.34) act as simple permutation symmetry within the two sets $\{s_1, s_2, s_3, s_4, s_5, s_6\}$ and $\{s_7, s_8, s_9\}$. The momentum (or α')–expansion of the six functions (3.50), which will be the subject of subsection 4.5, is given¹⁰ in terms of these new invariants:

$$\begin{aligned}
\Phi_1 &= \tilde{F} \begin{bmatrix} -1, -1, -2, 0, 0 \\ 0, 0, 0, 0 \end{bmatrix} = 1 - 3 s_1 - s_2 + s_3 + s_5 - s_6 + s_7 - s_8 + s_9 \\
&\quad + (s_1 - s_3 - s_4 - s_5) \zeta(2) + (s_1 + s_4 - s_7 - s_9) \zeta(3) + \dots \quad , \\
\Phi_2 &= \tilde{F} \begin{bmatrix} -1, -1, 0, 0, 0 \\ -2, 0, 0, 0 \end{bmatrix} = (1 + s_1 + s_4 - s_7 - s_9) \zeta(2) \\
&\quad - (2 s_1 + s_2 + s_3 + 2 s_4 + s_5 + s_6 - s_7 + s_8 - s_9) \zeta(3) + \dots \quad , \\
\Phi_3 &= \tilde{F} \begin{bmatrix} -1, 0, -2, 0, -1 \\ 0, 0, 0, 0 \end{bmatrix} = (1 - s_1 + s_2 + s_3 + 3 s_4 + s_5 + s_7 - s_8 - s_9) \zeta(2) \\
&\quad + (s_1 - s_2 - 2 s_3 - 4 s_4 - s_5 - s_7 + 2 s_8 + 2 s_9) \zeta(3) + \dots \quad , \\
\Phi_4 &= \tilde{F} \begin{bmatrix} -1, 0, -2, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} = -1 + \zeta(2) + s_1 - s_2 - s_4 + s_5 + 3 s_6 + s_7 - s_8 - s_9 \\
&\quad + (s_2 - s_3 - s_4 - s_5 - 2 s_6 + s_8 + 2 s_9) \zeta(2) \\
&\quad - (s_1 + s_2 - s_3 - 2 s_4 + s_7 + s_8 + 2 s_9) \zeta(3) \dots \quad , \\
\Phi_5 &= \tilde{F} \begin{bmatrix} 0, -2, -1, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} = -1 + \zeta(2) + s_1 - s_2 - 2 s_3 - 2 s_5 - s_6 + s_7 + 3 s_8 + s_9 \\
&\quad + (s_2 + s_3 - s_4 + s_5 + s_6 - 2 s_8) \zeta(2) \\
&\quad - (s_1 + s_2 - s_4 + s_6 + s_7 + s_9) \zeta(3) + \dots \quad ,
\end{aligned} \tag{3.54}$$

⁹ We may easily switch from the basis (3.23) to this new basis through: $s_1 \rightarrow s_8 - s_2 - s_3$, $s_2 \rightarrow s_3 + s_6 - s_8 - s_9$, $s_3 \rightarrow -s_1 - s_6 + s_9$, $s_4 \rightarrow s_3$, $s_5 \rightarrow s_9 - s_3 - s_4$, $s_6 \rightarrow s_1 + s_4 - s_7 - s_9$, $s_7 \rightarrow s_4$, $s_8 \rightarrow -s_4 - s_5 + s_7$, $s_9 \rightarrow s_5$.

¹⁰ For the function (3.49) we obtain:

$$\tilde{F} \begin{bmatrix} -1, -1, -1, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} = \zeta(3) - \frac{1}{4} (s_1 + 4 s_2 + 3 s_3 + 2 s_4 + 3 s_5 + 4 s_6 + s_7 + 4 s_8 + s_9) \zeta(4) + \dots \quad (3.53)$$

$$\begin{aligned}
\Phi_6 = \tilde{F} \begin{bmatrix} -2, -1, -1, 0, 0 \\ -1, 0, 0, 0 \end{bmatrix} &= -1 + \zeta(2) + s_1 + 3 s_2 + s_3 - s_4 - s_6 - s_7 - s_8 + s_9 \\
&- (2 s_2 + s_3 + s_4 + s_5 - s_6 - 2 s_7 - s_8) \zeta(2) \\
&- (s_1 - 2 s_4 - s_5 + s_6 + 2 s_7 + s_8 + s_9) \zeta(3) + \dots .
\end{aligned}$$

The higher orders, which are denoted by the dots, may be depicted from subsection 4.5. With these six functions the six–gluon superstring S –matrix (3.1) may be (formally) written for the group structure $\text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5 \lambda^6)$:

$$\boxed{\mathcal{A}^{(1,2,3,4,5,6)} = \sum_{j=1}^6 \mathcal{P}^j(s_1, \dots, s_9) \Phi_j .} \tag{3.55}$$

Here $\mathcal{P}^j(s_1, \dots, s_9)$ are meromorphic polynomials in the invariants s_i , which capture the set of kinematics $(\xi\xi)(\xi\xi)(\xi\xi)$, $(\xi\xi)(\xi\xi)(\xi k)(\xi k)$ and $(\xi\xi)(\xi k)(\xi k)(\xi k)(\xi k)$. The polynomials \mathcal{P}^j are quite huge expressions, to be compared with the corresponding five–gluon case in *Eqs.* (A.28) and (A.29). It is this form (3.55) of the six–gluon string S –matrix (3.1), which allows to arrange in the effective action the higher order α' gauge interaction terms with six external legs (like $F^6, D^4 F^4, D^2 F^5, D^6 F^4, D^2 F^6, \dots$) thanks to the properties of the functions Φ_j and coefficients \mathcal{P}^j . While the latter \mathcal{P}^j keep track about the pole structure of the reducible diagrams, the functions Φ_j organize the various terms according to their zeta–values [21,18].

Let us now present our final result for the six gluon string S –matrix (3.1), listed according to the kinematics A, B and C . As explained before, we have focused on the group structure $\text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5 \lambda^6)$, *i.e.* on the piece (3.29), since all other 59 group orderings may be obtained by relabeling the gauge vertices, *i.e.* the polarization vectors ξ_i and momenta k_i .

Kinematics $(\xi\xi) (\xi\xi) (\xi\xi)$

We start with the kinematics A , *i.e.* self–contractions of all six polarization vectors. Their expressions are given in *Eq.* (3.39). There we have also explained, that from the set (3.10) we have to display only the five kinematics $\Xi_1, \Xi_2, \Xi_5, \Xi_7$ and Ξ_8 as a result

of cyclic invariance. We obtain the following α' -expansion for that part of the six-gluon string S -matrix (3.1):

$$\begin{aligned}
& \Xi_1 \left\{ -\frac{s_9 s_2}{s_1 s_3 s_5} - \frac{s_2}{s_1 s_3} + \frac{s_2}{s_1 s_5} + \frac{s_2}{s_3 s_5} - \frac{s_2}{s_1 s_7} - \frac{s_4 s_2}{s_1 s_5 s_7} - \frac{s_6 s_2}{s_3 s_5 s_8} - \frac{s_2}{s_3 s_8} + \frac{s_4}{s_1 s_3} + \frac{s_6}{s_1 s_3} \right. \\
& - \frac{s_6}{s_1 s_5} + \frac{s_6}{s_3 s_5} - \frac{s_6 s_7}{s_1 s_3 s_5} - \frac{s_7}{s_1 s_5} + \frac{s_7 s_8}{s_1 s_3 s_5} - \frac{s_4 s_8}{s_1 s_3 s_5} - \frac{s_8}{s_3 s_5} + \frac{s_7 s_9}{s_1 s_3 s_5} + \frac{s_8 s_9}{s_1 s_3 s_5} - \frac{s_9}{s_1 s_3} \\
& + \frac{1}{s_1} + \frac{1}{s_3} + \frac{s_4}{s_1 s_5} - \frac{s_4}{s_3 s_5} + \frac{1}{s_5} - \frac{s_4}{s_5 s_7} - \frac{1}{s_7} - \frac{s_6}{s_5 s_8} - \frac{1}{s_8} - \frac{s_4}{s_3 s_9} - \frac{s_4 s_6}{s_1 s_3 s_9} - \frac{s_6}{s_1 s_9} - \frac{1}{s_9} \\
& + \left(\frac{s_2^2}{s_7} - \frac{s_2^2}{s_5} + \frac{s_4 s_2^2}{s_5 s_7} + \frac{s_6 s_2^2}{s_5 s_8} + \frac{s_2^2}{s_8} + \frac{s_9^2 s_2}{s_1 s_3} + \frac{s_3 s_2}{s_1} - \frac{s_4 s_2}{s_1} - \frac{s_5 s_2}{s_1} - \frac{s_5 s_2}{s_3} - \frac{s_6 s_2}{s_3} + \frac{s_1 s_6 s_2}{s_3 s_5} \right. \\
& - \frac{s_6 s_2}{s_5} + \frac{s_7 s_2}{s_1} - \frac{s_3 s_7 s_2}{s_1 s_5} + \frac{s_8 s_2}{s_3} - \frac{s_1 s_8 s_2}{s_3 s_5} + \frac{s_5 s_9 s_2}{s_1 s_3} + \frac{s_7 s_9 s_2}{s_1 s_5} + \frac{s_8 s_9 s_2}{s_3 s_5} - \frac{s_9 s_2}{s_1} - \frac{s_9 s_2}{s_3} \\
& - \frac{s_9 s_2}{s_5} + \frac{s_1 s_2}{s_3} + \frac{s_3 s_4 s_2}{s_1 s_5} - \frac{s_4 s_2}{s_5} + \frac{s_4^2 s_2}{s_1 s_7} + \frac{s_1 s_2}{s_7} + \frac{s_4 s_5 s_2}{s_1 s_7} + \frac{s_1 s_4 s_2}{s_5 s_7} + \frac{s_6^2 s_2}{s_3 s_8} + \frac{s_3 s_2}{s_8} \\
& + \frac{s_5 s_6 s_2}{s_3 s_8} + \frac{s_3 s_6 s_2}{s_5 s_8} - 2s_2 - \frac{s_4^2}{s_1} - \frac{s_6^2}{s_3} + \frac{s_6 s_7^2}{s_1 s_5} + \frac{s_3 s_7^2}{s_1 s_5} - \frac{s_7 s_8^2}{s_3 s_5} + \frac{s_4 s_8^2}{s_3 s_5} + \frac{s_1 s_8^2}{s_3 s_5} + \frac{s_5 s_9^2}{s_1 s_3} \\
& - \frac{s_7 s_9^2}{s_1 s_3} - \frac{s_8 s_9^2}{s_1 s_3} - \frac{s_1 s_4}{s_3} - 2s_4 + \frac{s_4 s_5}{s_3} - \frac{s_3 s_6}{s_1} - \frac{s_4 s_6}{s_1} - \frac{s_4 s_6}{s_3} + \frac{s_4 s_5 s_6}{s_1 s_3} + \frac{s_5 s_6}{s_1} + \frac{s_1 s_6}{s_5} \\
& - \frac{s_3 s_6}{s_5} - 2s_6 - \frac{s_3 s_7}{s_1} - \frac{s_6 s_7}{s_1} - \frac{s_6 s_7}{s_3} + \frac{s_3 s_6 s_7}{s_1 s_5} - \frac{s_6 s_7}{s_5} - \frac{s_3 s_7}{s_5} - \frac{s_3 s_4 s_7}{s_1 s_5} + \frac{s_4 s_7}{s_5} \\
& - \frac{s_7^2 s_8}{s_1 s_5} - \frac{s_4 s_8}{s_1} - \frac{s_4 s_8}{s_3} - \frac{s_1 s_6 s_8}{s_3 s_5} + \frac{s_6 s_8}{s_5} + \frac{s_6 s_7 s_8}{s_3 s_5} + \frac{s_7 s_8}{s_1} + \frac{s_7 s_8}{s_3} + \frac{s_4 s_7 s_8}{s_1 s_5} + \frac{2s_7 s_8}{s_5} \\
& - \frac{s_1 s_8}{s_3} - \frac{s_1 s_8}{s_5} + \frac{s_1 s_4 s_8}{s_3 s_5} - \frac{s_4 s_8}{s_5} - \frac{s_7^2 s_9}{s_1 s_5} - \frac{s_8^2 s_9}{s_3 s_5} + \frac{s_4 s_9}{s_3} - \frac{s_4 s_5 s_9}{s_1 s_3} - \frac{s_5 s_9}{s_1} - \frac{s_5 s_9}{s_3} \\
& - \frac{s_5 s_6 s_9}{s_1 s_3} + \frac{s_6 s_9}{s_1} + \frac{s_6 s_7 s_9}{s_1 s_3} + \frac{2s_7 s_9}{s_1} + \frac{s_7 s_9}{s_3} + \frac{s_7 s_9}{s_5} + \frac{s_4 s_8 s_9}{s_1 s_3} - \frac{s_7 s_8 s_9}{s_1 s_3} - \frac{s_7 s_8 s_9}{s_1 s_5} \\
& - \frac{s_7 s_8 s_9}{s_3 s_5} + \frac{s_8 s_9}{s_1} + \frac{2s_8 s_9}{s_3} + \frac{s_8 s_9}{s_5} - \frac{s_1 s_4}{s_5} + \frac{s_3 s_4}{s_5} + \frac{s_4^2}{s_7} + \frac{s_4 s_5}{s_7} + \frac{s_6^2}{s_8} + \frac{s_5 s_6}{s_8} + \frac{s_4^2}{s_9} \\
& + \frac{s_4 s_6^2}{s_3 s_9} + \frac{s_6^2}{s_9} + \frac{s_3 s_4}{s_9} + \frac{s_4^2 s_6}{s_1 s_9} + \frac{s_1 s_6}{s_9} + \frac{s_3 s_4 s_6}{s_1 s_9} + \frac{s_1 s_4 s_6}{s_3 s_9} \left. \right) \zeta(2) \Big\} \\
& + \Xi_2 \left\{ \frac{s_2}{s_1 s_7} - \frac{1}{s_1} + \frac{1}{s_7} + \frac{s_6}{s_1 s_9} + \frac{1}{s_9} \right. \\
& + \left(-\frac{s_2^2}{s_7} - \frac{s_3 s_2}{s_1} + \frac{s_5 s_2}{s_1} - \frac{s_1 s_2}{s_7} - \frac{s_4 s_5 s_2}{s_1 s_7} + s_2 + s_1 + s_4 + \frac{s_3 s_6}{s_1} - \frac{s_5 s_6}{s_1} + s_6 + \frac{s_3 s_7}{s_1} \right. \\
& \left. - s_7 + \frac{s_4 s_8}{s_1} - \frac{s_7 s_8}{s_1} + \frac{s_5 s_9}{s_1} - \frac{s_8 s_9}{s_1} - s_9 - \frac{s_4 s_5}{s_7} - \frac{s_6^2}{s_9} - \frac{s_3 s_4}{s_9} - \frac{s_1 s_6}{s_9} - \frac{s_3 s_4 s_6}{s_1 s_9} \right) \zeta(2) \Big\} \\
& + \Xi_5 \left\{ -\frac{1}{s_7} + \left(\frac{s_2 s_1}{s_7} - s_1 - s_2 - s_4 - s_5 + s_7 + s_8 + s_9 + \frac{s_4 s_5}{s_7} \right) \zeta(2) \right\}
\end{aligned} \tag{3.56}$$

$$\begin{aligned}
& + \Xi_7 \left\{ -\frac{s_1}{s_2 s_7} - \frac{s_4 s_1}{s_2 s_5 s_7} + \frac{s_9}{s_2 s_5} + \frac{1}{s_2} + \frac{1}{s_5} - \frac{s_4}{s_5 s_7} - \frac{1}{s_7} - \frac{s_3}{s_2 s_8} - \frac{s_3 s_6}{s_2 s_5 s_8} - \frac{s_6}{s_5 s_8} - \frac{1}{s_8} \right. \\
& + \left[\frac{s_4 s_1^2}{s_5 s_7} + \frac{s_1^2}{s_7} - \frac{s_4 s_1}{s_2} + \frac{s_6 s_1}{s_2} + \frac{s_6 s_1}{s_5} + \frac{s_4 s_8 s_1}{s_2 s_5} - \frac{s_8 s_1}{s_5} - \frac{s_9 s_1}{s_5} + \frac{s_3 s_1}{s_5} - \frac{s_4 s_1}{s_5} + \frac{s_4^2 s_1}{s_2 s_7} \right. \\
& + \frac{s_2 s_1}{s_7} + \frac{s_4 s_5 s_1}{s_2 s_7} + \frac{s_2 s_4 s_1}{s_5 s_7} - s_1 - s_2 - s_3 + \frac{s_3 s_4}{s_2} - s_4 - s_5 - \frac{s_3 s_6}{s_2} + \frac{s_4 s_6}{s_2} - \frac{s_3 s_6}{s_5} - s_6 \\
& - \frac{s_6 s_7}{s_2} + \frac{s_3 s_6 s_7}{s_2 s_5} - \frac{s_3 s_7}{s_5} + s_7 - \frac{s_4 s_8}{s_2} + s_8 - \frac{s_4 s_9}{s_2} - \frac{s_5 s_9}{s_2} - \frac{s_6 s_9}{s_2} + \frac{s_7 s_9}{s_2} + \frac{s_7 s_9}{s_5} - \frac{s_7 s_8 s_9}{s_2 s_5} \\
& + \frac{s_8 s_9}{s_2} + \frac{s_8 s_9}{s_5} - \frac{s_2 s_9}{s_5} - \frac{s_3 s_9}{s_5} + \frac{s_3 s_4}{s_5} + \frac{s_4^2}{s_7} + \frac{s_4 s_5}{s_7} + \frac{s_3^2}{s_8} + \frac{s_3 s_6^2}{s_2 s_8} + \frac{s_6^2}{s_8} + \frac{s_2 s_3}{s_8} + \frac{s_3 s_5 s_6}{s_2 s_8} \\
& \left. + \frac{s_5 s_6}{s_8} + \frac{s_3^2 s_6}{s_5 s_8} + \frac{s_2 s_3 s_6}{s_5 s_8} \right] \zeta(2) \Big\} \\
& + \Xi_8 (s_1 + s_2 + s_3 + s_4 + s_5 + s_6 - s_7 - s_8 - s_9) \zeta(2) + \mathcal{O}(k^4) .
\end{aligned}$$

To keep the expressions short we do not write the orders beyond k^4 . Clearly, the momentum expansions of the kinematics Ξ_8 shows cyclic invariance. There are no k^0 -contributions, a result, which has impact on the possible reducible diagrams contributing to these kinematics (*cf.* section 5).

Kinematics $(\xi\xi)$ $(\xi\xi)$ (ξk) (ξk)

Let us now present the B -kinematics. As outlined in subsection 3.1. there are many independent contractions of the form $(\xi\xi)$ $(\xi\xi)$ (ξk) (ξk) even after making use of the cyclic symmetry (3.34). It is not possible to write down all of them. Instead we pick up a representative example. *E.g.* in the string S -matrix (3.1) after eliminating $\xi_5 k_6$ and $\xi_6 k_5$ on-shell the kinematics $(\xi_1 \xi_2)(\xi_3 \xi_4)(\xi_5 k_2)(\xi_6 k_1)$ comes along with the momentum

dependent expression:

$$\begin{aligned}
& (\xi_1 \xi_2)(\xi_3 \xi_4)(\xi_5 k_2)(\xi_6 k_1) \\
& \times \left\{ -\frac{s_2}{s_3 s_5 s_8} - \frac{s_2}{s_3 s_6 s_8} - \frac{s_7}{s_1 s_3 s_5} + \frac{1}{s_1 s_3} - \frac{1}{s_1 s_5} + \frac{1}{s_3 s_5} + \frac{1}{s_3 s_6} - \frac{1}{s_5 s_8} - \frac{1}{s_6 s_8} \right. \\
& - \frac{s_4}{s_1 s_3 s_9} - \frac{1}{s_1 s_9} - \frac{s_4}{s_3 s_6 s_9} - \frac{1}{s_6 s_9} \\
& + \left(\frac{s_2^2}{s_5 s_8} + \frac{s_2^2}{s_6 s_8} + \frac{s_9 s_2}{s_3 s_6} - \frac{s_2}{s_3} + \frac{s_1 s_2}{s_3 s_5} - \frac{s_2}{s_5} - \frac{s_2}{s_6} + \frac{s_5 s_2}{s_3 s_8} + \frac{s_6 s_2}{s_3 s_8} + \frac{s_3 s_2}{s_5 s_8} + \frac{s_3 s_2}{s_6 s_8} \right. \\
& + \frac{s_7^2}{s_1 s_5} - \frac{s_3}{s_1} - \frac{s_4}{s_1} - \frac{s_4}{s_3} + \frac{s_4 s_5}{s_1 s_3} + \frac{s_5}{s_1} - \frac{s_6}{s_3} - \frac{s_7}{s_1} - \frac{s_7}{s_3} + \frac{s_3 s_7}{s_1 s_5} - \frac{s_7}{s_5} + \frac{s_7 s_8}{s_3 s_5} - \frac{s_1 s_8}{s_3 s_5} \\
& + \frac{s_8}{s_5} + \frac{s_4 s_8}{s_3 s_6} + \frac{s_8}{s_6} - \frac{s_5 s_9}{s_1 s_3} + \frac{s_7 s_9}{s_1 s_3} - \frac{s_8 s_9}{s_3 s_6} + \frac{s_9}{s_1} + \frac{s_9}{s_6} + \frac{s_1}{s_5} - \frac{s_3}{s_5} - \frac{s_3}{s_6} - \frac{s_4}{s_6} + \frac{s_5}{s_8} \\
& + \frac{s_6}{s_8} + \frac{s_4^2}{s_1 s_9} + \frac{s_1}{s_9} + \frac{s_3 s_4}{s_1 s_9} + \frac{s_1 s_4}{s_3 s_9} + \frac{s_4 s_6}{s_3 s_9} + \frac{s_6}{s_9} + \frac{s_4^2}{s_6 s_9} + \frac{s_3 s_4}{s_6 s_9} - 2 \left. \right) \zeta(2) \\
& + \left(-\frac{s_2^3}{s_5 s_8} - \frac{s_2^3}{s_6 s_8} - \frac{2s_3 s_2^2}{s_5 s_8} - \frac{2s_3 s_2^2}{s_6 s_8} - \frac{s_9^2 s_2}{s_3 s_6} + \frac{3s_7 s_2}{s_5} + \frac{s_8 s_2}{s_3} - \frac{s_1 s_8 s_2}{s_3 s_5} + \frac{s_8 s_2}{s_5} \right. \\
& + \frac{s_8 s_2}{s_6} - \frac{s_8 s_9 s_2}{s_3 s_6} + \frac{s_9 s_2}{s_3} - \frac{s_9 s_2}{s_6} + \frac{s_1 s_2}{s_3} - \frac{s_1 s_2}{s_5} + \frac{2s_3 s_2}{s_5} - \frac{s_1^2 s_2}{s_3 s_5} + \frac{2s_3 s_2}{s_6} \\
& + \frac{3s_4 s_2}{s_6} - \frac{s_5^2 s_2}{s_3 s_8} - \frac{s_6^2 s_2}{s_3 s_8} - \frac{2s_5 s_6 s_2}{s_3 s_8} - \frac{s_3^2 s_2}{s_5 s_8} - \frac{s_3^2 s_2}{s_6 s_8} - 2s_2 - \frac{s_7^3}{s_1 s_5} + \frac{s_2^3}{s_1} \\
& - \frac{s_4 s_5^2}{s_1 s_3} - \frac{s_5^2}{s_1} + \frac{s_6^2}{s_3} - \frac{2s_3 s_7^2}{s_1 s_5} - \frac{s_7 s_8^2}{s_3 s_5} + \frac{s_1 s_8^2}{s_3 s_5} - \frac{s_8^2}{s_5} - \frac{s_4 s_8^2}{s_3 s_6} - \frac{s_8^2}{s_6} + \frac{s_5 s_9^2}{s_1 s_3} \\
& - \frac{s_7 s_9^2}{s_1 s_3} + \frac{s_8 s_9^2}{s_3 s_6} - \frac{s_9^2}{s_1} - \frac{s_9^2}{s_6} + s_1 - 2s_3 + \frac{2s_3 s_4}{s_1} - 2s_4 - \frac{s_4 s_5}{s_1} + \frac{s_4 s_5}{s_3} - \frac{s_1 s_5}{s_3} \\
& + s_5 + 2s_6 + \frac{2s_3 s_7}{s_1} + \frac{3s_4 s_7}{s_1} + \frac{s_5 s_7}{s_1} + \frac{s_5 s_7}{s_3} + \frac{2s_6 s_7}{s_3} + \frac{s_1 s_7}{s_3} - \frac{s_3^2 s_7}{s_1 s_5} + \frac{s_1 s_7}{s_5} \\
& + \frac{2s_3 s_7}{s_5} + s_7 + \frac{s_4 s_8}{s_3} - \frac{s_1 s_7 s_8}{s_3 s_5} - \frac{s_7 s_8}{s_5} - \frac{s_1 s_8}{s_5} + \frac{s_1^2 s_8}{s_3 s_5} - \frac{s_4 s_8}{s_6} + 2s_8 + \frac{s_5^2 s_9}{s_1 s_3} \\
& + \frac{s_8^2 s_9}{s_3 s_6} + \frac{s_4 s_9}{s_1} + \frac{s_4 s_9}{s_3} - \frac{s_4 s_5 s_9}{s_1 s_3} - \frac{s_5 s_9}{s_1} - \frac{s_5 s_7 s_9}{s_1 s_3} - \frac{s_7 s_9}{s_1} - \frac{s_8 s_9}{s_3} - \frac{s_4 s_8 s_9}{s_3 s_6} \\
& - \frac{s_8 s_9}{s_6} + \frac{s_4 s_9}{s_6} + 2s_9 - \frac{s_1^2}{s_5} + \frac{s_3^2}{s_5} + \frac{s_3^2}{s_6} + \frac{2s_3 s_4}{s_6} - \frac{s_5^2}{s_8} - \frac{s_6^2}{s_8} - \frac{2s_5 s_6}{s_8} \\
& - \frac{s_4^3}{s_1 s_9} - \frac{s_1^2}{s_9} - \frac{2s_3 s_4^2}{s_1 s_9} - \frac{s_4 s_6^2}{s_3 s_9} - \frac{s_6^2}{s_9} - \frac{s_3^2 s_4}{s_1 s_9} - \frac{s_1^2 s_4}{s_3 s_9} - \frac{2s_1 s_6}{s_9} - \frac{2s_1 s_4 s_6}{s_3 s_9} \\
& \left. - \frac{s_4^3}{s_6 s_9} - \frac{2s_3 s_4^2}{s_6 s_9} - \frac{s_3^2 s_4}{s_6 s_9} \right) \zeta(3) \left. \right\} + \mathcal{O}(k^6).
\end{aligned} \tag{3.57}$$

Again, there are no k^0 -order contributions for this type of kinematics.

Kinematics $(\xi\xi)$ (ξk) (ξk) (ξk) (ξk)

Let us now present the C -kinematics. Here again, as outlined in subsection 3.1, there are many independent contractions of the form $(\xi\xi)$ (ξk) (ξk) (ξk) (ξk) even after applying the cyclic symmetry (3.34). Therefore, it is impossible to write down all of them. Instead, we pick up two representative examples: One with triple poles in the kinematic invariants and an other without any poles. *E.g.* in the string S -matrix (3.1) after eliminating $\xi_1 k_2$, $\xi_4 k_2$, $\xi_5 k_2$ and $\xi_6 k_2$ on-shell the kinematics $(\xi_2 \xi_3)(\xi_1 k_3)(\xi_4 k_1)(\xi_5 k_1)(\xi_6 k_1)$ comes along with the momentum dependent expression:

$$\begin{aligned}
 (\xi_2 \xi_3)(\xi_1 k_3)(\xi_4 k_1)(\xi_5 k_1)(\xi_6 k_1) & \left\{ \frac{1}{s_2 s_6 s_8} - \left(\frac{s_4}{s_2 s_6} + \frac{s_5}{s_2 s_8} + \frac{s_3}{s_6 s_8} \right) \zeta(2) \right. \\
 & + \left(\frac{s_4}{s_2} + \frac{s_5}{s_2} + \frac{s_3}{s_6} + \frac{s_4}{s_6} + \frac{s_4^2}{s_2 s_6} - \frac{s_7}{s_2} + \frac{s_5^2}{s_2 s_8} + \frac{s_2 s_3}{s_6 s_8} \right. \\
 & \left. \left. + \frac{s_3^2}{s_6 s_8} + \frac{s_5 s_6}{s_2 s_8} + \frac{s_4 s_8}{s_2 s_6} - \frac{s_9}{s_6} \right) \zeta(3) \right\} + \mathcal{O}(k^6) .
 \end{aligned} \tag{3.58}$$

Similarly, the kinematics $(\xi_1 \xi_2)(\xi_3 k_6)(\xi_4 k_2)(\xi_5 k_2)(\xi_6 k_2)$ appears with the momentum dependent expansion

$$\begin{aligned}
 (\xi_1 \xi_2)(\xi_3 k_6)(\xi_4 k_2)(\xi_5 k_2)(\xi_6 k_2) & \left\{ \zeta(3) - \left(\frac{1}{4} s_1 + s_2 + \frac{3}{4} s_3 + \frac{1}{2} s_4 + \frac{3}{4} s_5 + s_6 \right. \right. \\
 & \left. \left. + \frac{1}{4} s_7 + s_8 + \frac{1}{4} s_9 \right) \zeta(4) \right\} + \mathcal{O}(k^8) ,
 \end{aligned} \tag{3.59}$$

after eliminating $\xi_3 k_1$, $\xi_4 k_1$, $\xi_5 k_1$ and $\xi_6 k_1$ on-shell.

We have evaluated all kinematical configurations A , B and C up to including the k^6 -order, *i.e.* α'^4 in the string tension. The expressions become rather long. The interested reader may obtain more results from the authors upon request.

4. Multiple hypergeometric functions, generalized Kampé de Fériet function and Euler–Zagier sums

The integrals, which show up in any scattering process of (only) open strings are quite generic. Just as any four–point open string amplitude is given by the Euler–Beta function $B(a, b)$, higher point amplitudes are described by hypergeometric functions or generalizations thereof. The latter are also known as *multiple Gaussian hypergeometric functions*. A multiple Gaussian hypergeometric series is a hypergeometric series in two or more variables, which boils down to the famous Gaussian hypergeometric series ${}_2F_1$ if only one variable is non–zero.

4.1. Generalized hypergeometric functions ${}_pF_q$

Before moving to the six–point amplitude we review the structure of the integrals appearing for four– and five–point open string amplitudes.

The Euler–Beta function $B(a, b)$, relevant for four open string processes, is given by

$$\begin{aligned} B(a, b) &= \int_0^1 dx x^a (1-x)^b = \frac{1}{1+a} {}_2F_1[1+a, -b, 2+a; 1] \\ &= \frac{\Gamma(1+a) \Gamma(1+b)}{\Gamma(2+a+b)} \quad , \quad \text{Re } a > -1 \quad , \quad \text{Re } b > -1 \quad . \end{aligned} \quad (4.1)$$

Moreover, a scattering of five open strings boils down to the integral¹¹:

$$C(a, b, c, d, e) := \int_0^1 dx \int_0^1 dy x^a y^b (1-x)^c (1-y)^d (1-xy)^e \quad . \quad (4.2)$$

It is straightforward to evaluate this integral by making use of the integral representation of the generalized hypergeometric function ${}_pF_q$ [23]:

$$\begin{aligned} {}_2F_1[-c, 1+a, 2+a+b; y] \frac{\Gamma(1+a) \Gamma(1+b)}{\Gamma(2+a+b)} &= \int_0^1 dx x^a (1-x)^b (1-xy)^c \quad , \\ {}_{p+1}F_{q+1} \left[\begin{matrix} 1+a, a_1, \dots, a_p \\ 2+a+b, b_1, \dots, b_q \end{matrix} ; \lambda \right] \frac{\Gamma(1+a) \Gamma(1+b)}{\Gamma(2+a+b)} &= \int_0^1 dx x^a (1-x)^b {}_pF_q \left[\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} ; \lambda x \right] \quad , \end{aligned} \quad (4.3)$$

¹¹ In that case, we have five kinematic invariants, encoded in the powers a, b, c, d and e (cf. Eq. (A.15) for more details)

with $\text{Re } a > -1$, $\text{Re } b > -1$, and $p \leq q + 1$. In addition $|\lambda| < 1$ for the case $p = q + 1$. To this end one obtains

$$C(a, b, c, d, e) = \frac{\Gamma(1+a) \Gamma(1+b) \Gamma(1+c) \Gamma(1+d)}{\Gamma(2+a+c) \Gamma(2+b+d)} {}_3F_2 \left[\begin{matrix} 1+a, 1+b, -e \\ 2+a+c, 2+b+d \end{matrix} ; 1 \right], \quad (4.4)$$

with $\text{Re}(a), \text{Re}(b), \text{Re}(c), \text{Re}(d) > -1$. Later, we will need the power series representation of the hypergeometric functions ${}_pF_q$ [24]:

$${}_pF_q \left[\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} ; x \right] = \sum_{s=0}^{\infty} \frac{1}{s!} \frac{(a_1, s) \cdots (a_p, s)}{(b_1, s) \cdots (b_q, s)} x^s. \quad (4.5)$$

Here we have introduced the Pochhammer symbol $(a, m) = \frac{\Gamma(a+m)}{\Gamma(a)}$. The parameters a_i, b_j and the variable x assumed complex values, except $b_j \neq 0, -1, -2, \dots$. We shall be only concerned with the case $p = q + 1$. For that case, the series is convergent for $|x| < 1$. With

$$\omega := \text{Re} \left(\sum_{i=1}^q b_i - \sum_{i=1}^p a_i \right), \quad (4.6)$$

at $|x| = 1$ the series is absolutely convergent for $\omega > 0$ and divergent for $\omega \leq -1$.

4.2. Generalized Kampé de Fériet function $F_{C:D}^{A:B}$ and triple hypergeometric function $F^{(3)}$

After this warm up we now turn our attention to the integral (3.31):

$$F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] := \int_0^1 dx \int_0^1 dy \int_0^1 dz \\ \times x^a y^b z^c (1-x)^d (1-y)^e (1-z)^f (1-xy)^g (1-yz)^h (1-xyz)^j. \quad (4.7)$$

Any tree-level scattering process with six open strings involves such type of integral. Hence, the latter is as elementary as the expressions (4.1) or (4.2) are generic to open string four or five-point amplitudes, respectively. The function, which eventually will turn to represent some generalized Kampé de Fériet function shows among others the symmetry:

$$F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] = F \left[\begin{matrix} c, b, f, e, h \\ a, d, g, j \end{matrix} \right]. \quad (4.8)$$

To proceed, we use the series expansion:

$$(1-xy)^g (1-yz)^h = \sum_{m,n=0}^{\infty} \frac{(-g, m) (-h, n)}{(1, m) (1, n)} x^m z^n y^{m+n}, \quad |x| < 1; |y| < 1; |z| < 1. \quad (4.9)$$

After replacing in (4.7) the polynomials $(1 - xy)^g (1 - yz)^h$ by the above double sum we are able to perform the integrals with the help of the formulas (4.3), provided $\text{Re}(d), \text{Re}(e), \text{Re}(f) > -1$ and $m + \text{Re}(a) > -1$, $n + \text{Re}(c) > -1$, $\text{Re}(b) + m + n > -1$:

$$\begin{aligned}
F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] &= \Gamma(1+d) \Gamma(1+e) \Gamma(1+f) \sum_{m,n=0}^{\infty} \frac{(-g, m) (-h, n)}{(1, m) (1, n)} \\
&\times \frac{\Gamma(1+m+n+b) \Gamma(1+m+a) \Gamma(1+n+c)}{\Gamma(2+m+n+b+e) \Gamma(2+m+a+d) \Gamma(2+n+c+f)} \\
&\times {}_4F_3 \left[\begin{matrix} 1+m+n+b, 1+m+a, 1+n+c, -j \\ 2+m+n+b+e, 2+m+a+d, 2+n+c+f \end{matrix} ; 1 \right].
\end{aligned} \tag{4.10}$$

Evidently, the result (4.10) shows the symmetry (4.8).

Before we undertake to write (4.10) as a single analytic function we shall discuss the three special cases (i) $h = 0 = g$, (ii) $h = 0$ and (iii) $g = 0$. For the special case (i), *i.e.* $g = 0 = h$, the sum becomes trivial. In that case, only the terms with $m = 0 = n$ give a non-vanishing contribution. This can be seen from the identity $\frac{(-g, m)}{(1, m)} \rightarrow \delta_m$ for $g \rightarrow 0$ and similarly for $\frac{(-h, n)}{(1, n)}$. Hence we find:

$$\begin{aligned}
F \left[\begin{matrix} a, b, d, e, 0 \\ c, f, 0, j \end{matrix} \right] &= \frac{\Gamma(1+a) \Gamma(1+b) \Gamma(1+c) \Gamma(1+d) \Gamma(1+e) \Gamma(1+f)}{\Gamma(2+b+e) \Gamma(2+a+d) \Gamma(2+c+f)} \\
&\times {}_4F_3 \left[\begin{matrix} 1+b, 1+a, 1+c, -j \\ 2+b+e, 2+a+d, 2+c+f \end{matrix} ; 1 \right].
\end{aligned} \tag{4.11}$$

Furthermore, for the case (ii), with $h = 0$:

$$\begin{aligned}
F \left[\begin{matrix} a, b, d, e, g \\ c, f, 0, j \end{matrix} \right] &= \Gamma(1+d) \Gamma(1+e) \Gamma(1+f) \sum_{m=0}^{\infty} \frac{(-g, m)}{(1, m)} \\
&\times \frac{\Gamma(1+m+b) \Gamma(1+m+a) \Gamma(1+c)}{\Gamma(2+m+b+e) \Gamma(2+m+a+d) \Gamma(2+c+f)} \\
&\times {}_4F_3 \left[\begin{matrix} 1+m+b, 1+m+a, 1+c, -j \\ 2+m+b+e, 2+m+a+d, 2+c+f \end{matrix} ; 1 \right].
\end{aligned} \tag{4.12}$$

With the power series representation for the hypergeometric function (4.5) we may express (4.12) through a generalized Kampé de Fériet function $F_{C:D}^{A:B}$ [25]. These are functions in N variables x_1, \dots, x_N and have the power series representation:

$$\begin{aligned}
&F_{C:D}^{A:B} \left[\begin{matrix} a_1, \dots, a_A : b_{1,1}, \dots, b_{1,B}; b_{2,1}, \dots, b_{2,B}; \dots ; b_{N,1}, \dots, b_{N,B} \\ c_1, \dots, c_C : d_{1,1}, \dots, d_{1,D}; d_{2,1}, \dots, d_{2,D}; \dots ; d_{N,1}, \dots, d_{N,D} \end{matrix} ; x_1, \dots, x_N \right] \\
&= \sum_{m_1, \dots, m_N=0}^{\infty} \frac{\prod_{j=1}^A (a_j, m_1 + \dots m_N) \prod_{j=1}^B (b_{1,j}, m_1) \cdot \dots \cdot (b_{N,j}, m_N)}{\prod_{j=1}^C (c_j, m_1 + \dots m_N) \prod_{j=1}^D (d_{1,j}, m_1) \cdot \dots \cdot (d_{N,j}, m_N)} \frac{x_1^{m_1} \cdot \dots \cdot x_N^{m_N}}{m_1! \cdot \dots \cdot m_N!}.
\end{aligned} \tag{4.13}$$

Kampé de Fériet functions are generalizations of the four Lauricella hypergeometric functions¹² F_A, F_B, F_C and F_D . The latter are extensions of Appell's hypergeometric functions F_2, F_3, F_4 and F_1 from two variables to N variables [26], respectively. To this end we may rewrite (4.12):

$$F \left[\begin{matrix} a, b, d, e, g \\ c, f, 0, j \end{matrix} \right] = \frac{\Gamma(1+a) \Gamma(1+b) \Gamma(1+c) \Gamma(1+d) \Gamma(1+e) \Gamma(1+f)}{\Gamma(2+a+d) \Gamma(2+b+e) \Gamma(2+c+f)} \quad (4.15)$$

$$\times F_{2:2}^{2:1} \left[\begin{matrix} 1+a, 1+b : 1+c, -j ; -g, 1 \\ 2+a+d, 2+b+e : 2+c+f ; 1 \end{matrix} ; 1, 1 \right].$$

We may derive a similar expression for the third case (*iii*).

After having discussed the three special cases we now want to turn to the general case (4.10)

$$F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] = \frac{\Gamma(1+d) \Gamma(1+e) \Gamma(1+f)}{\Gamma(-g) \Gamma(-h) \Gamma(-j)} \sum_{m_i=0}^{\infty} \frac{\Gamma(-g+m_1) \Gamma(-h+m_2) \Gamma(-j+m_3)}{m_1! m_2! m_3!}$$

$$\times \frac{\Gamma(1+m_1+m_2+m_3+b)}{\Gamma(2+m_1+m_2+m_3+b+e)} \frac{\Gamma(1+m_1+m_3+a)}{\Gamma(2+m_1+m_3+a+d)} \frac{\Gamma(1+m_2+m_3+c)}{\Gamma(2+m_2+m_3+c+f)}. \quad (4.16)$$

A convenient expression for the triple sum (4.16) may be also

$$F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] = \frac{\Gamma(1+d) \Gamma(1+e) \Gamma(1+f)}{\Gamma(-g) \Gamma(-h) \Gamma(-j)}$$

$$\times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{n_1} \sum_{n_3=n_1-n_2}^{n_1} \frac{\Gamma(-g+n_1-n_3) \Gamma(-h+n_1-n_2) \Gamma(-j-n_1+n_2+n_3)}{\Gamma(1+n_1-n_3) \Gamma(1+n_1-n_2) \Gamma(1-n_1+n_2+n_3)}$$

$$\times \frac{\Gamma(1+n_1+b)}{\Gamma(2+n_1+b+e)} \frac{\Gamma(1+n_2+a)}{\Gamma(2+n_2+a+d)} \frac{\Gamma(1+n_3+c)}{\Gamma(2+n_3+c+f)} \quad (4.17)$$

¹² The latter correspond to the following four special cases of $F_{C:D}^{A:B}$:

$$F_A : A = 1, B = 1, C = 0, D = 1,$$

$$F_B : A = 0, B = 2, C = 1, D = 0,$$

$$F_C : A = 2, B = 0, C = 0, D = 1,$$

$$F_D : A = 1, B = 1, C = 1, D = 0. \quad (4.14)$$

with two terminating sums. We want to express the triple sum (4.16) through a generalized Kampé de Fériet function. In fact, Srivastava has introduced a general *triple hypergeometric function* $F^{(3)}[x, y, z]$ depending on three variables, *cf.* last *Ref.* of [25]

$$F^{(3)}[x, y, z] \equiv F^{(3)} \left[\begin{matrix} (a) :: (b); (b'); (b'') : (c); (c'); (c'') \\ (e) :: (g); (g'); (g'') : (h); (h'); (h'') \end{matrix} ; x, y, z \right] = \sum_{m, n, p=0}^{\infty} \Lambda(m, n, p) \frac{x^m}{m!} \frac{y^n}{n!} \frac{z^p}{p!}, \quad (4.18)$$

with

$$\begin{aligned} \Lambda(m, n, p) &= \frac{\prod_{j=1}^A (a_j, m+n+p) \prod_{j=1}^B (b_j, m+n) \prod_{j=1}^{B'} (b'_j, n+p) \prod_{j=1}^{B''} (b''_j, m+p)}{\prod_{j=1}^E (e_j, m+n+p) \prod_{j=1}^G (g_j, m+n) \prod_{j=1}^{G'} (g'_j, n+p) \prod_{j=1}^{G''} (g''_j, m+p)} \\ &\quad \times \frac{\prod_{j=1}^C (c_j, m) \prod_{j=1}^{C'} (c'_j, n) \prod_{j=1}^{C''} (c''_j, p)}{\prod_{j=1}^H (h_j, m) \prod_{j=1}^{H'} (h'_j, n) \prod_{j=1}^{H''} (h''_j, p)}, \end{aligned} \quad (4.19)$$

with (a) denoting a set of A parameter a_1, \dots, a_A and similarly for $(b), (b'), \dots$. With this nice expression we may write our integral (4.7), which is expressed through (4.16), in the following closed form:

$$\begin{aligned} F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] &= \frac{\Gamma(1+a) \Gamma(1+b) \Gamma(1+c) \Gamma(1+d) \Gamma(1+e) \Gamma(1+f)}{\Gamma(2+a+d) \Gamma(2+b+e) \Gamma(2+c+f)} \\ &\quad \times F^{(3)} \left[\begin{matrix} 1+b :: 1; 1+c; 1+a : -g, 1; -h, 1; -j, 1 \\ 2+b+e :: 1; 2+c+f; 2+a+d : 1; 1; 1 \end{matrix} ; 1, 1, 1 \right]. \end{aligned} \quad (4.20)$$

4.3. Harmonic Number and Euler sums

To analyze the string S -matrix results, we need their momentum expansion. Hence, we need to expand the hypergeometric series (4.5) w.r.t. small entries a_i, b_i . A formal expression for the generalized hypergeometric function (4.5) may be given

$$\begin{aligned} {}_pF_q \left[\begin{matrix} a_1 + \epsilon \alpha_1, \dots, a_p + \epsilon \alpha_p \\ b_1 + \epsilon \beta_1, \dots, b_q + \epsilon \beta_q \end{matrix} \right] &= \sum_{s=0}^{\infty} \frac{1}{s!} \frac{\prod_{i=1}^p (a_i + \epsilon \alpha_i, s)}{\prod_{j=1}^q (b_j + \epsilon \beta_j, s)} \\ \frac{(m + \alpha \epsilon, s)}{(m, s)} &= e^{-\sum_{k=1}^{\infty} \frac{(-\alpha \epsilon)^k}{k} (H_{m+s-1, k} - H_{m-1, k})}. \end{aligned} \quad (4.21)$$

Strictly speaking, in the last formula, m has to be an integer positive number, with $m > 1$. This condition limits the range of parameters a_i, b_j . However, by analytic continuation, the above expression keeps its form. At any rate, from this formula we see, that quite generically a class of infinite sums appears. These sums are known as *Euler sums* and involve the *harmonic number* H_n or the *generalized harmonic number* $H_{n,a}$. The harmonic number H_n and the generalized harmonic number $H_{n,a}$ are defined:

$$H_n = \sum_{k=1}^n \frac{1}{k} \quad , \quad H_{n,a} = \sum_{k=1}^n \frac{1}{k^a} . \quad (4.22)$$

Clearly, we have $H_{n,1} = H_n$ and $H_n = \gamma_E + \psi(n+1)$. The harmonic number is related to the polygamma function ψ :

$$\begin{aligned} \psi(n) &= -\gamma_E + H_{n-1} , \\ \psi^{(1)}(n) &= \zeta(2) - H_{n-1,2} . \end{aligned} \quad (4.23)$$

The sums, which appear after expanding the exponential (4.21) w.r.t. small ϵ , generically lead to Euler sums. The following two classes of Euler sums are relevant to us:

$$\begin{aligned} s_h(m, n) &= \sum_{k=1}^{\infty} \frac{H_k^m}{(k+1)^n} , \\ \sigma_h(m, n) &= \sum_{k=1}^{\infty} \frac{H_{k,m}}{(k+1)^n} . \end{aligned} \quad (4.24)$$

for $m \geq 1, n \geq 2$. We have the following relations:

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{H_{k,m}}{k^n} &= \sigma_h(m, n) + \zeta(m+n) , \\ \sum_{k=1}^{\infty} \frac{H_k^m}{k^n} &= s_h(m, n) - \sum_{j=0}^{m-1} \binom{m}{j} \sum_{k=1}^{\infty} (-1)^{m-j} \frac{H_k^j}{k^{m+n-j}} . \end{aligned} \quad (4.25)$$

From the last equation we obtain the identity:

$$\sum_{k=1}^{\infty} \frac{H_k^2}{k^n} = s_h(2, n) + 2 s_h(1, n+1) + \zeta(2+n) . \quad (4.26)$$

It has been already pointed out by Euler, that in some cases the sums may be reduced to a rational linear combination of product of single sums. There is recent work on the explicit

evaluation of some Euler sums in *Refs.* [27,28,29]. In the following let us report some of their results:

$$\begin{aligned}
\sigma_h(1, 2) &= \zeta(3) , \\
\sigma_h(1, 3) &= \frac{3}{2} \zeta(4) - \frac{1}{2} \zeta(2)^2 , \\
\sigma_h(2, 2) &= \frac{1}{2} \zeta(2)^2 - \frac{1}{2} \zeta(4) , \\
\sigma_h(1, 4) &= 2 \zeta(5) - \zeta(2) \zeta(3) , \\
\sigma_h(2, 3) &= -\frac{11}{2} \zeta(5) + 3 \zeta(2) \zeta(3) , \\
\sigma_h(2, 4) &= -6 \zeta(6) + \frac{8}{3} \zeta(2) \zeta(4) + \zeta(3)^2 , \\
\sigma_h(1, 5) &= \frac{5}{2} \zeta(6) - \zeta(2) \zeta(4) - \frac{1}{2} \zeta(3)^2 , \\
\sigma_h(4, 2) &= 5 \zeta(6) - \frac{5}{3} \zeta(2) \zeta(4) - \zeta(3)^2 ,
\end{aligned} \tag{4.27}$$

and:

$$\begin{aligned}
s_h(2, 2) &= \frac{3}{2} \zeta(4) + \frac{1}{2} \zeta(2)^2 = \frac{11}{4} \zeta(4) , \\
s_h(3, 2) &= \frac{15}{2} \zeta(5) + \zeta(2) \zeta(3) , \\
s_h(2, 4) &= \frac{2}{3} \zeta(6) - \frac{1}{3} \zeta(2) \zeta(4) + \frac{1}{3} \zeta(2)^3 - \zeta(3)^2 .
\end{aligned} \tag{4.28}$$

In addition, we have [28]:

$$s_h(1, n) = \sigma_h(1, n) = \frac{1}{2} n \zeta(n+1) - \frac{1}{2} \sum_{k=1}^{n-2} \zeta(n-k) \zeta(k+1) . \tag{4.29}$$

An useful identity is the *reflection formula* [29]

$$\sigma_h(s, t) + \sigma_h(t, s) = \zeta(s) \zeta(t) - \zeta(s+t) \quad , \quad s, t \geq 2 , \tag{4.30}$$

which leads to

$$\sigma_h(a, a) = \frac{1}{2} \zeta(a)^2 - \frac{1}{2} \zeta(2a) \quad , \quad a \geq 2 , \tag{4.31}$$

or:

$$\sum_{n=1}^{\infty} \frac{H_{n,a}}{n^a} = \frac{1}{2} \zeta(a)^2 + \frac{1}{2} \zeta(2a) \quad , \quad a \geq 2 . \tag{4.32}$$

So far, we have reviewed very useful identities about Euler sums from the literature. Essentially we have quoted results from [27,28,29]. Let us now apply these results to

infinite sums involving the harmonic number series (4.22) as they appear in the expansion of hypergeometric series: First we shall need the class of sums:

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{H_n}{n^2} &= 2 \zeta(3) \quad , \quad \sum_{n=1}^{\infty} \frac{H_n^2}{n^2} = \frac{11}{2} \zeta(4) - \frac{1}{2} \zeta(2)^2 = \frac{17}{4} \zeta(4) \quad , \quad \sum_{n=1}^{\infty} \frac{H_n}{n^3} = \frac{5}{4} \zeta(4) \quad , \\ \sum_{n=1}^{\infty} \frac{H_n}{n^4} &= 3 \zeta(5) - \zeta(2) \zeta(3) \quad , \quad \sum_{n=1}^{\infty} \frac{H_n}{n^5} = \zeta(2) \zeta(4) - \frac{1}{2} \zeta(3)^2 \quad . \end{aligned} \tag{4.33}$$

Later, we shall need the following series involving the generalized harmonic number. They may be easily derived from the above identities:

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{H_{n,2}}{n^4} &= \sigma_h(2, 4) + \zeta(6) = -5 \zeta(6) + \frac{8}{3} \zeta(2) \zeta(4) + \zeta(3)^2 \quad , \\ \sum_{n=1}^{\infty} \frac{H_{n,2}}{n^3} &= \sigma_h(2, 3) + \zeta(5) = -\frac{9}{2} \zeta(5) + 3 \zeta(2) \zeta(3) \quad , \\ \sum_{n=1}^{\infty} \frac{H_{n,4}}{n^2} &= \sigma_h(4, 2) + \zeta(6) = 6 \zeta(6) - \frac{5}{3} \zeta(2) \zeta(4) - \zeta(3)^2 \quad , \end{aligned} \tag{4.34}$$

In the following we shall be concerned to extend the above list to our needs in the subsections 4.4. and 4.5. First, using (4.32) let us evaluate the sum:

$$\sum_{k=1}^{\infty} \frac{1}{k^a} \psi^{(b)}(k) = (-1)^{b+1} b! \left[\zeta(a) \zeta(1+b) - \sigma_h(1+b, a) \right] \quad . \tag{4.35}$$

Here we have [23]:

$$\psi^{(b)}(x) = (-1)^{b+1} b! \sum_{k=0}^{\infty} \frac{1}{(x+k)^{b+1}} = (-1)^{b+1} b! \zeta(b+1, x) \quad , \tag{4.36}$$

which for $x \in \mathbf{N}$ may be also conveniently written:

$$\psi^{(b-1)}(n) = (-1)^b (b-1)! \left[\zeta(b) - H_{n-1,b} \right] \quad , \quad b > 1 \quad . \tag{4.37}$$

With (4.31) Eq. (4.35) boils down to

$$\sum_{k=1}^{\infty} \frac{1}{k^{1+b}} \psi^{(b)}(k) = \frac{1}{2} (-1)^{b+1} b! \left[\zeta(2b+2) + \zeta(b+1)^2 \right] \quad ,$$

for the case $a = 1 + b$.

Before we proceed, we shall present the series:

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)^\alpha} = \alpha - \sum_{i=2}^{\alpha} \zeta(i) . \quad (4.38)$$

This identity may be proven by induction, with noting: $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \psi(1) + \gamma_E + 1 = 1$; $\psi(1) = -\gamma_E$. Next, we present the identity:

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n(n+1)^\alpha} H_{n-1} &= \frac{1}{2} \alpha(\alpha+1) - \frac{1}{2} \sum_{i=2}^{\alpha} i \zeta(i+1) - \sum_{i=2}^{\alpha} (\alpha-i+1) \zeta(i) \\ &+ \frac{1}{2} \sum_{k=0}^{\alpha-3} \sum_{i \geq 0}^k \zeta(2+i) \zeta(2+k-i) . \end{aligned} \quad (4.39)$$

Again, this identity may be proven by induction, with noting $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} H_{n-1} = 1$. Some special cases are:

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{(n+1)^4} H_{n-1} &= -4 + \zeta(2) + \zeta(3) + \zeta(4) - \zeta(2) \zeta(3) + 2 \zeta(5) , \\ \sum_{n=1}^{\infty} \frac{1}{n(n+1)^4} H_{n-1} &= 10 - 3 \zeta(2) - 3 \zeta(3) - \frac{5}{4} \zeta(4) + \zeta(2) \zeta(3) - 2 \zeta(5) , \\ \sum_{n=1}^{\infty} \frac{1}{n^2(n+1)^4} H_{n-1} &= -20 + 6 \zeta(2) + 7 \zeta(3) + \frac{3}{2} \zeta(4) - \zeta(2) \zeta(3) + 2 \zeta(5) . \end{aligned} \quad (4.40)$$

For the power series expansion of the hypergeometric functions ${}_4F_3$ we shall need the two series:

$$\begin{aligned} (i) \quad \sum_{n=1}^{\infty} \frac{\psi^{(1)}(n)}{n(n+1)^3} &= 10 - 3 \zeta(2) - \frac{1}{2} \zeta(2)^2 - \zeta(3) - \frac{1}{2} \zeta(4) + 2 \zeta(2) \zeta(3) - \frac{11}{2} \zeta(5) , \\ (ii) \quad \sum_{n=1}^{\infty} \frac{H_{n-1}^2}{n(n+1)^3} &= 10 - \frac{1}{2} \zeta(2)^2 - 5 \zeta(3) - 2 \zeta(4) - \zeta(2) \zeta(3) + \frac{3}{2} \zeta(5) . \end{aligned} \quad (4.41)$$

Both series may be determined by making use of the decomposition:

$$\frac{1}{n(n+1)^3} = \frac{1}{n(n+1)} - \frac{1}{(n+1)^2} - \frac{1}{(n+1)^3} . \quad (4.42)$$

Let us first prove the series:

$$\sum_{n=1}^{\infty} \frac{H_{n,2}}{n(n+1)^3} = \zeta(3) - \frac{1}{2} \zeta(2)^2 + \frac{1}{2} \zeta(4) + \frac{11}{2} \zeta(5) - 3 \zeta(2) \zeta(3) . \quad (4.43)$$

According to the decomposition (4.42) we need the series:

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{H_{n,2}}{n(n+1)} &= \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right) H_{n,2} = \sum_{n=1}^{\infty} \left(\frac{H_{n,2}}{n} - \frac{H_{n-1,2}}{n} \right) = \sum_{n=1}^{\infty} \frac{1}{n^3} = \zeta(3) \ , \\
\sum_{n=1}^{\infty} \frac{H_{n,2}}{(n+1)^2} &= \sum_{n=1}^{\infty} \frac{H_{n-1,2}}{n^2} = \sum_{n=1}^{\infty} \frac{H_{n,2}}{n^2} - \sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{1}{2} \zeta(2)^2 - \frac{1}{2} \zeta(4) \ , \\
\sum_{n=1}^{\infty} \frac{H_{n,2}}{(n+1)^3} &= \sigma_h(2, 3) = -\frac{11}{2} \zeta(5) + 3 \zeta(2) \zeta(3) \ .
\end{aligned} \tag{4.44}$$

After putting together all three series along (4.42) we obtain (4.43).

To prove the series (i) of Eq. (4.41) we write:

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{\psi^{(1)}(n)}{n(n+1)^3} &= \sum_{n=1}^{\infty} \frac{1}{n(n+1)^3} \sum_{k=0}^{\infty} \frac{1}{(n+k)^2} = \sum_{n=1}^{\infty} \frac{1}{n(n+1)^3} \left(\sum_{k=1}^{\infty} \frac{1}{k^2} - \sum_{k=1}^{n-1} \frac{1}{k^2} \right) \\
&= \zeta(2) \sum_{n=1}^{\infty} \frac{1}{n(n+1)^3} - \sum_{n=1}^{\infty} \frac{H_{n-1,2}}{n(n+1)^3} \\
&= \zeta(2) [3 - \zeta(2) - \zeta(3)] - \sum_{n=1}^{\infty} \frac{H_{n,2}}{n(n+1)^3} + \sum_{n=1}^{\infty} \frac{1}{n^3(n+1)^3} \ .
\end{aligned} \tag{4.45}$$

The first term is obtained from (4.38). The second term corresponds to (4.43) and the last sum is straightforward to calculate:

$$\sum_{n=1}^{\infty} \frac{1}{n^3(n+1)^3} = 10 - \pi^2 \ .$$

Hence, after putting all pieces together we obtain the series (i) of Eq. (4.41).

Before we undertake to evaluate series (ii), we shall prove the series:

$$\sum_{n=1}^{\infty} \frac{H_n^2}{n(n+1)^3} = 3 \zeta(3) - \frac{3}{2} \zeta(4) - \frac{1}{2} \zeta(2)^2 + \frac{3}{2} \zeta(5) - \zeta(2) \zeta(3) \ . \tag{4.46}$$

Similar as before, we shall calculate the series:

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{H_n^2}{n(n+1)} &= \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+1} \right) H_n^2 = \sum_{n=1}^{\infty} \left(\frac{H_n^2}{n} - \frac{H_{n-1}^2}{n} \right) \\
&= 2 \sum_{n=1}^{\infty} \frac{H_n}{n^2} - \sum_{n=1}^{\infty} \frac{1}{n^3} = 3 \zeta(3) \ , \\
\sum_{n=1}^{\infty} \frac{H_n^2}{(n+1)^2} &= s_h(2, 2) = \frac{3}{2} \zeta(4) + \frac{1}{2} \zeta(2)^2 \ , \\
\sum_{n=1}^{\infty} \frac{H_n^2}{(n+1)^3} &= s_h(2, 3) \ .
\end{aligned} \tag{4.47}$$

The Euler sum $s_h(2, 3)$ may not be found in the list (4.28). Instead we may determine it by using some identities from [30]. In this work, the relation

$$\sum_{n=1}^{\infty} \frac{1}{n^a} H_{n-1,b} H_{n-1,c} = \zeta(a, b, c) + \zeta(a, c, b) + \sigma_h(b + c, a) , \quad (4.48)$$

involving the so-called *triple Euler sum*

$$\zeta(a, b, c) = \sum_{n=1}^{\infty} \sum_{m=1}^{n-1} \sum_{k=1}^{m-1} \frac{1}{n^a m^b k^c} , \quad (4.49)$$

has been proven. Furthermore, in *Ref.* [30] a list of values for $\zeta(a, b, c)$ has been given, in particular $\zeta(3, 1, 1) = 2 \zeta(5) - \zeta(2) \zeta(3)$. Hence we have:

$$s_h(2, 3) = \sum_{n=1}^{\infty} \frac{H_n^2}{(n+1)^3} = \sum_{n=1}^{\infty} \frac{H_{n-1}^2}{n^3} = 2 \zeta(3, 1, 1) + \sigma_h(2, 3) = -\frac{3}{2} \zeta(5) + \zeta(2) \zeta(3) . \quad (4.50)$$

From (4.47), together with the decomposition (4.42), we obtain (4.46).

Finally, we undertake to prove (ii) of *Eq.* (4.41). We first write:

$$\sum_{n=1}^{\infty} \frac{H_{n-1}^2}{n(n+1)^3} = \sum_{n=1}^{\infty} \frac{H_n^2}{n(n+1)^3} + \sum_{n=1}^{\infty} \frac{1}{n^3(n+1)^3} - 2 \sum_{n=1}^{\infty} \frac{H_n}{n^2(n+1)^3} . \quad (4.51)$$

The first two sums have been presented above, while for the last sum we have:

$$\sum_{n=1}^{\infty} \frac{H_n}{n^2(n+1)^3} = -3 \zeta(2) + 4 \zeta(3) + \frac{1}{4} \zeta(4) .$$

After inserting all sums into (4.51) we arrive at the final result (ii) of *Eq.* (4.41).

The relation (4.48) is very useful. For $b = c$ it boils down to:

$$\sum_{n=1}^{\infty} \frac{H_{n-1,b}^2}{n^a} = 2 \zeta(a, b, b) + \sigma_h(2b, a) . \quad (4.52)$$

From *Ref.* [30] we have

$$\zeta(a, a, a) = \frac{1}{6} \zeta(a)^3 - \frac{1}{2} \zeta(a) \zeta(2a) + \frac{1}{3} \zeta(3a) , \quad (4.53)$$

which allows to derive the following identity from (4.52):

$$\sum_{n=1}^{\infty} \frac{H_{n-1,2}^2}{n^2} = 2 \zeta(2, 2, 2) + \sigma_h(4, 2) = \frac{1}{3} \zeta(2)^3 - \zeta(3)^2 - \frac{8}{3} \zeta(2) \zeta(4) + \frac{17}{3} \zeta(6) . \quad (4.54)$$

4.4. Momentum expansion of hypergeometric functions ${}_3F_2$ and ${}_4F_3$

In the previous subsection we have compiled material, which one needs to expand Gaussian hypergeometric functions (4.5) w.r.t. to small parameter a_i and b_j . To obtain the α' -expansion of a five-point string S -matrix one needs the power series expansion of (4.2) w.r.t. the small parameter a, \dots, e . Let us present some important examples, which prove to be useful in any five-point scattering process on the disk (*cf.* appendix A). With the previous results and appendix B we may derive the following two expansions w.r.t. small parameter a, \dots, e

$$\begin{aligned}
C(a, b, c, d, e) &= \frac{\Gamma(1+a) \Gamma(1+b) \Gamma(1+c) \Gamma(1+d)}{\Gamma(2+a+c) \Gamma(2+b+d)} {}_3F_2 \left[\begin{matrix} 1+a, 1+b, -e \\ 2+a+c, 2+b+d \end{matrix} \right] \\
&= 1 - a - b - c - d - 2e + e \zeta(2) + a^2 + ab + b^2 + bc + c^2 + ad + cd + d^2 \\
&\quad + 3ae + 3be + 3ce + 3de + 3e^2 + 2ac + 2bd \\
&\quad - (ac + bd + ae + be + e^2) \zeta(2) - (ae + be + 2ce + 2de + e^2) \zeta(3) \\
&\quad - a^3 - a^2b - ab^2 - b^3 - 3a^2c - 2abc - b^2c - 3ac^2 - bc^2 - c^3 - a^2d - 2abd - 3b^2d \\
&\quad - 2acd - 2bcd - c^2d - ad^2 - 3bd^2 - cd^2 - d^3 - 4a^2e - 4abe - 4b^2e - 8ace \\
&\quad - 4bce - 4c^2e - 4ade - 8bde - 4cde - 4d^2e - 6ae^2 - 6be^2 - 6ce^2 - 6de^2 - 4e^3 \\
&\quad + (a^2c + abc + ac^2 + abd + b^2d + acd + bcd + bd^2 + a^2e + abe + b^2e + 3ace + 3bde \\
&\quad - cde + 2ae^2 + 2be^2 + e^3) \zeta(2) + (a^2c + ac^2 + b^2d + bd^2 + a^2e + abe + b^2e + 2ace \\
&\quad + 2bce + 2ade + 2bde + 2ae^2 + 2be^2 + 2ce^2 + 2de^2 + e^3) \zeta(3) \\
&\quad + \left(a^2e + abe + b^2e + 3ace + \frac{5bce}{4} + 3c^2e + \frac{5ade}{4} + 3bde + \frac{17cde}{4} + 3d^2e + \frac{ae^2}{4} \right. \\
&\quad \left. + \frac{be^2}{4} + 3ce^2 + 3de^2 + e^3 \right) \zeta(4) + \dots,
\end{aligned} \tag{4.55}$$

and

$$\begin{aligned}
C(a, b, c, d, e-1) &= \frac{\Gamma(1+a) \Gamma(1+b) \Gamma(1+c) \Gamma(1+d)}{\Gamma(2+a+c) \Gamma(2+b+d)} {}_3F_2 \left[\begin{matrix} 1+a, 1+b, 1-e \\ 2+a+c, 2+b+d \end{matrix} \right] \\
&= \zeta(2) - (a+b+2c+2d+e) \zeta(3) \\
&\quad + \left(a^2 + ab + b^2 + \frac{ac}{2} + \frac{5bc}{4} + 3c^2 + \frac{5ad}{4} + \frac{bd}{2} + 3d^2 + \frac{ae}{4} + \frac{be}{4} + 3ce \right. \\
&\quad \left. + \frac{17cd}{4} + 3de + e^2 \right) \zeta(4) + \dots,
\end{aligned} \tag{4.56}$$

for the two integrals $C(a, b, c, d, e)$ and $C(a, b, c, d, e - 1)$, defined in Eq. (4.4), respectively. We now want to propose an interesting identity

$$\begin{aligned}
{}_3F_2 \left[\begin{matrix} \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2 \end{matrix} \right] &= \frac{1}{\alpha_1 \alpha_2 (\alpha_1 - \beta_1) (\alpha_2 - \beta_2) (\alpha_1 + \alpha_2 + \alpha_3 - \beta_1 - \beta_2)} \\
&= \left\{ (1 - \alpha_3 + \beta_1) (1 - \alpha_3 + \beta_2) [\beta_1 \beta_2 (\alpha_1 + \alpha_2 + \alpha_3 - \beta_1 - \beta_2) - \alpha_1 \alpha_2 \alpha_3] \right. \\
&\times C(\alpha_1, \alpha_2, \beta_1 - \alpha_1, \beta_2 - \alpha_2, -\alpha_3) \\
&- \alpha_3 [\alpha_1^2 (\alpha_2 - \beta_1) (\alpha_2 - \beta_2) - \beta_1 \beta_2 (\alpha_2 + \alpha_3 - \beta_1 - \beta_2) (1 - \alpha_2 - \alpha_3 + \beta_1 + \beta_2) + \\
&+ \alpha_1 \alpha_2 \alpha_3 - \alpha_1 \alpha_2 \alpha_3^2 - \alpha_1 \alpha_2^2 \beta_1 + \alpha_1 \alpha_2 \beta_1^2 - \alpha_1 \alpha_2^2 \beta_2 - \alpha_1 \beta_1 \beta_2 \\
&+ 3 \alpha_1 \alpha_2 \beta_1 \beta_2 + 2 \alpha_1 \alpha_3 \beta_1 \beta_2 - 2 \alpha_1 \beta_1^2 \beta_2 + \alpha_1 \alpha_2 \beta_2^2 - 2 \alpha_1 \beta_1 \beta_2^2] \\
&\times C(\alpha_1, \alpha_2, \beta_1 - \alpha_1, \beta_2 - \alpha_2, -\alpha_3 - 1) \left. \right\} \frac{\Gamma(\beta_1) \Gamma(\beta_2)}{\Gamma(\alpha_1) \Gamma(\alpha_2) \Gamma(\beta_1 - \alpha_1) \Gamma(\beta_2 - \alpha_2)} \\
&= 1 - \frac{\alpha_1 \alpha_2 \alpha_3}{\beta_1 \beta_2 (\alpha_1 + \alpha_2 + \alpha_3 - \beta_1 - \beta_2)} \\
&\times \left\{ 1 + [(\alpha_2 - \beta_1) (\alpha_3 - \beta_1) + \alpha_1 (\alpha_2 + \alpha_3 - \beta_1 - \beta_2) - (\alpha_2 + \alpha_3 - \beta_1) \beta_2 + \beta_2^2] \zeta(2) \right. \\
&+ [\alpha_1^2 \alpha_2 + \alpha_1 \alpha_2^2 + \alpha_1^2 \alpha_3 + 4 \alpha_1 \alpha_2 \alpha_3 + \alpha_2^2 \alpha_3 + \alpha_1 \alpha_3^2 + \alpha_2 \alpha_3^2 - \alpha_1^2 \beta_1 - 4 \alpha_1 \alpha_2 \beta_1 \\
&- \alpha_2^2 \beta_1 - 4 \alpha_1 \alpha_3 \beta_1 - 4 \alpha_2 \alpha_3 \beta_1 - \alpha_3^2 \beta_1 + 3 \alpha_1 \beta_1^2 + 3 \alpha_2 \beta_1^2 + 3 \alpha_3 \beta_1^2 - 2 \beta_1^3 - \alpha_1^2 \beta_2 \\
&- 4 \alpha_1 \alpha_2 \beta_2 - \alpha_2^2 \beta_2 - 4 \alpha_1 \alpha_3 \beta_2 - 4 \alpha_2 \alpha_3 \beta_2 - \alpha_3^2 \beta_2 + 4 \alpha_1 \beta_1 \beta_2 + 4 \alpha_2 \beta_1 \beta_2 \\
&+ 4 \alpha_3 \beta_1 \beta_2 - 3 \beta_1^2 \beta_2 + 3 \alpha_1 \beta_2^2 + 3 \alpha_2 \beta_2^2 + 3 \alpha_3 \beta_2^2 - 3 \beta_1 \beta_2^2 - 2 \beta_2^3] \zeta(3) + \dots \left. \right\}, \tag{4.57}
\end{aligned}$$

which will be proven in section 5. In fact, it pops up automatically as a byproduct of our five-gluon scattering amplitude, once we perform the calculation in the way described in section 2. The above equation relates the hypergeometric function ${}_3F_2 \left[\begin{matrix} \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2 \end{matrix} \right]$ to the two expressions (4.55) and (4.56) from before. The power series of the hypergeometric function

$$\begin{aligned}
{}_3F_2 \left[\begin{matrix} \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2 \end{matrix} \right] &= \frac{\Gamma(\beta_1) \Gamma(\beta_2)}{\Gamma(\alpha_1) \Gamma(\alpha_2) \Gamma(\beta_1 - \alpha_1) \Gamma(\beta_2 - \alpha_2)} \\
&\times C(\alpha_1 - 1, \alpha_2 - 1, \beta_1 - \alpha_1 - 1, \beta_2 - \alpha_2 - 1, -\alpha_3) . \tag{4.58}
\end{aligned}$$

has meromorphic poles in β_1, β_2 and $\alpha_1 + \alpha_2 + \alpha_3 - \beta_1 - \beta_2$, while the functions (4.55) and (4.56) are finite. Hence the above identity yields a non-trivial relation between a singular hypergeometric function and two non-singular ones. From the previous results (4.55) and (4.56) we may easily deduce the power series expansion of the singular hypergeometric function ${}_3F_2 \left[\begin{matrix} \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2 \end{matrix} \right]$. In fact, from (4.21) one deduces, that in that case non-convergent sums would appear, if one performed a naive expansion. The techniques, how to deal with

such cases, have been pioneered in *Ref.* [31]. However, it is quite intriguing (*cf.* section 5), that from string–theory we find powerful identities to avoid this problem.

Finally, the momentum expansion of ${}_4F_3 \left[\begin{matrix} 1+b, 1+a, 1+c, -j \\ 2+b+e, 2+a+d, 2+c+f \end{matrix} \right]$, relevant to six open string amplitudes, will be given in appendix C.

4.5. Multiple zeta values, Euler–Zagier sums and momentum expansion of $F^{(3)}$

In this subsection we present the tools relevant for the momentum expansion of the triple hypergeometric function $F^{(3)}$, given in *Eq.* (4.16). Expanding the function $F^{(3)}$ w.r.t. the parameter a, \dots, j generically leads to triple infinite sums over positive integers. These sums represent generalizations of the multiple zeta values of length k :

$$\zeta(s_1, \dots, s_k) = \sum_{n_1 > \dots > n_k > 0} \prod_{j=1}^k \frac{1}{n_j^{s_j}} = \sum_{n_1, \dots, n_k=1}^{\infty} \prod_{j=1}^k \left(\sum_{i=j}^k n_i \right)^{-s_j}, \quad (4.59)$$

with $s_1 \geq 2$, $s_2, \dots, s_k \geq 1$. The multiple zeta value are related to the multiple polylogarithm

$$\mathcal{L}i_{s_1, \dots, s_k}(x_1, \dots, x_k) = \sum_{n_1 > \dots > n_k > 0} \prod_{j=1}^k \frac{x_j^{n_j}}{n_j^{s_j}}, \quad (4.60)$$

for $x_j = 1$, *i.e.* $\zeta(s_1, \dots, s_k) = \mathcal{L}i_{s_1, \dots, s_k}(1, \dots, 1)$. The latter reduces to the Riemann–Zeta function for $k = 1$, *i.e.* $\mathcal{L}i_a(1) = \zeta(a)$. The case $k = 3$ has already occurred in (4.49) and representative examples may be found in *Ref.* [30]. Furthermore, the multiple zeta values (4.59) have the integral representation [32]

$$\zeta(s_1, \dots, s_k) = \int_1^{\infty} \frac{dx_1}{x_1} \dots \int_1^{\infty} \frac{dx_k}{x_k} \prod_{j=1}^k \frac{1}{\Gamma(s_j)} \frac{(\ln x_j)^{s_j-1}}{\prod_{i=1}^j x_i - 1}. \quad (4.61)$$

Our case of interest is $k = 3$:

$$\zeta(s_1, s_2, s_3) = -\frac{(-1)^{s_1+s_2+s_3}}{\Gamma(s_1) \Gamma(s_2) \Gamma(s_3)} \int_0^1 dx \int_0^1 dy \int_0^1 dz x^2 y \frac{(\ln x)^{s_1-1} (\ln y)^{s_2-1} (\ln z)^{s_3-1}}{(1-x)(1-xy)(1-xyz)}. \quad (4.62)$$

It is this relation (4.62), which gives (formally) the link between the two expressions (3.31) and (4.16) after expanding them w.r.t. to small parameter a, \dots, j .

On the other hand, after looking closer at the nine polynomials showing up in (3.31) we realize, that (4.62) is not of considerable help. In fact, In the following we shall deal

with some generalizations of the sum (4.59). These sums are called *Euler–Zagier double series* or *Witten zeta function*¹³ [33,34]. One example is the following double sum:

$$W(r, s, t) = \sum_{m,n=1}^{\infty} \frac{1}{n^r m^s (m+n)^t} . \quad (4.63)$$

It enjoys the *Pascal triangle recurrence* relation

$$W(r, s, t) = W(r-1, s, t+1) + W(r, s-1, t+1) , \quad (4.64)$$

and the useful relations [35]:

$$2 W(a-2, 1, 1) - W(1, 1, a-2) = 2 \zeta(a) ,$$

$$W(1, 1, a-2) = (a-1) \zeta(a) - \sum_{i=2}^{a-2} \zeta(i) \zeta(a-i) , \quad (4.65)$$

$$W(a-2, 1, 1) = \frac{1}{2} W(1, 1, a-2) + \zeta(a) ,$$

$$W(1, 0, a-1) = \frac{1}{2} W(1, 1, a-2) .$$

From the latter relations we may deduce *e.g.*

$$W(1, 1, 1) = 2 \zeta(3) \quad , \quad W(1, 1, 2) = \frac{1}{2} \zeta(4) \quad (4.66)$$

among other remarkable identities. An immediate consequence of (4.65) is the identity ($\alpha \geq 1$):

$$\sum_{m,n=1}^{\infty} \frac{1}{m(1+n)(1+m+n)^\alpha} = -\alpha + (\alpha+1) \zeta(\alpha+2) + \sum_{i=2}^{\alpha} \zeta(i) - \sum_{i=2}^{\alpha} \zeta(i) \zeta(\alpha+2-i) , \quad (4.67)$$

which may be proven by noting $\sum_{m,n=1}^{\infty} \frac{1}{m(1+n)(1+m+n)^\alpha} = W(1, 1, \alpha) - \sum_{m=1}^{\infty} \frac{1}{m(m+1)^\alpha}$ and using (4.38). We may derive an other important series, similar to (4.67), namely

$$\begin{aligned} \sum_{m,n=1}^{\infty} \frac{1}{m(1+m+n)^\alpha} &= \sum_{\substack{m,n=1 \\ n>m}}^{\infty} \frac{1}{m(n+1)^\alpha} = \sum_{n=1}^{\infty} \frac{H_{n-1}}{(n+1)^\alpha} = \sum_{n=1}^{\infty} \frac{H_n}{(n+1)^\alpha} - \sum_{n=1}^{\infty} \frac{1}{n(n+1)^\alpha} \\ &= -\alpha + \frac{1}{2} \alpha \zeta(\alpha+1) + \sum_{i=2}^{\alpha} \zeta(i) - \frac{1}{2} \sum_{i=1}^{\alpha-2} \zeta(\alpha-i) \zeta(i+1) , \end{aligned} \quad (4.68)$$

¹³ These zeta functions represent sums over all finite-dimensional representations ρ of a semi-simple Lie algebra \mathfrak{g} : $\zeta_{\mathfrak{g}}(s) = \sum_{\rho} [\dim(\rho)]^{-s}$. *E.g.*: $\zeta_{sl(2)}(s) = \zeta(s)$ and $\zeta_{sl(3)}(s) = 2^s W(s, s, s)$. Special values of these zeta functions calculate the volume of certain moduli spaces of vector bundles of curves in analogy as $\zeta(2g)$ is proportional to the Euler characteristic of the moduli space of Riemann surfaces of genus g .

where we have used *Eqs.* (4.38) and (4.29). Similarly, we may deduce:

$$\sum_{m,n=1}^{\infty} \frac{1}{(1+m)(1+m+n)^{\alpha}} = 1 + \frac{1}{2} \alpha \zeta(\alpha+1) - \zeta(\alpha) - \frac{1}{2} \sum_{i=1}^{\alpha-2} \zeta(\alpha-i) \zeta(i+1). \quad (4.69)$$

After subtracting the two equations (4.68) and (4.69), we obtain:

$$\sum_{m,n=1}^{\infty} \frac{1}{m(1+m)(1+m+n)^{\alpha}} = -\alpha - 1 + 2 \zeta(\alpha) + \sum_{i=2}^{\alpha-1} \zeta(i). \quad (4.70)$$

The next type of sum we shall be concerned with is:

$$\sum_{m,n=1}^{\infty} \frac{1}{n(1+m)(1+m+n)(2+m+n)} = 2 \zeta(3) - \frac{9}{4}. \quad (4.71)$$

This equality may be proven after performing the partial fraction decomposition

$\frac{1}{(1+m+n)(2+m+n)} = \frac{1}{(1+m+n)} - \frac{1}{(2+m+n)}$ and determining the two series

$$\begin{aligned} \sum_{m,n=1}^{\infty} \frac{1}{n(1+m)(1+m+n)} &= -1 + 2 \zeta(3), \\ \sum_{m,n=1}^{\infty} \frac{1}{n(1+m)(2+m+n)} &= \sum_{m,n=1}^{\infty} \frac{1}{nm(1+m+n)} - \sum_{n=1}^{\infty} \frac{1}{n(n+2)} = 2 - \frac{3}{4} = \frac{5}{4}, \end{aligned} \quad (4.72)$$

which follow from (4.67) and:

$$\sum_{m,n=1}^{\infty} \frac{1}{nm(1+m+n)} = \sum_{m,n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{1+m+n} \right) \frac{1}{m(m+1)} = \sum_{m=1}^{\infty} \frac{H_{m+1}}{m(m+1)} = 2, \quad (4.73)$$

respectively. Furthermore, we present the identity

$$\sum_{m,n=1}^{\infty} \frac{1}{m n (1+m) (1+n) (1+m+n)} = 5 - 4 \zeta(3), \quad (4.74)$$

which may be proven after partial fraction decomposition

$$\frac{1}{m n (1+m) (1+n) (1+m+n)} = \left(\frac{1}{m} - \frac{1}{1+m} \right) \left(\frac{1}{n} - \frac{1}{1+n} \right) \frac{1}{1+m+n}$$

and determining the three sums:

$$\begin{aligned}
\sum_{m,n=1}^{\infty} \frac{1}{m n (1+m+n)} &= 2, \\
\sum_{m,n=1}^{\infty} \frac{1}{m (1+n) (1+m+n)} &= W(1,1,1) - \sum_{m=1}^{\infty} \frac{1}{m (m+1)} = 2 \zeta(3) - 1, \\
\sum_{m,n=1}^{\infty} \frac{1}{(1+m) (1+n) (1+m+n)} &= \sum_{m,n=1}^{\infty} \frac{1}{n (n+1)} \left(\frac{1}{m+1} - \frac{1}{m+n+1} \right) \\
&= \sum_{n=1}^{\infty} \frac{1}{n (n+1)} (H_{n+1} - 1) = 1.
\end{aligned} \tag{4.75}$$

The first sum follows from (4.73) and the second from (4.66) or (4.67). From *Eqs.* (4.75) we may also derive

$$\sum_{m,n=1}^{\infty} \frac{1}{(1+m) n (1+n) (1+m+n)} = 2 \zeta(3) - 2, \tag{4.76}$$

after performing the partial fraction decomposition:

$$\frac{1}{(1+m) n (1+n) (1+m+n)} = \left(\frac{1}{n} - \frac{1}{1+n} \right) \frac{1}{(1+m) (1+m+n)}.$$

Moreover, we prove the two identities:

$$\begin{aligned}
\sum_{m,n=1}^{\infty} \frac{1}{n m (1+m) (1+m+n)^2} &= 8 - 3 \zeta(2) - 2 \zeta(3) - \frac{1}{2} \zeta(4), \\
\sum_{m,n=1}^{\infty} \frac{1}{m (2+n) (1+m+n)^2} &= -\frac{3}{2} \zeta(2) + 3 \zeta(3) - 1.
\end{aligned} \tag{4.77}$$

After relabeling m and performing partial fraction decomposition the first sum may be casted into:

$$\sum_{\substack{n=1 \\ m \geq 2}}^{\infty} \frac{1}{n m (m-1) (m+n)^2} = \sum_{m,n=1}^{\infty} \frac{1}{m n (1+m+n)^2} - \sum_{m,n=1}^{\infty} \frac{1}{m n (m+n)^2} + \sum_{n=1}^{\infty} \frac{1}{n (n+1)^2}.$$

According to (4.66) and (4.38), the last two sums give $W(1,1,2) = \frac{1}{2}\zeta(4)$ and $2 - \zeta(2)$, respectively, while the first sum may be written:

$$\begin{aligned}
\sum_{m,n=1}^{\infty} \frac{1}{m n (1+m+n)^2} &= \sum_{m,n=1}^{\infty} \left(\frac{1}{m} + \frac{1}{n} \right) \frac{1}{(m+n) (1+m+n)^2} \\
&= 2 \sum_{m,n=1}^{\infty} \frac{1}{m (m+n) (1+m+n)^2} = 2 \sum_{n=1}^{\infty} \frac{H_{n-1}}{n(n+1)^2} = 2 [3 - \zeta(3) - \zeta(2)].
\end{aligned}$$

On the other hand, the second sum of (4.77) may be written:

$$\begin{aligned} \sum_{m,n=1}^{\infty} \frac{1}{m(2+n)(1+m+n)^2} &= 2 \sum_{m,n=1}^{\infty} \frac{1}{m(m+n)(m+n-1)^2} - \sum_{m=1}^{\infty} \frac{1}{2m(1+m)^2} \\ &- \sum_{m=1}^{\infty} \frac{1}{m^3} = -1 + \frac{1}{2} \zeta(2) - \zeta(3) + 2 \sum_{n=1}^{\infty} \frac{H_n}{n^2(1+n)} = -\frac{3}{2} \zeta(2) + 3 \zeta(3) - 1 . \end{aligned}$$

Besides, let us check the following sum:

$$\sum_{m,n=1}^{\infty} \frac{1}{(1+n)m^2(m+n)^2} = -\frac{1}{2} \zeta(4) + \frac{1}{2} \zeta(2)^2 + 5 \zeta(3) - 4 \zeta(2) . \quad (4.78)$$

After writing the numerator $1 = 1+n+m-m-n$ we may divide the sum into a combination of the following three pieces:

$$\begin{aligned} \sum_{m,n=1}^{\infty} \frac{1}{m^2(m+n)^2} &= \sum_{m<n} \frac{1}{m^2 n^2} = \zeta(2,2) = \frac{1}{2} \zeta(2)^2 - \frac{1}{2} \zeta(4) , \\ \sum_{m,n=1}^{\infty} \frac{1}{m(1+n)(m+n)^2} &= \sum_{m,n=1}^{\infty} \frac{1}{m(2+n)(1+m+n)^2} + \frac{1}{2} \sum_{m=1}^{\infty} \frac{1}{m(1+m)^2} \\ &= -2 \zeta(2) + 3 \zeta(3) , \\ \sum_{m,n=1}^{\infty} \frac{1}{m^2(1+n)(m+n)} &= \sum_{m,n=1}^{\infty} \frac{1}{m^2} \frac{1}{n(1+n)} - \sum_{n=1}^{\infty} \frac{1}{n(1+n)^2} \\ &- \sum_{m,n=1}^{\infty} \frac{1}{n(1+n)(1+m)(1+m+n)} = 2 \zeta(2) - 2 \zeta(3) . \end{aligned} \quad (4.79)$$

While the first sum of (4.79) is standard (*cf. Eq. (4.49)*), for the second sum the result (4.77) and for the third sum the identity (4.76) have been used in addition to *Eq. (4.38)*. After putting these sums together we prove (4.78).

So far we have listed all sums we shall need for expanding the six functions (3.50) up to first order in the momenta s_i . To go beyond this order we have also calculated the following three triple sums:

$$\begin{aligned} \sum_{m_i=1}^{\infty} \frac{m_3}{m_1 m_2 (m_1+m_3)(m_2+m_3)(m_1+m_2+m_3)} &= \frac{7}{4} \zeta(4) , \\ \sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (1+m_1+m_3)(m_2+m_3)(m_1+m_2+m_3)} &= \frac{19}{4} \zeta(4) - 4 \zeta(3) , \\ \sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (m_1+m_3)(m_2+m_3)(1+m_1+m_2+m_3)} &= \frac{17}{4} \zeta(4) + 2 \zeta(3) - \zeta(2) - 5 . \end{aligned} \quad (4.80)$$

These two sums may be proven, by first computing the following somewhat simpler triple sums:

$$\begin{aligned}
\sum_{m_i=1}^{\infty} \frac{1}{(1+m_1+m_3)(m_2+m_3)(m_1+m_2+m_3)^2} &= -\frac{13}{4} \zeta(4) - 3 \zeta(3) + 2 \zeta(2) + \frac{3}{2} \zeta(2)^2, \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1(1+m_1+m_3)(m_2+m_3)(m_1+m_2+m_3)} &= -2 \zeta(4) - 2 \zeta(3) + \zeta(2) + \frac{3}{2} \zeta(2)^2, \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (1+m_1+m_3)(m_2+m_3)} &= 2 \zeta(3) + \zeta(2), \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (m_2+m_3)(m_1+m_2+m_3)} &= 8 \zeta(4) - 2 \zeta(2)^2, \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (m_2+m_3)(m_1+m_3)} &= \frac{11}{2} \zeta(4) - \zeta(2)^2, \\
\sum_{m_i=1}^{\infty} \frac{1}{m_2 (m_1+m_3)(m_2+m_3)(m_1+m_2+m_3)} &= -\frac{5}{2} \zeta(4) + \frac{3}{2} \zeta(2)^2, \\
\sum_{m_i=1}^{\infty} \frac{1}{(m_1+m_3)(m_2+m_3)(m_1+m_2+m_3)^2} &= \frac{17}{4} \zeta(4) - 2 \zeta(2), \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (m_1+m_3)(m_2+m_3)(1+m_1+m_2+m_3)} &= \frac{17}{4} \zeta(4) + 2 \zeta(3) - \zeta(2) - 5, \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 m_2 (1+m_2)(m_2+m_3)(1+m_1+m_2+m_3)} &= 5 - 2 \zeta(3) - \zeta(2).
\end{aligned} \tag{4.81}$$

In addition, we have also derived the following sums involving the harmonic number

$$\begin{aligned}
\sum_{m_i=1}^{\infty} \frac{H_{m_1+m_3}}{m_1(1+m_1+m_3)^2} &= -\zeta(2) + \frac{1}{2} \zeta(2)^2 + \zeta(3) + \frac{3}{2} \zeta(4), \\
\sum_{m_i=1}^{\infty} \frac{H_{m_3}}{m_1(1+m_1+m_3)^2} &= \frac{9}{2} \zeta(4) - \zeta(2)^2, \\
\sum_{m_i=1}^{\infty} \frac{H_{m_1-1}}{m_1(1+m_1+m_3)^2} &= -3 + \zeta(2) + \zeta(3) + \zeta(4), \\
\sum_{m_i=1}^{\infty} \frac{H_{m_3}}{m_1(1+m_3)(1+m_1+m_3)(2+m_1+m_3)} &= 1 + \zeta(2) - 2 \zeta(3),
\end{aligned} \tag{4.82}$$

$$\begin{aligned}
\sum_{m_i=1}^{\infty} \frac{H_{m_1-1}}{m_1 (1+m_3) (1+m_1+m_3) (2+m_1+m_3)} &= \frac{15}{4} - \zeta(2) - 2 \zeta(3) , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_1+m_3}}{m_1 (1+m_3) (1+m_1+m_3) (2+m_1+m_3)} &= -\frac{3}{2} - \frac{5}{2} \zeta(2) + \frac{11}{2} \zeta(4) , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_1+m_2-1}}{m_1 (m_1+m_2)^2} &= \frac{3}{2} \zeta(4) + \frac{1}{2} \zeta(2)^2 , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_2}}{m_1 (m_1+m_2)^2} &= 5 \zeta(4) - \zeta(2)^2 , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_2}}{m_1 m_2 (m_1+m_2)} &= \frac{11}{2} \zeta(4) - \frac{1}{2} \zeta(2)^2 , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_2}}{m_2 (m_1+m_2)^2} &= \frac{1}{2} \zeta(4) + \frac{1}{2} \zeta(2)^2 , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_1+m_2-1}}{(1+m_1) (m_1+m_2)^2} &= \frac{7}{4} \zeta(4) + \frac{1}{2} \zeta(2)^2 - \zeta(3) , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_2}}{(1+m_1) (m_1+m_2)^2} &= 5 \zeta(4) - \zeta(2)^2 + 2 \zeta(3) - 2 \zeta(2) , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_3}}{m_2 (2+m_3) (1+m_2+m_3)^2} &= -2 \zeta(2) - \zeta(2)^2 + \frac{23}{4} \zeta(4) , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_2-1}}{m_2 (2+m_3) (1+m_2+m_3)^2} &= -\frac{3}{2} + \frac{1}{2} \zeta(2) - \zeta(2)^2 - \frac{5}{2} \zeta(3) + 6 \zeta(4) , \\
\sum_{m_i=1}^{\infty} \frac{H_{m_2+m_3}}{m_2 (2+m_3) (1+m_2+m_3)^2} &= \frac{23}{4} \zeta(4) - \frac{5}{2} \zeta(2) - \frac{3}{2} \zeta(3) ,
\end{aligned}$$

and finally the following triple sums:

$$\begin{aligned}
\sum_{m_i=1}^{\infty} \frac{1}{m_1 (1+m_1+m_3)^3} &= -3 + \frac{3}{2} \zeta(4) - \frac{1}{2} \zeta(2)^2 + \zeta(2) + \zeta(3) , \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 (1+m_3) (1+m_1+m_3)^2} &= -2 + 3 \zeta(4) - \zeta(2)^2 + \zeta(2) , \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 (1+m_3) (1+m_1+m_3) (2+m_1+m_3)} &= -\frac{9}{4} + 2 \zeta(3) , \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 (1+m_3)^2 (1+m_1+m_3) (2+m_1+m_3)} &= \frac{11}{4} - \frac{\pi^2}{6} + \frac{\pi^4}{72} - 2 \zeta(3) ,
\end{aligned}$$

(4.83)

$$\begin{aligned}
\sum_{m_i=1}^{\infty} \frac{1}{m_1 (1+m_3) (1+m_1+m_3) (2+m_1+m_3)^2} &= -\frac{29}{4} + \frac{\pi^2}{4} + 4 \zeta(3) , \\
\sum_{m_i=1}^{\infty} \frac{1}{m_1 (1+m_3) (1+m_1+m_3)^2 (2+m_1+m_3)} &= \frac{1}{4} + 3 \zeta(4) + \zeta(2) - \zeta(2)^2 - 2 \zeta(3) , \\
\sum_{m_i=1}^{\infty} \frac{1}{m_2 (2+m_3)^2 (1+m_2+m_3)^3} &= -\frac{1}{2} - \frac{15}{4} \zeta(2) + \frac{1}{2} \zeta(2)^2 + 4 \zeta(3) + \frac{1}{2} \zeta(4) , \\
\sum_{m_i=1}^{\infty} \frac{1}{m_2 (2+m_3) (1+m_1+m_2+m_3)^3} &= -\frac{3}{2} + \frac{5}{2} \zeta(2) - \frac{1}{2} \zeta(2)^2 - \frac{7}{2} \zeta(3) + \frac{11}{4} \zeta(4) , \\
\sum_{m_i=1}^{\infty} \frac{1}{(1+m_3) m_2 (2+m_3) (1+m_2+m_3)^2} &= -1 + 3 \zeta(4) + \frac{5}{2} \zeta(2) - 3 \zeta(3) - \zeta(2)^2 .
\end{aligned}$$

These results may be proven through multiple applying partial fractioning and using relations derived above and in subsection 4.3.

Let us move on to the power series expansion w.r.t. small parameter a, \dots, j of the function $F \left[\begin{smallmatrix} a+m_{24}, b+m_{25}, d+m_{26}, e+m_{34}, g+m_{35} \\ c+m_{36}, f+m_{45}, h+m_{46}, j+m_{56} \end{smallmatrix} \right]$, given in Eq. (4.7). However, the parameter m_{ij} are some integers in the range $m_{ij} \in \{0, \pm 1, \pm 2\}$. According to the definition (3.24), this corresponds to the integral (3.31), with the parameter a, \dots, b being some linear combination of kinematic invariants s_i and n_{ij} , given by Eq. (3.32). We shall expand the triple sum (4.16) w.r.t. small parameter a, \dots, j . Generically this leads to a power series in the parameter a, \dots, j with its coefficients given by some triple, double or single Euler sum of the type we have discussed above. Let demonstrate, how it works for the function $F \left[\begin{smallmatrix} a, b, d, e, g \\ c, f, h, j-2 \end{smallmatrix} \right]$. To keep the expressions short, we shall only write up, how the g and h -dependent terms appear up to the second order. With this example the procedure should become clear, how the power series expansion of (4.7) is obtained and how at each order generically some multiple Euler–Zagier sum appears. Of course working out all the dependence in all parameter a, \dots, j is quite cumbersome, since at each order some triple

sums have to be evaluated. For the case at hand we obtain from Eq. (4.16):

$$\begin{aligned}
F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j - 2 \end{matrix} \right] &= \frac{\Gamma(1+d) \Gamma(1+e) \Gamma(1+f)}{\Gamma(-g) \Gamma(-h) \Gamma(2-j)} \sum_{m_i=0}^{\infty} \frac{\Gamma(-g+m_1) \Gamma(-h+m_2) \Gamma(2-j+m_3)}{m_1! m_2! m_3!} \\
&\times \frac{\Gamma(1+m_1+m_2+m_3+b)}{\Gamma(2+m_1+m_2+m_3+b+e)} \frac{\Gamma(1+m_1+m_3+a)}{\Gamma(2+m_1+m_3+a+d)} \frac{\Gamma(1+m_2+m_3+c)}{\Gamma(2+m_2+m_3+c+f)} \\
&= 1 + \underbrace{\sum_{m_3=1}^{\infty} \frac{1}{(1+m_3)^2}}_{=\zeta(2)-1} - (g+h) \left[\underbrace{\sum_{m_1, m_3=1}^{\infty} \frac{1}{m_1 (1+m_1+m_3)^2}}_{=-2+\zeta(2)+\zeta(3)} + \underbrace{\sum_{m_1=1}^{\infty} \frac{1}{m_1 (1+m_1)^2}}_{=2-\zeta(2)} \right] \\
&+ gh \underbrace{\sum_{m_i=1}^{\infty} \frac{m_3}{m_1 m_2 (m_1+m_3) (m_2+m_3) (m_1+m_2+m_3)}}_{=\frac{7}{4} \zeta(4)} \\
&+ (g^2+h^2) \left[\underbrace{\sum_{m_1, m_3=1}^{\infty} \frac{H_{m_1-1}}{m_1 (m_1+m_3)^2}}_{=\zeta(2,1,1)=\zeta(4)} \right] + (a, b, c, d, e, f, j) - \text{dependent terms} + \dots \\
&= \zeta(2) - (g+h) \zeta(3) + \left(g^2 + h^2 + \frac{7}{4} gh \right) \zeta(4) + (a, b, c, d, e, f, j) - \text{dep. terms} + \dots .
\end{aligned} \tag{4.84}$$

We now present the full expansions for the functions (3.54), which we need to obtain the α' -expansion of the six-gluon amplitude. Above we have compiled all the sums necessary to perform this work. We obtain the following results:

$$\begin{aligned}
F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j \end{matrix} \right] &= 1 - a - b - c - d - e - f - 2g - 2h - 3j + (g+h+j) \zeta(2) + j \zeta(3) \\
&+ a^2 + ab + b^2 + ac + bc + c^2 + 2ad + bd + cd + d^2 + ae + 2be + ce \\
&+ de + e^2 + af + bf + 2cf + df + ef + f^2 + 3ag + 3bg + 2cg + 3dg \\
&+ 3eg + 2fg + 3g^2 + 2ah + 3bh + 3ch + 2dh + 3eh + 3fh + 5gh \\
&+ 3h^2 + 4aj + 4bj + 4cj + 4dj + 4ej + 4fj + 8gj + 8hj + 6j^2 \\
&- (ad + be + cf + ag + bg + cg + fg + g^2 + ah + bh + ch + dh + h^2 \\
&+ aj + bj + cj + 3gj + 3hj + 2j^2) \zeta(2) \\
&- [g^2 + 2dg + 2eg + 4hg + h^2 + 2j^2 + 2eh + 2fh \\
&+ 2(d+e+f+g+h)j + a(g+j) + c(h+j) + b(g+h+j)] \zeta(3) \\
&- [j(a+b+c) + \frac{j}{4} (5d+5e+5f+2g+2h+j)] \zeta(4) + \dots .
\end{aligned} \tag{4.85}$$

The other sums encountered above appear in the expansion of the following functions:

$$\begin{aligned}
F \left[\begin{array}{c} a, b, d, e, g \\ c, f, h, j-2 \end{array} \right] &= (1+j) \zeta(2) - (a+b+c+2d+2e+2f+g+h+2j) \zeta(3) \\
&+ j^2 \zeta(2) - j(a+b+c+2d+2e+2f+g+h+2j) \zeta(3) \\
&+ \left\{ a^2 + ba + ca + b^2 + c^2 + 3d^2 + 3e^2 + 3f^2 + g^2 + h^2 + 3dg \right. \\
&+ 3eg + 3eh + 3fh + bc + \frac{da}{2} + \frac{5ea}{4} + \frac{5fa}{4} + \frac{ga}{4} + \frac{ha}{2} + \frac{5ja}{4} \\
&+ \frac{5j^2}{4} + \frac{5bd}{4} + \frac{5cd}{4} + \frac{be}{2} + \frac{5ce}{4} + \frac{17de}{4} + \frac{5bf}{4} + \frac{cf}{2} + \frac{17df}{4} \\
&+ \frac{17ef}{4} + \frac{bg}{4} + \frac{cg}{2} + \frac{5fg}{2} + \frac{bh}{4} + \frac{ch}{4} + \frac{5dh}{2} + \frac{7gh}{4} + \frac{5bj}{4} \\
&+ \left. \frac{5cj}{4} + \frac{17dj}{4} + \frac{17ej}{4} + \frac{17fj}{4} + \frac{5gj}{2} + \frac{5hj}{2} + \frac{5j^2}{4} \right\} \zeta(4) + \dots, \\
F \left[\begin{array}{c} a+1, b, d, e, g \\ c, f, h, j-1 \end{array} \right] &= -1 + \zeta(2) + 3a - b - c + 3d + 2g + j \\
&- (2a - b - c + d - e - f - 2h) \zeta(2) \\
&- (b + c + 2d + 2e + 2f + 2g + 3h + j) \zeta(3) \\
&- 6a^2 + 4(b + c - 3d - 2g - j)a - b^2 - c^2 - 6d^2 + \frac{31g^2}{8} - j^2 \\
&+ 4cd + 3cg - 8dg + 4fg + cj - 4dj + gj + b(-c + 4d + 2g + j) \\
&+ \left\{ 3a^2 + [-3b - 3c + 5d - 2(e + f - g + 2h) + j]a + b^2 + c^2 \right. \\
&+ d^2 - g^2 - 2cd + ce - 2de + 2cf - 2df - cg - 2eg + 2ch \\
&- 4dh + b(c - 2d + 2e + f + 2h) - (e + f - g + 2h)j \left. \right\} \zeta(2) \\
&+ \left\{ a^2 + [b + c + 4d + 3e + 3f + 4(g + h) + 2j]a - b^2 - c^2 + 2d^2 \right. \\
&- 2e^2 - 2f^2 - 2g^2 - 3h^2 - bc - bd - cd - 3be - 2ce - 2bf - 3cf \\
&- 3ef - 2bg - 2cg + 2dg - 2eg - 4fg - 4bh - 4ch + dh - 6eh \\
&- 6fh - 4gh - bj - cj + dj - ej - fj - 2gj \left. \right\} \zeta(3) \\
&+ \left\{ b^2 + cb + \frac{5db}{4} + \frac{eb}{2} + \frac{5fb}{4} + \frac{gb}{2} + \frac{3hb}{2} + \frac{jb}{4} + c^2 + 3d^2 + 3e^2 \right. \\
&+ 3f^2 + \frac{7h^2}{2} + j^2 - \frac{3ad}{4} + \frac{5cd}{4} + \frac{5ce}{4} + \frac{17de}{4} + \frac{cf}{2} + \frac{17df}{4} + \frac{17ef}{4} \\
&+ \frac{5cg}{4} + 6dg + 6eg + \frac{5fg}{4} + \frac{7ah}{4} + \frac{3ch}{2} + \frac{11dh}{2} + \frac{29eh}{4} + \frac{29fh}{4} \\
&+ \left. \frac{19gh}{4} + \frac{cj}{4} + 3dj + 3ej + 3fj + \frac{13hj}{4} \right\} \zeta(4) + \dots.
\end{aligned} \tag{4.86}$$

$$\begin{aligned}
F \left[\begin{matrix} a, b, d, e, g-1 \\ c, f, h, j \end{matrix} \right] &= \zeta(2) - (c + f - h + j) \zeta(2) \\
&\quad - (a + b + 2d + 2e + g + 2h - j) \zeta(3) + \dots, \\
F \left[\begin{matrix} a, b, d, e, g \\ c+1, f, h, j-1 \end{matrix} \right] &= -1 + \zeta(2) - a - b + 3c + 3f + 2h + j \\
&\quad + (a + b - 2c + d + e - f + 2g) \zeta(2) \\
&\quad - (a + b + 2d + 2e + 2f + 3g + 2h + j) \zeta(3) + \dots, \\
F \left[\begin{matrix} a, b+1, d, e, g \\ c, f, h, j-1 \end{matrix} \right] &= -1 + \zeta(2) - a + 3b - c + 3e + 2g + 2h + j \\
&\quad + (a - 2b + c + d - e + f) \zeta(2) \\
&\quad - (a + c + 2d + 2e + 2f + 2g + 2h + j) \zeta(3) + \dots, \\
F \left[\begin{matrix} a, b, d, e, g \\ c, f, h, j-1 \end{matrix} \right] &= \zeta(3) - \frac{1}{4} (4a + 4b + 4c + 5d + 5e + 5f + 2g + 2h + j) \zeta(4) + \dots.
\end{aligned} \tag{4.87}$$

The function $F \left[\begin{matrix} a, b, d, e, g \\ c+1, f, h, j-1 \end{matrix} \right]$ may be obtained from $F \left[\begin{matrix} a+1, b, d, e, g \\ c, f, h, j-1 \end{matrix} \right]$ thanks to the symmetry (4.8), *i.e.* through the permutations $a \leftrightarrow c$, $d \leftrightarrow f$, $g \leftrightarrow h$ on the latter.

4.6. Momentum expansion of singular triple hypergeometric functions $F^{(3)}$

In this subsection we shall address the case when the integrand of (4.7) has poles. One example is the integral

$$\begin{aligned}
F \left[\begin{matrix} a-1, b-1, d, e, g \\ c, f, h, j \end{matrix} \right] &= \int_0^1 dx \int_0^1 dy \int_0^1 dz x^{a-1} y^{b-1} z^c \\
&\quad \times (1-x)^d (1-y)^e (1-z)^f (1-xy)^g (1-yz)^h (1-xyz)^j,
\end{aligned} \tag{4.88}$$

whose integrand diverges for $x, y \rightarrow 0$. Hence we should expect a double pole $\frac{1}{ab}$ in the expansion of $F \left[\begin{matrix} a-1, b-1, d, e, g \\ c, f, h, j \end{matrix} \right]$ w.r.t. to small parameter a, \dots, j . For the case at hand the sums in the general expression (4.16) are not convergent. However, this problem is only a manifestation of the chosen integral representation (4.16) and other representation would be well-behaved after proper analytic continuation. For the case at hand we may not expand (4.16) due to the lack of absolute convergence. We do not undertake to obtain the appropriate convergent expression of (4.16) for that case rather we follow the methods anticipated in [31]. We add and subtract a similar singular piece to $F \left[\begin{matrix} a-1, b-1, d, e, g \\ c, f, h, j \end{matrix} \right]$, such

that we obtain two expressions, which may be handled easily. The integral $F\left[\begin{smallmatrix} a-1, b-1, d, e, g \\ c, f, h, j \end{smallmatrix}\right]$ is rearranged as a sum of two pieces: $F\left[\begin{smallmatrix} a-1, b-1, d, e, g \\ c, f, h, j \end{smallmatrix}\right] = \mathcal{I}_I^a + \mathcal{I}_I^b$, with:

$$\begin{aligned} \mathcal{I}_I^a &= \int_0^1 dx \int_0^1 dy \int_0^1 dz x^{a-1} y^{b-1} z^c (1-x)^d (1-y)^e (1-z)^f (1-yz)^h \\ &\quad \times \{ (1-xy)^g (1-xyz)^j - 1 \} \\ \mathcal{I}_I^b &= \underbrace{\left(\int_0^1 dx x^{a-1} (1-x)^d \right)}_{= B(a-1, d)} \underbrace{\left(\int_0^1 dy \int_0^1 dz y^{b-1} z^c (1-y)^e (1-z)^f (1-yz)^h \right)}_{= C(b-1, c, e, f, h)}. \end{aligned} \quad (4.89)$$

The integrand of the first integral \mathcal{I}_I^a stays finite for $x, y \rightarrow 0$. Hence we may easily expand this integral w.r.t. to the parameter a, \dots, j . On the other hand, the two integrals of the second expression \mathcal{I}_I^b have poles. However, the latter may be treated easily by using material of subsections 4.1. and 4.4. The double integral $C(b-1, c, e, f, h)$ may be obtained from the results of appendix A. More precisely, the latter may be expressed as a linear combination of Φ_1 and Φ_2 , given in (4.55) and (4.56), respectively:

$$C(a-1, b, c, d, e) = \frac{(1-a+b+d+e)(1+a+c+e)\Phi_1 - e(1+c+d+e)\Phi_2}{a(1-a+b+d)}.$$

To this end, we find

$$\begin{aligned} \mathcal{I}_I^a &= j(\zeta(2) - 1) - (g+j)\zeta(3) + \dots, \\ \mathcal{I}_I^b &= \left(\frac{1}{a} - d\zeta(2) + \dots \right) \left(\frac{1}{b} - \frac{c+f}{b} + h + \frac{(c+f)^2}{b} - \zeta(2)\left(e + \frac{cf}{b} + h\right) + \dots \right), \end{aligned} \quad (4.90)$$

and altogether:

$$F\left[\begin{smallmatrix} a-1, b-1, d, e, g \\ c, f, h, j \end{smallmatrix}\right] = \frac{1}{ab} - \frac{c+f}{ab} + \frac{(c+f)^2 + bh}{ab} - \left(\frac{d}{b} + \frac{cf}{ab} + \frac{e+h}{a} \right) \zeta(2) + \mathcal{O}(\epsilon). \quad (4.91)$$

Our next example is the integral:

$$\begin{aligned} F\left[\begin{smallmatrix} a-1, b, d, e, g \\ c, f, h, j \end{smallmatrix}\right] &= \int_0^1 dx \int_0^1 dy \int_0^1 dz x^{a-1} y^b z^c \\ &\quad \times (1-x)^d (1-y)^e (1-z)^f (1-xy)^g (1-yz)^h (1-xyz)^j, \end{aligned} \quad (4.92)$$

which again may be split into two contributions:

$$\begin{aligned}
\mathcal{I}_{II}^a &= \int_0^1 dx \int_0^1 dy \int_0^1 dz x^{a-1} y^b z^c (1-x)^d (1-y)^e (1-z)^f (1-yz)^h \\
&\quad \times \{ (1-xy)^g (1-xyz)^j - 1 \} \\
\mathcal{I}_{II}^b &= \underbrace{\left(\int_0^1 dx x^{a-1} (1-x)^d \right)}_{= B(a-1,d)} \underbrace{\left(\int_0^1 dy \int_0^1 dz y^b z^c (1-y)^e (1-z)^f (1-yz)^h \right)}_{= C(b,c,e,f,h)} .
\end{aligned} \tag{4.93}$$

The first integrand of the first integral \mathcal{I}_{II}^a stays finite. Moreover, the second integral of \mathcal{I}_{II}^b has not any poles either and may be evaluated by the results of section 4.4. It is equivalent to Φ_1 , given in (4.55). With this information, we find:

$$\begin{aligned}
\mathcal{I}_{II}^a &= g + 3j - (g + 2j) \zeta(2) + \dots , \\
\mathcal{I}_{II}^b &= \left(\frac{1}{a} - d \zeta(2) + \dots \right) (1 - b - c - e - f - 2h + \zeta(2) h + \dots) .
\end{aligned} \tag{4.94}$$

Finally, we obtain

$$F \left[\begin{matrix} a-1, b, d, e, g \\ c, f, h, j \end{matrix} \right] = \frac{1}{a} - \frac{b+c+e+f+2h}{a} + \frac{h}{a} \zeta(2) + \mathcal{O}(\epsilon) . \tag{4.95}$$

As a third example we want to discuss the following integral:

$$\begin{aligned}
F \left[\begin{matrix} a-1, b-1, d-1, e-1, g \\ c-1, f-1, h, j \end{matrix} \right] &= \int_0^1 dx \int_0^1 dy \int_0^1 dz x^{a-1} y^{b-1} z^{c-1} (1-x)^{d-1} \\
&\quad \times (1-y)^{e-1} (1-z)^{f-1} (1-xy)^g (1-yz)^h (1-xyz)^j .
\end{aligned} \tag{4.96}$$

The integrand has meromorphic poles at $x, y, z = 0, 1$. Therefore, this integral implies various triple poles after expanding w.r.t. to small a, \dots, j . There is not a simple way to disentangle the singular part of this integral in the way applied before. On the other hand, from our system of equations for $F^{(3)}$ we may easily deduce the momentum expansion:

$$\begin{aligned}
F \left[\begin{matrix} a-1, b-1, d-1, e-1, g \\ c-1, f-1, h, j \end{matrix} \right] &= \frac{(a+d)(c+f)}{abcd f} + \frac{(a+d)(d+e)+dg}{acde(d+e+g)} + \frac{e+f}{aef(e+f+h)} \\
&\quad + \frac{(d+e)(e+f)(d+e+f+g) + [(d+e)(e+f)+eg]h}{def(d+e+g)(e+f+h)(d+e+f+g+h+j)} + \mathcal{O}(\epsilon^{-1}) .
\end{aligned} \tag{4.97}$$

5. Reducible diagrams and contact interactions

We have found, that thanks to the relations *Eq.* (3.48) the six-gluon string S -matrix (3.1) at the disk-level may be expressed as a linear combination of six triple hypergeometric functions. The latter encode the α' -dependence of the full string amplitude. In that linear combination the coefficients in front of those six functions are meromorphic polynomials in the kinematical invariants s_i . In *Eqs.* (3.56), (3.57) and (3.59) the string S -matrix is given as a power series in α' up to the order α'^4 . In order to correctly interpret this result, we look on the latter from the field theory side. More precisely, the string S -matrix comprises many field-theoretical Feynman diagrams with six external legs. Out of this set of diagrams we are only interested in the irreducible diagrams as they only represent new interaction terms in the low-energy effective action. The reducible diagrams may be built by the Feynman rules applied to the known terms in the effective action. The string S -matrix result is organized as a power series in the momenta of the external states (*cf.* *Eqs.* (3.56), (3.57) and (3.59)). Each power in the momenta comprises a certain class of reducible and irreducible Feynman diagrams. All terms in the expansions are weighted with special values of the Riemann zeta-function. Formally, the six-gluon string S -matrix assumes the momentum (α')-expansion:

$$\mathcal{A}_6(k) \sim k^{-2} + 0 k^0 + \zeta(2) k^2 + \zeta(3) k^4 + \zeta(4) k^6 + \mathcal{O}(k^8) . \quad (5.1)$$

Up to k^6 (α'^4), there occurs the set of zeta-values: $\zeta(2)$, $\zeta(3)$, $\zeta(4)$ and $\zeta(2)^2$, together with 1 accounting for pure Yang-Mills diagrams (field-theory). But only $\zeta(2)^2$ and $\zeta(4)$ appear at the order α'^4 (*cf.* *Eq.* (3.59)). This has important consequences on the number of reducible diagrams, which the string S -matrix may comprise at this order in α' . Each α' -order brings a new set of zeta functions and combinations thereof. *E.g.* at the α'^5 -order $\zeta(5)$ and $\zeta(2) \zeta(3)$ show up, while up to the α'^6 -order $\zeta(6)$, $\zeta(2) \zeta(4)$, $\zeta(2)^3$ and $\zeta(3)^2$ show up in addition to the previous zeta-values. As we shall see, each zeta-value in the α' -expansion gives information about a class of underlying reducible and irreducible diagrams.

Since we want to extract the six-point interactions¹⁴ of the terms F^6 , $D^2 F^5$ and $D^4 F^4$ from the string S -matrix (3.1), we shall be only interested in the irreducible diagrams at

¹⁴ Note, that the $D^4 F^4$ terms are in principle already known from the four- and five-gluon string S -matrix. Hence, its four-vertex \mathcal{V}_4^8 with four external gluons and its five-vertex \mathcal{V}_5^7 with five external gluons will also appear in the reducible diagrams. The same is true for $D^2 F^5$: Its five-vertex \mathcal{V}_5^7 is already determined from a five gluon scattering amplitude. However, the six-vertex \mathcal{V}_6 with six external gluons, build from $D^4 F^4$ and $D^2 F^5$, is determined by our six-gluon amplitude at α'^4 .

momentum order $\mathcal{O}(k^6)$, *i.e.* the contact interactions of six gluons, as only those represent new interaction terms in the effective action at order α'^4 . Therefore, we shall study the α'^4 -order of *Eqs.* (3.56), (3.57) and (3.59) in more detail. Of course, the lower orders of the latter are also interesting as they should correctly reproduce Yang–Mills interactions of Table 1 up to the order α'^3 as a consequence of unitarity. The lower momentum orders k^{-2} , k^2 , k^4 of the string S -matrix (3.1), represent reducible diagrams with six external legs. Because of unitarity they must be completely determined by the Feynman rules applied to the effective action, valid up to $\mathcal{O}(\alpha'^3)$. On the other hand, momentum powers of (3.1) beyond k^6 represent gluon interactions such as $D^2 F^6, D^4 F^5, D^6 F^4$, which only become relevant at α'^5 or higher. To conclude, we shall only be interested in the reducible and irreducible diagrams of momentum order k^6 .

Let P denote the number of internal propagators inside a reducible Feynman graph and V_n^k the number of vertices \mathcal{V}_n^k with n legs and energy dimension k . We are then able to compute the total number of external lines N of a given Feynman graph

$$N = \sum_{\substack{n \geq 3 \\ k \geq 0}} n V_n^k - 2P, \quad (5.2)$$

with N the total number of external legs. Notice, that the sum starts at $n = 3$, in accordance that a vertex cannot have less than three legs. If two vertices are glued together, two legs are converted into an propagator. This way one has to subtract the double number of propagators P . Let us now take into account, that we only consider tree-level diagrams. This restricts the number of propagators for a given diagram to

$$P = \sum_{\substack{n \geq 3 \\ k \geq 0}} V_n^k - 1, \quad (5.3)$$

where the sum runs over the whole number of vertices. After eliminating the number of propagators from (5.2) we arrive at:

$$N = \sum_{\substack{n \geq 3 \\ k \geq 0}} (n - 2) V_n^k + 2. \quad (5.4)$$

When studying a six-point string S -matrix, the vertices of their reducible (field-theory) diagrams have at most five legs. In addition, the string S -matrix may contain contact interactions with six legs. Hence, we shall only be concerned about vertices with at most six external legs and we may explicitly evaluate the sum over n in (5.4):

$$6 = 2 + \sum_{k \geq 0} (V_3^k + 2V_4^k + 3V_5^k + 4V_6^k). \quad (5.5)$$

There are five possibilities to fulfill the constraint (5.5). The cases range from diagrams comprising only four vertices \mathcal{V}_3^k with three legs up to diagrams of only one vertex \mathcal{V}_6^k with six legs. Only the latter diagram, representing a contact interaction, gives rise to an irreducible diagram. Hence, those are the interactions to be added to the effective action at α'^4 , while the other diagrams involving vertices with fewer than six legs are reducible and may be already obtained by the Feynman rules of the effective action, valid up to α'^3 . More concretely, Table 2 displays the possibilities of how to meet the condition (5.5).

	V_3^k	V_4^k	V_5^k	V_6^k	P
<i>a</i>	4	0	0	0	3
<i>b</i>	2	1	0	0	2
<i>c</i>	1	0	1	0	1
<i>d</i>	0	2	0	0	1
<i>e</i>	0	0	0	1	0

Table 2: Number of vertices V_n^k to meet the condition (5.5).

For the the five cases *a*)–*e*), shown in Table 2, their corresponding Feynman diagrams are schematically drawn in Figure 1 and 2.




Figure 1: Feynman diagrams for the three cases *a*), *b*) and *c*).




Figure 2: Feynman diagrams for the two cases *d*) and *e*).

The vertices \mathcal{V}_n^k with at most six external legs following from the F^n - and $D^m F^n$ -terms in the effective action (*cf.* Table 1) are displayed in Table 3.

1	α'^0	F^2	\mathcal{V}_3^1	\mathcal{V}_4^0	
$\zeta(2)$	α'^2	F^4	\mathcal{V}_4^4	\mathcal{V}_5^3	\mathcal{V}_6^2
$\zeta(3)$	α'^3	F^5	\mathcal{V}_5^5	\mathcal{V}_6^4	
$\zeta(3)$	α'^3	$D^2 F^4$	\mathcal{V}_4^6	\mathcal{V}_5^5	\mathcal{V}_6^4
$\zeta(4)$	α'^4	F^6	\mathcal{V}_6^6		
$\zeta(4)$	α'^4	$D^4 F^4$	\mathcal{V}_4^8	\mathcal{V}_5^7	\mathcal{V}_6^6
$\zeta(4)$	α'^4	$D^2 F^5$	\mathcal{V}_5^7	\mathcal{V}_6^6	
$\zeta(5)$	α'^5	$D^6 F^4$	\mathcal{V}_4^{10}	\mathcal{V}_5^9	\mathcal{V}_6^8
$\zeta(5)$	α'^5	$D^2 F^6$	\mathcal{V}_6^8		

Table 3: Possible vertices \mathcal{V}_n^k with at most six external legs

Of course, \mathcal{V}_3^1 and \mathcal{V}_4^0 are just the standard non-Abelian Yang-Mills vertices. Note, that we do not show any vertices (namely \mathcal{V}_3^3 , \mathcal{V}_4^2 and \mathcal{V}_5^1) stemming from $\text{Tr}F^3$ as this term is absent in the superstring. This fact leads to somewhat fewer diagrams relevant to us than in the bosonic string.

So far, we have only discussed the number and kind of vertices, which appear in a tree-level Feynman diagram with six external legs. We are interested in diagrams, which have momentum power k^6 . The latter correspond to order α'^4 and mix in the α' -expansion of (3.1) with contact interactions of the same order. Similarly as for the number of legs (cf. Eq. (5.2)) we may find an expression for the energy-dimension of a given graph. With the total energy-dimension K given by $K = \sum_{n \geq 3, k \geq 0} k V_n^k - 2P$, we find:

$$K = \sum_{\substack{n \geq 3 \\ k \geq 0}} (k-2) V_n^k + 2. \quad (5.6)$$

Demanding $K = 6$ and using the fact, that there is only one three-vertex \mathcal{V}_3^1 , which follows from the standard Yang-Mills interaction $\text{Tr}(\text{tr}F^2)$, we obtain the condition:

$$4 = 4 V_6^6 - V_3^1 + \sum_{k \geq 0} (k-2) V_4^k + \sum_{k \geq 0} (k-2) V_5^k. \quad (5.7)$$

Now, we have to verify, which of the five cases, shown in Table 2, fulfill the constraint (5.7). There is a fast answer to the two cases *a*) and *e*): Any Feynman tree-level diagram with only four three-vertices \mathcal{V}_3^1 leads to a total momentum power k^{-2} and never meets the constraint (5.7). Among others this diagram contributes to the k^{-2} powers in Eqs.




Figure 3: *Diagram $\sim \zeta(4) k^6$ with one four-vertex \mathcal{V}_4^8 and two three-vertices \mathcal{V}_3^1*

(3.56), (3.57) and (3.59). On the other hand, any single contact vertex \mathcal{V}_6^6 of momentum order k^6 automatically fulfills the constraint. These vertices give rise to a new interaction term in the effective action at α^4 . Next, the case *b*) may only fulfill the requirement (5.7) with the four-vertex \mathcal{V}_4^8 involved. This is shown in Figure 3.

Furthermore, the case *c*) needs the five-vertex \mathcal{V}_5^7 , shown in Figure 4.




Figure 4: *Diagram $\sim \zeta(4) k^6$ with one five-vertex \mathcal{V}_5^7 and one three-vertex \mathcal{V}_3^1*

Finally, the case *d*) allows for two four-vertices, with their energies k, \tilde{k} fulfilling the constraint: $k + \tilde{k} = 8$. This leads to two classes of possibilities: one possibility involving two times the same vertex \mathcal{V}_4^4 or an other with two different four-vertices \mathcal{V}_4^k and $\mathcal{V}_4^{\tilde{k}}$, with $k + \tilde{k} = 8$. According to Table 3 we have four different four-vertices \mathcal{V}_4^k , with $k = 0, 4, 6, 8$. Hence, the case *d*) allows for the following two pairs of four-vertices \mathcal{V}_4^k and $\mathcal{V}_4^{\tilde{k}}$, with $(k, \tilde{k}) \in \{(4, 4), (0, 8)\}$, shown in Figure 5 and 6.




Figure 5: *Diagram $\sim \zeta(2)^2 k^6$ with two four-vertices \mathcal{V}_4^4 and one internal line*




Figure 6: Diagram $\sim \zeta(4) k^6$ with the two four-vertices \mathcal{V}_4^0 and \mathcal{V}_4^8

To conclude, Figures Nr. 3–6 show the four reducible Feynman diagrams, which lead to total energy dimension six, *i.e.* order k^6 in the external momenta k_i . These diagrams correspond to the cases *b)–d)* of Table 2. Furthermore, the two vertices \mathcal{V}_6^6 from F^6 and D^4F^4 give rise to two contact interactions of momentum order k^6 , accounting for the case *e)* of Table 2. All these six diagrams are contained in the k^6 –part of our string S –matrix (3.1). The latter is given in *Eqs.* (3.56), (3.57) and (3.59). Hence, to extract from the string S –matrix the information about the new interaction terms F^6 and D^4F^4 , to be written at α'^4 in the effective action, one has to take from our string S –matrix results the k^6 –part and subtract the contribution of the above four diagrams, displayed in Figures Nr. 3–6. Hence the work to be still done is the following [18]: With the vertices from Table 3 we have to calculate the above described four reducible diagrams in field–theory. Then we subtract the result from the k^6 order of the string S –matrix (3.1). The result, which has no poles, then accounts for the contact six–point interactions following from F^6 , D^2F^5 and D^4F^4 after inserting the linearized field strength $F_{\mu\nu} = \xi_\mu k_\nu - \xi_\nu k_\mu$. To obtain their precise form with the right coefficients one makes an ansatz for the terms F^6 , D^2F^5 and D^4F^4 and matches this ansatz with the relevant piece of the string S –matrix. This gives a system of linear equations for the coefficients. Solving these equations allows to write down the complete α'^4 –order of the effective gluon action (1.3), *cf.* [18]. The calculation of the Feynman graphs, shown in Figure 3–6, consists of two parts: First extracting the Feynman rules for the vertices from either string–theory or field–theory. Second we have to calculate the exchange diagram under consideration with respecting permutation symmetries. At this point we should remind about the ambiguity of relating string S –matrix results with corresponding terms in the effective action. It is notorious (see *e.g.* *Ref.* [36]), that the string S –matrix reproduces given terms in the effective action only up to field redefinitions.

Hence it is a lot more efficient, if we extract the Feynman rules for the vertices from known string amplitudes with fewer than six external legs rather than extracting them from field–theory terms, which are ambiguous.

Finally, let us point out, that we do not encounter reducible Feynman diagrams at order k^6 involving vertices from the α'^3 –terms. This fact is related to the absence of vertices from F^3 –terms due to supersymmetry. Otherwise *e.g.* the diagram *c* of Figure 1 with the two vertices \mathcal{V}_5^5 and \mathcal{V}_3^3 connected by an internal line, would give rise to k^6 . In other words, since in the string amplitude at the α'^4 –order we do not encounter any odd zeta–functions or combinations thereof, like $\zeta(3)$, $\zeta(3)\zeta(2)$, no reducible diagrams involving the vertices from F^5 or D^2F^4 have to be considered at this order in α' . This may be of deeper origin related to the fact, that interaction terms of odd powers in α' appear with the transcendental ζ –values: $\zeta(3)$, $\zeta(5)$, \dots . As it can be seen from Table 3 any diagram involving a vertex from the α'^3 –terms would be proportional to $\zeta(3)$. However, our expansions *Eqs.* (3.56), (3.57) and (3.59) do not have $\zeta(3)$ at α'^4 . Hence, these vertices do not show up in the string S –matrix (3.1) at the α'^4 order, neither in reducible nor in irreducible diagrams. However, the vertex \mathcal{V}_6^4 from F^5 and D^2F^4 contributes at the order α'^3 as terms proportional to $\zeta(3)k^4$ in the expansions (3.56), (3.57) and (3.59). On the other hand, since the α'^4 –order of the six–gluon string S –matrix contains only $\zeta(4)$ and $\zeta(2)^2$, any reducible diagram contributing at this order contains either two vertices from F^4 or one vertex from D^4F^4 together with some vertices from the Yang–Mills interaction F^2 . This is precisely, what we have found above (*cf.* Figure 3-6). Hence, in the α' –expansion of a given gluon string S –matrix there seems to be an intriguing disentangling of terms between $\zeta(2s)$ and $\zeta(2s + 1)$, respectively at each order. This means, that for a given α' –order only a certain class of reducible and irreducible diagrams contributes. In addition there is a crucial difference between even and odd F^n –terms in the effective action, which relies in the difference of even and odd zeta–values. In the α' –expansion of a string S –matrix, *e.g.* a term $\zeta(2)\zeta(3)$ may be immediately traced back to a contribution from a reducible diagram involving one vertex from the F^5, D^2F^4 terms and an other from the F^4 –terms. Similarly for $\zeta(3)^2$ or $\zeta(3)\zeta(5)$. Hence the effective action terms involving odd zeta–values may be much easier extracted from a given string expansion than the terms proportional to even zeta–values or combinations thereof. In fact due to the identity

$$\zeta(2)^2 = \frac{5}{2} \zeta(4) \tag{5.8}$$

the contribution to the α'^4 -order of the string S -matrix of *e.g.* Figure 5 may not be disentangled from the contribution from Figure 6. Similar statements hold at the α'^6 -order due to the relations:

$$\begin{aligned}\zeta(2)^3 &= \frac{35}{8} \zeta(6) , \\ \zeta(2) \zeta(4) &= \frac{7}{4} \zeta(6) .\end{aligned}\tag{5.9}$$

6. Concluding remarks

In this work we have calculated the open superstring S -matrix with four, five and six external gluons on the disk by introducing a powerful method. We have found a systematic and efficient way to calculate tree-level string amplitudes by equating seemingly different expressions for one and the same string S -matrix. An intriguing way of using world-sheet supersymmetry generates a system of non-trivial equations for string tree-level amplitudes. These equations represent algebraic identities between different multiple hypergeometric functions or generalizations thereof. Their solutions give the ingredients of the string S -matrix. Generically, one- or higher-loop string amplitudes give much more constraints on the form of the effective action as a result of world-sheet supersymmetry manifested through Riemann identities in the string S -matrix calculation. The latter impose quite strong conditions on the allowed heavy string states running the loops. On the other hand, what we find here is, that it is the superdiffeomorphism invariance on the string world-sheet, which imposes constraints on the effective superstring tree-level action: by writing the tree-level string S -matrix in different, but equivalent forms, we obtain a system of equations, whose solutions capture the pieces of the full string S -matrix. Hence, in all cases, both string tree-level and string loop, it is world-sheet supersymmetry, which may be used to generate various non-trivial equations between different amplitudes. Especially consult *Ref.* [37] for highlighting this method at two-loop. It would be very interesting, whether in the pure spinor formalism similar relations may be obtained. See *Ref.* [38] for further information on this interesting question.

The full string S -matrix can be expressed by six (generalized) hypergeometric functions as in *Eq.* (3.55), which in the effective action play an important rôle in arranging the higher order α' gauge interaction terms (such as $F^6, D^2 F^6, D^4 F^4, D^6 F^4, \dots$) with six external legs. The final expression (3.55) is given by six triple hypergeometric functions Φ_j , which encode the full α' -dependence and some polynomials \mathcal{P}^j , which keep track about

the pole structure of the reducible diagrams. The functions Φ_j organize the various terms according to their zeta-values [21,18].

The general complication in constructing the effective action from string S -matrices is, that one has to expand the result for the string amplitude w.r.t. to the kinematical invariants (or α'), in order to subtract the terms accounting for the reducible diagrams. Since generically any tree-level string amplitude is given by some multiple hypergeometric functions (or generalizations thereof, *cf.* section 4), with the latter comprising the α' -dependence of the full amplitude, this problem turns to the mathematical task of expanding hypergeometric functions (or their generalizations) w.r.t. small parameters. In the case, when these functions have poles, this is a very challenging work (*cf.* subsection 4.6). However, with the formalism developed in this article, it is generically possible to express those hypergeometric functions, which have poles, as a linear combination of other hypergeometric functions, which are simpler to expand (*cf.* section 4 for explicit results). *Cf. Eq. (3.48)* for the six-gluon amplitude: There, the triple hypergeometric functions Φ_j comprise a basis of finite functions and the poles simply appear as coefficients Λ^j in front of those simpler hypergeometric functions. In fact, as we have demonstrated in subsection 3.3, these poles follow from solving algebraic equations.

So far, the heterotic-type I duality [39] has been checked up to certain α'^2 -terms in the effective action [40,41,42]. The terms tested represent 1/2 BPS saturated couplings, *i.e.* through supersymmetry they are related to eight fermion terms in the effective action. This duality maps certain (anomaly related) α'^2 -terms to each other. More precisely, the heterotic one-loop corrections to the terms F^4 , F^2R^2 , $(R^2)^2$, R^4 reproduce the analogous terms of the type I $SO(32)$ at tree-level [40,41]. Hence, there is a map between the integral over the complex structure modulus of the heterotic one-loop world-sheet torus to the single position integration of the disk-calculation of those α'^2 -terms. It is certainly important, to go beyond that order to verify this duality, in particular extending the checks to non-BPS saturated amplitudes. Though heterotic-type I duality is a strong-weak coupling duality, there have been given arguments due to non-renormalization theorems and underlying BPS-structure of certain amplitudes [43] (see also *Refs.* [44]), that this duality should also work perturbatively, *i.e.* perturbative heterotic calculations map to analogous perturbative type I results. In this line we expect, that a perturbative heterotic two-loop calculation of the six gluon string S -matrix in $D = 10$, captures the tree-level six gluon string S -matrix in the type I superstring [37,43]. Hence, the irreducible part of the α'^4 -order of *Eqs.* (3.56), (3.57) and (3.59) should agree with the two-loop results derived

in [37,43,45]. This check will be performed in *Ref.* [45]. At any rate, since a certain (BPS protected) kinematics of the heterotic two-loop amplitude (*cf.* the discussions in [37,43]) is given by an integral over the fundamental domain of the modular group of the $g = 2$ torus [46]

$$\int_{\mathcal{F}_2} \frac{d^3\Omega d^3\bar{\Omega}}{\det\text{Im}(\Omega)} = \frac{1}{3\pi} \zeta(4) , \quad (6.1)$$

the duality seems to work, since the relevant terms from (3.59), indeed start with $\zeta(4)$ [45]. Hence, as in the heterotic one-loop case, there seems to be a link between the integral over the moduli space of the heterotic two-loop torus (6.1) and the integral over positions of three vertex operators on the disk. Qualitatively for a kinematics $(\xi\xi)$ $(\xi\xi)$ $(\xi k)(\xi k)$ the latter integral may be mimicked by:

$$\int_0^1 dx \int_0^1 dy \int_0^1 dz \frac{(1-xy)^{\alpha'}}{(1-xyz)^2} \Big|_{\alpha'^2} = \zeta(4) . \quad (6.2)$$

The duality then relates the two integrals (6.1) and (6.2). Understanding this relation also for the non-BPS protected terms would be certainly interesting for a deeper understanding of the heterotic-type I duality. In that case, the integrand is of the generic form (3.31), while for the heterotic two-loop result the integrand over the $g = 2$ fundamental region represents some gauged heterotic two-loop partition function, given by some combination of $g = 2$ modular functions. Therefore, the duality turns into the non-trivial mathematical statement, that there is a link between the integral (3.31) and a two-loop torus integral over a certain combination of $g = 2$ modular functions. Perhaps the various identities between hypergeometric functions, that we have found (*cf.* subsection 3.3), may have a direct translation into two-loop Riemann identities on the heterotic side. Certainly this would lead to quite non-trivial relations between generalized hypergeometric functions and theta-functions, initiated by the duality between open and closed strings.

In perturbative gauge theories the maximally helicity violating (MHV) N -point amplitudes, with $N - 2$ gluons of one helicity and two gluons of the other helicity, show a remarkable simplicity, which is not apparent from the Feynman rules [47]. This result has a nice explanation after Fourier transforming the momentum space amplitudes to twistor space [48]. Based on this observation recently there has been made progress in perturbative gauge theory by studying the underlying twistor structure of scattering amplitudes. At tree-level this investigation has provided recursion relations for on-shell amplitudes [49], which give new and simple forms for many amplitudes in field-theory. The question one

may ask is, whether after including α' -corrections to the MHV-amplitudes would motivate a possible generalization of the duality between perturbative gauge theory and twistor string theory [21].

Acknowledgements: We wish to thank Tom Taylor and Christoph Sieg for many valuable and useful discussions. Furthermore, we are very grateful to Wolfgang Lerche and Dieter Lüst for useful discussions. D.O. thanks the Ludwig-Maximilians-Universität München for hospitality.

Appendix A. Five gluon open superstring S -matrix

In section 2 we have presented a powerful technique to calculate tree-level string amplitudes. As an example we outlined the procedure at the four gluon string S -matrix. In section 3 this technique considerably helped to derive the full six-gluon string S -matrix. For the reader, who still might not be convinced about the tremendous simplifications occurring once this method is used, we now apply this recipe to the five gluon string S -matrix. Though this amplitude has been already calculated in *Ref.* [6], we find it quite enlightening to redo this calculation within the frame established in this article. Moreover, this calculation allows to derive striking identities for the hypergeometric function ${}_3F_2\left[\begin{smallmatrix} a, b, c \\ c, d \end{smallmatrix}\right]$. In particular, we find interesting relations, which allows to express one ${}_3F_2$, which has a complicated pole structure, as a linear combination of two others, which do not have poles. Hence, in this linear combination the poles are completely captured by the coefficients in front of those hypergeometric functions. This is in lines of *Eq.* (3.48), where one needs a basis of six functions, to express a triple hypergeometric function $\tilde{F}\left[\begin{smallmatrix} n_{24}, n_{25}, n_{26}, n_{34}, n_{35} \\ n_{36}, n_{45}, n_{46}, n_{56} \end{smallmatrix}\right]$ in terms of others.

We study the string S -matrix describing the string tree-level scattering of five gauge bosons on the disk:

$$\begin{aligned} \mathcal{A}_5(k_1, \xi_1, a_1; k_2, \xi_2, a_2; k_3, \xi_3, a_3; k_4, \xi_4, a_4; k_5, \xi_5, a_5) \\ = V_{\text{CKG}}^{-1} \prod_{r=1}^5 \int d^2 z_r \langle V_{A^{a_1}}^{(-1)}(z_1) V_{A^{a_2}}^{(-1)}(z_2) V_{A^{a_3}}^{(0)}(z_3) V_{A^{a_4}}^{(0)}(z_4) V_{A^{a_5}}^{(0)}(z_5) \rangle . \end{aligned} \quad (\text{A.1})$$

We have total momentum conservation $\sum_{i=1}^5 k_i = 0$. Furthermore, we have chosen two gauge vertex operators in the (-1) -ghost picture in order to guarantee a total ghost charge of -2 on the disk. The explicit evaluation of the correlator in *Eq.* (A.1) leads to a sum of

integrals with the two classes of (space–time) kinematic contractions: $(\xi\xi)(\xi\xi)(\xi k)$ and $(\xi\xi)(\xi k)(\xi k)(\xi k)$. The first kinematics comprises the following 15×3 contractions

$$\begin{aligned}
\Xi_1 &:= (\xi_1\xi_2) (\xi_3\xi_4) (\xi_5k_s) \quad , \quad \Xi_2 := (\xi_1\xi_3) (\xi_2\xi_4) (\xi_5k_s) \quad , \quad \Xi_3 := (\xi_1\xi_4) (\xi_2\xi_3) (\xi_5k_s), \\
\Xi_4 &:= (\xi_1\xi_2) (\xi_3\xi_5) (\xi_4k_s) \quad , \quad \Xi_5 := (\xi_1\xi_3) (\xi_2\xi_5) (\xi_4k_s) \quad , \quad \Xi_6 := (\xi_1\xi_5) (\xi_2\xi_3) (\xi_4k_s), \\
\Xi_7 &:= (\xi_1\xi_2) (\xi_4\xi_5) (\xi_3k_s) \quad , \quad \Xi_8 := (\xi_1\xi_4) (\xi_2\xi_5) (\xi_3k_s) \quad , \quad \Xi_9 := (\xi_1\xi_5) (\xi_2\xi_4) (\xi_3k_s), \\
\Xi_{10} &:= (\xi_1\xi_3) (\xi_4\xi_5) (\xi_2k_s) \quad , \quad \Xi_{11} := (\xi_1\xi_4) (\xi_3\xi_5) (\xi_2k_s) \quad , \quad \Xi_{12} := (\xi_1\xi_5) (\xi_3\xi_4) (\xi_2k_s), \\
\Xi_{13} &:= (\xi_2\xi_3) (\xi_4\xi_5) (\xi_1k_s) \quad , \quad \Xi_{14} := (\xi_2\xi_4) (\xi_3\xi_5) (\xi_1k_s) \quad , \quad \Xi_{15} := (\xi_2\xi_5) (\xi_3\xi_4) (\xi_1k_s),
\end{aligned} \tag{A.2}$$

with k_j accounting for three space–time momenta as a result of applying the on–shell constraint $\xi_i k_i = 0$ and momentum conservation.

In Eq. (A.1) we have chosen the first two gauge vertex operators in the (-1) –ghost picture. As a matter of $PSL(2, \mathbf{R})$ –invariance on the disk, we are free to chose which pair (a, b) of the six gauge vertex operators we put into the (-1) –ghost picture. Hence, instead of the particular choice in (A.1), for a given group structure π , we may generically consider the expression

$$\begin{aligned}
&\text{Tr}(\lambda^{\pi(1)}\lambda^{\pi(2)}\lambda^{\pi(3)}\lambda^{\pi(4)}\lambda^{\pi(5)}) A^\pi(a, b, i, j, k) \\
&= V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^5 d^2 z_r \langle V_{A^{a_a}}^{(-1)}(z_a) V_{A^{a_b}}^{(-1)}(z_b) V_{A^{a_i}}^{(0)}(z_i) V_{A^{a_j}}^{(0)}(z_j) V_{A^{a_k}}^{(0)}(z_k) \rangle ,
\end{aligned} \tag{A.3}$$

with some permutation $(a, b, i, j, k) \in \overline{(1, 2, 3, 4, 5)}$ of vertex operators. According to (2.1) from this piece we may obtain the whole string S –matrix (A.1). After performing the Wick contractions and with the correlators (2.8) we may evaluate (A.3) and obtain:

$$\begin{aligned}
A^\pi(a, b, i, j, k) &= V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^5 d^2 z_r \sum_{\substack{(i_1, i_2, i_3) \\ \in (i, j, k)}} \left\{ \frac{1}{2} \mathcal{A}_1(a, b, i_1, i_2, i_3) (\xi_a \xi_b) (\xi_{i_1} \xi_{i_2}) \right. \\
&+ \mathcal{A}_2(a, i_1, b, i_2, i_3) (\xi_a \xi_{i_1}) (\xi_b \xi_{i_2}) \\
&+ \frac{1}{2} \mathcal{A}_3(a, i_1, i_2, i_3, b) (\xi_a \xi_{i_1}) (\xi_{i_2} \xi_{i_3}) + \frac{1}{2} \mathcal{A}_3(b, i_1, i_2, i_3, a) (\xi_b \xi_{i_1}) (\xi_{i_2} \xi_{i_3}) \\
&+ \frac{1}{6} \mathcal{C}_1(a, b, i_1, i_2, i_3) (\xi_a \xi_b) + \frac{1}{2} \mathcal{C}_3(i_1, i_2, a, b, i_3) (\xi_{i_1} \xi_{i_2}) \\
&\left. + \frac{1}{2} \mathcal{C}_2(a, i_1, b, i_2, i_3) (\xi_a \xi_{i_1}) + \frac{1}{2} \mathcal{C}_2(b, i_1, a, i_2, i_3) (\xi_b \xi_{i_1}) \right\} .
\end{aligned} \tag{A.4}$$

In the following, we shall need the integrated expressions

$$A_i^\pi = V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^6 z_r \mathcal{A}_i \quad , \quad C_i^\pi = V_{\text{CKG}}^{-1} \int_{\mathcal{I}_\pi} \prod_{r=1}^6 d^6 z_r \mathcal{C}_i \quad , \quad (\text{A.5})$$

which enter the expression (A.4). In front of the contractions $(\xi_a \xi_b)$ $(\xi_i \xi_j)$, $(\xi_a \xi_i)$ $(\xi_b \xi_j)$ and $(\xi_a \xi_i)$ $(\xi_j \xi_k)$ there appear the three basic functions $\mathcal{A}_1(a, b, i, j, k)$, $\mathcal{A}_2(a, i, b, j, k)$ and $\mathcal{A}_3(a, i, j, k, b)$, respectively

$$\begin{aligned} \mathcal{A}_1(a, b, i, j, k) &= i \mathcal{E} \left\{ \frac{1 - k_i k_j}{z_{ab}^2 z_{ij}^2} \left[(\xi_k k_a) \frac{z_{ja}}{z_{ak} z_{jk}} + (\xi_k k_b) \frac{z_{jb}}{z_{bk} z_{jk}} + (\xi_k k_i) \frac{z_{ji}}{z_{ik} z_{jk}} \right] \right. \\ &\quad \left. + \frac{1}{z_{ab}^2 z_{ik} z_{jk} z_{ij}} [(k_j k_k) (\xi_k k_i) - (k_i k_k) (\xi_k k_j)] \right\} \quad , \\ \mathcal{A}_2(a, i, b, j, k) &= i \mathcal{E} \left\{ \frac{k_i k_j}{z_{ab} z_{ai} z_{bj} z_{ij}} \left[(\xi_k k_a) \frac{z_{ja}}{z_{ak} z_{jk}} + (\xi_k k_b) \frac{z_{jb}}{z_{bk} z_{jk}} + (\xi_k k_i) \frac{z_{ji}}{z_{ik} z_{jk}} \right] \right. \\ &\quad \left. + \frac{1}{z_{ab} z_{ai} z_{bj} z_{ik} z_{jk}} [(k_i k_k) (\xi_k k_j) - (k_j k_k) (\xi_k k_i)] \right\} \quad , \\ \mathcal{A}_3(a, i, j, k, b) &= \frac{i \mathcal{E}}{z_{ab} z_{ai} z_{jk}} \left\{ \frac{(k_j k_k) (\xi_b k_i)}{z_{bi} z_{jk}} - \frac{(k_i k_k) (\xi_b k_j)}{z_{bj} z_{ik}} + \frac{(k_i k_j) (\xi_b k_k)}{z_{bk} z_{ij}} - \frac{(\xi_b k_i)}{z_{bi} z_{jk}} \right\} \quad , \end{aligned} \quad (\text{A.6})$$

with:

$$\mathcal{E} = \prod_{r < s}^5 |z_{rs}|^{k_r k_s} \quad .$$

Obviously, the first two integrals A_1^π and A_2^π account for the two cases whether the two vertices a, b in the (-1) -ghost picture are contracted among themselves or with other vertices, respectively. The integral A_3^π describes the case if only the polarization ξ_a of the vertex a is contracted with an other polarization ξ_i , while the polarization ξ_b is contracted with momenta. The function $\mathcal{A}_1(a, b, i_1, i_2, i_3)$ is invariant under permutations of the indices i_1, i_2 and a, b , while the function \mathcal{A}_2 enjoys the property $\mathcal{A}_2(a, i_1, b, i_2, i_3) = \mathcal{A}_2(b, i_2, a, i_1, i_3)$. Furthermore, the function $\mathcal{A}_3(a, i_1, i_2, i_3, b)$ is symmetric under permutations of i_2 and i_3 . Due to those symmetries it is obvious, that in the amplitude $A^\pi(a, b, i, j, k)$ any permutation of the three indices i, j, k , which label the three vertices in the zero-ghost picture, yields the same expression (3.4) as a result of Bose-symmetry of the NS -operators. The same is true for the permutation of a and b . In each amplitude $A^\pi(a, b, i, j, k)$, as given in Eq. (A.4), all 15 space-time kinematical contractions (A.2) show up. For the kinematics $(\xi\xi)(\xi k)(\xi k)(\xi k)$, which in the string S -matrix (A.4) may appear in the combinations

$C_1(a, b, i, j, k)$ ($\xi_a \xi_b$), $C_2(a, i, b, j, k)$ ($\xi_a \xi_i$) and $C_3(i, j, a, b, k)$ ($\xi_i \xi_j$), we have the corresponding integrands

$$\begin{aligned}
C_1(a, b, i, j, k) &= i \mathcal{E} \left\{ \frac{(\xi_j k_b)(\xi_k k_b)}{z_{ai} z_{bj} z_{bk}} \left(\frac{(\xi_i k_b) z_{ab}}{z_{aj} z_{ak} z_{bi}} - \frac{(\xi_i k_j)}{z_{ak} z_{ij}} - \frac{(\xi_i k_k)}{z_{aj} z_{ik}} \right) \right. \\
&\quad + \frac{(\xi_j k_i)(\xi_k k_b)}{z_{aj} z_{bk} z_{ij}} \left(\frac{(\xi_i k_b)}{z_{ak} z_{bi}} - \frac{(\xi_i k_k)}{z_{ab} z_{ik}} \right) + \frac{(\xi_j k_k)(\xi_k k_b)}{z_{ai} z_{bk} z_{jk}} \left(\frac{(\xi_i k_j)}{z_{ab} z_{ij}} - \frac{(\xi_i k_b)}{z_{aj} z_{bi}} + \frac{(\xi_i k_k) z_{ak}}{z_{ab} z_{aj} z_{ik}} \right) \\
&\quad + \frac{(\xi_i k_b)(\xi_k k_i)}{z_{aj} z_{bi} z_{ik}} \left(\frac{(\xi_j k_b)}{z_{ak} z_{bj}} - \frac{(\xi_j k_i) z_{ai}}{z_{ab} z_{ak} z_{ij}} - \frac{(\xi_j k_k)}{z_{ab} z_{jk}} - \frac{(\xi_i k_j)(\xi_j k_b)(\xi_k k_j)}{z_{ab} z_{ak} z_{bj} z_{ij} z_{ik}} \right) \\
&\quad \left. + \frac{(\xi_j k_b)(\xi_k k_j)}{z_{ai} z_{bj} z_{jk}} \left(\frac{(\xi_i k_b)}{z_{aj} z_{ak} z_{bi}} - \frac{(\xi_i k_j) z_{aj}}{z_{ab} z_{ak} z_{ij}} - \frac{(\xi_i k_k)}{z_{ab} z_{ik}} \right) - \frac{(\xi_i k_b)(\xi_j k_i)(\xi_k k_j)}{z_{ab} z_{ak} z_{bi} z_{ij} z_{jk}} \right\}, \\
C_2(a, i, b, j, k) &= i \mathcal{E} \left\{ \frac{(\xi_b k_i)(\xi_k k_b)}{z_{aj} z_{bi} z_{bk}} \left(\frac{(\xi_j k_k)}{z_{ai} z_{jk}} - \frac{(\xi_j k_b) z_{ab}}{z_{ai} z_{ak} z_{bj}} - \frac{(\xi_j k_i)}{z_{ak} z_{ij}} \right) \right. \\
&\quad - \frac{(\xi_j k_i)(\xi_k k_i)}{z_{ab} z_{ij} z_{ik}} \left(\frac{(\xi_b k_i) z_{ai}}{z_{aj} z_{ak} z_{bi}} + \frac{(\xi_b k_j)}{z_{ak} z_{bj}} + \frac{(\xi_b k_k)}{z_{aj} z_{bk}} \right) - \frac{(\xi_j k_b)(\xi_k k_c)}{z_{aj} z_{bj} z_{ik}} \left(\frac{(\xi_b k_i)}{z_{ak} z_{bi}} + \frac{(\xi_b k_k)}{z_{ai} z_{bk}} \right) \\
&\quad + \frac{(\xi_j k_k)(\xi_k k_i)}{z_{ab} z_{ik} z_{jk}} \left(\frac{(\xi_b k_i)}{z_{aj} z_{bi}} + \frac{(\xi_b k_j)}{z_{ai} z_{bj}} + \frac{(\xi_b k_k) z_{ak}}{z_{ai} z_{aj} z_{bk}} \right) - \frac{(\xi_b k_j)(\xi_j k_i)(\xi_k k_b)}{z_{ai} z_{ak} z_{bj} z_{bk} z_{ij}} \\
&\quad \left. - \frac{(\xi_j k_i)(\xi_k k_j)}{z_{ab} z_{ij} z_{jk}} \left(\frac{(\xi_b k_i)}{z_{ak} z_{bi}} + \frac{(\xi_b k_j) z_{aj}}{z_{ai} z_{ak} z_{bj}} + \frac{(\xi_b k_k)}{z_{ai} z_{bk}} \right) - \frac{(\xi_b k_i)(\xi_j k_b)(\xi_k k_j)}{z_{ai} z_{ak} z_{bi} z_{bj} z_{jk}} \right\}, \\
C_3(i, j, a, b, k) &= i \mathcal{E} \left\{ \frac{(\xi_k k_b)}{z_{ak} z_{bk} z_{ij}} \left(\frac{(\xi_a k_i)(\xi_b k_j)}{z_{ai} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_i)}{z_{aj} z_{bi}} \right) \right. \\
&\quad - \frac{(\xi_k k_i)}{z_{ab} z_{ak} z_{ij} z_{ik}} \left(\frac{(\xi_a k_i)(\xi_b k_j)}{z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_i) z_{ai}}{z_{aj} z_{bi}} \right) + \frac{(\xi_a k_i)(\xi_k k_j)}{z_{ab} z_{ai} z_{ij} z_{jk}} \left(\frac{(\xi_b k_j) z_{aj}}{z_{ak} z_{bj}} + \frac{(\xi_b k_k)}{z_{bk}} \right) \\
&\quad \left. - \frac{(\xi_b k_i)(\xi_k k_j)}{z_{ab} z_{bi} z_{ij} z_{jk}} \left(\frac{(\xi_a k_j)}{z_{ak}} + \frac{(\xi_a k_k)}{z_{ak}} \right) + \frac{(\xi_k k_i)}{z_{ak} z_{ij} z_{ik}} \left(\frac{(\xi_a k_k)(\xi_b k_j)}{z_{ak} z_{bj}} - \frac{(\xi_a k_j)(\xi_b k_k)}{z_{aj} z_{bk}} \right) \right\}, \tag{A.7}
\end{aligned}$$

respectively.

Exchanging in the amplitude $A^\pi(a, b, i, j, k)$ indices from the two sets $\{a, b\}$ and $\{i, j, k\}$, *i.e.* putting instead of the operators a, b other operators in the (-1) -ghost picture, leads to seemingly different expressions. In other words, putting different vertex operators in the (-1) -ghost picture, provides different function A_r^π, C_s^π in front of a given kinematics. The five-point amplitude (A.3) allows for $\binom{5}{2} = 10$ possibilities to choose which vertex pair (a, b) to put into the (-1) -ghost picture. Therefore, for a given group structure π and the given kinematics

$$(\xi_A \xi_B)(\xi_C \xi_D)(\xi_E k_s) \tag{A.8}$$

we obtain ten *a priori*-different expressions A_r^π , when we examine (A.4) for the choices $(a, b) \in \{(A, B), (A, C), (A, D), (A, E), (B, C), (B, D), (B, E), (C, D), (C, E), (D, E)\}$.

$$\tag{A.9}$$

The latter may be read off from (A.4):

$$\begin{aligned}
& A_1^\pi(A, B, C, D, E) \quad , \quad A_1^\pi(C, D, A, B, E) \quad , \\
& A_2^\pi(A, B, C, D, E) \quad , \quad A_2^\pi(B, A, C, D, E) \quad , \quad A_2^\pi(B, A, D, C, E) \quad , \quad A_2^\pi(A, B, D, C, E) \quad , \\
& A_3^\pi(A, B, C, D, E) \quad , \quad A_3^\pi(B, A, C, D, E) \quad , \quad A_3^\pi(C, D, A, B, E) \quad , \quad A_3^\pi(D, C, A, B, E) \quad .
\end{aligned} \tag{A.10}$$

They all have to be equated, which gives rise to 9×3 equations for each kinematics (A.8) under consideration. The additional factor of three arises, since $s = A, B, C$ due to the on-shell constraint $\xi_E k_E = 0$ and momentum conservation. *E.g.* the kinematics Ξ_1 , *i.e.* $A = 1, B = 2, C = 3, D = 4$ and $E = 5$ yields the system of equations:

$$\begin{aligned}
& A_1^\pi(1, 2, 3, 4, 5) = A_1^\pi(3, 4, 1, 2, 5) = A_2^\pi(1, 2, 3, 4, 5) = A_2^\pi(2, 1, 3, 4, 5) = A_2^\pi(2, 1, 4, 3, 5) \\
& = A_2^\pi(1, 2, 4, 3, 5) = A_3^\pi(1, 2, 3, 4, 5) = A_3^\pi(2, 1, 3, 4, 5) = A_3^\pi(3, 4, 1, 2, 5) = A_3^\pi(4, 3, 1, 2, 5) \quad .
\end{aligned} \tag{A.11}$$

Taking into account all 15×3 kinematics (A.2), we obtain $27 \times 15 = 405$ equations, which give rise to non-trivial relations among the integrals of the type (A.7) in the same way, as we have seen before in the case of the six- or four-gluon string S -matrix. However, again many equations are equal. Actually, we are left with only 291 equations.

Similarly, let us discuss the ten different expressions for a kinematics of the structure:

$$(\xi_A \xi_B) \quad (\xi_C k_r) \quad (\xi_D k_s) \quad (\xi_E k_t) \quad . \tag{A.12}$$

From (A.4) we read off the ten possibilities

$$\begin{aligned}
& C_1^\pi(A, B, C, D, E) \quad , \quad C_2^\pi(A, B, C, D, E) \quad , \quad C_2^\pi(B, A, C, D, E) \quad , \quad C_2^\pi(A, B, D, C, E) \quad , \\
& C_2^\pi(B, A, D, C, E) \quad , \quad C_2^\pi(A, B, E, C, D) \quad , \quad C_2^\pi(B, A, E, C, D) \quad , \\
& C_3^\pi(A, B, C, D, E) \quad , \quad C_3^\pi(A, B, C, E, D) \quad , \quad C_3^\pi(A, B, D, E, C)
\end{aligned} \tag{A.13}$$

to extract the kinematics (A.12). They all have to lead to the same result. After equating them we obtain nine equations for each kinematics (A.12). On-shell, there are $10 \times 3^3 = 270$ kinematics (A.12). Hence we obtain 2,430 equations. However, many of them are equal and in fact this system reduces to only 151 equations.

So far, we have not yet fixed any of the vertex operators. Due to the $PSL(2, \mathbf{R})$ invariance on the disk, we may fix three positions of the vertex operators. A convenient choice in (A.1) or in (A.7) is (3.21), which implies the ghost factor $\langle c(z_1)c(z_2)c(z_3) \rangle = -z_\infty^2$.

A five-point amplitude involving massless external particles, *i.e.* $k_i^2 = 0$, allows for five independent kinematic invariants s_i . We may choose¹⁵ the scalar products:

$$\begin{aligned} s_1 &= k_2 k_4 \quad , \quad s_2 = k_2 k_5 \quad , \quad s_3 = k_3 k_4 \quad , \\ s_4 &= k_3 k_5 \quad , \quad s_5 = k_4 k_5 \quad . \end{aligned} \tag{A.15}$$

The general structure of the integrals (A.7) is then given by:

$$\tilde{F}[n_{24}, n_{25}, n_{34}, n_{35}, n_{45}] := \int_{\mathcal{I}_\pi} dz_4 dz_5 |z_4|^{\alpha_{24}} |z_5|^{\alpha_{25}} |1 - z_4|^{\alpha_{34}} |1 - z_5|^{\alpha_{35}} |z_4 - z_5|^{\alpha_{45}} . \tag{A.16}$$

Here, the powers α_{ij} are real numbers, which are given by

$$\begin{aligned} \alpha_{24} &= s_1 + n_{24} \quad , \quad \alpha_{25} = s_2 + n_{25} \quad , \quad \alpha_{34} = s_3 + n_{34} \quad , \\ \alpha_{35} &= s_4 + n_{35} \quad , \quad \alpha_{45} = s_5 + n_{45} \quad , \end{aligned} \tag{A.17}$$

with some integers $n_{ij} \in \{-2, -1, 0\}$. The integrals (A.16) may be transformed into the form (4.2). Of course, this depends on the range of integration \mathcal{I}_π chosen in (A.16). For instance, if we concentrate on the group contractions $\text{Tr}(\lambda^1 \lambda^2 \lambda^3 \lambda^4 \lambda^5)$, which we shall do in the following, in *Eq.* (A.16) we have to perform the integral over the region:

$$\mathcal{I}_\pi = \{ z_4, z_5 \in \mathbf{R} \mid 1 < z_4 < z_5 < \infty \} . \tag{A.18}$$

For this case we obtain:

$$\begin{aligned} \tilde{F}[n_{24}, n_{25}, n_{34}, n_{35}, n_{45}] &= \int_1^\infty dz_4 \int_{z_4}^\infty dz_5 |z_4|^{\alpha_{24}} |z_5|^{\alpha_{25}} |1 - z_4|^{\alpha_{34}} |1 - z_5|^{\alpha_{35}} |z_4 - z_5|^{\alpha_{45}} \\ &= \int_0^1 dx \int_0^1 dy x^{-3 - \alpha_{24} - \alpha_{25} - \alpha_{34} - \alpha_{35} - \alpha_{45}} y^{-2 - \alpha_{25} - \alpha_{35} - \alpha_{45}} (1 - x)^{\alpha_{34}} (1 - y)^{\alpha_{45}} (1 - xy)^{\alpha_{35}} \\ &= \frac{\Gamma(-2 - \alpha_{24} - \alpha_{25} - \alpha_{34} - \alpha_{35} - \alpha_{45}) \Gamma(-1 - \alpha_{25} - \alpha_{35} - \alpha_{45}) \Gamma(1 + \alpha_{34}) \Gamma(1 + \alpha_{45})}{\Gamma(-1 - \alpha_{24} - \alpha_{25} - \alpha_{35} - \alpha_{45}) \Gamma(-\alpha_{25} - \alpha_{35})} \\ &\times {}_3F_2 \left[\begin{matrix} -2 - \alpha_{24} - \alpha_{25} - \alpha_{34} - \alpha_{35} - \alpha_{45}, & -1 - \alpha_{25} - \alpha_{35} - \alpha_{45}, & -\alpha_{35} \\ -1 - \alpha_{24} - \alpha_{25} - \alpha_{35} - \alpha_{45}, & -\alpha_{25} - \alpha_{35} \end{matrix} \right] . \end{aligned} \tag{A.19}$$

¹⁵ The remaining scalar products of momenta are expressed by these nine invariants:

$$\begin{aligned} k_1 k_2 &= s_3 + s_4 + s_5 \quad , \\ k_1 k_3 &= s_1 + s_2 + s_5 \quad , \\ k_1 k_4 &= -s_1 - s_3 - s_5 \quad , \\ k_1 k_5 &= -s_2 - s_4 - s_5 \quad , \\ k_2 k_3 &= -s_1 - s_2 - s_3 - s_4 - s_5 \quad . \end{aligned} \tag{A.14}$$

For the group structure π under consideration we may cast the integrals (A.7) into the form

$$\begin{aligned}
A_1^\pi(a, b, i, j, k) &= \sigma_{ij}\sigma_{jk} \left\{ (1 - k_i k_j) \left(\sigma_{ij}\sigma_{ja}\sigma_{ak} (\xi_k k_a) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ij} = -2 \\ \bar{n}_{jk}, \bar{n}_{ak} = -1, \bar{n}_{ja} = 1 \end{array} \right] \right. \right. \\
&\quad \left. \left. + \sigma_{ij}\sigma_{jb}\sigma_{bk} (\xi_k k_b) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ij} = -2 \\ \bar{n}_{jk}, \bar{n}_{bk} = -1, \bar{n}_{jb} = 1 \end{array} \right] \right) \right. \\
&\quad \left. + \sigma_{ik} [(\xi_k k_i)(k_j k_k) - (\xi_k k_j)(k_i k_k) - (1 - k_i k_j)(\xi_k k_i)] \tilde{F} \left[\begin{array}{l} \bar{n}_{ab} = -2 \\ \bar{n}_{ij}, \bar{n}_{ik}, \bar{n}_{jk} = -1 \end{array} \right] \right\}, \\
A_2^\pi(a, i, b, j, k) &= \sigma_{ab}\sigma_{ai}\sigma_{jk} \left\{ \sigma_{ij}\sigma_{bj}\sigma_{ja}\sigma_{ak} (\xi_k k_a)(k_i k_j) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ai}, \bar{n}_{ak} = -1 \\ \bar{n}_{bj}, \bar{n}_{ij}, \bar{n}_{jk} = -1, \bar{n}_{ja} = 1 \end{array} \right] \right. \\
&\quad \left. - \sigma_{bk}\sigma_{ij} (\xi_k k_b)(k_i k_j) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ai}, \bar{n}_{bk} = -1 \\ \bar{n}_{ij}, \bar{n}_{jk} = -1 \end{array} \right] \right. \\
&\quad \left. + \sigma_{bj}\sigma_{ik} [(\xi_k k_j)(k_i k_k) - (\xi_k k_i)(k_j k_k) - (\xi_k k_i)(k_i k_j)] \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ai}, \bar{n}_{bj} = -1 \\ \bar{n}_{ik}, \bar{n}_{jk} = -1 \end{array} \right] \right\}, \\
A_3^\pi(a, i, j, k, b) &= \sigma_{ab}\sigma_{ai}\sigma_{jk} \left\{ -\sigma_{bi}\sigma_{jk} (\xi_b k_i) (1 - k_j k_k) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ai}, \bar{n}_{bi} = -1 \\ \bar{n}_{jk} = -2, \end{array} \right] \right. \\
&\quad \left. - \sigma_{bj}\sigma_{ik} (\xi_b k_j) (k_i k_k) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ai}, \bar{n}_{bj} = -1 \\ \bar{n}_{ik}, \bar{n}_{jk} = -1 \end{array} \right] \right. \\
&\quad \left. + \sigma_{bk}\sigma_{ij} (\xi_b k_k) (k_i k_j) \tilde{F} \left[\begin{array}{l} \bar{n}_{ab}, \bar{n}_{ai}, \bar{n}_{bk} = -1 \\ \bar{n}_{ij}, \bar{n}_{jk} = -1 \end{array} \right] \right\}.
\end{aligned} \tag{A.20}$$

with $\sigma_{ij} = \text{sign}(i - j)$, and $\bar{n}_{ij} := \begin{cases} n_{ij}, & i < j \\ n_{ji}, & i > j \end{cases}$. The entries \bar{n}_{ij} in the functions \tilde{F} are to be understood such, that they are only of relevance, if they contribute in (A.16) or (A.17). Otherwise they are meaningless as a matter of the choice (3.21). Altogether, there appear 77 different functions \tilde{F} in the expressions (A.10) and (A.13).

Let us come back to the system of equations (A.11), written down for all 3×15 kinematics $(\xi\xi)(\xi\xi)(\xi k)$. With (A.20), it gives rise to 291 relations among the functions \tilde{F} , very much in the sense we have encountered in (2.22) for the case of four-gluon S -matrix or in subsection 3.3 for the six-gluon case: These relations are to be compared with *Eqs.* (3.47), *i.e.* they may be proven through partial integration. In addition, the 151 equations from the kinematics $(\xi\xi)(\xi k)(\xi k)(\xi k)$ give rise to simpler identities comparable to *Eqs.* (3.41) and (3.43), *i.e.* they follow from simple polynomial relations. Let us present

a selection of these relations:

$$\begin{aligned}
\tilde{F}[-2, 0, 0, -2, 1] &= \tilde{F}[-2, 0, 0, -1, 0] - \tilde{F}[-2, 0, 1, -2, 0] , \\
\tilde{F}[-2, -1, 0, -1, 0] &= \tilde{F}[-1, -1, 0, 0, -1] - \tilde{F}[-2, 0, 1, -1, -1] , \\
\tilde{F}[-1, -1, -1, -1, 0] &= \tilde{F}[-1, 0, -1, -1, 0] - \tilde{F}[-1, -, 1, -1, 0, 0] , \\
\tilde{F}[-1, 0, 0, -1, 0] &= \tilde{F}[-1, 0, 0, 0, -1] - \tilde{F}[-1, 0, 1, -1, -1] , \\
\tilde{F}[0, -2, -2, 0, 1] &= \tilde{F}[0, -2, -2, 1, 0] - \tilde{F}[0, -2, -1, 0, 0] .
\end{aligned} \tag{A.21}$$

After inserting for the \tilde{F} 's their definition (A.19) these equations appear like the theorems derived in *Ref.* [13].

As in the case of the six-gluon amplitude we may first solve for all equations following from the kinematics $(\xi\xi)(\xi k)(\xi k)(\xi k)$ and insert its solution into the equations derived from the kinematics $(\xi\xi)(\xi\xi)(\xi k)$. In doing so, solving for all 151 equations provides eight solutions, *i.e.* eight functions \tilde{F} out of the 77 may be expressed linearly in terms of others as in *Eq.* (A.21). Eliminating those eight functions in the second system of 291 equations leaves 275 equations to be solved. Of course, now these equations are more involved than in the four-gluon case and they describe various relations between hypergeometric functions ${}_3F_2$. This way we obtain a whole set of identities relating various functions \tilde{F} . These relations are similar to the identities for the hypergeometric function ${}_3F_2$, presented in [13]. Recall, that in the four-gluon scattering case (2.23) we have twelve equations and eleven functions F_j . However not all of these twelve equations are linear independent from each other. In effect, the system could be solved by introducing one single function F_0 and expressing all other eleven functions through F_0 , *cf.* *Eq.* (2.24). For the case at hand we encounter similar properties: Though the system of 275 equations is overdetermined, many of them are linear dependent on each other and it is possible to express 75 functions \tilde{F} in terms of a basis of two functions. In the following, the latter shall be denoted by Φ_1 and Φ_2 .

In the four-gluon it proved to be convenient, to express all eleven functions F_j ($j = 1, \dots, 11$) through one single function F_0 with special properties. The latter has no poles in the Mandelstam variables s, t or u , *cf.* the expansion (2.30). Similarly, in the six-gluon case we express 1,264 functions in terms of a basis of six finite functions Φ_j (3.54). All pole structure of the amplitudes has appeared through the algebraic equations (2.23) or (3.48). Similarly here, out of the total 77 functions ${}_3F_2$ only a few of them share the same

properties, namely they do not have any poles in the kinematic invariants s_i . *E.g.* we may single out the following two integrals

$$\begin{aligned}
\Phi_1 &:= \tilde{F}[-1, -2, 0, 0, 0] = \int_0^1 dx \int_0^1 dy x^{-s_1-s_2-s_3-s_4-s_5} y^{-s_2-s_4-s_5} \\
&\quad \times (1-x)^{s_3} (1-y)^{s_5} (1-xy)^{s_4}, \\
\Phi_2 &:= \tilde{F}[-1, -1, 0, -1, 0] = \int_0^1 dx \int_0^1 dy x^{-s_1-s_2-s_3-s_4-s_5} y^{-s_2-s_4-s_5} \\
&\quad \times (1-x)^{s_3} (1-y)^{s_5} (1-xy)^{s_4-1}
\end{aligned} \tag{A.22}$$

from the 77 functions \tilde{F} . The functions Φ_1, Φ_2 do not have poles in the invariants s_i . In fact, their expansion w.r.t. small s_i may be easily obtained from the expressions (4.55) and (4.56), *cf.* Eq. (A.27). According to (A.19) they may be also written as hypergeometric functions:

$$\begin{aligned}
\Phi_1 &= \frac{\Gamma(1-s_1-s_2-s_3-s_4-s_5) \Gamma(1-s_2-s_4-s_5) \Gamma(1+s_3) \Gamma(1+s_5)}{\Gamma(2-s_1-s_2-s_4-s_5) \Gamma(2-s_2-s_4)} \\
&\quad \times {}_3F_2 \left[\begin{matrix} 1-s_1-s_2-s_3-s_4-s_5, 1-s_2-s_4-s_5, -s_4 \\ 2-s_1-s_2-s_4-s_5, 2-s_2-s_4 \end{matrix} \right], \\
\Phi_2 &= \frac{\Gamma(1-s_1-s_2-s_3-s_4-s_5) \Gamma(1-s_2-s_4-s_5) \Gamma(1+s_3) \Gamma(1+s_5)}{\Gamma(2-s_1-s_2-s_4-s_5) \Gamma(2-s_2-s_4)} \\
&\quad \times {}_3F_2 \left[\begin{matrix} 1-s_1-s_2-s_3-s_4-s_5, 1-s_2-s_4-s_5, 1-s_4 \\ 2-s_1-s_2-s_4-s_5, 2-s_2-s_4 \end{matrix} \right].
\end{aligned} \tag{A.23}$$

In complete analogy to *Eqs.* (2.24) and (3.48) we may solve all remaining 275 equations through introducing the two-dimensional basis of functions Φ_1 and Φ_2 . All other 75 functions are given in terms of the latter

$$\tilde{F}[n_{24}, n_{25}, n_{34}, n_{35}, n_{45}] = \Lambda_{\{n_{ij}\}}^1(s_i) \tilde{F}[-1, -2, 0, 0, 0] + \Lambda_{\{n_{ij}\}}^2(s_i) \tilde{F}[-1, -1, 0, -1, 0]$$

(A.24)

with some polynomials $\Lambda_{\{n_{ij}\}}^1(s_i)$, $\Lambda_{\{n_{ij}\}}^2(s_i)$, which depend non-trivially on the five kinematic invariants s_i . Note, that due to the finiteness of Φ_1, Φ_2 , the pole structure of each of the 75 hypergeometric functions ${}_3F_2$ is determined by those prefactors $\Lambda_{\{n_{ij}\}}^j(s_i)$. The latter follow algebraically from solving the system of 275 equations. To demonstrate the

power of our method, let us present a few of those relations (A.24), we have found:

$$\begin{aligned}
\tilde{F}[-1, -1, 0, 0, -1] &= \frac{1-s_2}{s_5} \Phi_1 - \frac{s_4}{s_5} \Phi_2 , \\
\tilde{F}[-1, -1, -1, 0, 0] &= -\frac{(1-s_2)(1-s_1-s_2-s_5)}{s_3(s_2+s_4+s_5)} \Phi_1 + \frac{(1-s_2+s_3)s_4}{s_3(s_2+s_4+s_5)} \Phi_2 , \\
\tilde{F}[-2, 0, 0, 0, -1] &= \frac{(1-s_2)[(1-s_1)s_2+s_4(1-s_1-s_5)]}{(1-s_1)s_5(s_2+s_4+s_5)} \Phi_1 \\
&\quad - \frac{s_4[(1-s_1)s_2+s_4(1-s_1-s_5)-s_5(s_1+s_3+s_5)]}{(1-s_1)s_5(s_2+s_4+s_5)} \Phi_2 , \\
\tilde{F}[0, -2, -1, 0, 0] &= \frac{(1-s_1-s_2-s_5)}{s_3} \Phi_1 - \frac{s_4}{s_3} \Phi_2 , \\
\tilde{F}[0, -2, 0, -1, 0] &= \frac{1-s_1-s_2-s_5}{1+s_1+s_3+s_5} \Phi_1 + \frac{s_1-s_4}{1+s_1+s_3+s_5} \Phi_2 , \\
\tilde{F}[-1, -1, -1, -1, 1] &= -\frac{(1-s_2)(1-s_1-s_2-s_5)}{s_3(s_2+s_4+s_5)} \Phi_1 + \frac{s_4-s_2(s_3+s_4)-s_3s_5}{s_3(s_2+s_4+s_5)} \Phi_2 .
\end{aligned} \tag{A.25}$$

In addition, let us present the somewhat more involved relation:

$$\begin{aligned}
&\tilde{F}[0, 0, -1, 0, -1] \\
&= \frac{(1-s_2)(1-s_1-s_2-s_5)[s_1s_3(s_2+s_4)+s_1s_2s_5+s_2(s_3+s_5)(s_2+s_4+s_5)]}{s_3s_5(s_2+s_4+s_5)(s_3+s_4+s_5)(s_1+s_2+s_3+s_4+s_5)} \Phi_1 \\
&\quad - \left\{ \frac{(1-s_1-s_2)(s_1+s_2)(s_2+s_4)}{s_5(s_2+s_4+s_5)(s_3+s_4+s_5)(s_1+s_2+s_3+s_4+s_5)} \right. \\
&\quad \left. + \frac{(1-s_2)s_2-s_1s_3}{s_3(s_2+s_4+s_5)(s_3+s_4+s_5)} \right\} s_4 \Phi_2 .
\end{aligned} \tag{A.26}$$

The importance of this identity becomes clear, once we insert (A.19) into this equation. According to Eq. (A.19) the function $\tilde{F}[0, 0, -1, 0, -1]$ corresponds to the integral

$$\begin{aligned}
\tilde{F}[0, 0, -1, 0, -1] &= \int_0^1 dx \int_0^1 dy x^{-1-s_1-s_2-s_3-s_4-s_5} y^{-1-s_2-s_4-s_5} \\
&\quad \times (1-x)^{-1+s_3} (1-y)^{-1+s_5} (1-xy)^{s_4} ,
\end{aligned}$$

which has several double poles in the invariants s_i . However, since the functions Φ_1, Φ_2 in the expansion (A.26) do not have any poles, all the poles of $\tilde{F}[0, 0, -1, 0, -1]$ are encoded in the fractions in front of those functions and we may easily obtain the full power series of $\tilde{F}[0, 0, -1, 0, -1]$ by only knowing the series expansions of Φ_1, Φ_2 . Furthermore, this equation becomes (4.57). Hence, we have proven the non-trivial equation (4.57) just from relating equivalent parts of the five-gluon string S -matrix. According to (A.19) the

above identities (A.21), (A.25) and (A.26) give rise to non-trivial relations among three hypergeometric functions ${}_3F_2$, which only partly may be found in the literature [13]. We should stress, that the relations listed above are just twelve out of 75 identities we obtain from our method.

After having solved all equations, the full α' -dependence of the five gluon amplitude is captured by the two functions Φ_1, Φ_2 , given in *Eqs.* (A.22) or (A.23). Hence, to obtain the α' -expansion of (A.1), we only have to know their power series expansions w.r.t. to the kinematical invariants s_i . With the material, derived in section 4, in particular setting $a = -s_1 - s_2 - s_3 - s_4 - s_5$, $b = -s_2 - s_4 - s_5$, $c = s_3$, $d = s_5$ and $e = s_4$ in (4.55) and (4.56), we obtain:

$$\begin{aligned}
\Phi_1 &= 1 + s_1 + 2 s_2 + s_5 + s_4 \zeta(2) + s_1^2 + 3 s_1 s_2 + 3 s_2^2 + 2 s_1 s_5 + 3 s_2 s_5 + s_5^2 \\
&\quad + (s_1 s_3 + s_2 s_3 + s_3^2 + s_1 s_4 + 2 s_2 s_4 + 2 s_3 s_4 + s_4^2 + s_2 s_5 + s_3 s_5 + 3 s_4 s_5 + s_5^2) \zeta(2) \\
&\quad + (s_1 s_4 + 2 s_2 s_4 - s_3 s_4 + s_4^2) \zeta(3) + \dots , \\
\Phi_2 &= \zeta(2) + (s_1 + 2 s_2 - s_3 + s_4) \zeta(3) \\
&\quad + \left(s_1^2 + 3 s_1 s_2 + 3 s_2^2 + \frac{3}{2} s_1 s_3 + \frac{5}{4} s_2 s_3 + \frac{7}{2} s_3^2 + \frac{11}{4} s_1 s_4 + \frac{11}{2} s_2 s_4 + 4 s_3 s_4 + \right. \\
&\quad \left. + \frac{7}{2} s_4^2 + \frac{7}{4} s_1 s_5 + \frac{17}{4} s_2 s_5 + \frac{17}{4} s_3 s_5 + \frac{27}{4} s_4 s_5 + \frac{17}{4} s_5^2 \right) + \dots .
\end{aligned} \tag{A.27}$$

These two functions Φ_1, Φ_2 have no poles in the kinematic invariants s_i and their momentum expansion may be obtained by the methods presented in section 4. This property is just in lines of (2.30) in the four-gluon case and (3.54) in the six-gluon case.

E.g. in the string S -matrix (A.1) after eliminating $\xi_5 k_4$ on-shell the kinematics $(\xi_1 \xi_2)(\xi_3 \xi_4)(\xi_5 k_1)$ comes along with the momentum dependent expression:

$$\begin{aligned}
&(\xi_1 \xi_2)(\xi_3 \xi_4)(\xi_5 k_1) \frac{1}{s_3 s_5 (s_2 + s_4 + s_5) (s_3 + s_4 + s_5)} \\
&\times \left\{ (1 - s_2) (1 - s_1 - s_2 - s_5) [s_1 s_3 (s_2 + s_4) + s_1 s_2 s_5 + s_2 (s_3 + s_5) (s_2 + s_4 + s_5)] \Phi_1 \right. \\
&\quad + s_4 \{ (-1 + s_1 + s_2) (s_1 + s_2) s_3 (s_2 + s_4) - [(1 - s_2) s_2 - s_1 s_3] (s_1 + s_2 + s_3 + s_4) s_5 \\
&\quad \left. - [(1 - s_2) s_2 - s_1 s_3] s_5^2 \} \Phi_2 \right\} .
\end{aligned} \tag{A.28}$$

According to *Eq.* (A.8) the above expression may be derived from $A_1(1, 2, 3, 4, 5)$. On the other hand, in the string S -matrix (A.1) after eliminating $\xi_5 k_4, \xi_4 k_5$ and $\xi_3 k_5$ on-shell the kinematics $(\xi_1 \xi_2)(\xi_3 k_1)(\xi_4 k_2)(\xi_5 k_1)$ comes along with the momentum dependent

expression:

$$\begin{aligned}
& (\xi_1 \xi_2)(\xi_3 k_1)(\xi_4 k_2)(\xi_5 k_1) \frac{1}{s_5 (s_3 + s_4 + s_5)} \\
& \times \{ (1 - s_2) (1 - s_1 - s_2 - s_5) \Phi_1 - [s_4 (1 - s_1 - s_2 - s_5) - s_5 (s_1 + s_3 + s_5)] \Phi_2 \} .
\end{aligned} \tag{A.29}$$

According to *Eq.* (A.12) the above expression may be derived from $C_1(1, 2, 3, 4, 5)$. All other kinematics may be expressed similarly. The result (A.28) is to be compared with the final result (3.55) of the six-gluon case. To conclude, we may express the full five gluon kinematics (A.10) through the two functions Φ_1, Φ_2 , which are rather easy to handle due to their pole structure. In fact, after expanding (A.28) and (A.29) up to second order in the Mandelstam variables, we find agreement with the results of [6]. In addition, we shall point out, that in the beautiful works [7], it has also been found, that the five-gluon string S -matrix may be expressed by two single hypergeometric functions ${}_3F_2$.

Appendix B. Series involving the Harmonic Number

For the power series expansion of the hypergeometric function ${}_3F_2$ we shall need the nine series:

$$\begin{aligned}
(i) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \psi^{(1)}(n+1) = 1 , \\
(ii) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \psi^{(1)}(n+2) = \zeta(2) - \zeta(3) , \\
(iii) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \psi^{(1)}(n+3) = -3 + 2 \zeta(2) , \\
(iv) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} H_{n+1} \psi(n+1) = (1 - \gamma_E) \zeta(2) + \zeta(3) , \\
(v) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} H_{n+2} \psi(n+1) = 3 - 2 \gamma_E ,
\end{aligned} \tag{B.1}$$

$$\begin{aligned}
(vi) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} H_{n+2} \psi(n+2) = -2 \gamma_E + \zeta(2) + 2 \zeta(3) , \\
(vii) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \psi(n+1)^2 = (1 - \gamma_E)^2 + \zeta(2) , \\
(viii) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \psi(n+2)^2 = \gamma_E^2 - 2 \gamma_E \zeta(2) + 3 \zeta(3) , \\
(ix) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)} \psi(n+3)^2 = 3 - 4 \gamma_E + \gamma_E^2 + \zeta(2) .
\end{aligned}$$

These identities may be proven by using (4.23) and applying formulae shown in subsection

4.3. Furthermore, we prove the three series:

$$\begin{aligned}
(i) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)^2} \psi(n+1)^2 = \frac{11}{4} \zeta(4) - 2 \gamma_E \zeta(3) + \gamma_E^2 \zeta(2) , \\
(ii) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)^2} \psi(n+2)^2 = \frac{17}{4} \zeta(4) - 4 \gamma_E \zeta(3) + \gamma_E^2 \zeta(2) , \\
(iii) \quad & \sum_{n=0}^{\infty} \frac{1}{(n+1)^2} \psi^{(1)}(n+1)^2 = \frac{1}{3} \zeta(2)^3 - \zeta(3)^2 - \frac{5}{3} \zeta(2) \zeta(4) + \frac{17}{3} \zeta(6) .
\end{aligned} \tag{B.2}$$

After rewriting the sums with (4.23), these identities may be proven by using various

relations presented in subsection 4.3, in particular *Eq.* (4.54).

Appendix C. Momentum expansion of ${}_4F_3 \left[\begin{matrix} 1+b, 1+a, 1+c, -j \\ 2+b+e, 2+a+d, 2+c+f \end{matrix} \right]$

$$\begin{aligned}
& \frac{\Gamma(1+d) \Gamma(1+e) \Gamma(1+f) \Gamma(1+b) \Gamma(1+a) \Gamma(1+c)}{\Gamma(2+b+e) \Gamma(2+a+d) \Gamma(2+c+f)} {}_4F_3 \left[\begin{matrix} 1+b, 1+a, 1+c, -j \\ 2+b+e, 2+a+d, 2+c+f \end{matrix} \right] \\
&= 1 - a - b - c - d - e - f - 3j + a^2 + ba + ca + 2da + ea + fa + 4ja + b^2 + c^2 + d^2 \\
&+ e^2 + f^2 + 6j^2 + bc + bd + cd + 2be + ce + de + bf + 2cf + df + ef + 4bj + 4cj + 4dj \\
&+ 4ej + 4fj - a^3 - ba^2 - ca^2 - 3da^2 - ea^2 - fa^2 - 5ja^2 - b^2a - c^2a - 3d^2a - e^2a \\
&- f^2a - 10j^2a - bca - 2bda - 2cda - 2bea - cea - 2dea - bfa - 2cfa - 2dfa - efa \\
&- 5bja - 5cja - 10dja - 5eja - 5fja - b^3 - c^3 - d^3 - e^3 - f^3 - 10j^3 - bc^2 - bd^2 - cd^2 \\
&- 3be^2 - ce^2 - de^2 - bf^2 - 3cf^2 - df^2 - ef^2 - 10bj^2 - 10cj^2 - 10dj^2 - 10ej^2 - 10fj^2 \\
&- b^2c - b^2d - c^2d - bcd - 3b^2e - c^2e - d^2e - 2bce - 2bde - cde - b^2f - 3c^2f - d^2f \\
&- e^2f - 2bcf - bdf - 2cdf - 2bef - 2cef - def - 5b^2j - 5c^2j - 5d^2j - 5e^2j - 5f^2j \\
&- 5bcj - 5bdj - 5cdj - 10bej - 5cej - 5dej - 5bfj - 10cfj - 5dfj - 5efj \\
&+ \zeta(2) \{ j - 2j^2 - aj - bj - cj - ad - be - cf + 3j^3 + 3aj^2 + 3bj^2 + 3cj^2 + a^2j + b^2j \\
&+ c^2j + abj + acj + bcj + 4adj + 4bej - dej + 4cfj - dfj - efj + ad^2 + be^2 + cf^2 + a^2d \\
&+ abd + acd + b^2e + abe + bce + ade + bde + c^2f + acf + bcf + adf + cdf + bef + cef \} \\
&+ \frac{1}{4} \zeta(4) \{ -j^2 - 4aj - 4bj - 4cj - 5dj - 5ej - 5fj + 5j^3 + 5aj^2 + 5bj^2 + 5cj^2 + 17dj^2 \\
&+ 17ej^2 + 17fj^2 + 4a^2j + 4b^2j + 4c^2j + 12d^2j + 12e^2j + 12f^2j + 4abj + 4acj + 4bcj \\
&+ 2adj + 5bdj + 5cdj + 5aej + 2bej + 5cej + 17dej + 5afj + 5bfj + 2cfj + 17dfj + 17efj \} \\
&+ \zeta(3) \{ j - 2j^2 - aj - bj - cj - 2dj - 2ej - 2fj + 3j^3 + 3aj^2 + 3bj^2 + 3cj^2 + 4dj^2 + 4ej^2 \\
&+ 4fj^2 + a^2j + b^2j + c^2j + abj + acj + bcj + 2adj + 2bdj + 2cdj + 2aej + 2bej + 2cej \\
&+ 2afj + 2bfj + 2cfj + ad^2 + be^2 + cf^2 + a^2d + b^2e + c^2f \} \\
&+ \zeta(2) \zeta(3) \{ -j^3 - aj^2 - bj^2 - cj^2 + d^2j + e^2j + f^2j + adj - bdj - cdj - aej + bej - cej \\
&- dej - afj - bfj + cfj - dfj - efj \} \\
&+ \frac{1}{2} \zeta(5) \{ 4j^3 + 4aj^2 + 4bj^2 + 4cj^2 + dj^2 + ej^2 + fj^2 + 2a^2j + 2b^2j + 2c^2j - d^2j - e^2j \\
&- f^2j + 2abj + 2acj + 2bcj - 3adj + 6bdj + 6cdj + 6aej - 3bej + 6cej + 7dej + 6afj \\
&+ 6bfj - 3cfj + 7dfj + 7efj \} + \dots
\end{aligned} \tag{C.1}$$

Appendix D. Some group theoretical facts

The field strength $F_{\mu\nu}^a$ is defined as:

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c , \quad (\text{D.1})$$

with the structure constants f^{abc} for an $U(N)$ gauge group. Let us recall some basic facts about Lie Algebras. The commutation relations of a representation T^a of the Lie Algebra read:

$$[T^a, T^b] = i f^{abc} T^c , \quad (\text{D.2})$$

with the structure constants f^{abc} . We impose the standard normalization condition

$$\text{Tr}(T^a T^b) = C(r) \delta^{ab} , \quad (\text{D.3})$$

with $C(r)$ being a constant for each representation r . Then we have the relation

$$f^{abc} = -\frac{i}{C(r)} \text{Tr} ([T^a, T^b] T^c) , \quad (\text{D.4})$$

which implies, that f^{abc} is totally anti-symmetric.

The adjoint representation $r = G$ is given by the matrices

$$(T^a)_{bc} = -i f^{abc} , \quad (\text{D.5})$$

which obviously satisfies (D.2) and (D.4). The covariant derivative D_μ^{adj} acting on fields in the adjoint representation is introduced as:

$$(D_\lambda^{\text{adj}})_{ab} = \partial_\lambda \delta_{ab} - i g A_\lambda^m (T^m)_{ab} = \partial_\lambda \delta_{ab} - g f_{mab} A_\lambda^m . \quad (\text{D.6})$$

Hence, we have:

$$D_\lambda F_{\mu\nu}^a = \partial_\lambda F_{\mu\nu}^a - i g A_\lambda^m (T^m)_{ab} F_{\mu\nu}^b . \quad (\text{D.7})$$

In addition, we derive:

$$\begin{aligned} D_\kappa D_\lambda F_{\mu\nu}^a &= \partial_\kappa \partial_\lambda F_{\mu\nu}^a + g f^{and} A_\kappa^n \partial_\lambda F_{\mu\nu}^d - g f^{mac} \partial_\kappa (A_\lambda^m F_{\mu\nu}^c) \\ &+ g^2 f^{amc} f^{mnd} A_\kappa^n A_\lambda^d F_{\mu\nu}^c + g^2 f^{amc} f^{cnd} A_\lambda^m A_\kappa^n F_{\mu\nu}^d . \end{aligned} \quad (\text{D.8})$$

Finally, for $F_{\mu\nu} \equiv T^a F_{\mu\nu}^a$ in the adjoint representation, we may write (D.1) as:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i g [A_\mu, A_\nu] , \quad (\text{D.9})$$

and (D.7) as:

$$D_\lambda = \partial_\lambda - i g [A_\lambda, \star] . \quad (\text{D.10})$$

Furthermore, Eq. (D.8) gives rise to:

$$\begin{aligned} D_\kappa (D_\lambda F_{\mu\nu}) &= \partial_\kappa \partial_\lambda F_{\mu\nu} - i g [A_\kappa, \partial_\lambda F_{\mu\nu}] - i g [\partial_\kappa A_\lambda, F_{\mu\nu}] \\ &- i g [A_\lambda, \partial_\kappa F_{\mu\nu}] - g^2 [A_\kappa, [A_\lambda, F_{\mu\nu}]] \end{aligned} \quad (\text{D.11})$$

To this end, we may prove:

$$[D_\mu, D_\nu] F_{\rho\sigma} = -i g [F_{\mu\nu}, F_{\rho\sigma}] . \quad (\text{D.12})$$

References

- [1] M.B. Green and J.H. Schwarz, “Supersymmetrical Dual String Theory. 2. Vertices And Trees,” Nucl. Phys. B **198**, 252 (1982);
J.H. Schwarz, “Superstring Theory,” Phys. Rept. **89**, 223 (1982).
- [2] D.J. Gross and E. Witten, “Superstring Modifications Of Einstein’s Equations,” Nucl. Phys. B **277**, 1 (1986);
H. Kawai, D.C. Lewellen and S.H.H. Tye, “A Relation Between Tree Amplitudes Of Closed And Open Strings,” Nucl. Phys. B **269**, 1 (1986).
- [3] A.A. Tseytlin, “Vector Field Effective Action In The Open Superstring Theory,” Nucl. Phys. B **276**, 391 (1986) [Erratum-ibid. B **291**, 876 (1987)].
- [4] A.A. Tseytlin, “On non-abelian generalisation of the Born-Infeld action in string theory,” Nucl. Phys. B **501**, 41 (1997) [arXiv:hep-th/9701125].
- [5] A. Bilal, “Higher-derivative corrections to the non-abelian Born-Infeld action,” Nucl. Phys. B **618**, 21 (2001) [arXiv:hep-th/0106062].
- [6] R. Medina, F.T. Brandt and F.R. Machado, “The open superstring 5-point amplitude revisited,” JHEP **0207**, 071 (2002) [arXiv:hep-th/0208121].
- [7] L.A. Barreiro and R. Medina, “5-field terms in the open superstring effective action,” JHEP **0503**, 055 (2005) [arXiv:hep-th/0503182];
F. Machado and R. Medina, “The open superstring and the non-abelian Born-Infeld theory,” Nucl. Phys. Proc. Suppl. **127**, 166 (2004);
O. Chandia and R. Medina, “4-point effective actions in open and closed superstring theory,” JHEP **0311**, 003 (2003) [arXiv:hep-th/0310015];
F. Brandt, F. Machado and R. Medina, “Open and closed superstring five-point amplitudes at tree-level,” Int. J. Mod. Phys. A **18**, 2127 (2003).
- [8] M. de Roo and M.G.C. Eenink, “The effective action for the 4-point functions in abelian open superstring theory,” JHEP **0308**, 036 (2003) [arXiv:hep-th/0307211];
A. Collinucci, M. De Roo and M.G.C. Eenink, “Supersymmetric Yang-Mills theory at α'^3 ,” JHEP **0206**, 024 (2002) [arXiv:hep-th/0205150]; “Derivative corrections in 10-dimensional super-Maxwell theory,” JHEP **0301**, 039 (2003) [arXiv:hep-th/0212012].
- [9] M.R. Garousi and R.C. Myers, “World-volume potentials on D-branes,” JHEP **0011**, 032 (2000) [arXiv:hep-th/0010122].
- [10] D. Lüst, P. Mayr, R. Richter and S. Stieberger, “Scattering of gauge, matter, and moduli fields from intersecting branes,” Nucl. Phys. B **696**, 205 (2004) [arXiv:hep-th/0404134].
- [11] M.D. Fisk, “Five Point Tree Amplitude For The Heterotic String,” Nucl. Phys. B **320**, 377 (1989).
- [12] Y. Kitazawa, “Effective Lagrangian For Open Superstring From Five Point Function,” Nucl. Phys. B **289**, 599 (1987).

- [13] W.N. Bailey, *Generalised Hypergeometric Series*, Cambridge, England: Cambridge University Press, 1935.
- [14] A. Hashimoto and W.I. Taylor, “Fluctuation spectra of tilted and intersecting D-branes from the Born-Infeld action,” Nucl. Phys. B **503**, 193 (1997) [arXiv:hep-th/9703217].
- [15] A. Sevrin and A. Wijns, “Higher order terms in the non-Abelian D-brane effective action and magnetic JHEP **0308**, 059 (2003) [arXiv:hep-th/0306260];
P. Koerber and A. Sevrin, “The non-abelian D-brane effective action through order α'^4 ,” JHEP **0210**, 046 (2002) [arXiv:hep-th/0208044]. “Testing the α'^3 term in the non-abelian open superstring effective JHEP **0109**, 009 (2001) [arXiv:hep-th/0109030]. “The non-Abelian Born-Infeld action through order α'^3 ,” JHEP **0110**, 003 (2001) [arXiv:hep-th/0108169];
A. Sevrin, J. Troost and W. Troost, “The non-abelian Born-Infeld action at order F^6 ,” Nucl. Phys. B **603**, 389 (2001) [arXiv:hep-th/0101192];
L. De Fosse, P. Koerber and A. Sevrin, “The uniqueness of the Abelian Born-Infeld action,” Nucl. Phys. B **603**, 413 (2001) [arXiv:hep-th/0103015];
P. Koerber, “Abelian and non-Abelian D-brane effective actions,” Fortsch. Phys. **52**, 871 (2004) [arXiv:hep-th/0405227].
- [16] W. Taylor, “Lectures on D-branes, tachyon condensation, and string field theory,” arXiv:hep-th/0301094;
W. Taylor and B. Zwiebach, “D-branes, tachyons, and string field theory,” arXiv:hep-th/0311017.
- [17] P. Koerber and A. Sevrin, “The non-abelian D-brane effective action through order α'^4 ,” JHEP **0210**, 046 (2002) [arXiv:hep-th/0208044];
A. Sevrin and A. Wijns, “Higher order terms in the non-Abelian D-brane effective action and magnetic background fields,” JHEP **0308**, 059 (2003) [arXiv:hep-th/0306260].
- [18] Work in progress.
- [19] R.C. Myers, “Dielectric-branes,” JHEP **9912**, 022 (1999) [arXiv:hep-th/9910053].
- [20] Z. Koba and H.B. Nielsen, “Reaction Amplitude For N Mesons: A Generalization Of The Veneziano-Bardakci-Ruegg-Virasora Model,” Nucl. Phys. B **10**, 633 (1969);
“Manifestly Crossing Invariant Parametrization Of N Meson Amplitude,” Nucl. Phys. B **12**, 517 (1969).
- [21] S. Stieberger and T.R. Taylor, work to appear.
- [22] M.L. Mangano and S.J. Parke, “Multiparton Amplitudes In Gauge Theories,” Phys. Rept. **200**, 301 (1991);
M.L. Mangano, S.J. Parke and Z. Xu, “Duality And Multi - Gluon Scattering,” Nucl. Phys. B **298**, 653 (1988).

- [23] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, Academic Press 1994.
- [24] L.J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, 1966.
- [25] P. Appell, *Sur le fonctions hypergéométriques de plusieurs variables*, Paris: Gauthier-Villars, 1925;
 J. Kampé de Fériet, *La fonction hypergéométrique*, Paris: Gauthier-Villars, 1937;
 H. Exton, *Multiple Hypergeometric Functions and Applications*, Chichester, England: Ellis Horwood, 1976;
 H.M. Srivastava, and P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Chichester, England: Ellis Horwood, 1985.
- [26] P. Appell, and J. Kampé de Fériet, *Fonctions hypergéométriques et hypersphériques: polynomes d'Hermite*, Paris: Gauthier-Villars, 1926.
- [27] D. Borwein, and J.M. Borwein, *On an Intriguing Integral and Some Series Related to $\zeta(4)$* , Proc. Amer. Math. Soc. 123, 1191 (1995).
- [28] D.H. Bailey, J.M. Borwein, and R. Girgensohn, *Experimental evaluation of Euler sums*, Experiment. Math. 3 (1994), 17-30.
- [29] D. Borwein, J.M. Borwein, and R. Girgensohn, *Explicit evaluation of Euler sums*, Proc. Edinburgh Math. Soc. 38 (1995), 277-294.
- [30] J.M. Borwein and R. Girgensohn, *Evaluation of triple Euler sums*, with appendix *Euler sums in quantum field theory* by D. J. Broadhurst, Electronic J. Combinatorics 3 (1996), R23.
- [31] N.E. Nörlund, *Vorlesungen über Differenzenrechnung*, Berlin: Springer, 1924.
- [32] J.M. Borwein, D.M. Bradley, D.J. Broadhurst, and P. Lisonek, *Special values of multidimensional polylogarithms*, Trans. Amer. Math. Soc. 353 (2001), 907-941;
 D.J. Broadhurst, J. M. Borwein, and D. M. Bradley, *Evaluation of k -fold Euler/Zagier sums: a compendium of results for arbitrary k* , Electronic J. Combinatorics 4(2) (1997), R5.
- [33] D. Zagier, *Values of zeta functions and their applications*, First European Congress of Mathematics, Paris, 1992, Vol. II, Ed. A. Joseph et. al, Birkhäuser, Basel, 1994, pp. 497–512.
- [34] R.E. Crandall and J.P. Buhler *On the evaluation of Euler sums*, Experiment. Math. 3 (1994) 275;
 K. Boyadzhiev, *Evaluation of Euler-Zagier sums*, Internat. J. Math. Math. Sci. 27 (2001), 407-412.
- [35] L. Tornheim, *Harmonic double series*, Amer. J. Math. 72 (1950), 303-314;
 O. Espinosa and V. H. Moll, *The evaluation of Tornheim double sums. Part 1*, arXiv: math-ca/0505647.
- [36] D.J. Gross and J.H. Sloan, “The Quartic Effective Action For The Heterotic String,” Nucl. Phys. B **291**, 41 (1987).

- [37] S. Stieberger and T.R. Taylor, “Non-Abelian Born-Infeld action and type I - heterotic duality (I): Heterotic F^6 terms at two loops,” Nucl. Phys. B **647**, 49 (2002); [arXiv:hep-th/0207026].
- [38] N. Berkovits, “Multiloop amplitudes and vanishing theorems using the pure spinor formalism for the superstring,” JHEP **0409**, 047 (2004) [arXiv:hep-th/0406055]; “Covariant multiloop superstring amplitudes,” arXiv:hep-th/0410079.
- [39] E. Witten, “String theory dynamics in various dimensions,” Nucl. Phys. B **443** (1995) 85 [arXiv:hep-th/9503124];
J. Polchinski and E. Witten, “Evidence for Heterotic - Type I String Duality,” Nucl. Phys. B **460** (1996) 525 [arXiv:hep-th/9510169].
- [40] A.A. Tseytlin, “On SO(32) heterotic - type I superstring duality in ten dimensions,” Phys. Lett. B **367** (1996) 84 [arXiv:hep-th/9510173]; “Heterotic - type I superstring duality and low-energy effective actions,” Nucl. Phys. B **467** (1996) 383 [arXiv:hep-th/9512081].
- [41] C. Bachas and E. Kiritsis, “ F^4 terms in $N = 4$ string vacua,” Nucl. Phys. Proc. Suppl. **55B**, 194 (1997) [arXiv:hep-th/9611205];
C. Bachas, C. Fabre, E. Kiritsis, N.A. Obers and P. Vanhove, “Heterotic/type-I duality and D-brane instantons,” Nucl. Phys. B **509**, 33 (1998) [arXiv:hep-th/9707126];
E. Kiritsis and N.A. Obers, “Heterotic/type-I duality in $D < 10$ dimensions, threshold corrections and D-instantons,” JHEP **9710**, 004 (1997) [arXiv:hep-th/9709058].
- [42] W. Lerche and S. Stieberger, “Prepotential, mirror map and F-theory on K3,” Adv. Theor. Math. Phys. **2**, 1105 (1998) [Erratum-ibid. **3**, 1199 (1999)] [arXiv:hep-th/9804176];
W. Lerche, S. Stieberger and N.P. Warner, “Quartic gauge couplings from K3 geometry,” Adv. Theor. Math. Phys. **3**, 1575 (1999) [arXiv:hep-th/9811228]; “Prepotentials from symmetric products,” Adv. Theor. Math. Phys. **3**, 1613 (1999) [arXiv:hep-th/9901162];
K. Foerger and S. Stieberger, “Higher derivative couplings and heterotic-type I duality in eight dimensions,” Nucl. Phys. B **559**, 277 (1999) [arXiv:hep-th/9901020].
- [43] S. Stieberger and T.R. Taylor, “Non-Abelian Born-Infeld action and type I - heterotic duality (II): Nonrenormalization theorems,” Nucl. Phys. B **648**, 3 (2003) [arXiv:hep-th/0209064].
- [44] E. D’Hoker and D.H. Phong, “Two-loop superstrings. VI: Non-renormalization theorems and the 4-point function,” arXiv:hep-th/0501197; “Two-loop superstrings. V: Gauge slice independence of the N-point function,” arXiv:hep-th/0501196.
- [45] S. Stieberger and T.R. Taylor, work to appear.
- [46] C.L. Siegel, *Symplectic Geometry*, Am. Jour. Math. 65 (1943), 1-86.
- [47] S.J. Parke and T.R. Taylor, “An Amplitude For N Gluon Scattering,” Phys. Rev. Lett. **56**, 2459 (1986).

- [48] E. Witten, “Perturbative gauge theory as a string theory in twistor space,” *Commun. Math. Phys.* **252**, 189 (2004) [arXiv:hep-th/0312171].
- [49] F. Cachazo, P. Svrcek and E. Witten, “MHV vertices and tree amplitudes in gauge theory,” *JHEP* **0409**, 006 (2004) [arXiv:hep-th/0403047];
R. Britto, F. Cachazo, B. Feng and E. Witten, “Direct proof of tree-level recursion relation in Yang-Mills theory,” arXiv:hep-th/0501052;
R. Britto, F. Cachazo and B. Feng, “New recursion relations for tree amplitudes of gluons,” arXiv:hep-th/0412308;
F. Cachazo and P. Svrcek, “Lectures on twistor strings and perturbative Yang-Mills theory,” arXiv:hep-th/0504194.