## Generalized coordinate gauge and nonabelian Stokes theorem

V.I.Shevchenko\*and Yu.A.Simonov<sup>†</sup>

Institute for Theoretical and Experimental Physics 117218, B.Cheremushkinskaya, 25, Moscow, Russia

## Abstract

A contour gauge of general type is analysed where 1-form (vector potential) is expressed as a contour integral of the 2-form (field strength) along an arbitrary contour C. For a special class of contours the gauge condition reduces to  $k_{\mu}(x)A_{\mu}(x) = 0$  where  $k_{\mu}(x)$  is a tangent vector to the contour C. It is shown, that this gauge is advantageous to give a simple proof of the nonabelian Stokes theorem.

<sup>\*</sup>shevchen@heron.itep.ru †simonov@vxitep.itep.ru

The procedure of the gauge fixing is an essential part of QCD [1] and however final results do not depend on the gauge, different forms of gauge conditions are useful in different settings of physical problems. For example, in high energy scattering in QCD the axial gauge has proved to be useful [2], while in the OPE analysis [3] the Fock-Schwinger [4] (sometimes called the coordinate or radial) gauge was applied (for discussions and derivation see [5] and also [6]).

In another physical situation, where the time axis is singled out, as e.g. in the heavy quarkonium theory, the modified coordinate gauge [7] can be convenient. This gauge was used recently in the context of equations for the quark [8] and gluon [9] Green's functions, displaying the property of chiral symmetry breaking and confinement.

There is another set of studies where an emphasis is made on formulation of gauge theory without gauge-dependent degrees of freedom from the very beginning, and the role of dynamical variables is played by 2-forms [10] or loop variables [11]. These completely gauge-invariant approaches encountered their own difficulties and as a matter of fact many gauge-invariant observables are easier to calculate using gauge-dependent diagrammatic rules.

Both in the coordinate gauge [4] and in its modified form [7] the shape of the contour C(x), in the integral, connecting vector potential and the field strength,

$$A_{\mu}(x) = \int_{C(x)} dz_{\nu} \alpha_{\rho\mu}(z) F_{\nu\rho}(z)$$
(1)

is fixed and consists of straight lines. Inessential for physical results, it may be inconvenient in the course of computations. In particular, in the confining phase of QCD, when the QCD string is formed between two colour charges it would be advantageous to choose the contours C lying on the world sheet of the string; in this case one could do simplifying approximations as in [8, 9], namely to keep only Gaussian field correlator. The decoupling of ghosts, known to occur for the gauges (1) (see [12], [13] and references therein) is also an attractive feature, which suggests to look for generalizations of (1) with arbitrary contours C.

The gauge condition of the type we are interested in was introduced for the first time in [13]. In the present paper we give a refined treatment of this gauge, paying special attention to some important details, missing in the original paper. Let us briefly mention them. To define this gauge condition correctly, the set of contours C, determining the gauge must satisfy some additional requirement (eq.(3) of the present paper). This condition is essential for the representation (1) to hold true. With this requirement we are also able to formulate the gauge condition in the local form (eq.(12) of the present paper). An immediate use of the generalized contour gauge which also has not yet been discussed in the literature is the ability to give a short and direct proof of the nonabelian Stokes theorem [14, 15] as we do below in this paper.

Let us proceed with the definition of the generalized contour gauge. Let M be a d-dimensional connected Euclidean manifold. We choose some subspace  $M_0$ ,  $M_0 \subset M$  which in general may be disconnected and of lower dimension than d. For our purposes it is sufficient to take  $M_0$  consisting of the only one point  $x_0$ . If  $M_0$  is of more complicated structure some specific features appear, which we plan to discuss elsewhere (see e.g. the remark before the eq.(16)).

For each point  $x \in M \setminus M_0$  we define the unique smooth contour  $C^x_{x_0}$ ,  $x_0 \in M_0$  connecting points x and  $x_0$ . The contours are parametrized as follows:

$$C_{x_0}^x: z_\mu = z_\mu(s, x); \ s \in [0, 1]; \ z_\mu(0, x) = x_{0\mu}; \ z_\mu(1, x) = x_\mu$$
 (2)

The map  $M \setminus M_0 \to M_0$  defined above is naturally extended to  $M \to M_0$ by setting  $C_{x_0}^{x_0}$  to be the unit contour:  $z_{\mu}(s, x_0) \equiv x_{0\mu}$ . The resulting map  $M \to M_0$  is assumed to be smooth. In the particular case when  $M_0$  consists of the only point  $x_0$  it means, that the manifold M should be contractible.

Let us choose two arbitrary points  $z_{\mu}(s, x)$  and  $z_{\mu}(s', x)$  on the given contour C in such a way that the point  $z_{\mu}(s', x)$  lies between points  $z_{\mu}(s, x)$ and  $z_{\mu}(1, x) = x_{\mu}$  (if s is natural parameter, it simply means that s < s'). We assume the following condition - for any s, s' there exists s'' such that

$$z_{\mu}(s,x) = z_{\mu}(s'', z(s',x)) \tag{3}$$

The geometrical meaning of (3) is simple: for any point z lying on some contour  $C_{x_0}^x$  its own contour  $C_{x_0}^z$  coincides with the corresponding part of the contour  $C_{x_0}^x$ . The eq.(3) does not mean, generally speaking, that contours  $C_{x_0}^{x_1}$  and  $C_{x_0}^{x_2}$  from different points  $x_1 \neq x_2$  have no common points except  $x_0$ . The condition that contours  $C_{x_0}^x$  should not selfintersect (the only condition discussed in [13]) is necessary but not sufficient to guarantee (3) and therefore

to derive (8) and (12) below. The defined set of contours forms an oriented tree graph without closed cycles according to (3).

Let us now start with the gauge potential  $A_{\mu}(x)$  taken in some arbitrary gauge and perform the gauge rotation

$$A'_{\mu}(x) = \Omega^{+}(x)A_{\mu}(x)\Omega(x) + \frac{i}{g}\Omega^{+}(x)\partial_{\mu}\Omega(x)$$
(4)

with

$$\Omega(x) = U(x, x_0) = Pexp(ig \int_{x_0}^x A_\mu(z) dz_\mu)$$

and integration goes along the contour  $C_{x_0}^x$ . The important point is the differentiation of the phase factors [10, 11] which is a well defined procedure for our choice of contours since the function  $z_{\mu}(s, x)$  is given:

$$\partial_{\mu}\Omega(x) = igA_{\mu}(x)\Omega(x) + ig\int_{0}^{1} ds \frac{\partial z_{\nu}(s,x)}{\partial s} \alpha_{\rho\mu}(z)U(x,z(s))F_{\rho\nu}(z(s))U(z(s),x_{0})$$
(5)

where

$$\alpha_{\rho\mu}(z) = \frac{\partial z_{\rho}(s, x)}{\partial x_{\mu}} \tag{6}$$

Substituting (5) into the (4) one gets:

$$A'_{\mu}(x) = -U(x_0, x) \int_{0}^{1} ds \frac{\partial z_{\nu}(s, x)}{\partial s} \alpha_{\rho\mu}(z) U(x, z(s)) F_{\rho\nu}(z(s)) U(z(s), x_0)$$
(7)

Taking into account the condition (3) and the gauge transformation property

$$U(x_0, x)U(x, z(s))F_{\rho\nu}(z(s))U(z(s), x_0) \to F'_{\rho\nu}(z(s))$$

we arrive to the final result (omitting primes in what follows)

$$A_{\mu}(x) = \int_{0}^{1} ds \frac{\partial z_{\nu}(s,x)}{\partial s} \frac{\partial z_{\rho}(s,x)}{\partial x_{\mu}} F_{\nu\rho}(z(s))$$
(8)

This formula was proposed in [13] and used without derivation in [9]. Note at the same time, that the condition (3), is crucial for proceeding from (7) to (8).

The eq.(8) leads to important local condition for vector-potential. To this end, note, that solving (3) with respect to s' we find:

$$s' = f(s, s'', x); f(s, s, x) = 1$$

Substituting s' = f(s, s'', x) into (3) and differentiating with respect to s one gets:

$$\frac{\partial z_{\mu}(s,x)}{\partial s} = \frac{\partial z_{\mu}(s'',z(s',x))}{\partial z_{\rho}(s',x)} \frac{\partial z_{\rho}(s',x)}{\partial s'} \frac{\partial f(s,s'',x)}{\partial s}$$
(9)

By putting s' to be equal to unity (9) reads:

$$\frac{\partial z_{\mu}(s,x)}{\partial s} = \frac{\partial z_{\mu}(s,x)}{\partial z_{\rho}(1,x)} \left(\frac{\partial z_{\rho}(s',x)}{\partial s'}\right)\Big|_{s'=1} \cdot g(s,x)$$
(10)

with  $g(s,x) = (\partial f(s,s'',x)/\partial s)_{s''=s}$ . We can now multiply both sides of (8) by  $k_{\mu}(x) = (\partial z_{\mu}(s,x)/\partial s)_{s=1}$  and get

$$A_{\mu}(x) \cdot k_{\mu}(x) =$$

$$= \int_{0}^{1} ds \frac{\partial z_{\nu}(s,x)}{\partial s} \frac{\partial z_{\rho}(s,x)}{\partial x_{\mu}} \left( \frac{\partial z_{\mu}(s,x)}{\partial s} \right) \Big|_{s=1} F_{\nu\rho}(z(s)) =$$

$$= \int_{0}^{1} ds \frac{\partial z_{\nu}(s,x)}{\partial s} \frac{\partial z_{\rho}(s,x)}{\partial s} (g(s,x))^{-1} F_{\nu\rho}(z(s)) = 0 \qquad (11)$$

where the second equality holds by virtue of (10) and the third due to antisymmetry of  $F_{\rho\nu}$ . The condition (11) can be easily understood taking into account, that phase factors along the contours  $C_{x_0}^x$  specifying the gauge are equal to unity:

$$U(x, x_0) = Pexp(ig \int_{x_0}^x A_\mu(z)dz_\mu) = 1$$

Since (11) holds for any x, one gets:

$$A_{\mu}(x) k_{\mu}(x) = 0 \tag{12}$$

Specific examples of the gauge condition discussed in the present article are known in the literature. The best studied is the radial or Fock-Schwinger gauge [4]. In this gauge the set of contours is defined by

$$z_{\mu}(s,x) = s \cdot x_{\mu} \tag{13}$$

and (12) reads:

$$x_{\mu}A_{\mu}(x) = 0 \tag{14}$$

The relation (8) becomes

$$A_{\mu}(x) = \int_{0}^{1} dssx_{\nu} F_{\nu\mu}(z(s))$$
(15)

Note that due to the topological restrictions stated above this gauge condition is already defined in some neighbourhood of the point  $x_0 = 0$  but might not be well defined globally. This was noticed in different respect also in [12].

Another example is the gauge condition introduced in [7]. It singles out not a point as the Fock-Schwinger gauge but a line. One can take this line to be defined by  $z_1 = z_2 = z_3 = 0$ . Then the contours C(x) are made of two straight paths:

$$z_i(s,x) = q(s) \cdot x_i, \ i = 1, 2, 3; \ z_4(s,x) = x_4 + p(s,x) \cdot n_4$$

where  $n_4$  is the unit vector in the forth direction, q(s) is the linear function, satisfying q(1) = 1,  $q(s_0) = 0$  and the function p(s, x) is such that  $p(s, x) \equiv 0$ for  $1 \ge s \ge s_0$  and  $p(s, x) \to \infty$  if  $s \to 0$ . The choice of  $s_0$  is arbitrary.

Piecewise nature of C leads to two different conditions depending on whether the point x lies on the singled out line or not. By imposing an additional requirement

$$\left(\partial p(s,x)/\partial s\right)\big|_{s=s_0} = 1$$

we obtain from (12):

$$A_4(x_i = 0) = 0; \ A_i(x) \cdot x_i = 0$$

This gauge was proved to be useful in the studies of heavy quarkonium dynamics [7, 16]. A natural generalization of the above conditions is a gauge, which could be called planar and it was actually used without derivation in [9]. It is constructed by choosing a plane  $z_1 = z_2 = 0$  and the contours C to be orthogonal to this plane. Then (12) reads:

$$A_1(x)x_1 + A_2(x)x_2 = 0$$

and additional gauge freedom for the potentials on the plane itself still remains to be fixed in an appropriate way.

It should be obvious from the derivation and considered examples that the gauge conditions of the type (12) are generally not enough to fix the gauge modulo global transformations (see discussion of this point for the radial gauge in [12]). But if the base manifold M satisfies some requirements (contractible if  $M_0$  consists of the only point, as it is, for example, in the case of radial gauge) the additional gauge freedom is absent. We plan to discuss these questions in detail in subsequent publication.

As an illustrative example let us consider the use of the generalized gauge condition for the nonabelian Stokes theorem. There are different proofs of this theorem in the literature [14, 15], but what we are going to present is perhaps the simplest one. It is close in spirit to the paper [14]. Namely, we define the gauge condition in such a way that potential  $A_{\mu}(x)$  on the contour is expressed as a function of field strength  $F_{\mu\nu}(u)$  defined on the (arbitrary) surface, bound by the contour. Then rewriting gauge-invariant Wilson loop in this gauge we obtain a relation, valid in the chosen gauge and as the last step put it into gauge-covariant form. It was done in [14] for the completely fixed axial gauge condition, which is a convenient choice in two dimensions (or for planar surfaces in higher dimensional case). Our procedure allows one to choose an arbitrary surface S bound by the simple contour  $C = \partial S$  and therefore the gauge condition we use entirely depends on the shape of S.

We parametrize the surface by  $w_{\mu}(s,t)$ ;  $s,t \in [0,1]$ . We choose an arbitrary point  $x_{0\mu}$  on the surface in such a way, that  $w_{\mu}(0,t) \equiv x_{0\mu}$ . If s = 1 then  $w_{\mu}(1,t)$  goes along the contour C and  $w_{\mu}(1,0) = w_{\mu}(1,1)$  according to  $\partial C = 0$ .

The following important remark is in order. It is usually assumed that S has the disk topology, in this simplest case we are free in our choice of  $M_0$ , which may consist of only one point, what we actually have used. For this topology it is always possible to define a set of contours obeying (3)

by continuous deformation of the planar disk with the radial contours (13). Indeed, the continuous, without cuts and gluings, deformation of the surface leaves (3) intact for the contours, defined on this surface. But if S is onehole surface of nontrivial genus, the proof should be modified, because in this case S cannot be retracted to a point. In other words, it is impossible to define the smooth set of contours, obeying (3) on a higher genus surface if  $M_0$  consists of only one point. Instead  $M_0$  must be taken as 1-cycle. So the validity of the nonabelian Stokes theorem in this case depends on the possibility to retract arbitrary surface with a hole to  $S^1$ . We plan to discuss the subtleties of nonabelian Stokes theorem for the surfaces of higher genus in subsequent publication.

According to (8) the gauge potential  $A_{\mu}(z)$  is related to  $F_{\mu\nu}(z)$  in the following way:

$$A_{\mu}(z(s,t)) = \int_{0}^{1} ds' \frac{\partial z_{\nu}(s',x(t))}{\partial s'} \frac{\partial z_{\rho}(s',x(t))}{\partial x_{\mu}(t)} F_{\nu\rho}(z(s',t))$$
(16)

Equation (16) is actually nothing else than the Stokes theorem in its infinitesimal form. It is well known that the generalization to finite contours is nontrivial in the nonabelian case, in particular the integral  $\int_S F_{\mu\nu} d\sigma_{\mu\nu} de$ pends on the surface even if the contour  $C = \partial S$  is closed. But this integral does not enter by itself in the nonabelian Stokes theorem. Instead the quantity which should be considered here is a *P*-ordered exponent, by definition it reads:

$$Pe^{ig \int_{C} A_{\mu}(x)dx^{\mu}} = \hat{1} + \sum_{n=1}^{\infty} (ig)^{n} \int ... \int dx^{(1)}_{\mu 1}...dx^{(n)}_{\mu n} A_{\mu_{n}}(x^{(n)})...A_{\mu_{1}}(x^{(1)}) \theta(x^{(1)} > x^{(2)} > ... > x^{(n)})$$
(17)

The  $\theta$ -function in (17) orders the points  $x^{(i)}$  along the contour C.

Substitution of (16) into (17) leads to the expression:

$$Pe^{ig \int_{C} A_{\mu}(x)dx_{\mu}} =$$

$$= 1 + \sum_{n=1}^{\infty} (ig)^{n} \int ... \int d\sigma_{\mu\nu}(w^{(1)}(s_{1}, t_{1}))...d\sigma_{\rho\phi}(w^{(n)}(s_{n}, t_{n}))$$

$$F_{\rho\phi}(w^{(n)}(s_{n}, t_{n}))..F_{\mu\nu}(w^{(1)}(s_{1}, t_{1})) \theta (t_{1} > t_{2} > ... > t_{n})$$
(18)

Note, that ordering procedure in (18) is the same as for the original Wilson loop – only the points along the contour C are ordered with respect one to another, i.e. ordered in parameters  $t_i$ , while the integrals over  $s_i$  are taken independently for each  $t_i$ .

To bring (18) to the gauge covariant form we introduce phase factors along the *s*-direction on the surface, which are equal to unity due to (12), i.e. we replace

$$F_{\mu\nu}(w(s,t)) \to G_{\mu\nu}(w(s,t)) = U(x_0; w(s,t))F_{\mu\nu}(w(s,t))U(w(s,t); x_0)$$

Under arbitrary gauge rotations the l.h.s. of (17) transforms as:

$$Pe^{ig \int\limits_{C_{x^*x^*}} A_{\mu}(x)dx_{\mu}} \to \Omega^+(x^*)Pe^{ig \int\limits_{C_{x^*x^*}} A_{\mu}(x)dx_{\mu}} \Omega(x^*)$$
(19)

where  $x^*$  is an arbitrary fixed point on the contour C (lower limit in all integrals in (17)). If the point  $x_0$  does not lie on the contour C, then the final gauge-covariant answer reads:

$$Pe^{ig \int_{C_{x^*x^*}} A_{\mu}dz_{\mu}} = U(x^*, x_0) \mathcal{P}e^{ig \int_{S} d\sigma_{\mu\nu}(z)G_{\mu\nu}(z)} U(x_0, x^*)$$
(20)

where the meaning of the ordering simbol  $\mathcal{P}$  is explained in (18). Under the gauge rotations both sides of (20) are transformed in the same way which finishes the proof. The more often used gauge-invariant form of (20) is

$$Tr P e^{ig \int_{C} A_{\mu} dz_{\mu}} = Tr \mathcal{P} e^{ig \int_{S} d\sigma_{\mu\nu}(z)G_{\mu\nu}(z)}$$
(21)

We stress again, that the exact meaning of the symbol  $\mathcal{P}$  is completely determined by the choice of the set of contours, defining the gauge which may be done as the most convenient one for a given application of the nonabelian Stokes theorem.

The advantages of the discussed gauge condition are not exhausted by the simple and transparent proof of Stokes theorem given above, in particular see [13] for some applications. In our opinion, the most interesting development has not yet been investigated. Namely, in the partition function one could introduce the integration over the set of contours defining the gauge (8) in addition to the integration over field strengths. This contour integration is

natural to associate with the sum over the surfaces, bounded by the Wilson contour C. This yields a choice of integration variables alternative to what is usually discussed in the literature [17]. We plan to develop this issue in future publications.

## Acknowledgments

The work was supported by the RFFI grants 96-02-19184a, 96-15-96740, and partially by RFFI-DFG grant 96-02-00088G. The authors are grateful to A.Levin, M.Olshanetsky, V.Novikov for valuable discussions and to G.Korchemsky for pointing authors attention to the Ref.[13].

## References

- L.D. Faddeev, "Gauge Fields: An Intro to Quantum Theory", Addison-Wesley, 1991.
   F. Yndurain, "The Theory of Quark and Gluon Interactions", 2-nd ed., Springer-Verlag, New-York, 1993.
- [2] L.V. Gribov, E.V. Levin, M.G. Ryskin, Phys.Rep., **100** (1983) 1.
- [3] V. Novikov, M. Shifman, A. Vainshtein, V. Zakharov, Nucl. Phys. B249 445 (1985).
- [4] V.A. Fock, Sov.Phys., 12 (1937) 404.
  J. Schwinger, Phys.Rev., 82 (1952) 684, see also "Particles, Sources and Fields", vols.1 and 2, Addison-Wesley, Reading, 1970 and 1973.
- [5] M. Shifman, Nucl.Phys.**B173** 13 (1980).
- [6] M.S. Dubovikov, A.V. Smilga, Nucl. Phys., **B** 185 (1981) 109.
- [7] I.Balitskii, Nucl.Phys.**B** 254 (1985) 166.
- [8] Yu.A. Simonov, hep-ph/9704301
   Yu.A. Simonov, hep-th/9712250
- [9] Yu.A. Simonov, hep-ph/9712248

- [10] S.Mandelstam, Phys.Rev. bf 175 (1968) 1580;
   Ann.Phys. 19, 1 (1962) 25.
- [11] Yu.M. Makeenko, A.A. Migdal, Nucl.Phys.B188 (1981) 269, for a review see A.A. Migdal, Phys.Rep.102 (1983) 199.
- [12] S. Leupold, H. Weigert, hep-th/9604015.
- [13] S.V. Ivanov, G.P. Korchemsky, Phys.Lett. B154 (1985) 197.
   S.V. Ivanov, G.P. Korchemsky, A.V. Radyushkin, Sov.J.Nucl.Phys. 44 (1986) 145.
- [14] M. Halpern, Phys.Rev.**D19** (1979) 517.
- [15] Y. Aref'eva, Theor.Math.Phys. 43 (1980) 353;
  N.Bralic, Phys.Rev. D22 (1980) 3090;
  Yu.A. Simonov, Sov.J.Nucl.Phys. 50 (1989) 134;
  M.Hirayama, S.Matsubara, hep-th/9712120.
- [16] N. Brambilla, A. Vairo, Phys.Lett. **B** 407 (1997) 167.
- [17] E. Akhmedov, M. Chernodub, M. Polykarpov, M. Zubkov, Phys.Rev. D 53 (1996) 20.
   A.M. Polyakov, Nucl.Phys., B486 (1997) 23.