MEMBRANES AND FIVEBRANES WITH LOWER SUPERSYMMETRY AND THEIR AdS SUPERGRAVITY DUALS

S. Ferrara, A. Kehagias, H. Partouche and A. Zaffaroni[†]

Theory Division, CERN, 1211 Geneva 23, Switzerland

Abstract

We consider superconformal field theories in three and six dimensions with eight supercharges which can be realized on the world-volume of M-theory branes sitting at orbifold singularities. We find that they should admit a $\mathcal{N} = 4$ and $\mathcal{N} = 2$ supergravity dual in AdS_4 and AdS_7 , respectively. We discuss the characteristics of the corresponding gauged supergravities.

[†] e-mail: Sergio.Ferrara, Alexandros.Kehagias, Herve.Partouche, Alberto.Zaffaroni@cern.ch CERN-TH-98-79 March 1998 In recent papers, the close connection between AdS_{p+2} and the dynamics on the worldvolume of p-branes was explored [1, 2, 3, 4, 5, 6]. The explicit proposal of Maldacena [7] that the large N limit of certain conformal field theories can be described in terms of supergravity paved the way for a novel approach to superconformal theories. In particular, it has been argued that the type IIB supergravity on $AdS_5 \times S^5$ is dual to D = 3, $\mathcal{N} = 4 U(N)$ SYM theory at large N. Similarly, it has been proposed that the (2,0), D = 6 superconformal field theory with N tensor multiplets is dual for large N to eleven-dimensional supergravity on $AdS_7 \times S^4$ with large radii. It has also been conjectured that these dualities can be elevated to field-theory/string-theory and field-theory/M-theory equivalence for finite N.

The proposed duality between large N field theories and anti-de Sitter supergravities has been tested by identifying massless excitations in the bulk with singletons composite operators on the anti-de Sitter boundary [8]. Extending further this relation, it has been argued that in a suitable limit, the generating functional for the boundary correlators of singleton composite field is reproduced by the anti-de Sitter supergravity action [9, 10]. The conjecture was further explored in [11, 12, 13, 14, 15, 16, 17, 18]. Moreover, models with lower world-volume supersymmetries started to be explored [19, 20, 21, 22]. We could ask if the field-theory/string-theory correspondence can be extended in such a way that for any given $\mathcal{N} = 0, 1, 2$ superconformal model in four dimensions there exist a supergravity theory in AdS_5 , and, similarly, if the field-theory/M-theory duality gives a supergravity theory on AdS_4 or AdS_7 for any superconformal theory in three and six dimensions, respectively. The conjecture, in the form in which it has been formulated up to now, requires the realization of the superconformal theory in terms of branes, in a setting which is compatible with the limits considered in [7]. Fortunately enough, quite a lot of superconformal models can be realized in this way. It is one of the purpose of this paper, to show superconformal models in D = 3, 6 which fulfil the previous requirements.

The maximally supersymmetric theories in three and six dimensions have been explored

recently [23, 24, 25, 26]. Here, we will consider D = 3 and D = 6 theories with lower supersymmetries, in particular, $\mathcal{N} = 4$ and $\mathcal{N} = 1$ supersymmetric theories realized on the world-volume of M2- and M5-branes. Such theories can be obtained by appropriate orbifolds of the transverse space. For the M2-brane, the transverse space is \mathbf{C}^4 and one may consider orbifolds of the form $\mathbf{C}^2/\Gamma \times \mathbf{C}^2/\Gamma'$, $\mathbf{C} \times \mathbf{C}^3/\Gamma$, $or \mathbf{C}^4/\Gamma$, corresponding to M-theory compactifications on $K3 \times K3$, CY_3 and CY_4 , respectively, giving rise to theories with $\mathcal{N} = 4$, $\mathcal{N} = 2$ and $\mathcal{N} = 1$. For the M5-brane, the transverse space is \mathbf{R}^5 and one may consider orbifolds $\mathbf{C}^2/\Gamma \times \mathbf{R}$ that lead to (1,0) theories on the world-volume of the M5brane. A similar procedure was followed in [19] and further elaborated in [21] for theories on the D3-brane of the type IIB string theory.

Let us start with D = 6. The M5-brane breaks half of the 32 supersymmetries leaving a (2,0) theory on the M5 world-volume. There exist five scalar fields representing fluctuations transverse to the brane and three additional degrees of freedom coming as collective coordinates associated to the three-form of the eleven-dimensional supergravity. All together, we have 8 degrees of freedom which fill the unique $\mathcal{N} = 2$, D = 6 tensor multiplet of the chiral (2,0), D = 6 supersymmetry. The eleven-dimensional supergravity solution for N M5-branes is

$$ds^{2} = f^{-1/3} \left(-dt^{2} + dx^{a} dx^{a} \right) + f^{2/3} dy^{a} dy^{a} ,$$

$$f = 1 + \frac{\pi N \ell_{p}^{3}}{r^{3}} , \quad r^{2} = y^{a} y^{a} , \qquad (1)$$

where ℓ_p is the eleven-dimensional Planck length, x^a, y^a (a = 1, ..., 5) are coordinates on the five-brane and transverse to it, respectively. There is a horizon at r = 0. The near horizon geometry is given by $AdS_7 \times S^4$, namely as the product of a seven-dimensional anti-de Sitter space with a four-sphere of radii $R_{S^4} = R_{AdS_7}/2 = \ell_p (\pi N)^{1/3}$. The M-theory five-brane interpolates between flat Minkowski space-time at $r \to \infty$ and $AdS_7 \times S^4$ at the horizon. The decoupling limit considered in [7] is $\ell_p \to 0$ and r/ℓ_p^3 finite, while supergravity is valid for N >> 1.

In order to describe a gauge theory in six dimensions, we should further break the (2,0) supersymmetry to the (1,0) one. The massless sector of the latter contains tensors, vectors and hypermultiplets. In particular, the (2,0) tensor multiplet gives rise to a tensor and a hyper of the (1,0) superalgebra. A (1,0) supersymmetry on the M5-brane can be obtained by taking an appropriate orbifold in the transverse space. Since we want to keep in the near-horizon geometry the AdS_7 structure, the orbifolding should be such that it acts on S^4 only. Now at $t, x^a = const.$, the transverse space is topologically \mathbb{R}^5 with metric

$$ds_{\perp} = \left(1 + \frac{\pi N \ell_p^3}{r^3}\right)^{2/3} dy^a dy^a \,. \tag{2}$$

We may act with a discrete group $\Gamma \subset SU(2)$ on \mathbb{R}^5 to form $\mathbb{R}^5/\Gamma = \mathbb{C}^2/\Gamma \times \mathbb{R}$ where the modding by Γ identifies points which are at equal distance from the origin. Thus, the transverse space to the M5-branes is the product of a flat sixth direction \mathbb{R} and the ALE space \mathbb{C}^2/Γ . We will consider here the cases $\Gamma = \mathbb{Z}_k$ $(k \ge 2)$ and $\Gamma = \mathcal{D}_k$ $(k \ge 4)$ where \mathcal{D}_k is the binary extension of the dihedral group. The former gives an A_{k-1} ALE space and the latter a D_k one. The generators of Γ act on \mathbb{C}^2 with coordinates z^1, z^2 as

$$\mathbf{Z}_{k} : \qquad z^{1} \to e^{\frac{2i\pi}{k}} z^{1}, \quad z^{2} \to e^{-\frac{2i\pi}{k}} z^{2}, \tag{3}$$

$$\mathcal{D}_{k} : \qquad z^{1} \to e^{i\pi/(k-2)} z^{1}, \quad z^{2} \to e^{-2i\pi/(k-2)} z^{2} \quad \text{and} \quad z^{1} \to z^{2}, \quad z^{2} \to -z_{1}.$$

Since these ALE spaces are of SU(2) holonomy, they break half of the original 32 supersymmetries while half more are broken by the M5-branes. Thus, the eight remaining supercharges lead to a (1,0) world-volume theory in six dimensions. The (2,0) tensor multiplet gives a (1,0) tensor and a hyper. The tensor contains an antiself-dual two-form and a scalar representing fluctuations of the five-brane along the sixth direction **R**, which is not affected by the orbifold. The hyper contains two complex scalars Z^1, Z^2 corresponding to the position of the M5-branes in the ALE space. In addition, there exist vector multiplets coming from the wrapping of M-theory membranes along the non-vanishing two-cycles of the ALE space. We will first consider the case of N M5-branes with world-volume (0, 1, 2, 3, 4, 5) located at N points in the sixth direction **R** and at the same point in the A_{k-1} ALE space. To analyze the spectrum, it is more convenient to consider our system in the context of type IIA theory. Let us recall that a metric of the form

$$ds_{TN}^{2} = V^{-1}(dX_{11} + \vec{\omega}) + Vd\vec{X}d\vec{X},$$

$$V = 1 + \sum_{i=1}^{k} \frac{R_{11}}{2|\vec{X} - \vec{X}_{i}|}, \quad \vec{\nabla} \times \vec{\omega} = -\vec{\nabla}V,$$
(4)

where \vec{X} is a three vector and X_{11} has periodicity $2\pi R_{11}$ describes a Taub-NUT space. In the limit $R_{11} \to \infty$ we can ignore the 1 in the expression of V and the Taub-NUT space approaches the ALE space $\mathbf{C}^2/\mathbf{Z}_k$. By identifying R_{11} with the type IIA string coupling $g^A = (R_{11}/\ell_p)^{3/2}$, we may replace the ALE space in M-theory with a Taub-NUT one. This can be interpreted as M-theory with k KK monopoles, which is type IIA with k D6-branes located at \vec{X}_i [27, 28, 29, 30]. Thus, the spectrum of N M5-branes with transverse space $\mathbf{R} \times \mathbf{C}^2/\mathbf{Z}_k$ in M-theory for large R_{11} can be analyzed in the type IIA side by considering k D6-branes and N NS5-branes at strong string coupling. One should expect that the spectrum can be evaluated at weak coupling and followed into strong coupling by rescaling appropriate parameters and moduli.

The spectrum of the above system at weak coupling can be analyzed by standard brane technology [31, 32, 33]. We consider a system of N NS5-branes with world-volume (0, 1, 2, 3, 4, 5) and k D6-branes stretched between them with world-volume (0, 1, 2, 3, 4, 5, 6). This configuration breaks the ten-dimensional Lorentz-invariance to $SO(1, 5) \times SO(3)$ and has a (1, 0), D=6 supersymmetry living on the world-volume of the NS5-branes. The latter has an Sp(1) R-symmetry which is geometrically realized as rotations in the threedimensional space transverse to the D6-brane (the SO(3) group above.) We choose the N NS5-branes to be located at $(x_{\alpha}^6, x^7, x^8, x^9)$, $\alpha = 1, ..., N$, and coincident with k D6branes at the same fixed coordinate (x^7, x^8, x^9) in the transverse space. If the NS5-branes were absent, we would have a system of k coincident D6-branes, which in the M-theory language would be interpreted as k KK-monopoles. In this case, there would be an U(k) gauge symmetry. The presence of the NS5-branes implies that the D6-branes are in fact divided in N-1 pieces of finite length along the sixth direction, each of them stretched between two adjacent NS5-branes. In addition, there are k semi-infinite D6-branes on the left and right of the first and last NS5-branes, respectively. This has the effect to give rise to the gauge group $U(k)^{N-1}$ on the NS5-branes world-volume as follows from the KK reduction of the D6 world-volume theory along x^6 . There exist charged hypermultiplets in the representation

$$(\mathbf{k}, \bar{\mathbf{k}}, 1, ..., 1) \oplus (1, \mathbf{k}, \bar{\mathbf{k}}, 1, ..., 1) \oplus \cdots \oplus (1, ..., 1, \mathbf{k}, \bar{\mathbf{k}}) \oplus$$

 $k(\bar{\mathbf{k}}, 1, ..., 1) \oplus k(1, ..., 1, \mathbf{k})$ (5)

of the gauge group, where the two last terms arise from the semi-infinite D6-branes. In addition to the massless spectrum comming from the D6-branes, each of the NS5-branes provides a (2,0) tensor multiplet. In addition to the self-dual two-forms, these N tensor multiplets of the (2,0) superalgebra contain five scalars ϕ_{α}^{I} (I = 6, ..., 10). Under the unbroken (1,0) supersymmetry, these multiplets decompose into tensor multiplets that contain the scalars ϕ_{α}^{6} , while $\phi_{\alpha}^{7,8,9,10}$ fill hypermultiplets. The KK reduction to (5 + 1) dimensions shows that ϕ_{α}^{6} appears as the effective coupling of each of the N - 1 U(k)factors

$$\frac{1}{g_{\alpha}^{2}}F_{\mu\nu}^{\alpha\,2} + (\partial\phi_{\alpha}^{6})^{2} + \sqrt{c}\phi_{\alpha}^{6}F_{\mu\nu}^{\alpha\,2}.$$
(6)

The bare coupling g_{α} can be absorbed into ϕ_{α}^{6} so that the effective coupling is indeed

$$\frac{1}{g_{\alpha}^{eff^2}} = \sqrt{c}\phi_{\alpha}^6\,,\tag{7}$$

where c is the anomaly coefficient [34]. $\phi_{\alpha}^{7,8,9}$ are SO(3)-triplets of FI terms for the diagonal U(1)'s of the gauge group factors. The hypermultiplets they belong to are exactly what is needed to cancel the U(1) anomalies and, as a consequence of a Green-Schwarz mechanism,

all the diagonal U(1) gauge bosons are massive. As a result, the (5 + 1) world-volume theory we obtain is the $\mathcal{N} = 1$, $SU(k)^{N-1}$ gauge theory with one tensor multiplet and khypermultiplets in the $\mathbf{k} \oplus \bar{\mathbf{k}}$ representation for each of the gauge group factors.

Six-dimensional gauge theories are restricted by the gauge anomaly, which is given by

$$Tr_{adj}F^4 - Tr_RF^4 = a \ trF^4 + \frac{c}{d^2}(trF^2)^2 \tag{8}$$

for a simple group, where Tr_{adj} , Tr_R and tr are traces in the adjoint, R and fundamental representations, respectively, and d is the dimension of the fundamenantal. When the theory is not coupled to gravity, it makes sence when $\alpha = 0$ and $c \geq 0$. For c = 0, the theory is anomaly free, while for c > 0 the anomaly can be canceled by a tensor multiplet [35, 34]. In our case, $c = 3k^2$ for each SU(k) factor and the anomaly can indeed be canceled by the tensor multiplet. In fact, as follows from eq.(7), the theory may have a non-trivial fixed point at $\phi_{\alpha}^6 = 0$ where it is strongly coupled. After absorbing the bare coupling into ϕ_{α}^6 , the terms in (6) are scale invariant, supporting the existence of this non-trivial fixed point. In fact, in the large N limit, the six-dimensional (1,0) $SU(k)^{N-1}$ SYM theory with one tensor and k hypermultiplets in $\mathbf{k} \oplus \bar{\mathbf{k}}$ for each SU(k) factor should be dual to eleven-dimensional supergravity on $AdS_7 \times S^4/\mathbf{Z}_k$ of radii $R_{S^4} = R_{AdS_7}/2 = \ell_p (\pi N)^{1/3}$ and therefore conformal invariant. Then, it is natural to conjure that for all N, the above six-dimensional (1,0) $SU(k)^{N-1}$ SYM theory is M-theory on $AdS_7 \times S^4/\mathbf{Z}_k$.

We will now consider the case where Γ is the binary dihedral group which corresponds to a D_k ALE space. It is known that M-theory on such spaces is type IIA with 2k D6branes sitting at an orientifold O6-plane with charge -4. Therefore, the system we consider is as in the A_{k-1} case, namely N NS5-branes with world-volume (0, 1, 2, 3, 4, 5) and D6-branes stretched between them, with world-volume (0, 1, 2, 3, 4, 5, 6), together with an orientifold six-plane. The system is located at a single point in the transverse space. Charge conservation imposes N to be even and the gauge group turns out to be [31, 32, 33] $Sp(k-4) \times (SO(2k) \times Sp(k-4))^{N/2-1}$. Each factor of the gauge group is coupled to a (1,0) tensor multiplet as in eq. (6) and the hypermultiplet content is

$$\frac{1}{2} \Big\{ (\mathbf{2k} - \mathbf{8}, \mathbf{2k}, 1, ..., 1) \oplus (1, \mathbf{2k}, \mathbf{2k} - \mathbf{8}, 1, ..., 1) \oplus \cdots \oplus (1, ..., 1, \mathbf{2k} - \mathbf{8}, \mathbf{2k}, 1) \oplus (1, ..., 1, \mathbf{2k}, \mathbf{2k} - \mathbf{8}) \oplus 2k (\mathbf{2k} - \mathbf{8}, 1, ..., 1) \oplus 2k (1, ..., 1, \mathbf{2k} - \mathbf{8}) \Big\}.$$
(9)

Effectively, each SO(2k) factor of the gauge group is coupled to 2k - 8 hypers in the vectorial representation 2k, while each of the Sp(k - 4) factor is coupled to 2k hypers in the fundamental 2k - 8. In this case the anomaly eq.(8) has a = 0 and $c = 12k^2$, $c = 12(k - 4)^2$ for each SO(2k) and Sp(k - 4) factor, respectively. Thus, the anomaly can be cancelled by the same mechanism as before, namely by the coupling of the scalars in the tensor multiplets with the gauge fields. The theory flows then to a non-trivial fixed point in the IR which can be described by the supergravity on $AdS_7 \times S^5/\mathcal{D}_k$.

Branes at orbifold singularities generally give rise to six-dimensional superconformal fixed points [36]. They can be realized using variations of the previous construction [32, 33]. The ones which admit a consistent description in M-theory (in some cases defined on R/Z_2 -see [20] for a related example) are likely to have a dual description as a supergravity in AdS_7 .

The $\mathcal{N} = 2$ gauged supergravity, coupled to matter, on AdS_7 was constructed in [38]. The gauge group is $SU(2) \times H$, where the SU(2) vector fields, gauging the R-symmetry of the superconformal theory on the boundary, live in the supergraviton multiplet, while the vectors in the adjoint of H, gauging the flavour symmetries of the boundary theory, are provided by additional vector multiplets. At the six-dimensional superconformal point discussed above (considering, for simplicity, the A_k singularity), the global symmetry is H = SU(k). The scalars in the supergravity theory parametrize the manifold,

$$R^+ \times \frac{O(3,n)}{O(3) \times O(n)},\tag{10}$$

with $n = \dim H$ and coordinates a real scalar ϕ_0 belonging to the graviton multiplet, and by the scalars ϕ_i^{Λ} in the adjoint of SU(2) and Lie-algebra valued in H. The $\mathcal{N} = 2$ supergravity has a potential which admits a stable [37] SU(2) invariant anti-de Sitter vacuum at $\phi_0 = \hat{\phi}_0, \phi_i^{\Lambda} = 0$ [38].

It would be interesting to understand better how the previous theory can be obtained as a KK reduction of the eleven-dimensional supergravity to seven dimensions [39], since the KK states would give information on the spectrum of conformal operator of the superconformal theory living at the boundary [8, 10, 13, 23, 24]. The relevant superconformal algebra is OSp(6, 2/2), which also classifies the particle states in AdS_7 . Here we simply note that singleton representations of the algebra, corresponding to degrees of freedom living on the boundary of AdS_7 , would correspond to the *fields* of the superconformal theory, while the other representations correspond to composite operators [8].

Similar constructions can be repeated for the case of superconformal theories in three dimensions, by orbifolding the $\mathcal{N} = 8$ example discussed in [7]. The theory of N coincident membranes in M theory is supposed to be dual for large N to the eleven-dimensional supergravity on $AdS_4 \times S^7$. We can consider orbifolds of M-theory in which the AdS_4 part is not affected by the projection. This corresponds to projecting by a suitable discrete group the transverse space of the membranes R^8 .

 $\mathcal{N} = 4$ superconformal theories in three dimensions have a OSp(4/4) symmetry. Nontrivial superconformal theories can be obtained in the IR limit of three dimensional Yang-Mills theories [40, 41]. We will now determine a large class of three dimensional Yang-Mills theories, which have a brane realization and whose IR limit should admit a dual description in terms of a $\mathcal{N} = 4$ supergravity in AdS_4 .

The original $\mathcal{N} = 8$ example in [7] can be also understood in the following way. Let us start with N D2-branes in type IIA which give rise to the maximal supersymmetric Yang-Mills theory in three dimensions. The theory is not conformal, having a dimensionful gauge coupling but flows in the IR to a superconformal fixed point which is the same as the one discussed in [7]. The gauge coupling for the D2-branes is determined by the IIA string coupling in such a way that it becomes very large, and the theory therefore flows to the IR superconformal fixed point, exactly when the M-theory description takes over and the system is better described with N M-theory membranes. Note also that the SO(7)global symmetry of the D2-branes theory, which rotates the transverse direction in type IIA, is promoted at the superconformal point to an SO(8) symmetry [42], which rotates the directions transverse to the membranes, and which is the appropriate R-symmetry of the $\mathcal{N} = 8$ superconformal algebra.

Let us now consider the $\mathcal{N} = 4$ case. The simplest example is obtained by putting N membranes near an orbifold singularity of the form $R^4/Z_k \times R^4/Z_n$. The spectrum for this theory was computed in [28]. In a type IIA description, we have a theory of D2-branes sitting at an orbifold singularity in the presence of n D6-branes, whose world-volume is, in part, parallel to the D2 world-volume and, for the remaining part, wrapped around a four-dimensional singular space R^4/Z_k . The corresponding Yang-Mills theory realized on the world-volume of the D2-branes can be easily derived to be [43, 28] a $SU(N)^k$ theory with hypermultiplets in the representations:

$$(\mathbf{N}, \mathbf{N}, 1, ..., 1) \oplus (1, \mathbf{N}, \mathbf{N}, ..., 1) \oplus \cdots \oplus (\mathbf{N}, 1, ..., 1, \mathbf{N}) + n(\mathbf{N}, 1, ..., 1).$$
 (11)

In the IR, these theories flow to a superconformal fixed point which should have a supergravity dual description as eleven-dimensional supergravity on $AdS_4 \times S^7/(Z_k \times Z_n)$.

Since there are two ways to get a type IIA description starting from M theory on $R^4/Z_k \times R^4/Z_n$, which correspond to *compactify*, in one case, on R^4/Z_k and, in the second case, on R^4/Z_n , we could expect to obtain two Yang-Mills candidates (corresponding to the exchange of k and n) for the same IR fixed point (and the same supergravity description). However, and this was the original motivation of [28], these two theories are actually three dimensional mirror pairs in the sense of [41], and therefore flow in the IR to the same superconformal fixed point. The Yang-Mills theories described above indeed contain and generalize the example in [41]. The differences between the theory in (11) and its mirror,

with k and n interchanged, in fact disappear in the eleven-dimensional description which is the relevant one for capturing the structure of the superconformal fixed point.

The R-symmetry $SU(2) \times SU(2)$ of the superconformal point is already manifest in the Yang-Mills theory at finite coupling. In the $\mathcal{N} = 4$ case, it is the global flavour symmetry which is enhanced at the IR fixed point [41]. The manifest SU(n) flavour symmetry of the theories discussed above is enhanced in the IR to $SU(n) \times SU(k)$ (a symmetry which is manifest in the M-theory description).

The case in which the orbifold projection is performed with a dihedral discrete group is analogous. The Yang-Mills theories that can be obtained in this way can be found in [28].

These superconformal fixed points should have a description as a $\mathcal{N} = 4$ supergravity in AdS_4 , whose supergraviton multiplet contains the $SU(2) \times SU(2)$ vector fields gauging the R-symmetry, coupled to additional vector multiplets which gauge the flavour symmetries. Such gauged supergravity was constructed in [44]. If the gauge group is $SU(2) \times SU(2) \times H$, the scalars in the theory parametrize the manifold,

$$\frac{SU(1,1)}{U(1)} \times \frac{SO(6,n)}{SO(6) \times SO(n)},$$
(12)

with $n = \dim H$. The coset SU(1,1)/U(1) is parametrized by a $SU(2) \times SU(2)$ singlet complex scalar Z in the supergraviton multiplet. Since we are looking for an anti-de Sitter $SU(2) \times SU(2)$ invariant vacuum, all the remaining scalars, in the adjoint of $SU(2) \times SU(2)$ and Lie-algebra valued in H, can be set to zero. The potential for Z admits a stable AdS_4 vacuum when the coupling constants for the two SU(2) are equal [44], giving the symmetric gauged SO(4) supergravity. From the point of view of the Yang-Mills theory which flows to this fixed point, the symmetry between the two SU(2) is nothing else than the mirror symmetry of [41].

Also in these three-dimensional theories, it would be interesting to have a description in terms of an explicit KK reduction of the eleven-dimensional supergravity to four dimensions, since the KK states would give information on the spectrum of conformal operator of the superconformal theory living at the boundary [10]. The relevant superconformal algebra is OSp(4/4), which also classifies the particle states in AdS_4 . The irreducible representations of OSp(4/4) are discussed in [45]. The singleton representations of the algebra, in this case, correspond to a multiplet with a complex scalar and a fermion, transforming as (1/2, 0) + (0, 1/2) under $SU(2) \times SU(2)$, in the superconformal boundary theory, while the other representations correspond to composite operators [8].

In principle, one could construct, by changing the orbifold projection, three-dimensional theories with lower $\mathcal{N} = 2, 1$ supersymmetries, based on the superconformal algebras OSp(2/4) and OSp(1/4). Their supergravity duals would correspond to 4d $\mathcal{N} = 2, 1$ AdS_4 supergravities with some additional matter multiplets. The supermultiplets in AdS_4 would be composite operators of the singleton representations on the 3d boundary [8].

It would be interesting, both in three and six dimensions, to extend the results of the present paper to cases with lower supersymmetry and eventually to cases with $\mathcal{N} = 0$. This can be achieved by considering different types of orbifold projections. In the dual supergravity side, it is known that the scalar potential for the gauged supergravities has other critical points where supersymmetry is partially or completely broken. We may ask if we can identify a superconformal boundary theory with lower or zero supersymmetry, whose generating functional for composite operators is reproduced by the tree-level supergravity expanded around these other critical points. This would provide an explicit flow between superconformal theories with different supersymmetries.

Acknowledgements

This work is supported in part by the EEC under TMR contract ERBFMRX-CT96-0090. S.F. is supported in part by the DOE under grant DE-FG03-91ER40662, Task C, and by ECC Science Program SCI^{*} -CI92-0789 (INFN-Frascati).

References

- C. W. Gibbons and P. K. Townsend, Phys. Rev. Lett. 71 (1993) 3754; M. P. Blencowe and M. J. Duff, Phys. Lett. B203 (1988) 229; Nucl. Phys B310 (1988), 389; M. J. Duff, Class. Quantum Grav. 5 (1988) 189; E. Bergshoeff, M. J. Duff, C. N. Pope and E. Sezgin, Phys. Lett. B199 (1988) 69; H. Nicolai, E. Sezgin and Y. Tanii, Nucl. Phys B305 (1988) 483.
- [2] K. Sfetsos and K. Skenderis, "Microscopic derivation of the Bekenstein-Hawking entropy formula for non-extremal black holes", hep-th/9711138.
- [3] H. J. Boonstra, B. Peeters and K. Skenderis, "Branes and Anti-de Sitter Space-Times", hep-th/9801076.
- [4] P. Claus, R. Kallosh and A. Van Proeyen, "M5-brane and superconformal (0,2) tensor multiplet in 6 dimensions", hep-th/9711161; R. Kallosh, J. Kumar and A. Rajaraman, "Special Conformal Symmetry of World-volume Actions", hep-th/9712073; P. Claus, R. Kallosh, J. Kumar, P. Townsend and A. Van Proeyen, "Conformal theory of M2, D3, M5 and D1+D5 branes", hep-th/9801206.
- [5] M. Gunaydin and D. Minic, "Singletons, Doubletons and M-theory", hep-th/9802047.
- [6] L. Castellani, A. Ceresole, R. D'Auria, S. Ferrara, P. Fré and M. Trigiante, "G/H M-branes and AdS_{p+2} Geometries, hep-th/9803039.
- [7] J. Maldacena, "The Large N Limit of Superconformal Field Theories and Supergravity", hep-th/9711200.
- [8] S. Ferrara and C. Fronsdal, "Conformal Maxwell Theory as a Singleton Field Theory on AdS₅, IIB Three-Branes and Duality, hep-th/971223; "Gauge Fields as Composite Boundary Excitations", hep-th/9802126.

- S. S. Gubser, I. R. Klebanov and A. M. Polyakov, "Gauge Theory Correlators from Non-Critical String Theory", hep-th/9802109.
- [10] E. Witten, "Anti-de Sitter Space And Holography", hep-th/9802150.
- [11] G. T. Horowitz and H. Ooguri, "Spectrum of Large N Gauge Theory from Supergravity", hep-th/9802116.
- [12] N. Itzhaki, J.M. Maldacena, J. Sonnenschein and S. Yankielowicz, "Supergravity and the Large N Limit of Theories with Sixteen Supercharges", hep-th/9802042.
- [13] S. Ferrara, C. Fronsdal and A. Zaffaroni, "On N=8 Supergravity on AdS₅ and N=4 Superconformal Yang-Mills Theory", hep-th/9802203; S. Ferrara and A. Zaffaroni, "N=1, N=2 4-D Superconformal Field Theories and Supergravity in AdS₅", hep-th/9803060.
- [14] S. J. Rey and J. Yee, "Macroscopic Strings as Heavy Quarks of Large N Gauge Theory and Anti-de Sitter Supergravity", hep-th/9803001.
- [15] J. M. Maldacena, "Wilson loops in large N field theories", hep-th/9803002.
- [16] I. Ya. Aref'eva and I. V. Volovich, "Field Theories in Anti-de Sitter Space and Singletons", hep-th/9803028.
- [17] E. Bergshoeff and K. Behrndt, "D-Instantons and Asymptotic Geometries, hepth/9803090.
- [18] M.J. Duff, H. Lu, C.N. Pope "Gauge Theory \rightarrow IIB \rightarrow IIA \rightarrow M Duality", hep-th/9803061.
- [19] S. Kachru and E. Silverstein, "4d Conformal Field Theories and Strings on Orbifolds", hep-th/9802183.
- [20] M. Berkooz, "A Supergravity Dual of a (1,0) Field Theory in Six Dimensions, hepth/9802195.

- [21] A. Lawrence, N. Nekrasov and C. Vafa, "On Conformal Field Theories in Four-Dimensions", hep-th/9803015.
- [22] M. Bershadsky, Z. Kakushadze and C. Vafa, "String Expansion as Large N Expansion of Gauge Theories", hep-th/9803076
- [23] O. Aharony, Y. Oz and Z. Yin, "M-Theory on AdS_p × S^{11-p} and Superconformal Field Theories, hep-th/9803051.
- [24] R.G. Leigh and M. Rozali, "The Large N Limit of the (2,0) Superconformal Field Theory", hep-th/9803068.
- [25] S. Minwalla, "Particles on $AdS_{4/7}$ and Primary Operators on $M_{2/5}$ Brane Worldvolumes", hep-th/9803053.
- [26] E. Halyo, "Supergravity on $AdS_{4/7} \times S^{7/4}$ and M Branes", hep-th/9803077.
- [27] N. Seiberg, Phys.Lett. B384 (1996) 81, hep-th/9606017.
- [28] M. Porrati and A. Zaffaroni, Nucl. Phys. B490 (1997) 107.
- [29] E. Witten, Nucl.Phys. B500 (1997) 3, hep-th/9703166.
- [30] A. Sen, "A Note on Enhanced Gauge Symmetries in M- and String Theory", hepth/9707123.
- [31] I. Brunner and A. Karch, Phys. Lett. B409 (1997) 109, hep-th/9705022.
- [32] A. Hanany and A. Zaffaroni, "Branes and Six Dimensional Supersymmetric Theories, hep-th/9712145.
- [33] I. Brunner and A. Karch, "Branes at Orbifolds versus Hanany Witten in Six Dimensions", hep-th/9712143.
- [34] N. Seiberg, Phys. Lett. B390 (1997) 169, hep-th/9609161.

- [35] A. Sagnotti, Phys. Lett. B294 (1992) 196, hep-th/9210127.
- [36] K. Intriligator, Nucl.Phys. B496 (1997) 17, hep-th/9702038; J. D. Blum and K. Intriligator, Nucl.Phys. B506 (1997) 22, hep-th/9705033; Nucl.Phys. B506 (1997) 199, hep-th/9705044.
- [37] P. Breitenlohner and D.Z. Freedman, Ann. Phys. 144 (1982) 197.
- [38] L. Mencizescu, P. K. Townsend and P. van Nieuwenhuizen, Phys. Lett. 143B (1984)584; E. Bergshoeff, I. G. Koh and E. Sezgin, Phys. Rev. D32 (1985) 1357.
- [39] P. van Nieuweinhuizen, Class. Quantum Grav. 2 (1985) 1.
- [40] N. Seiberg and E. Witten, "Gauge Dynamics And Compactification To Three Dimensions", hep-th/9607163.
- [41] K. Intriligator and N. Seiberg, Phys. Lett. B387 (1996) 513.
- [42] S. Sethi and L. Susskind, Phys. Lett. B400 (1997) 265; T. Banks and N. Seiberg, Nucl. Phys. B497 (1997) 41, hep-th/9702187; N. Seiberg, "Notes on Theories with 16 Supercharges", hep-th/9705117.
- [43] M.R. Douglas and G. Moore, "D-Branes, Quivers, and ALE Instantons", hepth/9603167; M.R. Douglas, B.R. Greene and D.R. Morrison, "Orbifold resolution by D-Branes", Nucl. Phys. B506 (1997) 84.
- [44] D. Z. Freedman and J. H. Schwarz, Nucl. Phys. B137 (1978) 333; E. Bergshoeff, I. G. Koh and E. Sezgin, Phys. Lett. 155B (1985) 71; M. de Roo and P. Wagemans, Nucl. Phys. B262 (1985) 644.
- [45] M. Flato and C. Fronsdal, J. Math. Phys. 22 (1981) 1100; Phys. Lett. 172B (1986)
 412; Lett. Math. Phys. 2 (1978) 421; Phys. Lett. 97B (1980) 236; A. Angelopoulos, M. Flato, C. Fronsdal and D. Sternheimer, Phys. Rev.D23 (1981) 1278.