M apping H awking into Unruh Therm al Properties S.Deser and Orit Levin

Physics Department, Brandeis University, Waltham, Massachusetts 02454-9110, USA

A bstract: By globally embedding curved spaces into higher dimensional at ones, we show that Hawking therm alproperties map into their Unruh equivalents: The relevant curved space detectors become R indler ones, whose temperature and entropy reproduce the originals. Speci c illustrations include Schwarzschild, Schwarzschild-(anti)deSitter, Reissner-N ordstrom and BTZ spaces.

1.Introduction

It is well-understood that, for both Hawking and Unruh e ects, tem perature em erges from information loss associated with real and accelerated-observer horizons, respectively. Given that any D-dimensional geometry has a higher-dimensional global embedding M inkowskian (possibly with more than one timelike coordinate) spacetime (GEMS) [1], it is natural to ask whether these m appings can unify the two e ects, by associating the relevant detectors of the curved spaces and their horizons with (constant acceleration) R indler detectors and their horizons. C on m ation of these ideas was recently given in an analysis of deSitter (dS) [2] and anti-deSitter (AdS) [3] geom etries and their GEMS. There, constantly accelerated observers were mapped into similar ones in the GEM S.The resulting Unruh tem peratures associated with these R indlerm otions agreed with those in the originaldS and A dS spaces. A ctually, A dS has no realhorizon, but tem perature is wellde ned for su ciently large accelerations and the two methods agree both as to the range where T exists and to its m agnitude.] In the present paper¹, we will show that the GEM S approach indeed provides a uni ed derivation of tem perature for a wide variety of curved spaces, including general rotating BTZ, Schwarzschild together with its dS and AdS extensions, and Reissner{Nordstrom. In each case the usual black-hole (BH) detectors are mapped into R indler observers with the correct temperature as determined from their (constant) accelerations. Conversely, we will also connect surface gravity and U nruh tem peratures, for both R indler observers in at space and various accelerated observers in de-Sitter (dS) and anti de-Sitter (AdS) spaces, thereby establishing the equivalence principle between constant acceleration and \true" gravity e ects. We will also consider the associated extensive quantity, the entropy, and again show the mapping correctly matches the area of the GEMS R indler motion and \true" horizons, thereby con ming the equivalence for entropy as well.

We will rst review how temperature measured by an accelerated detector in dS/AdS geometries, say in D = 4, is just its U nruh temperature (i.e., R indler acceleration divided by 2) in the D = 5 GEM S, by relating the corresponding 4- and 5-accelerations. In this connection we will also explicitly relate surface gravity to the associated temperatures. Next we shall treat rotating and non-rotating D = 3 BTZ spaces [5], [6]. Since BTZ is obtained from AdS through geodesic identication, we will show that we can use the treatment of U nruh observers in AdS to calculate the BH temperature here as well, in agreement with earlier results. Our nal applications will be to Schwarzschild, Schwarzschild-dS, Schwarzschild-AdS and Reissner-Nordstrom spacetimes, where the same connections are made, this time the required GEMS extensions having D 6. M ore generally, it will be seen that for any geometry admitting a group of constantly accelerated observers

deser@ binah.cc.brandeis.edu, olevin@ binah.cc.brandeis.edu

¹A brief sum m ary of part of this work was given in [4].

which encounter a horizon as they follow a \bifurcate" tim elike K illing vector eld, the tem perature measured by each observer is simply 2 T = a_{c} when a_{c} is their acceleration as mapped into the GEM S. Finally, we will establish equivalence of entropies using the Unruh de nition in terms of the "transverse" R indler area [7], together with the fact that horizons m ap into horizons.

2. Surface gravity-Unruh e ect connection in dS/AdS

We begin with a brief summary of the GEMS approach to temperature given in [3], for dS/AdS spaces of cosm ological constant $3R^{2}$; these are hyperboloids in the D = 5 GEMS $ds^{2} = \frac{1}{AB} (dz^{A})^{2} (dz^{B})^{2}$,

$$_{AB} (z^{A})^{2} (z^{B})^{2} = R^{2}$$
: (1)

Here A; B = 0:::4; $_{AB}$ = diag(1; 1; 1; 1; 1); throughout, upper/lower signs refer to dS/AdS respectively. We speci cally consider $z^{1} = z^{2} = 0$ and $z^{4} = Z = const$ trajectories, obeying $(z^{1})^{2}$ $(z^{0})^{2} = R^{2}$ Z^{2} a_{5}^{2} . Now the Unruh e ect states that at space detectors with constant acceleration a along the x direction, whose motions are thus on x^{2} $t^{2} = a^{2}$, measure temperature 2 T = a. Since our embedding space detectors follow precisely such trajectories i.e., have a R indler-like motion with constant acceleration a_{5} , they measure

2 T =
$$a_5 = (R^2 Z^2)^{1=2} (R^2 + a^2)^{1=2}$$
: (2)

The last equality expresses the tem perature in term of the D = 4 quantities, using $a_5^2 = R^2 + a^2$.

The relation between H aw king-B ekenstein horizon surface gravity $k_{_{\rm H}}$ and the B H tem perature (originally found for Schwarzschild B H) [8], [9]

$$T = \frac{1}{2} \frac{k_{H}}{p_{00}};$$
(3)

where x^0 is the time-like K illing vector of a detector in its rest frame, holds also for Schwarzschild-A dS and BTZ spacetimes [10]. For these latter two, the local temperature vanishes at in nity, and no Hawking particles are present far from the BH: created at the horizon, they do not have enough energy to escape to in nity (where the \e ective potential" becomes in nite). The connection (3) between temperature and surface gravity also holds [11] for R indler motions, reinforcing the connection between the Hawking and Unruh e ects as being based on the existence of horizons, whether \real" or just seen by accelerated observers. In both cases, inserting the horizon surface gravity in (3) will give the temperature. To calculate T, it is convenient to use the detector rest fram e.² The simplest example is the at space R indler observer, best described by R indler coordinates (;)

$$ds^2 = L^2 \exp(2) (d^2 d^2) (dy^2 + dz^2)$$
: (4)

A = const detector (following the time-like Killing vector = @) has a constant acceleration a = L¹ exp(). This group of accelerated observers sees an event horizon at = 1. Since is perpendicular to the horizon (and therefore null) we can calculate the surface gravity using its de nition [11]

$$k_{H}^{2} = \frac{1}{2} (r) (r)$$
 (5)

 $^{^{2}}$ The vacuum states in these timelike K illing coordinate systems are Schwarzschild-like. Therefore, determining the temperature by the (lowest order) transition rate obtained from the W ightman function for these vacua gives zero temperature, while the same calculation for H awking-H artle and K ruskal-like vacua gives the temperature (3).

where the right side is to be evaluated at the horizon. For us

$$k_{H}^{2} = k^{2} (= 1) = 1;$$
 (6)

Inserting k_{μ} in (3) gives the desired result

$$2 T = L^{-1} \exp() = a;$$
 (7)

Let us show that use of surface gravity to calculate temperature also works for dS/AdS. Consider rst dS with its real horizon, expressed in the static coordinates (t;r; ;) related to the z^{A} according to³, p _____ p ____

$$z^{0} = \stackrel{p}{R^{2}} \stackrel{r^{2}}{r^{2}} \sinh(t=R) \quad z^{1} = \stackrel{p}{R^{2}} \stackrel{r^{2}}{r^{2}} \cosh(t=R)$$

$$z^{2} = r \sin \cos z^{3} = r \sin \sin z^{4} = r \cos z^{3} = r \sin z^{4} = r \cos z^{4}$$

The metric

$$ds^{2} = 1 \frac{r^{2}}{R^{2}} dt^{2} 1 \frac{r^{2}}{R^{2}} dr^{2} r^{2} (d^{2} + \sin^{2} d^{2})$$
(9)

has an intrinsic horizon at r = R. It is seen by \static" detectors (r; ; const), or equivalently (choosing = 0, as is allowed by symmetry) $z^1 = z^2 = 0$ and $z^4 = r = \frac{Z}{P} = \frac{Z}{R^2 - r^2}$. They follow the time-like K illing vector e_t and have constant acceleration $a = r = (R - \frac{R^2}{R^2 - r^2})$. Hence by (5), we have

$$k_{_{\rm H}} = 1 = R$$
 (10)

and the tem perature m easured by these detectors agrees with the known results of [2],

$$T = \frac{1}{2} \frac{p}{R^2 r^2} = \frac{1}{2} \frac{1}{R^2 + a^2}$$
(11)

In AdS,

$$ds^{2} = 1 + \frac{r^{2}}{R^{2}} dt^{2} \qquad 1 + \frac{r^{2}}{R^{2}} dr^{2} \qquad r^{2} (d^{2} + \sin^{2} d^{2}); \qquad (12)$$

there is no intrinsic horizon. So although r = const detectors have constant acceleration $a = r = (R^{-1}R^{-2} + r^2) < R^{-1}$, they will not measure any temperature. The intrinsic horizon of dS causes even inertial detectors to measure temperature, while in AdS the absence of a real horizon causes su ciently slow ly (a < R^{-1}) accelerated detectors not to measure one. There is no contradiction with the Unruh picture: as we will see, the GEM S acceleration a_5^2 becomes negative for them $^{4;5}$. Indeed the \GEM S temperature" was obtained only for $(z^4)^2 = const^2 > R^2$ (a > R^{-1}) trajectories there [3]. Using the formula for time-like trajectories with a < R^{-1} (not $(z^4)^2 > R^2$ trajectories, but for example the $z^1 = const$, or the r = const case we discussed above) would lead to imaginary T : the detector will not measure any temperature because it sees no event horizon, hence no loss

 $^{^{3}}$ A lthough this coordinate transform ation covers only part of the space, it is easy to extend it continuously to the whole dS, resulting in a global embedding.

⁴ If we take the in aginary point $r_{H} = iR$ to de ne the AdS \horizon" and calculate the surface gravity at that point, (3) will give, as expected, an in aginary temperature 2 T = $i(R^{2} + r^{2})^{1=2} = \frac{P}{R^{2} + a^{2}}$, but (by the last equality) the correct temperature form ula for AdS [3].

⁵ It is also possible to get the AdS result from that of Schwarzschild-AdS [10], not by taking the limit m ! 0 but only by setting m = 0 initially. This is exactly like the impossibility of reaching at space by taking the m ! 0 limit of the H awking tem perature form ula for Schwarzschild.

of inform ation. To calculate the tem perature using (3) when $a > R^{-1}$ it is convenient to use a new coordinate system (the one in [3] is not suitable here since its x^0 is not the time-like K illing vector followed by the observers). Instead we introduce an \accelerated" coordinate system obtained by the GEM S coordinates de ned from the D = 4 covering of A dS

$$ds^{2} = \frac{R^{2} + 2}{R^{2}} d^{2} \frac{R^{2}}{R^{2} + 2} d^{2} \frac{R^{2}}{R^{2} + 2} d^{2} \frac{R^{2}}{R^{2} + 2} d^{2} (d^{2} + \sinh^{2} d^{2})$$
(13)

as follow s:

$$z^{0} = \frac{q}{R^{2} + 2} \sinh(=R);; z^{1} = \frac{q}{R^{2} + 2} \cosh(=R)$$
(14)

$$z^2 = \sinh \cos z^3 = \sinh \sin z^4 = \cosh z^4$$

Here 1 < ; < 1 ; < < ; while this coordinate patch only covers the region > R, it can be extended to the entire space. Since we are interested in $z^1 = z^2 = 0$ $z^4 = const$ trajectories, is set to zero, and to a constant Z; their accelerations are $a^2 = Z^2 R^2 (Z^2 R^2)^{-1} > R^{-2}$. For AdS, the horizon appears in this \accelerated" fram e exactly as it did upon transform ing from

M inkow ski to R indler coordinates in at space. These trajectories follow the tim e-like K illing vector eld @ which is null at the event horizon = R, so (5) gives

$$k_{\mu} = R^{-1}$$
 (15)

The corresponding tem perature, from (3), is

Ν

2 T =
$$(R^2 + Z^2)^{1=2} = (R^2 + a^2)^{1=2}$$
 (16)

which is exactly the result obtained using the kinematical behavior of these trajectories in the $G \in M$ S, as well as by calculating the transition rate in the non-accelerated coordinate system.

3.BTZ spaces

In the previous section, we demonstrated the feasibility of using surface gravity (or equivalently Hawking{Bekenstein temperature) to calculate the temperature measured in dS/AdS, in agreement with that obtained by purely kinematicalUnruh considerations. This immediately raises the converse question: calculate Hawking temperature entirely from GEMS kinematics when \real", mass-related, horizons are present. The simplest candidate for this would seem to be the BTZ black hole solution, due to its relation to AdS; we now use ourmethod to calculate BTZ temperature, at least for some observers, and compare with previous calculations using surface gravity [5],[12].

The general rotating BTZ black hole is described by the 3-m etric

$$ds^{2} = N^{2}dt^{2} N^{2}dr^{2} r^{2} (d + N dt)^{2}$$

$$^{2} (r^{2} r_{+}^{2})(r^{2} r^{2}) = (r^{2}R^{2}) ; N r_{+}r = (r^{2}R):$$
(17)

It arises from AdS upon making the geodesic identication = +2. The coordinate transformations to the (2+2) AdS GEMS $ds^2 = (dz^0)^2 (dz^1)^2 (dz^2)^2 + (dz^3)^2$ are for, r r₊ (the extension to r < r₊ is given in [6])

$$z^{\circ} = R \frac{r^{2}}{r_{+}^{2}} \frac{r_{+}^{2}}{r_{+}^{2}} \sinh \frac{r_{+}}{R^{2}} t \frac{r}{R} z^{1} = R \frac{r^{2}}{r_{+}^{2}} \frac{r_{+}^{2}}{r_{+}^{2}} \cosh \frac{r_{+}}{R^{2}} t \frac{r}{R}$$
(18)

$$z^{2} = R = \frac{r^{2}}{r_{+}^{2}} \frac{r^{2}}{r^{2}} \sinh \frac{r_{+}}{R} = \frac{r}{R^{2}}t = R = \frac{r^{2}}{r_{+}^{2}} \frac{r^{2}}{r^{2}} \cosh \frac{r_{+}}{R} = \frac{r}{R^{2}}t$$

where the constants $(r_+; r_-)$ are related to the m ass and angularm on entum. This A dS G EM S can serve as the BTZ embedding space for our purpose. In spite of the fact that there is no longer a one to one mapping between it and the BTZ space due to the identication, following a detector motion with certain initial condition such as (t = 0) = 0 still gives a unique trajectory in the embedding space which is the basic requirement of our approach based on the observer's kinematical behavior in the GEM S: If the detector trajectory maps (without ambiguity) into an Unruh one in the GEM S, then we can use it for temperature calculation.

Consider rst non-rotating BTZ (r = 0) and focus on \static" detectors (;r = const). These detectors have constant 3-acceleration a = rR⁻¹ (r² r_{+}^2)¹⁼², and are describe by a (xed) point in the (z²;z³) plane (for example = 0 gives z² = 0 z³ = const), and constant accelerated motion in (z⁰;z¹) with a₄ = r₊ R⁻¹ (r² r_{+}^2)¹⁼². So in the GEM S we have a constant R indler-like accelerated motion and the temperature measured by the detector is

2 T =
$$a_4 = r_+ R^{-1} (r^2 - r_+^2)^{-1=2} = (R^{-2} + a^2)^{1=2}$$
 (19)

which is that obtained using (3), and agrees with the tem perature given by the response function of particle detectors [13]. In the asymptotic limit r! 1, BTZ tends to AdS, the acceleration a! R¹, which is of course the acceleration of a \static" detector at in nity in AdS; both detectors m easure zero tem perature⁶ (no H aw king particle at in nity). The rotating case is m ore com plicated. The Hawking temperature 2 T = (R r) (r² r₊²) ¹⁼² (r² r²) ¹⁼² k_H , k_H = (r₊² r²) (r₊ R²) ¹ was calculated [12], [10] for trajectories that follow the time-like K illing vector $= Q_{1} N Q$, i.e. observers that obey = N t r = const (and hence are static" at in nity). A lthough they have a constant D = 3 acceleration, a = $(r^4 r^2 r_+^2) = (r^2 R^2 (r^2 r_+^2) (r^2 r_+^2))$, these trajectories do not describe pure R indler m otion in the GEMS, combining accelerated m otion in the $(z^0; z^1)$ plane with a space-like motion in $(z^3; z^2)$. Therefore, we cannot use their kinematical behavior in this GEM S to calculate the tem perature they measure. Exactly the same problem would arise for any AdS detector with \oint const in (14). This particular case resembles AdS motion with = (r)t, = 0. Our method can be used only for a group of detectors that maps into a group of pure Unruh observers in the GEM S. Hence, it is only possible to use it for those observers for whom the m ap of the detector trajectory into the τ answerse" embedding space (for BTZ the z^2 ; z^3 plane) is time-independent, i.e., the detector motion at any time is described by a xed point in that plane. There is one group of time-like observers obeying

$$= \frac{r}{r_{+}R}t ; r = const$$
 (20)

that does allow us to use the above GEMS and hence to compare the two calculation of T. These detectors have a constant acceleration $a = (r^2 r^2)^{1=2} (r^2 r_+^2)^{1=2} R^{-1}$ in BTZ and a R indler-like motion in the GEMS with acceleration $a_4 = R^{-1} (r_+^2 r^2)^{1=2} (r^2 r_+^2)^{-1=2}$ and therefore measure

2 T =
$$a_4 = \frac{1}{R} \frac{r_+^2 r_-^2}{r_+^2 r_+^2} = \frac{p}{R^2 + a^2}$$
 (21)

 $^{{}^{6}}$ BTZ form ally becomes AdS in our coordinates by setting r = 0 and r₊ = iR; (17) and the D = 3 version of (12) are the same. This shows again that AdS has a hidden in aginary horizon which causes the threshold in the temperature (acceleration sm aller than R⁻¹ measures no temperature).

On the other hand, inserting = $\frac{r}{r+1}$ t into (17) gives

$$ds^{2} = \frac{(r^{2} - r_{+}^{2})(r_{+}^{2} - r_{-}^{2})}{r_{+}^{2}R^{2}}dt^{2} - \frac{r^{2}R^{2}}{(r^{2} - r_{+}^{2})(r^{2} - r_{-}^{2})}dr^{2}$$
(22)

which show us that they follow the time-like K illing vector eld = Q for this metric (or $= Q_t + r (r_+ R)^{-1}Q$ if we use (17)) and see an event horizon at the metric's own \real" event horizon $r = r_+$. The surface gravity is

$$k_{H} = (r_{+}^{2} r_{-}^{2}) = (r_{+} R^{2});$$
 (23)

which is the same as that calculated for the other group by using the other K illing vector. This equivalence exists since both have the same horizon $r = r_+$ and the K illing vectors they follow are the same there. Any scaling problem s are avoided since we used a common coordinate system. W hile surface gravity can be obtained from either of the metrics (22) or (17), the appropriate g_{00} must taken from (22) because only there is x^0 the time-like K illing vector followed by the observer. This gives

2 T =
$$\frac{1}{R} = \frac{1}{r_{+}^2} \frac{r_{+}^2}{r_{-}^2} r_{+}^2$$
; (24)

exactly the result obtained by using the GEMS. Finally, we note that a common alternative definition of BH temperature is to scale T by $p_{\overline{000}}$: $T_0 = p_{\overline{000}}T = k_{\mu} = 2$; as distinct from the local temperature T, it is T_0 that enters into the BH them odynamics relations. Since there is one observer (the $r = r_{+}$ one) that belongs to both of the di erent observer groups (= N t and $= r t_{-}(r_{+}R)$), and since T_0 is a global feature of all the members in the group, it is obvious that both groups should give the same temperature (this of course could be seen immediately from their surface gravity equivalence). On the other hand, it should be no surprise that detectors in the two di erent observer groups measure di erent temperatures even though their absolute accelerations are the same (the R indler relation 2 T = a_4 does not apply to the = N t group) because the temperature T is observer-dependent in general. Since BTZ is asymptotically AdS, both detectors will again measure zero temperature at r ! 1, where a ! R ¹.

4. Schwarzschild and related geom etries.

We now come to spaces with \m ore m anifest" real horizons. Once a GEM S has been found (they always exist [1]) for the desired physical space, it is a mechanical procedure, using the familiar embedding G auss-C odazzi-R icci equations to relate constant acceleration a_{c} in GEM S to the embedded space physics; this is also possible when (as for Schwarzschild) the GEM S is more than one dimension higher. The acceleration of detectors that follow a time-like K illing vector in the physical space is $[2]a = r = j^{2}j$ where j j is the norm of . It is related to a_{c} in the GEM S according to

$$a_{G}^{2} = a^{2} + {}^{2}jj^{4}$$
 (25)

where is the second fundamental form [1]. Thus the temperature should simply be 2 T = $a_{c} = [a^{2} + {}^{2}j j^{4}]^{1=2}$. One should not, however, assume from this form ula that there is always a temperature, since in fact 2 need not always be positive (it is ${}^{2}j j^{4} = R {}^{2}$ in AdS). After all, it is only when a_{c}^{2} is non-negative that the Unruh description itself is meaningful in a at space.

We apply these ideas rst to the three types of Schwarzschild (vacuum) spaces, beginning with the usual case without cosm ological constant; it can be globally embedded in at D = 6

$$ds^{2} = (dz^{0})^{2} (dz^{1})^{2} (dz^{2})^{2} (dz^{3})^{2} (dz^{4})^{2} (dz^{5})^{2}; \qquad (26)$$

using the coordinate transform ation [14],

$$z^{0} = 4m^{2} \frac{q}{1 - 2m = r} \sinh(t = 4m) \quad z^{1} = 4m^{2} \frac{q}{1 - 2m = r} \cosh(t = 4m)$$

$$z^{2} = dr^{2} \frac{q}{(2m r^{2} + 4m^{2}r + 8m^{3}) = r^{3}}$$

$$z^{3} = r \sin \sin z^{4} = r \sin \cos z^{5} = r \cos z^{5}$$
(27)

This transform ation can be extended to cover the r < 2m interior thanks to the analyticity of z^2 (r) in r > 0. Indeed, the extension is just the maximal K ruskalone [15]. The original H awking detectors (moving according to constant r, ,), are here U nruh detectors; their six-space motions are the now familiar hyperbolic trajectories

$$(z^{1})^{2}$$
 $(z^{0})^{2} = 16m^{2} (1 \ 2m = r) = a_{6}^{2}$: (28)

Hence we immediately infer the local Hawking and BH temperatures

$$T = a_6 = 2 = (8 \text{ m} \ 1 \ 2\text{m} = \text{r})^{-1}$$
; $T_0 = \ p \ \overline{g_{00}} T = (8 \text{ m})^{-1}$: (29)

It should be cautioned that use of incomplete embedding spaces, that cover only r > 2m (as for example in [16]), will lead to observers there for whom there is no event horizon, no loss of inform ation, and no temperature.

The above calculation is easily generalized to Schwarzschild-AdS spaces (where 1 2m = r is replaced by 1 $2m = r + r^2 = R^2$) using a D = 7 GEMS with an additional timelike dimension z^6 ,

$$z^{0} = k_{H}^{1} \frac{1}{1} \frac{2m}{2m} = r + r^{2} = R^{2} \sinh(k_{H} t); \quad z^{1} = k_{H}^{1} \frac{q}{1} \frac{2m}{2m} = r + r^{2} = R^{2} \cosh(k_{H} t)$$

$$z^{2} = \frac{Z}{R^{2} + 3r_{H}^{2}} \frac{R^{3} + Rr_{H}^{2}}{R^{2} + 3r_{H}^{2}} \frac{\frac{q}{r}}{r^{3}(r^{2} + rr_{H} + r_{H}^{2} + R^{2})} dr \qquad (30)$$

$$z^{6} = \frac{Z}{t} \frac{V_{H}^{0}}{r^{2} + rr_{H}^{2} + rr_{H}^{2} + R^{2}} \frac{dr}{r^{2} + rr_{H}^{2} + rr_{H}^{2} + R^{2}} \frac{dr}{R^{2} + 3r_{H}^{2}}$$

and $(z^3; z^4; z^5)$ as in (27); $k_{_H} = (R^2 + 3r_{_H}^2)=2r_{_H}R^2$ is the surface gravity at the root $r_{_H}$ of (1 $2m = r + r^2 = R^2) = 0$. Using this GEM S⁷, we obtain 2 $T = k_{_H}$ (1 $2m = r + r^2 = R^2$) ¹⁼², equal to that calculated in [10]. [It may seem that we have the freedom to choose an arbitrary constant rather than $k_{_H}$ in z^0 and z^1 and thereby get a di erent temperature. But for any other choice, z^2 and z^6 cannot be chosen so that both their integrands are nite at the horizon. Hence, such embedding spaces are not global, cover only the area outside the horizon and cannot be extended; they are therefore excluded.]

For Schwarzschild-dS, which di ers form ally from Schwarzschild-AdS by $R^2 ! R^2$, there are two real horizons $(r_+; r)$ in general, both of which could be seen by physical detectors (such as constant r, with $r < r < r_+$). This requires use of a GEMS that captures both horizons. A lthough we have not tried to de ne this bigger GEMS, we do reproduce the known results [17] for

⁷ It is easy to see that when R ! 1 (the Schwarzschild lim it), r_{μ} ! 2m, k_{μ} ! $(2r_{\mu})^{1}$ and z^{2} becomes identical to the Schwarzschild one while z^{6} vanishes, so that we indeed get back the Schwarzschild GEM S.W hen m = 0 (the AdS lim it), we have $r_{\mu}^{2} = k_{\mu}^{2} = R^{2}$ and both z^{2} and z^{6} vanish, leaving the AdS GEM S of (12).

tem perature of each separate horizon, by using (30), with $R^2 ! R^2$ and the respective $k_{_{H}}$ (r₊), $k_{_{H}}$ (r). [Durm ethod becomes meaningless for the extrem al (r₊ = r) case since the whole R indler wedge vanishes there.]

We turn now to an example with matter, the Reissner {Nordstrom solution with

$$ds^{2} = 1 \frac{2m}{r} + \frac{e^{2}}{r^{2}} dt^{2} = 1 \frac{2m}{r} + \frac{e^{2}}{r^{2}} dr^{2} = r^{2} (d^{2} + \sin^{2} d^{2}):$$
(31)

A lthough there are two horizons $(r = m - m^2 - e^2)$ in the nonextrem al case (m > e), it is still simple to calculate the temperature via the embedding space. As explained earlier, a reliable GEMS has to cover (or be extendable to cover) both sides of the horizon, or else there is no loss of inform ation for a detector in that space. But physical $(r > r_+) r = \text{const Reissner}\{N \text{ ordstrom} detectors are aware only of the existence of one horizon <math>r_+$, unlike the physical Schwarzschild-dS r = const detectors $(r_+ > r > r)$ that see two horizons. Therefore, it is enough to use as the embedding space, again with an added tim elike z^6 dimension,

$$z^{0} = k_{H}^{1} \frac{q}{1 - 2m = r + e^{2} = r^{2}} \sinh(k_{H} t); \quad z^{1} = k_{H}^{1} \frac{q}{1 - 2m = r + e^{2} = r^{2}} \cosh(k_{H} t)$$

$$z^{2} = \frac{z}{r^{2} (r_{+} + r_{-}) + r_{+}^{2} (r_{+} + r_{+})}{r^{2} (r_{-} r_{-})} dr \qquad (32)$$

$$z^{6} = \frac{z}{r^{4} (r_{+} + r_{-})^{2}} dr$$

with (z^3, z^4, z^5) as in (27), and $k_{_H} = k(r_+) = (r_+ r_-) = 2r_+^2$. [In the neutral, e = 0, limit, z^6 vanishes and this GEMS becomes the (D=6) Schwarzschild one.] Even though it does not reach down to r_-r_- , this embedding su ces, because it covers r_+ , for the purpose of calculating the Reissner{N ordstrom temperature in the nonextrem al case⁸. It is clear from (32) that the relevant D=7 acceleration $a_7 = ((z^1)^2 (z^0)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (2r_+^2 (z^2 - 1)^2)^{-1=2} = (r_+ r_-) = (r_+ r_+) = (r_+ r_+)^{-1=2} = (r_+ r_+)^{-1=2$

5. Entropy

We turn now to the \extensive" companion of temperature, the entropy. For those of our curved spaces with intrinsic horizons, and at our sem iclassical level, entropy is just one quarter of the horizon area. Entropy can also be de ned for a R indler wedge [7], using arguments similar to those used originally [19] for Schwarzschild and dS. Here the relevant area is that of the null surface $x^2 + t^2 = 0$. This \transverse" area is in general in nite for otherwise unrestricted R indler motion, being just the cartesian dy dz for D = 4, say. For our purposes, however, we must evaluate this area subject to the embedding constraints, and we shall see, the resulting integral becomes nite and agrees with that of the original horizon. [This is not a tautology: we are not initially writing the original horizon area in embedding coordinates, although the result is indeed that real and embedding horizon areas agree. Nor is it a surprise: we have insured that (when present) horizons m ap to horizons.]

 $^{^{8}}$ To be sure, our mapping approach has limitations: since the R indler horizon of the GEMS is K illing bifurcate, one can only map from spaces whose horizons also are; this excludes the strictly extrem al (m = e) R eissner-N ordstrom, which is also exceptional from the D = 4 point of view [18].

Let us begin with the dS case, where the R indler horizon condition is $(z^1)^2 (z^0)^2 = 0$ which was Z = R, and of course $(z^2)^2 + (z^3)^2 + (z^4)^2 = R^2$. Thus the integration over $dz^2 dz^3 dz^4$ is restricted to the surface of the sphere of radius R, precisely that of the true horizon. The AdS case di ers, (as expected from lack of an intrinsic horizon) and the corresponding restrictions are $(z^{1})^{2}$ $(z^{0})^{2} = 0$ which again implies Z = R, but now $(z^{2})^{2} + (z^{3})^{2}$ $(z^{4})^{2} = R^{2}$, and the area of this hyperboloidal surface diverges, having no further restrictions. For comparison with the BTZ case below, the cause of the in nity can be traced to the fact that the limits on the $\frac{1}{2}$ integral are

R sinh, with 1 < < 1.

We now see how the BTZ solution leads to a nite Unruh area due to the periodic identi cation of mod 2. The $(z^1)^2$ $(z^0)^2 = 0$ R indler horizon condition in plies $r = r_+$, while $(z^3)^2$ $(z^2)^2 = R^2 (r^2 r^2) = (r_+^2 r^2) = R^2$ still looks hyperbolic. However, the relevant bounds on z^3 due to the periodicity are R sinh ($r_{+} = R^2$) and R sinh ($r_{+} = R^2$) for the nonrotating case, so that one has the integral

(and lim its R sinh ($r_+ = R r t = R^2$) and R sinh ($r_+ = R + r t = R^2$) in the rotating case) and it yields the desired area integral 2 r+ . It is clear that the limits di er from the AdS ones precisely in having the $\angle's''$ bounds be nite here.

The Schwarzschild case, where there are two additional dimensions in the transverse area, dz^2 ::: dz^5 , is correspondingly subject to three constraints: $(z^1)^2$ $(z^0)^2 = 0$ leads to r = 2m, phorizon to horizon mapping) $z^2 = f(r)$ and $(z^3)^2 + (z^4)^2 + (z^5)^2 = r^2$. Thus the z^2 integral, dz^2 (z^2 f (r)), is unity, while the remaining integrals of course reproduce the area of the r = 2m sphere in D = 3. The R eissner{N ordstrom and the Schwarzschild-A dS calculations are essentially the same⁹, except that in these cases there are three (rather than two) additionald in ensions, and four constraints: $(z^1)^2$ $(z^0)^2 = 0$ leads to $r = r_+$, $z^2 = f_1(r)$, $z^6 = f_2(r)$ and $(z^3)^2 + (z^4)^2 + (z^5)^2 = r^2$. Thus the z^2 ; z^6 integrals, $dz^2 dz^6 (z^2 f_1(r)) (z^6 f_2(r))$, are unity, and the z^3 ; z^4 ; z^5 integrals gives the desired area, that of the $r = r_+$ sphere. Having two separate horizons, the SchwarzschilddS system is more delicate to handle, but just as for temperature, we can calculate entropy for each horizon separately, to obtain the corresponding D = 4 results [17].

6. Sum m ary

We have form ulated a uniform mechanism for reducing curved space BH horizon temperatures and entropies to those of the kinem atical Unruh e ect due to R indler m otion in their GEM S. The latter must of course include the horizon in question. This method has been applied to a variety of \true" BH spacetimes, both vacuum ones like BTZ, Schwarzschild and its dS/AdS extensions, as well as Reissner-Nordstrom . It would be interesting to consider other possible applications of GEMS, for example to superradiance in rotating geom etries.

W e thanks M. Banados for useful discussions of BTZ. This work was supported by NSF grant PHY-9315811 and by the Fishbach Foundation for OL.

References

 $^{^9}$ W e m ay still use the Reissner{N ordstrom partial GEM S for the area calculation, since it covers the original $r = r_{+}$ horizon.

- [1] Goenner H F in 1980 General Relativity and Gravitation Held A ed. (New York: Plenum) 441
- [2] Namhofer H, Peter I and Thirring W 1996 Int. J. Mod. Phys. B 10 1507
- [3] Deser S and Levin O 1997 Class. Quantum Grav. 14 L163
- [4] Deser S and Levin O 1998 hep-th 9806223
- [5] Banados M, Teitelboim C and Zanelli J 1992 Phys. Rev. Let. 69 1849
- [6] Banados M, Henneaux M, Teitelboim C and Zanelli J 1993 Phys. Rev. D 48 1506
- [7] La ammeR 1987 Phys. Lett. B 196 449
- [8] Hawking SW 1975 Commun.Math.Phys. 42 199
- [9] Bekenstein J D 1973 Phys. Rev. D 7 2333
- [10] Brown JD, Creighton J and M ann R B. 1994 Phys. Rev. D 50 6394
- [11] W ald R M 1994 Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics (Chicago: The University of Chicago Press)
- [12] Carlip S 1995 C lass. Q uantum G rav. 12 2853
- [13] Lifschytz G and Ortiz M 1994 Phys. Rev. D 49 1929
- [14] FronsdalC.1959 Phys. Rev. 116 778
- [15] KruskalM D 1960 Phys. Rev. 119 1743
- [16] Rosen J 1993 Rev. M od. Phys. 37 204
- [17] Gibbons G W and Hawking S W 1977 Phys. Rev. D 15 2738
- [18] Jacobson T 1994 Phys.Rev.D 50 R 6031 Hawking S W , Horowitz G T and Ross S F 1995 Phys.Rev.D 51 4302 M oretti V and SteidlS 1996 C lass.Quantum G rav.13 2121
- [19] Gibbons G W and Hawking S W 1977 Phys. Rev. D 15 2752