Extended Superconformal Algebras on AdS_3

Katsushi Ito

Yukawa Institute for Theoretical Physics Kyoto University, Kyoto 606-8502, Japan

Abstract

We study a supersymmetric extension of the Virasoro algebra on the boundary of the anti-de Sitter space-time AdS_3 . Using the free field realization of the currents, we show that the world-sheet affine Lie superalgebras $osp(1|2)^{(1)}$, $sl(1|2)^{(1)}$ and $sl(2|2)^{(1)}$ provide the boundary $\mathcal{N}=1,2$ and 4 extended superconformal algebras, respectively.

The duality between the type IIB string theory on $AdS_3 \times S^3 \times M^4$, where $M^4 = K3$ or T^4 , and two-dimensional $\mathcal{N}=4$ superconformal field theory on a symmetric product of M^4 [1, 2, 3], is one of interesting examples of the AdS/CFT correspondence[4]. Conformal symmetry on the boundary of AdS_3 , firstly introduced by Brown and Henneaux [5], is realized as the chiral algebra of the boundary conformal field theory associated with the SL(2,R) Chern-Simons theory[6]. On the other hand, Giveon et al. [3] constructed the boundary Virasoro algebra from the string theory on AdS_3 . The generators of the algebra are identified as the global charges associated with the world-sheet sl(2,R) current algebra, which is expressed in terms of the free fields[7]. They also constructed (a part of) the $\mathcal{N}=4$ superconformal algebra from the superstrings on $AdS_3 \times S^3 \times T^4$. Their boundary algebra, however, is not manifestly supersymmetric since the supercurrents are introduced by the bosonization of the world-sheet fermions and the algebra is defined up to the picture changing operator[8].

In the present paper, we study the boundary extended superconformal algebra realized in a manifestly supersymmetric way. By replacing the world-sheet affine Lie algebra sl(2,R) to an affine Lie superalgebra which includes the subalgebra sl(2,R) and employing the free field realization of the currents, we will obtain the extended superconformal algebra which acts on the boundary AdS_3 superspace. In particular, we will show that the world-sheet affine Lie superalgebras $osp(1|2)^{(1)}$, $sl(1|2)^{(1)}$ and $sl(2|2)^{(1)}$ provide the boundary $\mathcal{N}=1,2$ and 4 extended superconformal algebras, respectively.

We begin with reviewing the boundary Virasoro algebra associated with AdS_3 . The anti-de Sitter space AdS_3 is the hypersurface $-U^2-V^2+X^2+Y^2=-\ell^2$ embedded in the flat space $R^{2,2}$. In the coordinates (ρ, τ, ϕ) defined by $U=\ell \cosh \rho \sin \tau$, $V=\ell \cosh \rho \cos \tau$, $X=\ell \sinh \rho \cos \phi$, $Y=\ell \cosh \rho \sin \phi$, the metric is given by

$$\frac{ds^2}{\ell^2} = -\cosh^2 \rho d\tau^2 + \sinh^2 \rho d\phi^2 + d\rho^2. \tag{1}$$

In the AdS_3 space, there is $SL(2,R)_L \times SL(2,R)_R$ symmetry. The generators of $SL(2,R)_L$ read [1]

$$L_0 = i\partial_u, \quad L_{\pm 1} = ie^{\pm iu} \left(\coth 2\rho \partial_u - \frac{1}{\sinh 2\rho} \partial_v \mp \frac{i}{2} \partial_\rho \right),$$
 (2)

where $u = \tau + \phi$, $v = \tau - \phi$. $SL(2,R)_R$ is generated by $\bar{L}_{0,\pm 1}$, which are obtained by

 $u \leftrightarrow v$. Define the coordinates $(\tilde{\phi}, \gamma, \bar{\gamma})$ by

$$\tilde{\phi} = \ell \cosh \rho - i\tau,$$

$$\gamma = \tanh \rho e^{iu}, \quad \bar{\gamma} = \tanh \rho e^{iv}.$$
(3)

The metric (1) becomes

$$ds^2 = \ell^2 (d\tilde{\phi}^2 + e^{2\tilde{\phi}} d\gamma d\bar{\gamma}). \tag{4}$$

The $SL(2,R)_L$ generators are

$$L_{0} = \frac{1}{2}\partial\tilde{\phi} - \gamma\partial_{\gamma},$$

$$L_{-1} = -\partial_{\gamma},$$

$$L_{1} = \gamma\partial\tilde{\phi} - \gamma^{2}\partial_{\gamma} + \ell^{2}e^{-2\tilde{\phi}}\partial_{\bar{\gamma}}.$$
(5)

Thus in the limit $\tilde{\phi} \to \infty$, the derivative with respect to $\bar{\gamma}$ in L_1 , decouples and the SL(2,R) generators become holomorphic with respect to γ . ¹ These currents are regarded as the zero-mode part of the Wakimoto realization [7] of the affine Lie algebra sl(2,R) at level k:

$$H(z) = -i\sqrt{2}\alpha_{+}\partial\varphi + 2\gamma\beta,$$

$$J_{-}(z) = \beta,$$

$$J_{+}(z) = i\sqrt{2}\alpha_{+}\partial\varphi\gamma - \gamma^{2}\beta - k\partial\gamma,$$
(6)

where $\alpha_+ = \sqrt{-k+2}$. $(\beta(z), \gamma(z))$ is the bosonic ghost system with conformal weights (1,0) and has the operator product expansion (OPE) $\beta(z)\gamma(w) = 1/(z-w) + \cdots$. $\varphi(z)$ is a free boson with the OPE $\varphi(z)\varphi(w) = -\log(z-w) + \cdots$. Then

$$L_0 = -\frac{1}{2} \oint \frac{dz}{2\pi i} H(z), \quad L_{\pm 1} = \pm \oint \frac{dz}{2\pi i} J_{\pm}(z),$$
 (7)

generate sl(2,R). For integer n, it is shown in [3] that the generators

$$L_n = \oint \frac{dz}{2\pi i} \mathcal{L}_n(z),\tag{8}$$

where

$$\mathcal{L}_n(z) = -\frac{1-n^2}{2}H\gamma^n - \frac{n(n-1)}{2}J_-\gamma^{n+1} + \frac{n(n+1)}{2}J_+\gamma^{n-1},\tag{9}$$

Here we consider the Euclidean version of the theory, which is obtained by replacing τ by $-i\tau_E$.

satisfy the Virasoro algebra

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m+n,0}, \tag{10}$$

with central charge c = 6kp. Here the contour integral in (8) is taken around z = 0 and p is defined by

$$p \equiv \oint \frac{dz}{2\pi i} \frac{\partial \gamma}{\gamma}.$$
 (11)

We now consider an \mathcal{N} supersymmetric extension of this Virasoro algebra (10), which is realized in the superspace $(\gamma, \xi_1, \dots, \xi_{\mathcal{N}})$. Here $\xi_1, \dots, \xi_{\mathcal{N}}$ denote the Grassmann odd coordinates. The supercharges are realized as the conserved charges of the anti-commuting currents with spin one on the world-sheet, whose zero modes should obey the Lie superalgebras osp(1|2), osp(2|2) = sl(1|2) and sl(2|2) for $\mathcal{N} = 1, 2, 4$ supersymmetries respectively. Hence we expect that conformal field theory whose chiral algebra is the affine Lie superalgebra with a subalgebra sl(2, R), provides a natural supersymmetric generalization of the boundary conformal symmetry. In the following we will construct some boundary superconformal algebras explicitly. The affine Lie superalgebra \mathbf{g} is generated by the bosonic currents $J_{\alpha}(z)$ ($\alpha \in \Delta_0$), $H^i(z)$ ($i = 1, \dots, r$) and fermionic currents $j_{\beta}(z)$ ($\beta \in \Delta_1$). Here $H^i(z)$ are the Cartan currents and Δ_0 (Δ_1) the set of even (odd) roots.

The free field representation [9] of an affine Lie superalgebra with rank r is obtained by r free bosons $\phi^i(z)$ ($i=1,\dots,r$), pairs of bosonic ghosts ($\beta_{\alpha}(z),\gamma_{\alpha}(z)$) with conformal weights (1,0) for positive even roots α and pairs of fermionic ghosts ($\eta_{\alpha}(z),\xi_{\alpha}(z)$) with conformal weights (1,0) for positive odd roots α . The OPEs for these fields are $\phi^i(z)\phi^j(w) = -\delta_{ij}\log(z-w) + \cdots$, $\beta_{\alpha}(z)\gamma_{\alpha'}(w) = \delta_{\alpha,\alpha'}/(z-w) + \cdots$, $\eta_{\alpha}(z)\xi_{\alpha'}(w) = \delta_{\alpha,\alpha'}/(z-w) + \cdots$.

Firstly we consider the $\mathcal{N}=1$ superconformal symmetry. The Lie superalgebra osp(1|2) contains odd roots $\pm \alpha_1$ and even roots $\pm 2\alpha_1$ with $\alpha_1^2=1$. The free field realization of affine Lie superalgebra $osp(1|2)^{(1)}$ at level -k is

$$j_{\alpha_{1}}(z) = (-2k+2)\partial\xi + 2i\alpha_{+}\xi\partial\varphi + \gamma\eta - 2\xi\gamma\beta,$$

$$J_{2\alpha_{1}}(z) = \frac{-k}{2}\partial\gamma - (-k+2)\xi\partial\xi + i\alpha_{+}\gamma\partial\varphi - \gamma^{2}\beta - \gamma\xi\eta,$$

$$j_{-\alpha_{1}}(z) = \eta - 2\xi\beta, \quad J_{-2\alpha_{1}}(z) = \beta,$$

$$\alpha_{1} \cdot H(z) = -i\alpha_{+}\partial\varphi + 2\gamma\beta + \xi\eta,$$
(12)

where $(\eta, \xi) = (\eta_{\alpha_1}, \xi_{\alpha_1})$, $(\beta, \gamma) = (\beta_{2\alpha_1}, \gamma_{2\alpha_1})$ and $\alpha_+ = \sqrt{-k+3}$. This algebra contains the sl(2, R) current algebra which is generated by $H(z) = \alpha_1 \cdot H(z)$, $J_{\pm}(z) = J_{\pm 2\alpha_1}(z)$. Then the generators $\mathcal{L}_n(z)$ defined by (9) satisfy the Virasoro algebra (10). Concerning the supercurrent, let us introduce the conserved charges

$$G_{\pm \frac{1}{2}} = \frac{1}{\sqrt{2}} \oint \frac{dz}{2\pi i} j_{\pm \alpha_1}(z). \tag{13}$$

From the OPE for the currents, it is shown that L_n $(n = 0, \pm 1)$ and $G_{\pm 1/2}$ satisfy the Lie superalgebra osp(1|2). Higher modes of the supercurrent are obtained by extracting the simple pole term in the OPE between $\mathcal{L}_n(z)$ and $j_{-\alpha_1}(z)$, which is denoted by $\mathcal{G}_{n-1/2}(z)$. The result is

$$\mathcal{G}_{n-\frac{1}{2}}(z) = \frac{n}{\sqrt{2}} j_{\alpha_1} \gamma^{n-1} - \frac{n-1}{\sqrt{2}} j_{-\alpha_1} \gamma^n - \sqrt{2} n(n-1) \left(H \gamma^{n-1} \xi + J_+ \gamma^{n-2} \xi - J_- \gamma^n \xi \right). \tag{14}$$

We define the generators $G_{n-1/2}$ by

$$G_{n-\frac{1}{2}} = \oint \frac{dz}{2\pi i} \mathcal{G}_{n-\frac{1}{2}}(z). \tag{15}$$

It is shown that the generators L_n and G_r obey the $\mathcal{N}=1$ superconformal algebra in the NS sector:

$$[L_{m}, L_{n}] = (m-n)L_{m+n} + \frac{c}{12}(m^{3}-m)\delta_{m+n,0},$$

$$[L_{n}, G_{r}] = \left(\frac{n}{2}-r\right)G_{n+r},$$

$$\{G_{r}, G_{s}\} = 2L_{r+s} + \frac{c}{3}(r^{2}-\frac{1}{4})\delta_{r+s,0}, \quad (n, m \in \mathbf{Z}, r, s \in \mathbf{Z} + \frac{1}{2})$$

$$(16)$$

with the central charge c = 6kp. For example, the third eq. in (16) follows from OPEs

$$j_{\alpha_{1}}(z)j_{\alpha_{1}}(w) = \frac{4J_{2\alpha_{1}}(w)}{z-w} + \cdots, \quad j_{-\alpha_{1}}(z)j_{-\alpha_{1}}(w) = \frac{-4J_{-2\alpha_{1}}(w)}{z-w} + \cdots,$$

$$j_{\alpha_{1}}(z)j_{-\alpha_{1}}(w) = \frac{2k}{(z-w)^{2}} + \frac{-2\alpha_{1} \cdot H(w)}{z-w} + \cdots,$$

$$j_{\alpha_{1}}(z)\gamma(w) = \frac{-2\gamma\xi(w)}{z-w} + \cdots, \quad j_{\alpha_{1}}(z)\xi(w) = \frac{\gamma(w)}{z-w} + \cdots,$$

$$j_{-\alpha_{1}}(z)\gamma(w) = \frac{-2\xi(w)}{z-w} + \cdots, \quad j_{-\alpha_{1}}(z)\xi(w) = \frac{1}{z-w} + \cdots,$$

$$(17)$$

and $(H\gamma^{n-1} + J_+\gamma^{n-2} - J_-\gamma^n)\xi = -k/2\gamma^{n-2}\partial\gamma\xi$. Note that the third term in (14) is necessary to obtain the correct superconformal algebra, although in the bosonic case the

combination $-H\gamma - J_+ + J_-\gamma^2$ becomes the total derivative and may be regarded as the longitudinal part. Let us take the zero mode part of the supercurrent and the Virasoro generator. Regarding β and η as the derivatives $\frac{\partial}{\partial \gamma}$ and $\frac{\partial}{\partial \xi}$ respectively, we find

$$G_{n-1/2} \sim \frac{1}{\sqrt{2}} (\gamma^n \frac{\partial}{\partial \xi} - 2\gamma^n \xi \frac{\partial}{\partial \gamma}),$$

$$L_n \sim -\gamma^{n+1} \frac{\partial}{\partial \gamma} - \frac{n+1}{2} \gamma^n \xi \frac{\partial}{\partial \xi}.$$
(18)

Thus, in this classical limit, the $\mathcal{N}=1$ superconformal algebra is realized on the superspace with coordinates (γ, ξ) .

Now we generalize this result to the case of $\mathcal{N}=2,4$ extended superconformal algebras by considering affine Lie superalgebras $sl(1|2)^{(1)}$, $sl(2|2)^{(1)}$, respectively. The Lie superalgebra sl(N|M) has even roots $\pm(e_i-e_j)$ $(1 \le i < j \le N)$, $\pm(\delta_a-\delta_b)$ $(1 \le a < b \le M)$ and odd roots $\pm(e_i-\delta_a)$ $(i=1,\cdots,N,\ a=1,\cdots,M)$. Here e_i (δ_a) are the orthonormal basis with positive (negative) metric $e_i \cdot e_j = \delta_{ij}$ $(\delta_a \cdot \delta_b = -\delta_{ab})$. We take the simple roots as $\alpha_i = e_i - e_{i+1}$, $(i=1,\cdots,N-1)$, $\alpha_N = e_N - \delta_1$, $\alpha_{N+a} = \delta_a - \delta_{a+1}$ $(a=1,\cdots,M-1)$. The OPEs for the currents of the affine Lie superalgebra $sl(N|M)^{(1)}$ at level k are given by

$$\begin{split} J_{e_{i}-e_{j}}(z)J_{e_{k}-e_{l}}(w) &= \frac{k\delta_{j,k}\delta_{i,l}}{(z-w)^{2}} \\ &+ \frac{\delta_{j,k}(1-\delta_{i,l})J_{e_{i}-e_{l}}(w) - \delta_{i,l}(1-\delta_{k,i})J_{e_{k}-e_{i}}(w) + \delta_{j,k}\delta_{il}(e_{i}-e_{j}) \cdot H(w)}{z-w} + \cdots, \\ J_{\delta_{a}-\delta_{b}}(z)J_{\delta_{c}-\delta_{d}}(w) &= \frac{-k\delta_{b,c}\delta_{a,d}}{(z-w)^{2}} \\ &+ \frac{\delta_{d,a}(1-\delta_{c,b})J_{\delta_{c}-\delta_{b}}(w) - \delta_{b,c}(1-\delta_{a,d})J_{\delta_{a}-\delta_{d}}(w) - \delta_{b,c}\delta_{a,d}(\delta_{a}-\delta_{b}) \cdot H(w)}{z-w} + \cdots, \\ j_{e_{i}-\delta_{a}}(z)j_{\delta_{b}-e_{j}}(w) &= \frac{-k\delta_{i,j}\delta_{a,b}}{(z-w)^{2}} \\ &- \frac{\delta_{a,b}(1-\delta_{i,j})J_{e_{i}-e_{j}}(w) + \delta_{i,j}(1-\delta_{a,b})J_{\delta_{b}-\delta_{a}}(w) - \delta_{i,j}\delta_{a,b}(\delta_{a}-e_{i}) \cdot H(w)}{z-w} + \cdots, \\ J_{e_{i}-e_{j}}(z)j_{e_{k}-\delta_{a}}(w) &= \frac{\delta_{j,k}j_{e_{i}-\delta_{a}}(w)}{z-w} + \cdots, \quad J_{e_{i}-e_{j}}(z)j_{\delta_{a}-\delta_{k}}(w) &= \frac{-\delta_{i,k}j_{\delta_{a}-\delta_{j}}(w)}{z-w} + \cdots, \\ J_{\delta_{a}-\delta_{b}}(z)j_{e_{k}-\delta_{c}}(w) &= \frac{\delta_{a,c}j_{e_{k}-\delta_{b}}(w)}{z-w} + \cdots, \quad J_{\delta_{a}-\delta_{b}}(z)j_{\delta_{c}-e_{k}}(w) &= \frac{-\delta_{b,c}j_{\delta_{a}-e_{k}}(w)}{z-w} + \cdots, \\ H^{i}(z)H^{j}(w) &= \frac{k\delta^{ij}}{(z-w)^{2}} + \cdots, \end{split}$$

$$H^{i}(z)J_{\alpha}(w) = \frac{\alpha^{i}J_{\alpha}(w)}{z - w} + \cdots, \quad H^{i}(z)j_{\alpha}(w) = \frac{\alpha^{i}j_{\alpha}(w)}{z - w} + \cdots.$$
 (19)

For M = 2, $H(z) = \alpha_{N+1} \cdot H(z)$, $J_{\pm}(z) = J_{\pm \alpha_{N+1}}(z)$ become the currents of the sl(2, R) subalgebra. Then it will be shown that (9) for $\gamma = \gamma_{\alpha_{N+1}}$ obeys the Virasoro algebra (10). The other even subalgebra $u(N)^{(1)}$ ($sl(2)^{(1)}$ for N = 2) turns out to be an affine Lie algebra symmetry on the boundary.

Let us consider the $\mathcal{N}=2$ (sl(1|2)) case. The free field realization of the affine Lie superalgebra $sl(1|2)^{(1)}$ at level k is given by

$$j_{-\alpha_{1}}(z) = \eta_{\alpha_{1}}, \quad J_{-\alpha_{2}}(z) = \beta_{\alpha_{2}} - \xi_{\alpha_{1}}\eta_{\alpha_{1}+\alpha_{2}}, \quad j_{-\alpha_{1}-\alpha_{2}}(z) = \eta_{\alpha_{1}+\alpha_{2}},$$

$$j_{\alpha_{1}}(z) = k\partial\xi_{\alpha_{1}} + i\alpha_{+}\alpha_{1} \cdot \partial\varphi\xi_{\alpha_{1}} - (\xi_{\alpha_{1}+\alpha_{2}} + \xi_{\alpha_{1}}\gamma_{\alpha_{2}})\beta_{\alpha_{2}} - \xi_{\alpha_{1}}\xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{1}+\alpha_{2}},$$

$$J_{\alpha_{2}}(z) = -(k+1)\partial\gamma_{\alpha_{2}} - i\alpha_{+}\alpha_{2} \cdot \partial\varphi\gamma_{\alpha_{2}} - \gamma_{\alpha_{2}}^{2}\beta_{\alpha_{2}} - \xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{1}},$$

$$j_{\alpha_{1}+\alpha_{2}}(z) = k\partial\xi_{\alpha_{1}+\alpha_{2}} + (k+1)(\partial\gamma_{\alpha_{2}})\xi_{\alpha_{1}} + i\alpha_{+}(\alpha_{1}+\alpha_{2}) \cdot \partial\varphi\xi_{\alpha_{1}+\alpha_{2}}$$

$$+i\alpha_{+}\alpha_{2} \cdot \partial\varphi\gamma_{\alpha_{2}}\xi_{\alpha_{1}} + \xi_{\alpha_{1}}\xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{1}} + \xi_{\alpha_{1}+\alpha_{2}}\gamma_{\alpha_{2}}\beta_{\alpha_{2}} + \xi_{\alpha_{1}}\gamma_{\alpha_{2}}^{2}\beta_{\alpha_{2}},$$

$$H^{i}(z) = -i\alpha_{+}\partial\varphi^{i} + \alpha_{1}^{i}\xi_{\alpha_{1}}\eta_{\alpha_{1}} + \alpha_{2}^{i}\gamma_{\alpha_{2}}\beta_{\alpha_{2}} + (\alpha_{1}+\alpha_{2})^{i}\xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{1}+\alpha_{2}},$$

$$(20)$$

where $\alpha_{+} = \sqrt{k-1}$. Let us introduce the operator

$$\mathcal{L}_n(z) = \frac{1 - n^2}{2} \alpha_2 \cdot H \gamma^n - \frac{n(n-1)}{2} J_{-\alpha_2} \gamma^{n+1} + \frac{n(n+1)}{2} J_{\alpha_2} \gamma^{n-1}, \tag{21}$$

which corresponds to the Virasoro generator. Here $\gamma = \gamma_{\alpha_2}$. As in the case of $\mathcal{N} = 1$, from the OPEs between $\mathcal{L}_n(z)$ and $j_{\pm \alpha_1}(w)$, one may construct the operators corresponding to the supercurrents:

$$\mathcal{G}_{n-1/2}(z) = -(n-1)j_{-\alpha_1-\alpha_2}\gamma^n + nj_{-\alpha_1}\gamma^{n-1},
\bar{\mathcal{G}}_{n-1/2}(z) = -(n-1)j_{\alpha_1}\gamma^n - nj_{\alpha_1+\alpha_2}\gamma^{n-1}
-n(n-1)\left(-\alpha_2 \cdot H\gamma^{n-1}\xi_{\alpha_1+\alpha_2} + J_{-\alpha_2}\gamma^n\xi_{\alpha_1+\alpha_2} - J_{\alpha_2}\gamma^{n-2}\xi_{\alpha_1+\alpha_2}\right). (22)$$

By extracting the simple pole term in the OPE between $\mathcal{G}_{n-1/2}(z)$ and $\bar{\mathcal{G}}_{1/2}(w)$, one may read off the operator $\mathcal{T}_n(z)$, which corresponds to the U(1) current. The result is

$$\mathcal{T}_n(z) = -(\alpha_1 + 2\alpha_2) \cdot H\gamma^n - n\tilde{\xi}_{\alpha_1 + \alpha_2}\tilde{\eta}_{\alpha_1}\gamma^{n-1}, \tag{23}$$

where $\tilde{\xi}_{\alpha_1+\alpha_2} = \xi_{\alpha_1+\alpha_2} + \gamma_{\alpha_2}\xi_{\alpha_1}$, $\tilde{\eta}_{\alpha_1} = \eta_{\alpha_1} - \gamma_{\alpha_2}\eta_{\alpha_1+\alpha_2}$. After introducing these operators, we can show that the generators

$$L_n = \oint \frac{dz}{2\pi i} \mathcal{L}_n(z), \quad T_n = \oint \frac{dz}{2\pi i} \mathcal{T}_n(z), \quad (n \in \mathbf{Z})$$

$$G_r = \oint \frac{dz}{2\pi i} \mathcal{G}_r(z), \quad \bar{G}_r = \oint \frac{dz}{2\pi i} \bar{\mathcal{G}}_r(z), \quad (r \in \mathbf{Z} + 1/2)$$
(24)

satisfy the $\mathcal{N}=2$ superconformal algebra with central charge c=6kp:

$$[L_{m}, L_{n}] = (m-n)L_{m+n} + \frac{c}{12}(m^{3}-m)\delta_{m+n,0},$$

$$[L_{m}, G_{r}] = \left(\frac{m}{2}-r\right)G_{m+r}, \quad [L_{m}, \bar{G}_{r}] = \left(\frac{m}{2}-r\right)\bar{G}_{m+r},$$

$$[L_{m}, T_{n}] = -nT_{m+n},$$

$$[T_{m}, G_{r}] = G_{m+r}, \quad [T_{m}, \bar{G}_{r}] = \bar{G}_{m+r},$$

$$[T_{m}, T_{n}] = \frac{c}{3}m\delta_{m+n,0},$$

$$\{G_{r}, G_{s}\} = \{\bar{G}_{r}, \bar{G}_{s}\} = 0,$$

$$\{G_{r}, \bar{G}_{s}\} = L_{r+s} + \frac{1}{2}(r-s)T_{r+s} + \frac{c}{6}(r^{2} - \frac{1}{4})\delta_{r+s,0},$$
(25)

where $n, m \in \mathbf{Z}$ and $r, s \in \mathbf{Z} + 1/2$.

Similar construction can be done for $\mathcal{N}=4$ (sl(2|2)) case. In terms of free fields, the currents of the affine Lie superalgebra $sl(2|2)^{(1)}$ at level k are given by

$$J_{-\alpha_{1}}(z) = \beta_{\alpha_{1}}, \quad j_{-\alpha_{2}}(z) = \eta_{\alpha_{2}} - \gamma_{\alpha_{1}}\eta_{\alpha_{1}+\alpha_{2}},$$

$$J_{-\alpha_{3}}(z) = \beta_{\alpha_{3}} - \xi_{\alpha_{2}}\eta_{\alpha_{2}+\alpha_{3}} - \xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}}, \quad j_{-\alpha_{1}-\alpha_{2}}(z) = \eta_{\alpha_{1}+\alpha_{2}},$$

$$j_{-\alpha_{2}-\alpha_{3}}(z) = \eta_{\alpha_{2}+\alpha_{3}} - \gamma_{\alpha_{3}}\eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}}, \quad j_{-\alpha_{1}-\alpha_{2}-\alpha_{3}}(z) = \eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}},$$

$$J_{\alpha_{1}}(z) = k\partial\gamma_{\alpha_{1}} + i\alpha_{+}\alpha_{1} \cdot \partial\varphi\gamma_{\alpha_{1}} - \gamma_{\alpha_{1}}^{2}\beta_{\alpha_{1}} + \gamma_{\alpha_{1}}\xi_{\alpha_{2}}\eta_{\alpha_{2}} - \gamma_{\alpha_{1}}\xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{1}+\alpha_{2}}$$

$$+\gamma_{\alpha_{1}}\xi_{\alpha_{2}+\alpha_{3}}\eta_{\alpha_{2}+\alpha_{3}} - \gamma_{\alpha_{1}}\xi_{\alpha_{1}+\alpha_{2}+\alpha_{3}}\eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}} - \xi_{\alpha_{1}+\alpha_{2}}\eta_{\alpha_{2}} - \xi_{\alpha_{1}+\alpha_{2}+\alpha_{3}}\eta_{\alpha_{2}+\alpha_{3}},$$

$$j_{\alpha_{2}}(z) = (k+1)\partial\xi_{\alpha_{2}} + i\alpha_{+}\alpha_{2} \cdot \partial\varphi\xi_{\alpha_{2}} + \xi_{\alpha_{1}+\alpha_{2}}\beta_{\alpha_{1}} - \xi_{\alpha_{2}}\gamma_{\alpha_{3}}\beta_{\alpha_{3}}$$

$$-\xi_{\alpha_{2}}\xi_{\alpha_{2}+\alpha_{3}}\eta_{\alpha_{2}+\alpha_{3}} + \xi_{\alpha_{2}+\alpha_{3}}\beta_{\alpha_{3}},$$

$$J_{\alpha_{3}}(z) = -(k+2)\partial\gamma_{\alpha_{3}} - i\alpha_{+}\alpha_{3} \cdot \partial\varphi - \gamma_{\alpha_{3}}^{2}\beta_{\alpha_{3}} + \xi_{\alpha_{2}+\alpha_{3}}\eta_{\alpha_{2}} + \xi_{\alpha_{1}+\alpha_{2}+\alpha_{3}}\eta_{\alpha_{2}+\alpha_{3}},$$

$$H^{i}(z) = -i\alpha_{+}\partial\varphi^{i} + \alpha_{1}^{i}\gamma_{\alpha_{1}}\beta_{\alpha_{1}} + \alpha_{3}^{i}\gamma_{\alpha_{3}}\beta_{\alpha_{3}} + \alpha_{2}^{i}\xi_{\alpha_{2}}\eta_{\alpha_{2}} + (\alpha_{1}+\alpha_{2})^{i}\xi_{\alpha_{1}+\alpha_{2}+\alpha_{3}},$$

$$+(\alpha_{2}+\alpha_{3})^{i}\xi_{\alpha_{2}+\alpha_{3}}\eta_{\alpha_{2}+\alpha_{3}} + (\alpha_{1}+\alpha_{2}+\alpha_{3})^{i}\xi_{\alpha_{1}+\alpha_{2}+\alpha_{3}}\eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}},$$

$$(26)$$

where $\alpha_{+} = \sqrt{k}$. The currents for positive non-simple roots are obtained from those for the simple roots. Introduce the operators

$$\mathcal{L}_{n}(z) = \frac{1 - n^{2}}{2} \alpha_{3} \cdot H \gamma^{n} - \frac{n(n-1)}{2} J_{-\alpha_{3}} \gamma^{n+1} + \frac{n(n+1)}{2} J_{\alpha_{3}} \gamma^{n-1},
\mathcal{G}_{n-1/2}^{1}(z) = -(n-1) j_{-\alpha_{1}-\alpha_{2}} \gamma^{n} + n j_{-\alpha_{1}-\alpha_{2}-\alpha_{3}} \gamma^{n-1},
\mathcal{G}_{n-1/2}^{2}(z) = -(n-1) j_{-\alpha_{2}} \gamma^{n} + n j_{-\alpha_{2}-\alpha_{3}} \gamma^{n-1},
\bar{\mathcal{G}}_{n-1/2}^{1}(z) = -(n-1) j_{\alpha_{1}+\alpha_{2}} \gamma^{n} - n j_{\alpha_{1}+\alpha_{2}+\alpha_{3}} \gamma^{n-1}
-n(n-1) \left(\alpha_{3} \cdot H \gamma^{n-1} \tilde{\xi}_{\alpha_{1}+\alpha_{2}+\alpha_{3}} + J_{-\alpha_{3}} \gamma^{n} \tilde{\xi}_{\alpha_{1}+\alpha_{2}+\alpha_{3}} - J_{\alpha_{3}} \gamma^{n-2} \tilde{\xi}_{\alpha_{1}+\alpha_{2}+\alpha_{3}}\right),
\bar{\mathcal{G}}_{n-1/2}^{2}(z) = -(n-1) j_{\alpha_{2}} \gamma^{n} - n j_{\alpha_{2}+\alpha_{3}} \gamma^{n-1}
-n(n-1) \left(\alpha_{3} \cdot H \gamma^{n-1} \tilde{\xi}_{\alpha_{2}+\alpha_{3}} + J_{-\alpha_{3}} \gamma^{n} \tilde{\xi}_{\alpha_{2}+\alpha_{3}} - J_{\alpha_{3}} \gamma^{n-2} \tilde{\xi}_{\alpha_{2}+\alpha_{3}}\right),
\mathcal{T}_{n}^{+}(z) = J_{\alpha_{1}} \gamma^{n} + n \tilde{\xi}_{\alpha_{1}+\alpha_{2}+\alpha_{3}} \tilde{\eta}_{\alpha_{1}+\alpha_{2}} \gamma^{n-1},
\mathcal{T}_{n}^{-}(z) = J_{-\alpha_{1}} \gamma^{n} + n \tilde{\xi}_{\alpha_{2}+\alpha_{3}} \tilde{\eta}_{\alpha_{1}+\alpha_{2}} \gamma^{n-1},
\mathcal{T}_{n}^{0}(z) = \frac{1}{2} \alpha_{1} \cdot H \gamma^{n} + \frac{n}{2} \tilde{\xi}_{\alpha_{1}+\alpha_{2}+\alpha_{3}} \tilde{\eta}_{\alpha_{1}+\alpha_{2}} \gamma^{n-1} - \frac{n}{2} \tilde{\xi}_{\alpha_{2}+\alpha_{3}} \tilde{\eta}_{\alpha_{2}} \gamma^{n-1},$$
(27)

where $\gamma = \gamma_{\alpha_3}$ and

$$\tilde{\xi}_{\alpha_{2}+\alpha_{3}} = \xi_{\alpha_{2}+\alpha_{3}} + \gamma_{\alpha_{3}}\xi_{\alpha_{3}},$$

$$\tilde{\xi}_{\alpha_{1}+\alpha_{2}+\alpha_{3}} = \xi_{\alpha_{1}+\alpha_{2}+\alpha_{3}} + \gamma_{\alpha_{1}}\xi_{\alpha_{2}+\alpha_{3}} + \gamma_{\alpha_{1}}\gamma_{\alpha_{3}}\xi_{\alpha_{2}} + \gamma_{\alpha_{3}}\xi_{\alpha_{1}+\alpha_{2}},$$

$$\tilde{\eta}_{\alpha_{1}+\alpha_{2}} = \eta_{\alpha_{1}+\alpha_{2}} - \gamma_{\alpha_{3}}\eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}},$$

$$\tilde{\eta}_{\alpha_{2}} = \eta_{\alpha_{2}} - \gamma_{\alpha_{1}}\eta_{\alpha_{1}+\alpha_{2}} + \gamma_{\alpha_{1}}\gamma_{\alpha_{3}}\eta_{\alpha_{1}+\alpha_{2}+\alpha_{3}} - \gamma_{\alpha_{3}}\eta_{\alpha_{2}+\alpha_{3}}.$$
(28)

Then the generators

$$L_{n} = \oint \frac{dz}{2\pi i} \mathcal{L}_{n}(z),$$

$$G_{r}^{a} = \oint \frac{dz}{2\pi i} \mathcal{G}_{r}^{a}(z), \quad \bar{G}_{r}^{a} = \oint \frac{dz}{2\pi i} \bar{\mathcal{G}}_{r}^{a}(z),$$

$$T_{n}^{\pm} = T_{n}^{1} \pm i T_{n}^{2} = \oint \frac{dz}{2\pi i} \mathcal{T}_{n}^{\pm}(z), \quad T_{n}^{0} = \oint \frac{dz}{2\pi i} \mathcal{T}_{n}^{0}(z), \quad (n \in \mathbf{Z}, r \in \mathbf{Z} + \frac{1}{2}) \quad (29)$$

can be shown to satisfy $\mathcal{N}=4$ superconformal algebra with central charge c=6kp:

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m+n,0},$$

$$[L_m, T_n^i] = -nT_{m+n}^i,$$

$$[L_{m}, G_{r}^{a}] = \left(\frac{m}{2} - r\right) G_{m+r}^{a}, \quad [L_{m}, \bar{G}_{r}^{a}] = \left(\frac{m}{2} - r\right) \bar{G}_{m+r}^{a},$$

$$[T_{m}^{i}, T_{n}^{j}] = i\varepsilon^{ijk} T_{m+n}^{k} + \frac{c}{12} m \delta^{ij} \delta_{m+n,0},$$

$$[T_{m}^{i}, G_{r}^{a}] = -\frac{1}{2} (\sigma^{i})_{b}^{a} G_{m+r}^{b}, \quad [T_{m}^{i}, \bar{G}_{r}^{a}] = \frac{1}{2} (\sigma^{*i})_{b}^{a} G_{m+r}^{b},$$

$$\{G_{r}^{a}, G_{s}^{b}\} = \{\bar{G}_{r}^{a}, \bar{G}_{s}^{b}\} = 0,$$

$$\{G_{r}^{a}, \bar{G}_{s}^{b}\} = \delta^{ab} L_{r+s} - (r-s)(\sigma^{i})_{ab} T_{r+s}^{i} + \frac{c}{6} (4r^{2} - 1)\delta^{ab} \delta_{r+s,0}, \tag{30}$$

where $i, j, k = 1, 2, 3, a, b = 1, 2, n, m \in \mathbf{Z}$ and $r, s \in \mathbf{Z} + 1/2$.

In the present work, we have constructed boundary extended superconformal symmetry from the world-sheet affine Lie superalgebra with a subalgebra sl(2, R). The Wakimoto type free field realization [9] is shown to be useful for the construction of manifestly supersymmetric extended superconformal algebras. We have write down only the NS sector of superconformal algebras. The R-sector of the algebra is obtained by the spectral flow [10] in the case of $\mathcal{N}=2,4$ extended superconformal algebras. For $\mathcal{N}=1$ case, it is necessary to find the world-sheet current which represents the supercurrent in the R-sector.

The present construction would be generalized to other affine Lie superalgebras associated with AdS_3 Lie supergroups[11]. The corresponding conformal algebras would be linearly extended superconformal algebras. On the other hand, the Chern-Simons theories based on general AdS_3 Lie supergroups lead to non-linearly extended superconformal algebras constructed in [12]. Hence it is an interesting problem to examine the boundary conformal field theory for general AdS_3 supergroups. Finally, in order to relate the present construction to the type IIB string theory on $AdS_3 \times S_3 \times T^4$ [13] or K3, we need to clarify the hidden affine Lie superalgebra symmetry in the type IIB theory. These subjects will be discussed elsewhere.

This work is supported in part by the Grant-in-Aid from the Ministry of Education, Science and Culture, Priority Area: "Supersymmetry and Unified Theory of Elementary Particles" (#707).

References

- [1] J. Maldacena and A. Strominger, hep-th/9804085.
- [2] J. de Boer, hep-th/9806104
- [3] A. Giveon, D. Kutasov and N. Seiberg, hep-th/9806194.
- [4] J. Maldacena, hep-th/9711200; S. Gubser, I. Klebanov and A. Polyakov, Phys. Lett.
 B428 (1998) 105, hep-th/9802109; E. Witten, hep-th/9802150.
- [5] J.D. Brown and M. Henneaux, Comm. Math. Phys. **104** (1986) 207.
- [6] O. Coussaert, M. Henneaux and P. van Driel, Class. Quant. Grav. 12 (1995) 2961;
 M. Bañados, K. Bautier, O. Courssaert, M. Henneaux and M. Ortiz, Phys. Rev. D58 (1998) 085020, hep-th/9805162;
 M. Bañados, M. E. Ortiz, hep-th/980689;
 K. Behrndt, I. Brunner and I. Gaida, hep-th/9806195.
- [7] M. Wakimoto, Commun. Math. Phys. 92 (1984) 455; D.Bernard and G.Felder, Commun. Math. Phys. 127 (1990) 145.
- [8] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986) 93.
- [9] M. Bershadsky and H. Ooguri, Phys. Lett. B229 (1989) 374; K. Ito, Int. J. Mod. Phys. A (1992) 4885; Phys. Lett. B259 (1991) 73; J. Rasmussen, Nucl. Phys. B510 (1998) 688, hep-th/9706091.
- [10] A. Schwimmer and N. Seiberg, Phys. Lett. **B184** (1987) 191.
- [11] M. Günaydin, G. Sierra and P.K. Townsend, Nucl. Phys. **B274** (1986) 429.
- [12] V. Knizhnik, Theor. Math. Phys. 66 (1986) 68; M. Bershadsky, Phys. Lett. B174 (1986) 285; K. Schoutens, Nucl. Phys. B314 (1989) 519; P. Bowcock, Nucl. Phys. B381 (1992) 415; K. Ito, J.O. Madsen, and J.L. Petersen, Nucl. Phys. B398 (1993) 425; Phys. Lett. B318 (1993) 315; E.S. Fradkin and V. Ya Linetsky, Phys. Lett. B282 (1992) 352; A. Sevrin, K. Thielmans and W. Troost, Nucl. Phys. B407 (1993) 459.

 $[13]\,$ I. Pesando, hep-th/9809145; J. Rahmfeld and A. Rajaraman, hep-th/9809164