

Discrete symmetry breaking and restoration at finite temperature in 3D Gross-Neveu model

Bang-Rong Zhou*

Department of Physics, Graduate School at Beijing

University of Science and Technology of China, Academia Sinica, Beijing 100039, **China**

and

The Abdus Salam International Centre for Theoretical Physics, Trieste 34100, **Italy**

Abstract

Dynamical spontaneous breaking of some discrete symmetries including special parities and time reversal and their restoration at finite temperature T are researched in 3D Gross-Neveu model by means of Schwinger-Dyson equation in the real-time thermal field theory in the fermion bubble diagram approximation. When the momentum cut-off Λ is large enough, the equation of critical chemical potential μ_c and critical temperature T_c will be Λ -independent and identical to the one obtained by auxiliary scalar field approach. The dynamical fermion mass $m \equiv m(T, \mu)$, as the order parameter of symmetry breaking, has the same $(T_c - T)^{1/2}$ behavior at $T \lesssim T_c$ as one in 4D NJL-model and this shows the second-order phase transition feature of the symmetry restoration at $T > T_c$. It is also proven that no scalar bound state could exist in this model.

PACS numbers: 11.10.Kk, 11.10.Wx, 11.30.Er, 11.30.Qc

Key words: 3D Gross-Neveu model, discrete symmetry breaking and restoration, real-time thermal field theory, scalar bound state

*Regular Associate of The Abdus Salam ICTP.

The Gross-Neveu (GN) model [1] in 2+1 space-time dimension is a model with four-fermion interactions. As a well-known and applicable model to condensed matter physics, it has been extensively researched [2-6]. The conventional approach is to introduce an auxiliary scalar field, then to replace the four-fermion interactions by the Yukawa couplings between the auxiliary scalar field and the fermion fields. Alternatively, one can also use Schwinger-Dyson equation directly to research the fermion condensates induced by the four-fermion interactions and relevant dynamical behavior. In the latter approach, no effective scalar field is introduced and, as far as some problems, for example, the calculation of the propagators for the bound states composed of fermions at finite temperature, are concerned, this approach could be more straightforward and convenient. On the other hand, almost all the discussions about the finite temperature behavior of the model were conducted in the imaginary-time formalism of thermal field theory. However, we also want to examine the application of the real-time formalism of thermal field theory to the model. Therefore, in this letter, instead of using the auxiliary scalar field and the imaginary-time thermal field theory, we will research the 3D GN model at finite temperature by means of Schwinger-Dyson equation in the real-time thermal field theory. All the discussions will be made in the fermion bubble diagram approximation. This approximation amounts to the leading order of $1/N$ expansion and is sometimes also called the mean field approximation. Since 3D GN model is renormalizable in $1/N$ expansion [5] and in 2+1 dimension, there is no the kink effect which could appear in a 1+1 dimension system [7], the mean field approximation can be considered as a reliable one. The results will also be compared with the previous works' ones based on the auxiliary field and the imaginary-time thermal field theory.

The Lagrangian of the model can be written by

$$\mathcal{L}(x) = \sum_{k=1}^N \bar{\psi}^k(x) i \gamma^\mu \partial_\mu \psi_k(x) + \frac{g}{2N} \sum_{k=1}^N [\bar{\psi}^k(x) \psi_k(x)]^2. \quad (1)$$

In 3 dimensions, the coupling constant g has mass dimension -1 thus the model is perturbatively non-renormalizable. However, this fact does not affect our discussions based on the $1/N$ expansion. The γ^μ ($\mu = 0, 1, 2$) can be taken as 2×2 matrices and related to the Pauli matrices σ^i ($i = 1, 2, 3$) by

$$\gamma^0 = \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^1 = i\sigma^1 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad \gamma^2 = i\sigma^2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (2)$$

which satisfy the Dirac algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}, \quad \mu, \nu = 0, 1, 2. \quad (3)$$

In this representation $\psi_k(x)$ is two-component complex spinor with N "color" degrees of freedom. Notice that there is no 2×2 matrix which anti-commutes with all the three γ^μ , i.e. no " γ_5 " matrix exists in 3 dimensions. Consequently $\mathcal{L}(x)$ has no chiral symmetry, though $\psi_k(x)$ are zero-mass fields. However, it is easy to verify that the action of system is invariant for the discrete transformation (omitting the index k of ψ)

$$\psi^1(x) \xleftrightarrow{Z_2} \psi^2(x), \quad \text{if } \psi(x) = \begin{pmatrix} \psi^1(x) \\ \psi^2(x) \end{pmatrix}, \quad (4)$$

the special \mathcal{P}_1 and \mathcal{P}_2 parity transformations

$$\psi(t, x^1, x^2) \xrightarrow{\mathcal{P}_1} \gamma^1 \psi(t, -x^1, x^2), \quad (5)$$

$$\psi(t, x^1, x^2) \xrightarrow{\mathcal{P}_2} \gamma^2 \psi(t, x^1, -x^2), \quad (6)$$

and the time reversal

$$\psi(t, x^1, x^2) \xrightarrow{\mathcal{T}} \gamma^2 \psi(-t, x^1, x^2). \quad (7)$$

We indicate that the mass term $-m\bar{\psi}(x)\psi(x)$ (if it exists) is not invariant under Z_2 , \mathcal{P}_1 , \mathcal{P}_2 and \mathcal{T} . Hence, if the four-fermion interactions could lead to generation of a non-zero dynamical mass of the ψ -fields, then the above Z_2 , \mathcal{P}_1 , \mathcal{P}_2 and \mathcal{T} discrete symmetries will be spontaneously broken. Since the broken symmetries are discrete, no Goldstone bosons are expected to appear. In the following we will research generation of the dynamical mass of $\psi(x)$ fields and its finite temperature behavior in the fermionic bubble graph approximation. This approximation, as the leading order of the $1/N$ expansion, will be able to give us a clear insight into the breaking and restoration of above discrete symmetries.

At zero-temperature $T = 0$, assume the four-fermion scalar interactions $g \sum_{k=1}^N (\bar{\psi}^k \psi_k)^2 / 2N$ can lead to the fermion condensates $\sum_{k=1}^N \langle \bar{\psi}^k \psi_k \rangle \neq 0$, then we will get the dynamical fermion mass $m(0) = -(g/N) \sum_{k=1}^N \langle \bar{\psi}^k \psi_k \rangle$ which satisfies the gap equation

$$1 = 2g \int \frac{id^3l}{(2\pi)^3} \frac{1}{l^2 - m^2(0) + i\varepsilon} \quad (8)$$

After Wick rotation, angular integration and introduction of 3D Euclidean momentum cut-off Λ , Eq. (8) becomes

$$1 = \frac{g\Lambda}{\pi^2} \left[1 - \frac{m(0)}{\Lambda} \arctan \frac{\Lambda}{m(0)} \right]. \quad (9)$$

We indicate that since Λ corresponds to the square root of the 3d squared Euclidean momentum, its introduction does not explicitly break the discussed discrete symmetries including \mathcal{P}_1 , \mathcal{P}_2 and \mathcal{T} . It is seen from Eq.(9) that the necessary condition for formation of the fermion condensates is sufficiently strong four-fermion coupling g that $g\Lambda/\pi^2 > 1$, similar to the one in 4D NJL-model case [8]. In both cases, one can consider the four-fermion couplings as an effective interactions below some high momentum scale Λ . For the sake of satisfying the gap equation Eq. (8), the fine-tuning of the coupling constant g is also required.

When $T \neq 0$, we must replace the vacuum expectation value $\sum_{k=1}^N \langle \bar{\psi}^k \psi_k \rangle$ by the thermal expectation value $\sum_{k=1}^N \langle \bar{\psi}^k \psi_k \rangle_T$. In the real-time thermal field theory, this implies the substitution of the fermion propagator [9,10]

$$\frac{i}{\not{V} - m + i\varepsilon} \rightarrow \frac{i}{\not{V} - m + i\varepsilon} - 2\pi\delta(l^2 - m^2)(\not{V} + m) \sin^2 \theta(l^0, \mu) \quad (10)$$

with

$$\sin^2 \theta(l^0, \mu) = \frac{\theta(l^0)}{\exp[\beta(l^0 - \mu)] + 1} + \frac{\theta(-l^0)}{\exp[\beta(-l^0 + \mu)] + 1}, \quad (11)$$

where $m \equiv m(T, \mu)$ is the dynamical fermion mass at finite temperature T and finite chemical potential μ and the denotation $\beta = 1/T$. As a result, the gap equation at $T \neq 0$ becomes

$$1 = gI \quad (12)$$

$$I = 2 \int \frac{d^3l}{(2\pi)^3} \left[\frac{i}{l^2 - m^2 + i\varepsilon} - 2\pi\delta(l^2 - m^2) \sin^2 \theta(l^0, \mu) \right] \quad (13)$$

$$= \frac{\Lambda}{\pi^2} \left\{ 1 - \frac{m}{\Lambda} \arctan \frac{\Lambda}{m} - \frac{\pi T}{2\Lambda} [I_2(y, -r) + I_2(y, r)] \right\} \quad (14)$$

where we have used the denotations

$$I_2(y, \mp r) = \int_0^\infty \frac{dx}{\sqrt{x^2 + y^2}} \frac{1}{\exp(\sqrt{x^2 + y^2} \mp r) + 1} = \ln \left[1 + e^{-(y \mp r)} \right] \quad (15)$$

with

$$x = \beta | \vec{l} |, \quad y = \beta m \quad \text{and} \quad r = \beta \mu. \quad (16)$$

Considering the gap equation (9) at $T = 0$, Eq.(12) may be changed into

$$m(0) \arctan \frac{\Lambda}{m(0)} = m \arctan \frac{\Lambda}{m} + \frac{\pi}{2} f(T) \quad (17)$$

with

$$f(T) = T \ln \left\{ \left[1 + e^{-(m-\mu)/T} \right] \left[1 + e^{-(m+\mu)/T} \right] \right\}. \quad (18)$$

It is easy to see that no matter $m - \mu \geq 0$ or $m - \mu < 0$ we always have $f'(T) = df(T)/dT > 0$, i.e. $f(T)$ is a monotone increasing function of temperature T . To keep the left-handed side of Eq.(17) to be a constant, the first term in the right-handed side must decrease as T increases and finally has $m \rightarrow 0$ in it at a critical temperature T_c and a critical chemical potential μ_c . Hence taking $m = 0$ in Eq.(17) we will obtain the equation to determine T_c and μ_c

$$m(0) \arctan \frac{\Lambda}{m(0)} = \frac{\pi}{2} T_c \left[\ln(1 + e^{\mu_c/T_c}) + \ln(1 + e^{-\mu_c/T_c}) \right]. \quad (19)$$

Here the explicit momentum cut-off Λ is not important. In fact, as long as the ratio $\Lambda/m(0) \geq 10^4$, then the value of $\arctan[\Lambda/m(0)]$ is almost equal to $\pi/2$. Assume that is the case then the momentum cut-off Λ will disappear from the equation and we obtain

$$m(0) = T_c \left[\ln(1 + e^{\mu_c/T_c}) + \ln(1 + e^{-\mu_c/T_c}) \right]. \quad (20)$$

We indicate that the $\mu - T$ criticality equation (20) comes from the two-point function of fermions which corresponds to $O(1)$ order in $1/N$ expansion, hence no explicit N -dependence is contained in it. It follows from Eq.(20) that

$$\begin{aligned} T_c/m(0) &= 1/2 \ln 2, & \text{if } \mu_c &= 0 \\ \mu_c/m(0) &= 1, & \text{if } T_c &= 0 \end{aligned} \quad (21)$$

If scaled in $m(0)$ by defining $u_c = \mu_c/m(0)$ and $t_c = T_c/m(0)$ then the equation (20) of criticality curve of chemical potential and temperature can be rewritten as

$$u_c = 1 + 2t_c \ln \left[(1 + \sqrt{1 - 4e^{-1/t_c}}) / 2 \right]. \quad (22)$$

These results obtained under the assumption $\Lambda/m(0) \gg 1$ are identical to the ones given by effective scalar field approach in the leading order of $1/N$ expansion [5] and also consistent with the calculations of Ref. [4].

By means of the dynamical fermion mass, we may have more clear understanding of the feature of symmetry restoration phase transition. Assuming $\Lambda/m(0) \gg 1$ and in view of $m \leq m(0)$, we will have $\arctan[\Lambda/m(0)] \simeq \arctan[\Lambda/m] \simeq \pi/2$. Thus Eq.(17) will become Λ -independent, i.e.

$$m(0) = m + T \left\{ \ln \left[1 + e^{-(m-\mu)/T} \right] + \ln \left[1 + e^{-(m+\mu)/T} \right] \right\}. \quad (23)$$

When $T \lesssim T_c$, we have $m/T \ll 1$ thus Eq.(23) can be transformed into

$$\begin{aligned} m(0) &= T \left[\ln(1 + e^{\mu/T}) + \ln(1 + e^{-\mu/T}) \right] + T \ln \left[1 + (e^{\mu/T} - 1) + \frac{(e^{\mu/T} - 1)^2}{(1 + e^{\mu/T})(1 + e^{-\mu/T})} \right] - m \\ &\simeq T \left\{ \left[\ln(1 + e^{\mu/T}) + \ln(1 + e^{-\mu/T}) \right] + \frac{m^2}{2T^2[1 + \cosh(\mu/T)]} + \mathcal{O}\left(\frac{m^3}{T^3}\right) \right\}, \end{aligned} \quad (24)$$

where we have made the power series expansion of $[\exp(m/T) - 1]$ in m/T and kept only the terms to the order $\mathcal{O}(m^2/T^2)$. Combining Eq.(24) with Eq.(20) (in which μ_c is replaced by μ) we obtain

$$m^2 = 2T \left(1 + \cosh \frac{\mu}{T} \right) \left\{ T_c \ln \left[2 \left(1 + \cosh \frac{\mu}{T_c} \right) \right] - T \ln \left[2 \left(1 + \cosh \frac{\mu}{T} \right) \right] \right\}. \quad (25)$$

By means of the approximate expression at $T \lesssim T_c$

$$\begin{aligned} \ln \left[2 \left(1 + \cosh \frac{\mu}{T} \right) \right] &= \ln \left[2 \left(1 + \cosh \left\{ \frac{\mu}{T_c} + \frac{\mu(T_c - T)}{T_c T} \right\} \right) \right] \\ &\simeq \ln \left[2 \left(1 + \cosh \frac{\mu}{T_c} \right) \right] + \frac{\mu(T_c - T)}{T_c T} \tanh \frac{\mu}{2T_c} \end{aligned} \quad (26)$$

Eq.(25) may be changed into

$$m^2 = 2T \left(1 + \cosh \frac{\mu}{T} \right) \left\{ \ln \left[2 \left(1 + \cosh \frac{\mu}{T_c} \right) \right] - \frac{\mu}{T_c} \tanh \frac{\mu}{2T_c} \right\} (T_c - T) \quad \text{when } T \lesssim T_c \quad (27)$$

We see that the dynamical fermion mass m , as the only order parameter of the discrete symmetry breaking, will have $(T_c - T)^{1/2}$ behavior at $T \lesssim T_c$. This situation is similar to the one in 4 dimension NJL-model [10]. Hence, based on the same arguments given in Ref. [10], we can conclude that in the 3D Gross-Neveu model the discrete symmetry restoration at $T > T_c$ will also be a second-order phase transition.

In this model, no continuous symmetry breaking means no Goldstone boson appearing. An interesting question is that whether any (massive or massless) scalar bound state could exist. Such scalar bound state can be thought as the equivalent of a "Higgs" particle appearing in spontaneous breaking of a continuous symmetry. In present case, it could be the configuration $\sum_{k=1}^N (\bar{\psi}^k \psi_k)$. However, we will prove that the answer to the above question is negative in 3D Gross-Neveu model. Based on the method taken in Ref.[11] in the real-time formalism of thermal field theory, we can obtain from the scalar four-point function of fermions the formal propagator for the "scalar bound state" expressed by

$$\Gamma_S(p) = -i/N(p^2 - 4m^2 + i\varepsilon) \left[K(p) + H(p) - iS(p) - \frac{R^2(p)}{K(p) + H(p) + iS(p)} \right], \quad (28)$$

where the functions

$$\begin{aligned} K(p) &= \frac{1}{4\pi^2} \int_0^1 \frac{dx}{[m^2 - p^2 x(1-x)]^{1/2}} \arctan \frac{\Lambda}{[m^2 - p^2 x(1-x)]^{1/2}} \\ &\simeq \frac{1}{8\pi\sqrt{p^2}} \ln \frac{4m^2 + 4m\sqrt{p^2} + p^2}{4m^2 - p^2}, \quad \text{if } \arctan \frac{\Lambda}{[m^2 - p^2 x(1-x)]^{1/2}} \approx \frac{\pi}{2} \quad \text{and } 0 < p^2 < 4m^2, \end{aligned}$$

$$\begin{aligned}
H(p) &= 2\pi \int \frac{d^3l}{(2\pi)^3} \left\{ \frac{(l+p)^2 - m^2}{[(l+p)^2 - m^2]^2 + \varepsilon^2} + (p \rightarrow -p) \right\} \delta(l^2 - m^2) \sin^2 \theta(l^0, \mu), \\
S(p) &= 2\pi^2 \int \frac{d^3l}{(2\pi)^3} \delta(l^2 - m^2) \delta[(l+p)^2 - m^2] \left[\sin^2 \theta(l^0, \mu) \cos^2 \theta(l^0 + p^0, \mu) \right. \\
&\quad \left. + \cos^2 \theta(l^0, \mu) \sin^2 \theta(l^0 + p^0, \mu) \right], \\
R(p) &= \pi^2 \int \frac{d^3l}{(2\pi)^3} \delta(l^2 - m^2) \delta[(l+p)^2 - m^2] \sin 2\theta(l^0, \mu) \sin 2\theta(l^0 + p^0, \mu). \quad (29)
\end{aligned}$$

It is indicated that when deriving Eq. (28) we have used Eq. (12). As a result, the Λ -dependent divergent sectors in $\Gamma_S(p)$ have cancelled each other and this is just a peculiarity in the leading order of $1/N$ expansion. Formally $p^2 = 4m^2$ seems to be a mere simple pole of $\Gamma_S(p)$, but this is not true. First of all we notice that when $p^2 \rightarrow 4m^2$,

$$\lim_{p^2 \rightarrow 4m^2} K(p) \rightarrow \infty \quad (30)$$

and $K(p)$ has a logarithmic singularity. Next, we can write

$$\begin{aligned}
H(p) &= \frac{1}{8\pi^2} \int \frac{d^2l}{\omega_l} \left\{ \frac{p^2 + 2\omega_l p^0 - 2|\vec{l}||\vec{p}|\cos\varphi}{[p^2 + 2\omega_l p^0 - 2|\vec{l}||\vec{p}|\cos\varphi]^2 + \varepsilon^2} + \frac{p^2 - 2\omega_l p^0 + 2|\vec{l}||\vec{p}|\cos\varphi}{[p^2 - 2\omega_l p^0 + 2|\vec{l}||\vec{p}|\cos\varphi]^2 + \varepsilon^2} \right. \\
&\quad \left. \cdot \frac{1}{\exp[\beta(\omega_l - \mu)] + 1} + (p^0 \rightarrow -p^0, \mu \rightarrow -\mu) \right\}, \quad \omega_l = (\vec{l}^2 + m^2)^{1/2}. \quad (31)
\end{aligned}$$

The angular integrations in Eq. (31) are of the following general forms and results:

$$\int_0^{2\pi} d\varphi \frac{1}{b + c \cos \varphi \pm \varepsilon^2} = 0, \quad (32)$$

where b and c are constants. Hence the function $H(p) = 0$. As for $S(p)$ and $R(p)$, when $p^2 = 4m^2$ each of them may be generally expressed by

$$\begin{aligned}
A(p)|_{p^2=4m^2} &= \int d^3l \delta(l^2 - m^2) \delta[(l+p)^2 - m^2] f(l^0, p^0, \mu)|_{p^2=4m^2} \\
&= \int_0^\infty \frac{d|\vec{l}||\vec{l}|}{4\omega_l} \int_0^{2\pi} d\varphi \delta(2m^2 - \omega_l p^0 - |\vec{l}||\vec{p}|\cos\varphi) f(-\omega_l, p^0, \mu), \quad (33)
\end{aligned}$$

where $f(l^0, p^0, \mu)$ is a finite function. In Eq.(33), the condition $|\cos\varphi| \leq 1$ limits $|\vec{l}|$ to the only possible value $|\vec{l}| = |\vec{p}|/2$. Let $\tan(\varphi/2) = t$, then the integration of the angle φ in Eq. (33) can be written by

$$\begin{aligned}
B &= \left(\int_0^\pi + \int_\pi^{2\pi} \right) d\varphi \delta(2m^2 - \omega_l p^0 - |\vec{l}||\vec{p}|\cos\varphi) \\
&= 2 \int_0^\infty dt \left\{ \delta[(2m^2 - \omega_l p^0 + |\vec{l}||\vec{p}|)t^2 + (2m^2 - \omega_l p^0 - |\vec{l}||\vec{p}|)] \right. \\
&\quad \left. + \delta[(2m^2 - \omega_l p^0 - |\vec{l}||\vec{p}|)t^2 + (2m^2 - \omega_l p^0 + |\vec{l}||\vec{p}|)] \right\}. \quad (34)
\end{aligned}$$

Since the δ -functions have non-zero values only if $|\vec{l}| = |\vec{p}|/2$, we can set

$$|\vec{l}| = \left(\frac{1}{2} \pm \varepsilon\right) |\vec{p}|, \quad \varepsilon = 0_+ \quad (35)$$

and obtain

$$\begin{aligned} 2m^2 - \omega_l p^0 + |\vec{l}| |\vec{p}| &\simeq \frac{\varepsilon^2 \vec{p}^4}{\vec{p}^2 + 4m^2}, \\ 2m^2 - \omega_l p^0 - |\vec{l}| |\vec{p}| &\simeq -\vec{p}^2 \end{aligned} \quad (36)$$

which further lead to

$$\begin{aligned} B &= 2 \lim_{\varepsilon \rightarrow 0} \delta_{|\vec{l}|, |\vec{p}|/2} \left\{ \int_0^\infty dt \delta \left[\frac{\varepsilon^2 \vec{p}^4}{\vec{p}^2 + 4m^2} t^2 - \vec{p}^2 \right] + \int_0^\infty dt \delta \left[\frac{\varepsilon^2 \vec{p}^4}{\vec{p}^2 + 4m^2} - \vec{p}^2 t^2 \right] \right\} \\ &= 2 \frac{(\vec{p}^2 + 4m^2)^{1/2}}{\vec{p}^2} \lim_{\varepsilon \rightarrow 0} \frac{\delta_{|\vec{l}|, |\vec{p}|/2}}{\varepsilon |\vec{p}|} = 2 \frac{(\vec{p}^2 + 4m^2)^{1/2}}{\vec{p}^2} \lim_{|\vec{l}| \rightarrow |\vec{p}|/2} \frac{\delta_{|\vec{l}|, |\vec{p}|/2}}{||\vec{l}| - |\vec{p}|/2|} \\ &= 2 \frac{(\vec{p}^2 + 4m^2)^{1/2}}{\vec{p}^2} \delta(|\vec{l}| - |\vec{p}|/2). \end{aligned} \quad (37)$$

Substituting Eq. (37) into Eq.(33) we will have

$$A(p)|_{p^2=4m^2} = \frac{1}{2|\vec{p}|} f(-\sqrt{\vec{l}^2 + m^2}, p^0, \mu) \Big|_{|\vec{l}|=|\vec{p}|/2}. \quad (38)$$

This means that both $S(p)|_{p^2=4m^2}$ and $R(p)|_{p^2=4m^2}$ with the form of Eq. (38) are finite. To summarize the above results, we can obtain

$$\lim_{p^2 \rightarrow 4m^2} (p^2 - 4m^2) \Gamma_S(p) = 0, \quad (39)$$

hence $p^2 = 4m^2$ is not a simple pole of $\Gamma_S(p)$ and $\Gamma_S(p)$ does not represent the propagator for any scalar bound state. This proves that no scalar bound state could exist in this model. This result, seen from Eqs. (28) and (30), is true at $T = 0$, and also maintained at a finite temperature. We point out that the above result at $T = 0$ is different from the one obtained in Reference [5].

In conclusion, in 3 dimension Gross-Neveu model, the fermion condensates induced by the scalar four-fermion interactions will spontaneously break the discrete Z_2 , the special parity \mathcal{P}_1 , \mathcal{P}_2 and the time-reversal symmetries and lead to generation of dynamical fermion mass. By means of the real-time thermal field theory in the fermionic bubble diagram approximation, it is clearly shown that the dynamical fermion mass m , as the order parameter of symmetry breakings, will decrease as temperature increases and finally becomes zero at some critical temperature T_c . This implies restorations of these discrete symmetries at $T > T_c$. The equation of the criticality curve of chemical potential and temperature will be independent of the momentum cut-off and identical to one obtained by the auxiliary scalar field approach in the $1/N$ expansion if the ratio between the momentum cut-off and the dynamical fermion mass at $T = 0$ is sufficiently large. Similar to the case in 4D NJL-model, the dynamical mass m has the $(T_c - T)^{1/2}$ behavior at $T \lesssim T_c$ and this fact indicates the second-order phase transition feature of the symmetry restorations. We have also proven by explicit analysis of the fermionic scalar four-point

functions that there is no scalar bound state in this model.

The author is grateful to Prof. Zhao Bao-Heng for helpful discussions. This work was done (in part) in the frame of Associate Membership Programme of the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy and partially supported by National Natural Science Foundation of China and by Grant No. LWZ-1298 of Chinese Academy of Sciences.

References

- [1] D. J. Gross and A. Neveu, *Phys. Rev. D***10** (1974) 3235.
- [2] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, (Clarendon Press, Oxford, 1993).
- [3] K. G. Klimenko, *Z. Phys. C***37** (1988) 457.
- [4] A. Okopińska, *Phys. Rev. D***38** (1988) 2507.
- [5] B. Rosenstein, B. J. Warr and S. H. Park, *Phys. Rep.* **205** (1991) 59.
- [6] V. E. Rochev and P. A. Saponov, hep-th/9710006.
- [7] R. F. Dashen, S-K, Ma and R. Rajaraman, *Phys. Rev. D* **11** (1975) 1499.
- [8] Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* 122 (1961) 345; 124 (1961) 246.
- [9] N. P. Landsman and Ch. G. van Weert, *Phys. Rep.* **145** (1987) 141, and the refereces therein.
- [10] B. R. Zhou, *Phys. Rev. D***57** (1998) 3171; *Commun. Theor. Phys.*, to be published.
- [11] B. R. Zhou, Institution Report, AS-GS-TP-004 (1998).