

D-Branes and Hilbert Schemes

Yu. Malyuta and T. Obikhod

*Institute for Nuclear Research
National Academy of Sciences of Ukraine
252022 Kiev, Ukraine
e-mail: interdep@kinr.kiev.ua*

Abstract

The Nakamura's algorithm is applied to compute the Hilbert scheme for the D-brane model $\frac{1}{13}(1, 2, 10)$.

1 Introduction

The moduli space of a D-Brane localized at the singular point of the Calabi-Yau orbifold $\mathbb{C}^3/\mathbb{Z}_n$ is an interesting object to study, because it describes phases of models [1, 2]

$$\frac{1}{n}(a_1, a_2, a_3),$$

where characters a_1, a_2, a_3 are coprime to n and $\sum_{i=1}^3 a_i = 0 \pmod{n}$. The D-brane moduli space is isomorphic to the Hilbert scheme, which is the fine moduli space of \mathbb{Z}_n -clusters [3, 4, 5]. Nakamura [3] proposes an algorithm to compute Hilbert schemes as toric varieties given by fans.

The purpose of the present paper is to apply the Nakamura's algorithm to compute the Hilbert scheme for the model

$$\frac{1}{13}(1, 2, 10),$$

which has been studied in [4] by another method.

2 Constructions

In this section we give a sketch of Nakamura's constructions [3].

Proposition 1. For every \mathbb{Z}_n -cluster the defining equations can be written as equations in one of the two following forms: either

$$\begin{aligned} x^{a+d+1} &= \lambda y^b z^f & y^{b+1} z^{f+1} &= \mu \nu x^{a+d} \\ y^{b+e+1} &= \mu z^c x^d & z^{c+1} x^{d+1} &= \lambda \nu y^{b+e} & xyz &= \lambda \mu \nu & (\uparrow) \\ z^{c+f+1} &= \nu x^a y^e & x^{a+1} y^{e+1} &= \lambda \mu z^{c+f} \end{aligned}$$

for some $a, b, c, d, e, f \geq 0$; or

$$\begin{aligned}
x^{a+d} &= \beta\gamma y^{b-1} z^{f-1} & y^b z^f &= \alpha x^{a+d-1} \\
y^{b+e} &= \alpha\gamma z^{c-1} x^{d-1} & z^c x^d &= \beta y^{b+e-1} & xyz &= \alpha\beta\gamma & (\downarrow) \\
z^{c+f} &= \alpha\beta x^{a-1} y^{e-1} & x^a y^e &= \gamma z^{c+f-1}
\end{aligned}$$

for some $a, b, c, d, e, f \geq 1$.

Proposition 2. The fan of the Hilbert scheme is given by the cones with the vertexes (divisors)

$$\begin{aligned}
P &= (bc+bf+ef+b+c+e+f+1, ac+cd+df+d, ab+ae+de+a) \\
Q &= (bc+bf+ef+b, ac+cd+df+a+d+c+f+1, ab+ae+de+e) \\
R &= (bc+bf+ef+f, ac+cd+df+c, ab+ae+de+a+b+d+e+1)
\end{aligned}$$

for the equations (\uparrow), and by the cones with the vertexes (divisors)

$$\begin{aligned}
P &= (bc+bf+ef-b-c-e-f+1, ac+cd+df-d, ab+ae+de-a) \\
Q &= (bc+bf+ef-b, ac+cd+df-a-d-c-f+1, ab+ae+de-e) \\
R &= (bc+bf+ef-f, ac+cd+df-b, ab+ae+de-a-b-d-e+1)
\end{aligned}$$

for the equations (\downarrow).

3 Computations

In this section we compute the Hilbert scheme for the model $\frac{1}{13}(1, 2, 10)$.

Figure 1 shows the McKay quiver tessellated by tripods for $\frac{1}{13}(1, 2, 10)$. The corresponding monomial representation of this quiver is illustrated in Figure 2.

Comparing the hexagons of Figure 1 marked by identical numbers and using the corresponding monomials of Figure 2, we infer the following equations for \mathbb{Z}_{13} -clusters:

$$\begin{array}{lll}
x = y^2 z & y^4 = z^2 x & z^2 = xy^3 & (1) \\
x = y^2 z & y^5 = z & z^3 = y^2 & (2) \\
x^2 = y & y^4 = z^2 x & z^3 = y^2 & (3) \\
x^2 = y & y^5 = z & z^2 = xy^3 & (4) \\
x^2 = y & y^2 = z^3 & z^4 = x & (5) \\
x^2 = y & y^7 = x & z = y^5 & (6) \\
x = z^4 & y^2 = z^3 & z^8 = y & (7) \\
x^4 = z^3 & y = x^2 & z^4 = x & (8) \\
x^7 = z^2 & y = x^2 & z^3 = x^4 & (9) \\
x^{10} = z & y = x^2 & z^2 = x^7 & (10) \\
x^{13} = 1 & y = x^2 & z = x^{10} & (11) \\
x = y^7 & y^{13} = 1 & z = y^5 & (12) \\
x = z^4 & y = z^8 & z^{13} = 1 & (13)
\end{array}$$

Note that we write these equations omitting multiplicative constants $\lambda, \mu, \nu, \alpha, \beta, \gamma$. Note also that we require the agreement of our equations with the defining equations of Proposition 1. This requirement gives us the exponents of x^*, y^*, z^* :

$$\begin{array}{ll}
a = 1, & b = 2, & c = 2, & d = 1, & e = 3, & f = 1 & \text{for (1)} \\
a = 0, & b = 2, & c = 1, & d = 0, & e = 2, & f = 1 & \text{for (2)} \\
a = 0, & b = 1, & c = 2, & d = 1, & e = 2, & f = 0 & \text{for (3)} \\
a = 1, & b = 1, & c = 1, & d = 0, & e = 3, & f = 0 & \text{for (4)} \\
a = 1, & b = 1, & c = 3, & d = 0, & e = 0, & f = 0 & \text{for (5)} \\
a = 0, & b = 1, & c = 0, & d = 1, & e = 5, & f = 0 & \text{for (6)} \\
a = 0, & b = 0, & c = 3, & d = 0, & e = 1, & f = 4 & \text{for (7)} \\
a = 1, & b = 0, & c = 0, & d = 2, & e = 0, & f = 3 & \text{for (8)} \\
a = 4, & b = 0, & c = 0, & d = 2, & e = 0, & f = 2 & \text{for (9)} \\
a = 7, & b = 0, & c = 0, & d = 2, & e = 0, & f = 1 & \text{for (10)} \\
a = 10, & b = 0, & c = 0, & d = 2, & e = 0, & f = 0 & \text{for (11)}
\end{array}$$

$$\begin{array}{ll}
a = 0, & b = 7, & c = 0, & d = 0, & e = 5, & f = 0 & \text{for (12)} \\
a = 0, & b = 0, & c = 8, & d = 0, & e = 0, & f = 4 & \text{for (13)}
\end{array}$$

Using these exponents and Proposition 2, we obtain the following cones of the fan:

$$\begin{array}{llll}
P = (2, 4, 7) & Q = (7, 1, 5) & R = (8, 3, 2) & \text{for (1)} \\
P = (13, 0, 0) & Q = (8, 3, 2) & R = (7, 1, 5) & \text{for (2)} \\
P = (8, 3, 2) & Q = (3, 6, 4) & R = (2, 4, 7) & \text{for (3)} \\
P = (7, 1, 5) & Q = (2, 4, 7) & R = (1, 2, 10) & \text{for (4)} \\
P = (8, 3, 2) & Q = (4, 8, 1) & R = (3, 6, 4) & \text{for (5)} \\
P = (7, 1, 5) & Q = (1, 2, 10) & R = (0, 0, 13) & \text{for (6)} \\
P = (13, 0, 0) & Q = (4, 8, 1) & R = (8, 3, 2) & \text{for (7)} \\
P = (4, 8, 1) & Q = (0, 13, 0) & R = (3, 6, 4) & \text{for (8)} \\
P = (3, 6, 4) & Q = (0, 13, 0) & R = (2, 4, 7) & \text{for (9)} \\
P = (2, 4, 7) & Q = (0, 13, 0) & R = (1, 2, 10) & \text{for (10)} \\
P = (1, 2, 10) & Q = (0, 13, 0) & R = (0, 0, 13) & \text{for (11)} \\
P = (13, 0, 0) & Q = (7, 1, 5) & R = (0, 0, 13) & \text{for (12)} \\
P = (13, 0, 0) & Q = (0, 13, 0) & R = (4, 8, 1) & \text{for (13)}
\end{array}$$

The Hilbert scheme for $\frac{1}{13}(1, 2, 10)$ is illustrated in Figure 3.

Acknowledgement

The authors would like to thank Professor I. Nakamura for sending his preprint.




Figure 1




Figure 2




Figure 3

References

- [1] M.R. Douglas, B.R. Greene and D.R. Morrison, *Orbifold Resolution by D-Branes*, hep-th/9704151.
- [2] K. Mohri, *Kähler Moduli Space for a D-Brane at Orbifold Singularities*, hep-th/9806052.
- [3] I. Nakamura, *Hilbert Schemes of Abelian Group Orbits*, preprint 1998.
- [4] M. Reid, *McKay Correspondence*, alg-geom/9702016.
- [5] Y. Ito and H. Nakajima, *McKay Correspondence and Hilbert Schemes in Dimension Three*, math.AG/9803120.