

Comment on Two-Loop Free Energy in $N = 4$ Supersymmetric Yang-Mills Theory at Finite Temperature[?]

A. Fotopoulos and T. R. Taylor

Department of Physics, Northeastern University, Boston, MA 02115, U.S.A.

Abstract

The strong coupling behavior of finite temperature free energy in $N = 4$ supersymmetric $SU(N)$ Yang-Mills theory has been recently discussed by Gubser, Klebanov and Tseytlin in the context of AdS-SYM correspondence. In this note, we focus on the weak coupling behavior. As a result of a two-loop computation we obtain, in the large N 't Hooft limit, $F(g^2N \neq 0) = \frac{2}{6}N^2V_3T^4(1 - \frac{3}{2}g^2N)$. Comparison with the strong coupling expansion provides further indication that free energy is a smooth monotonic function of the coupling constant.

[?]Research supported in part by the National Science Foundation under grant PHY {96{02074.

Finite temperature effects break supersymmetry [1]. By switching on non-zero temperature one can interpolate between supersymmetric and non-supersymmetric theories. For instance in gauge theories, one can interpolate between the supersymmetric case and a theory which contains pure Yang-Mills (YM) as the massless sector, with some additional thermal excitations. In the infinite temperature limit, the time dimension decouples and, at least formally, one obtains a non-supersymmetric Euclidean gauge theory. If no phase transition occurs when the YM gas is heated up, then the dynamics of realistic gauge theories such as QCD is smoothly connected to their supersymmetric relatives.

Maldacena conjecture [2] which relates the large N limit of $N = 4$ supersymmetric $SU(N)$ Yang-Mills theory (SYM) to type IIB superstrings propagating on $AdS_5 \times S^5$, provides a very promising starting point towards QCD. On the superstring side, non-zero temperature can be simulated by including Schwarzschild black holes embedded in AdS geometry [3]. The classical geometry of black holes with Hawking temperature T does indeed encode correctly many qualitative features of large N gauge theory heated up to the same temperature. At the quantitative level though, the comparison between SYM and supergravity becomes rather subtle because the supergravity side merely provides the strong coupling expansion for physical quantities while most of finite temperature computations in SYM are limited to the perturbative, weak coupling expansion. In this note, we comment on the computation of free energy.

The AdS-SYM correspondence has been recently employed for computing the two leading terms in the strong coupling expansion [4] (valid in the $g^2 N \gg 1$ limit):

$$F(\infty) = \frac{2}{6} N^2 V_3 T^4 \left[\frac{3}{4} + \frac{45}{32} (3/2)^{3-2} \right]; \quad (1)$$

The comparison with the limiting free-theory value,

$$F(=0) = \frac{2}{6} N^2 V_3 T^4; \quad (2)$$

indicates that the exact answer has the form :

$$F(\tau) = \frac{2}{6} N^2 V_3 T^4 f(\tau); \tag{3}$$

where the function $f(\tau)$ interpolates smoothly between the asymptotic values $f(0) = 1$ and $f(1) = 3/4$. The sign of the subleading correction $O[(2\tau)^{-3=2}]$ in eq.(1) indicates that f decreases monotonically from 1 to 3/4.

The question whether free energy interpolates smoothly between weak and strong coupling limits deserves careful investigation, especially in view of the recent claim in favor of a phase transition at finite τ [5].¹ There is, however, a place to look for further hints on the properties of free energy: the subleading terms in the weak coupling expansion. Surprisingly enough, they cannot be found in the existing literature. In order to fill this gap, we calculated the two-loop correction to free energy. The result is:

$$F(\tau \neq 0) = \frac{2}{6} N^2 V_3 T^4 \left[1 - \frac{3}{2\tau^2} \right]; \tag{4}$$

The (relative) negative sign of the two-loop correction provides further indication that the free energy is a smooth, monotonic function of the 't Hooft coupling τ . In the remaining part of this note we present some details of the two-loop computation leading to eq.(4).

For the purpose of diagrammatic computations, it is convenient to use the $N = 1$ decomposition of $N = 4$ SYM, with one Majorana fermion corresponding to the gaugino, and the three remaining Majorana fermions combined with scalars in three $N = 1$ chiral multiplets. The two-loop diagrams are displayed in Figure 1, together with the combinatorial/statistics factors. The two-loop integrals can be readily performed by using techniques described in

¹ It is beyond the scope of this note to review the arguments of ref.[5], however we would like to point out that they involve certain assumptions on the convergence properties of perturbative expansions. The proposed 2^{-2} convergence radius does not seem realistic after one looks at the two-loop correction, see eq.(4).




Figure 1: Two-loop diagrams contributing to free energy. Gauge bosons are represented by wiggly lines, ghosts by dotted lines, fermionic (Majorana) components of chiral multiplets by dashed lines, scalars by solid lines and $N = 1$ gauginos by double-dashed lines.

refs.[6]. In the table below, we list results for individual diagrams²

A	B	C	D	E	F	G	H	I	J
3	$\frac{9}{4}$	$\frac{1}{4}$	$\frac{3}{4}$	$\frac{1}{4}$	$\frac{9}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	12	$\frac{15}{2}$

(5)

where

$$= g^2 c_A dV_3 \frac{T^4}{144} + \frac{T^2}{12(2)^3} \int \frac{d^3k}{k^j} ; \quad (6)$$

$$= g^2 c_A dV_3 \frac{5T^4}{144} - \frac{T^2}{3(2)^3} \int \frac{d^3k}{k^j} ; \quad (7)$$

with d denoting the dimension of the gauge group and c_A the Casimir operator in the adjoint representation. Note that individual diagrams contain ultraviolet divergences. After combining all contributions, we obtain the (finite) result:

$$F_{2-loop} = 16 + 4 = \frac{1}{4} g^2 c_A dV_3 T^4 ; \quad (8)$$

Specified to the case of $SU(N)$, with $d = N^2 - 1$ and $c_A = N$, in the leading large N order the above result yields eq.(4).

²Diagrams are computed in the Feynman gauge.

Acknowledgments

Most of this work was done while the authors were visiting Laboratoire de Physique Théorique et Hautes Energies at Université Paris-Sud, Orsay. We are grateful to Pierre Binétruy and all members of LPTHE for their kind hospitality.

References

- [1] L. Girardello, M. T. Grisaru and P. Salomonson, Nucl. Phys. B 178 (1981) 331.
- [2] J. Maldacena, hep-th/9711200.
- [3] E. Witten, hep-th/9803131.
- [4] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, hep-th/9805156.
- [5] Miao Li, hep-th/9807196
- [6] J. I. Kapusta, "Finite Temperature Field Theory," Cambridge University Press (1989); Nucl. Phys. B 148 (1979) 461.