# Generalised quantum anharmonic oscillator using an operator ordering approach

Anirban Pathak Department of Physics, Visva-Bharati, Santiniketan-731235, India

#### Abstract

We construct a generalised expression for the normal ordering of  $(a + a^{\dagger})^m$  for integral values of m and use the result to study the quantum anharmonic oscillator problem in the Heisenberg approach. In particular, we derive generalised expressions for energy eigen values and frequency shifts for the Hamiltonian  $H = \frac{x^2}{2} + \frac{\dot{x}^2}{2} + \frac{\lambda}{m} x^m$ . We also derive a closed form first order multi-scale perturbation theoretic operator solution of this Hamiltonian with a view to generalise some recent results of Bender and Bettencourt .

PACS numbers: 03.65.Ge, 02.30.Mv, 11.15.Bt, 11.15.Tk

#### 1 Introduction

The simple harmonic oscillator (SHO) model can be used to understand a wide variety of physical phenomena ranging from problems in Newtonian mechanics to those in quantum field theory. This solvable model describes small oscillation of a system about the mean position of equilibrium under the action of a linear restoring force. However, for a real physical problem one has to incorporate the anharmonicity in the model Hamiltonian. In this case the equation of motion is not exactly solvable and, in-fact, the anharmonic oscillators (AHO) provide one of the simplest examples of quantum mechanical systems which can not be solved without making use of approximation techniques. This was why AHO problem got considerable attention of the physicists and mathematicians from the very beginning of the subject. It is an outstanding curiosity to note that studies in the AHO problems have enriched the subject of large order perturbation theory [1-5], divergent expansion of quantum mechanics [6-7], Laplace transformation representation of energy eigen values [8-12], computational physics, Pade and Borel summation of perturbation series [13]. In the present paper we shall study the generalised quantum anharmonic oscillator problem in the Heisenberg representation by making use of two simple normal ordering theorems derived by us for the expansion of  $(a+a^{\dagger})^m$ . We shall see that the merit of the present paper is its simplicity. For example, the present method provides, on the one hand, a straight forward mathematical frame work to construct expressions for the energy eigen values as well as frequency shifts and, on the other hand, generalises some of the recent results on the topic by Bender and Bettencourt [14-15].

The Hamiltonian of a generalised anharmonic oscillator having unit mass and unit frequency is written as

$$H = H_0 + \frac{\lambda}{m} x^m$$
  
=  $\frac{x^2}{2} + \frac{\dot{x}^2}{2} + \frac{\lambda}{m} x^m$  (1)

 $\operatorname{with}$ 

$$x = \frac{1}{\sqrt{2}}(a^{\dagger} + a) \ . \tag{2}$$

Here we have chosen to work in units in which  $\hbar = 1$ . Equation of motion corresponding to (1) is

$$\ddot{x} + x + \lambda x^{m-1} = 0. \tag{3}$$

Equation (3) can not be solved exactly for m > 2. However, large number of approximation methods are available for solving (3) for particular values of m. These include perturbation technique [16], variation of parameters and Taylor series approach [17]. Ordinary perturbation technique leads to the unwanted secular terms that grows up rapidly with time. Secular terms are unwanted because they are in conflict with the physical requirement that the solution be finite. There are some methods which are successfully used to sum up the secular terms for all orders. Multiscale perturbation theory (MSPT) is one of those methods [14-16].

Anharmonic oscillator problem can be approached in two different ways in quantum mechanics. One way is the c-number approach or the Schroedinger approach and the other way is the operator approach or the Heisenberg approach. There is an extensive amount of literature on c-number approach [1-13,18] but the situation is different for the Heisenberg approach because the complicated operator algebra tends to pose serious mathematical problems. Our main objective in this work is to provide a theoretical frame work which is free from calculational difficulties. The first operator solution of a quantum oscillator was given by Bender and Bettencourt [14,15] using MSPT. They generalised the existing theory of MSPT into an operator approach to get the zeroth order solution involving a quantum operator analog of the classical first order frequency shift and interpreted it as an operator mass renormalization that expresses the first order shift of all energy levels. Somewhat later Mandal [19] approached the problem (up to first order) using Taylor series method. All these solutions were found for the quartic oscillator only. Application of the Taylor series method to sextic and octic oscillator problems are now in order [20].

The Schroedinger approach to AHO problem in quantum mechanics has made a significant contribution for the development of the perturbative techniques as we use today but the suitability of the other approach (Heisenberg approach) has not yet been studied in detail. Till now we don't have any operator solutions of higher anharmonic oscillators although these might be relevant to various physical problems. Keeping these in mind, we will suggest a novel method in which MSPT results for anharmonic oscillators in general can be found. To that end we prove in section 2 two theorems to construct a normal ordered expansion of  $(a + a^{\dagger})^m$ . We obtain a generalised expression for the energy eigen values in section 3. We devote section 4 to construct a generalised solution for the equation of motion (3) and specialise our result to reproduce some of the existing results as useful checks on the generalised solution obtained by us.

## 2 Operator Ordering theorems

On a very general ground one knows that in quantum mechanics and quantum field theory proper ordering of the operators plays a crucial role. Let  $f(a, a^{\dagger})$  be an arbitrary operator function of the usual bosonic annihilation and creation operators a and  $a^{\dagger}$  which satisfy the commutation relation

$$[a, a^{\dagger}] = 1. \tag{4}$$

One can write  $f(a, a^{\dagger})$  in such a way that all powers of  $a^{\dagger}$  always appear to the left of all powers of a. Then  $f(a, a^{\dagger})$  is said to be normal ordered. In this paper we want to write  $(a + a^{\dagger})^m$  in the normal ordered form for integral values of m. Traditionally for a given value of m this is achieved by using a very lengthy procedure which involves repeated application of (4). One of our objectives in this work is to construct a normal ordered expansion of  $(a+a^{\dagger})^m$  without taking recourse to such repeated applications.

We shall denote normal ordered form of f by  $f_N$ . On the other hand : f : will denote an operator obtained from f by arranging all powers of  $a^{\dagger}$  to the left of all powers of a without making use of the commutation relation in (4). Now if  $f = aa^{\dagger}$  then  $f_N = a^{\dagger}a + 1$  and :  $f := a^{\dagger}a$ . Therefore, we can write

$$: (a+a^{\dagger})^{m} :=: (a^{\dagger}+a)^{m} := a^{m} + {}^{m}C_{1}a^{\dagger}a^{m-1} + \dots + {}^{m}C_{r}a^{\dagger^{r}}a^{m-r} + \dots + a^{\dagger^{m}}.$$
 (5)

Thus in this notation :  $(a + a^{\dagger})^m$  : is simply a binomial expansion in which powers of the  $a^{\dagger}$  are always kept to the left of the powers of the a. To write  $(a^{\dagger} + a)_N^m$  we shall proceed by using the following theorems.

#### 2.1 Theorem 1.

$$: (a^{\dagger} + a)^{m} : (a^{\dagger} + a) =: (a^{\dagger} + a)^{m+1} : +m : (a^{\dagger} + a)^{m-1} : .$$
 (6)

Proof: From (5) :  $(a^{\dagger} + a)^m$  :  $(a^{\dagger} + a)$  can be written in the form

$$: (a^{\dagger} + a)^{m} : (a^{\dagger} + a) = \left[ a^{\dagger^{m+1}} + ({}^{m}C_{1} + {}^{m}C_{0})a^{\dagger^{m}}a + ({}^{m}C_{2} + {}^{m}C_{1})a^{\dagger^{m-1}}a^{2} + \dots + ({}^{m}C_{r} + {}^{m}C_{r-1})a^{\dagger^{m-r+1}}a^{r} + \dots \right] + ({}^{m}C_{1}a^{\dagger^{m-1}} + 2{}^{m}C_{2}a^{\dagger^{m-2}}a + \dots + r{}^{m}C_{r}a^{\dagger^{m-r}}a^{r-1} + \dots).$$
(7)

The theorem in (6) can be obtained by simply using the following identities in the above

$$\begin{array}{rcl}
(a^{r}a^{\dagger})_{N} &=& a^{\dagger}a^{r} + ra^{r-1}, \\
r^{n}C_{r} &=& n^{n-1}C_{r-1}, \\
and & {}^{m+1}C_{r+1} &=& ({}^{m}C_{r} + {}^{m}C_{r+1}).
\end{array}$$
(8)

Note that first identity in (8) can be proved with the help of the general operator ordering theorems [21] while the other two are trivial.

#### 2.2 Theorem 2 :

For any integral values of m

$$(a^{\dagger} + a)_N^m = \sum_{r=0,2,4..}^m t_r \,{}^m C_r : \, (a^{\dagger} + a)^{m-r} : \tag{9}$$

with

$$t_r = \frac{(r-1)!}{2^{(\frac{r}{2}-1)}(\frac{r}{2}-1)!} \text{ for } r \ge 4$$
(10)

 $\quad \text{and} \quad$ 

$$t_0 = t_2 = 1. (11)$$

Proof:

Using theorem 1, we can write

$$\begin{array}{rcl} (a^{\dagger}+a)_{N} & = & : (a^{\dagger}+a): \\ (a^{\dagger}+a)_{N}^{2} & = & : (a^{\dagger}+a)^{2}:+1 \\ (a^{\dagger}+a)_{N}^{3} & = & : (a^{\dagger}+a)^{3}:+3: (a^{\dagger}+a): \\ (a^{\dagger}+a)_{N}^{4} & = & : (a^{\dagger}+a)^{4}:+6: (a^{\dagger}+a)^{2}:+3 \\ (a^{\dagger}+a)_{N}^{5} & = & : (a^{\dagger}+a)^{5}:+10: (a^{\dagger}+a)^{3}:+15: (a^{\dagger}+a): \\ & \cdots & & \cdots & & \cdots \\ & \cdots & & \cdots & & \cdots \\ (a^{\dagger}+a)_{N}^{9} & = & : (a^{\dagger}+a)^{9}:+36: (a^{\dagger}+a)^{7}:+378: (a^{\dagger}+a)^{5}: \\ & + & 1260: (a^{\dagger}+a)^{3}:+945: (a^{\dagger}+a): \end{array}$$

From (12) we venture to identify the general form of the above expansion for a given value of m as

$$(a^{\dagger} + a)_N^m = \sum_{r=0,2,4..}^m t_r \, {}^m C_r : \, (a^{\dagger} + a)^{m-r} : \ . \tag{13}$$

It is easy to check that (13) gives all the expansions of (12) such that (13) is true for m = 1, 2, ..., 9. The general validity of (13) can be ensured by using the method of induction. This establishes that (13) gives the normal order expansion of  $(a^{\dagger} + a)^m$  for any arbitrary integer m. We shall now use this normal ordered expansion to study the anharmonic oscillator problem.

## 3 Energy eigen values

The first order energy eigen value  $E_1$  is  $\langle n|H|n \rangle$ , where  $|n\rangle$  is the number state. In the literature there are two lengthy procedures [22] to obtain  $E_1$ . The first one is the usual normal ordering method. This method involves iterative use of (4) and number of iteration increases very first as m increases. For a given value of m we need  $[2^m - (m+1)]$  iterations. This shows that for large mthe construction of the expansion  $(a^{\dagger} + a)_N^m$  becomes formidable. In the second procedure one proceeds by making repeated application of x operator on the number state. As in the normal ordering method this procedure is also equally lengthy. In the following we implement theorem 2 to derive an uncomplicated method to construct an expansion for  $E_1$  for the Hamiltonian (1).

Using the result in (9) we can write the expression for  $E_1$  for any integer value of m in the form

$$E_{1} = (n + \frac{1}{2}) + \frac{\lambda}{2^{\frac{m}{2}}m} < n | \sum_{r=0,2,4..}^{m} t_{r} \, {}^{m}C_{r} : (a^{\dagger} + a)^{m-r} : |n >$$

$$= (n + \frac{1}{2}) + \frac{\lambda}{2^{\frac{m}{2}}m} < n | \sum_{r=0,2,4..}^{m} t_{r} \, {}^{m}C_{r} \, {}^{m-r}C_{\frac{m-r}{2}} a^{\dagger \frac{m-r}{2}} a^{\frac{m-r}{2}} |n >$$

$$= (n + \frac{1}{2}) + \frac{\lambda}{2^{\frac{m}{2}}m} \sum_{r=0,2,4..}^{m} t_{r} \, {}^{m}C_{r} \, {}^{m-r}C_{\frac{m-r}{2}} \, {}^{n}C_{\frac{m-r}{2}} (\frac{m-r}{2})!$$

$$(14)$$

The expression (14) involves summations which are easy to evaluate and thereby avoids the difficulties associated with earlier iterative procedures. Although somewhat forced, it may be tempting to compare the simplicity sought in our approach with the use of logarithm in a numerical calculation or use of integral transforms in solving a partial differential equation. It is of interest to note that expression similar to (14) can also be constructed for the first order energy eigen value of a Hamiltonian in which the anharmonic term is a polynomial in x. This type of Hamiltonians are very frequently used in nonlinear optics.

# 4 MSPT solution of the generalised quantum anharmonic oscillator.

We want to obtain MSPT operator solution of equation (3) for arbitrary integral values of m. The essential idea behind our approach is that the quantum operator analog of the classical first order frequency shift is an operator function ( $\Omega(H_0)$ ) of the unperturbed Hamiltonian  $H_0$  and secondly a correct solution should reproduce the first order energy spectrum. Now the general form of the zeroth order solution of (3) is given by

$$x_0(t) = \frac{1}{G(n)} [x(0)cos(t + \lambda\Omega(H_0)t) + cos(t + \lambda\Omega(H_0)t)x(0) + \dot{x}(0)sin(t + \lambda\Omega(H_0)t) + sin(t + \lambda\Omega(H_0)t)\dot{x}(0)]$$
(15)

where G(n) is a normalization factor. So our task is to find out  $\Omega(H_0)$  and G(n) in general. From equation (14) we obtain the energy difference for two consecutive energy levels as

$$\begin{aligned}
\omega_{n,n-1} &= (E_1)_{m,n} - (E_1)_{m,n-1} \\
&= 1 + \lambda \omega(m,n) \\
&= 1 + \frac{\lambda}{2^{\frac{m}{2}}m} \sum_{r=0,2,4...}^{(m-2)} t_r^{m} C_r^{m-r} C_{\frac{m-r}{2}}^{n-1} C_{\frac{m-r-2}{2}} \left(\frac{m-r}{2}\right)!.
\end{aligned}$$
(16)

Since the correct quantum operator solution has to give the first order energy spectrum, we should have

$$< n-1|x_0(t)|n> = < n-1|x_0(0)|n> \cos[t+\lambda\omega(m,n)t] + < n-1|p_0(0)|n> \sin[t+\lambda\omega(m,n)t] + < (17)$$

Equation (15) and (17) impose restrictions on our unknown functions  $\Omega(H_0)$ and G(n). The condition imposed on  $\Omega(H_0)$  is

$$< n|\Omega(H_0)|n> + < n-1|\Omega(H_0)|n-1> = 2\omega(m,n)$$
 (18)

or,

$$\Omega(n+\frac{1}{2}) + \Omega(n-\frac{1}{2}) = 2\omega(m,n) .$$
(19)

For a particular *m* right hand side is a known polynomial in *n* and our job is simply to find out  $\Omega(n + \frac{1}{2})$ . We obtain this as

$$\Omega(n+\frac{1}{2}) = 2\left[\sum_{k=0}^{n} (-1)^{n-k} \omega(m,k)\right] + (-1)^{n+\frac{m}{2}} \frac{t_m}{2^{\frac{m-2}{2}}m}.$$
 (20)

Substituting the functional form of  $\Omega(n+\frac{1}{2})$  or  $\Omega(H_0)$  in (15) if we impose condition (17) we will get

$$G(n) = 2\cos\left[\frac{\lambda t}{2}\left(\Omega(n+\frac{1}{2}) - \Omega(n-\frac{1}{2})\right)\right].$$
(21)

The results in (20) and (21) when substituted in (15) solve the generalised quantum anharmonic oscillator problem.

# 4.1 Specific results and their comparison with the existing spectra:

Although we have solved the quantum anharmonic oscillator in general, it is not possible to compare our solution directly with other existing results since the present study happens to be the first operator solution of the generalised anharmonic oscillator. There are methods to calculate first order classical frequency shift for particular m in the appropriate classical limit  $(x(0) = a, \dot{x}(0) = 0)$ . Here we calculate some specific results from our general expressions and compare them with the existing results.

For m = 4 we have,

$$\Omega(n+\frac{1}{2}) = \frac{3n}{4} + \frac{3}{8} = \frac{3}{4}(n+\frac{1}{2}).$$
(22)

Therefore,

$$\Omega(H_0) = \frac{3}{4}H_0$$
  
and  $G(n) = 2\cos(\frac{3\lambda t}{8}).$  (23)

,

,

In terms of (23) the total solution for the quantum quartic anharmonic oscillator is

$$\begin{aligned} x(t)|_{m=4} &= \frac{1}{2\cos(\frac{3\lambda t}{8})} \left[ x(0)\cos[t + \frac{3\lambda t}{4}H_0] + \cos[t + \frac{3\lambda t}{4}H_0] x(0) \right. \\ &+ \dot{x}(0)\sin[t + \frac{3\lambda t}{4}H_0] + \sin[t + \frac{3\lambda t}{4}H_0] \dot{x}(0) \right] \end{aligned}$$
(24)

This exactly coincides with the solutions given in [14,15 and 19] and gives the correct classical frequency shift [23] in the limit x(0) = a,  $\dot{x}(0) = 0$ . Similarly we have

$$\begin{aligned} x(t)|_{m=6} &= \frac{1}{2\cos(\frac{5\lambda t}{4}n)} \left[ x(0)\cos[t + \frac{5\lambda}{4}(H_0^2 + \frac{1}{4})t] + \cos[t + \frac{5\lambda}{4}(H_0^2 + \frac{1}{4})t]x(0) \right. \\ &+ \dot{x}(0)\sin[t + \frac{5\lambda}{4}(H_0^2 + \frac{1}{4})t] + \sin[t + \frac{5\lambda}{4}(H_0^2 + \frac{1}{4})t]\dot{x}(0) \right] \end{aligned}$$

$$(25)$$

$$\begin{aligned} x(t)|_{m=8} &= \frac{1}{2cos[\frac{35\lambda}{64}(6n^2+3)t]} \left[ x(0)cos[t+\frac{35\lambda}{64}(4H_0^3+5H_0)t] + cos[t+\frac{35\lambda}{64}(4H_0^3+5H_0)t]x(0) \right. \\ &+ \dot{x}(0)sin[t+\frac{35\lambda}{64}(4H_0^3+5H_0)t] + sin[t+\frac{35\lambda}{64}(4H_0^3+5H_0)t]\dot{x}(0) \right] \end{aligned}$$

 $\operatorname{and}$ 

$$\begin{aligned} x(t)|_{m=10} &= \frac{1}{2\cos\left[\frac{63\lambda t}{8}(n^3+2n)\right]} \left[ x(0)\cos\left[t + \frac{63\lambda}{16}(H_0^4 + \frac{7}{2}H_0^2 + \frac{9}{16})t\right] \\ &+ \cos\left[t + \frac{63\lambda}{16}(H_0^4 + \frac{7}{2}H_0^2 + \frac{9}{16})t\right] x(0) + \dot{x}(0)\sin\left[t + \frac{63\lambda}{16}(H_0^4 + \frac{7}{2}H_0^2 + \frac{9}{16})t\right] \\ &+ \sin\left[t + \frac{35\lambda}{64}\frac{63\lambda}{16}(H_0^4 + \frac{7}{2}H_0^2 + \frac{9}{16})t\right] \dot{x}(0) \right] . \end{aligned}$$

$$(27)$$

The solutions for sextic and octic oscillator exactly coincides with the solution obtained by us using Taylor series approach [20] and all the solutions give the correct classical frequency shifts in the appropriate limit [24,25].

## 5 Summary and concluding remarks

We conclude by noting that depending on the nature of nonlinearity in a physical problem the treatment of higher anharmonic oscillators assumes significance. But studies in such oscillators (for m > 4) are not undertaken in the Heisenberg approach presumably because the existing methods tends to introduce inordinate mathematical complications in a detailed study. In the present work we contemplate to circumvent them by proving a theorem for the expansion of  $(a^{\dagger} + a)_N^m$ . Thus the results of the present work are expected to serve a useful purpose for physicists working in nonlinear mechanics, molecular physics, quantum optics and quantum field theory.

Acknowledgments: The author is thankful to the CSIR for the award of a Junior Research Fellowship. He is also grateful to Prof B K Talukdar and Dr S Mandal for their kind interest in this work.

#### References

- 1. Bender C M and Wu T T, Phys Rev., 184 (1969) 1231
- 2. Bender C M and Wu T T, Phys Rev. Let , 27 (1971) 461
- 3. Bender C M and Wu T T, Phys Rev. D, 7(1973) 1620
- 4. Arteca G A, Fernandez F M and Castro E A, (1990) Large Order Perturbation Theory and Summation Methods in Quantum Mechanics (Berlin: Springer-Verlag).
- Lowdin P O (Ed.) Int. J. Quant. Chem (1982) , (Proceedings of the Sanibel Workshop on Perturbation Theory at large order, Sanibel Conference, Florida, 1981)
- 6. Simon B, Int. J. Quant. Chem. 21 (1982) 3
- 7. Graffi S, Greechi V and Simon B, Phys Lett. 32B (1970) 631.
- Ivanov I A, J. Phys A **31** (1998) 5697; Ivanov I A, J. Phys A **31** (1998) 6995.
- 9. Weniger E J, Annals of Physics 246 (1996) 133.
- 10. Weniger E J, Cizek J and Vinette F, J. Math. Phys. 39 (1993) 571.
- 11. Vienette F and Cizek J, J. Math. Phys. 32 (1991) 3392
- 12. Fernandez F and Cizek J, Phys. Lett. 166A (1992) 173.
- 13. Graffi S, Greechi V and Turchetti G, IL Nuovo Cimento 4B (1971) 313
- 14. Bender C M, Bettencourt L M A, Phys. Rev. Lett. 77 (1996) 4114.

- 15. Bender C M, Bettencourt L M A, Phys. Rev. D 54 (1996) 1710.
- Nayfeh A H (1981) Introduction to Perturbation Techniques (Wiely, New York).
- 17. Margenau H and Murphy G M (1956) The Mathematics of Physics and Chemistry (D. Van Nostrand Company, Princeton).
- Bender C M, J. Math. Phys. 11 (1970) 796; Biswas S N etal J. Math. Phys. 14 (1973) 1190; Killingbeck J, Rep. Prog. Phys. 40 (1977) 963; ; Zamatsila J, Cizek J and Skala L Annals of Physics 276 (1998) 39.
- 19. Mandal S (1998), J.Phys. A 31 L501.
- 20. Pathak A and Mandal S, Communicated for publication.
- 21. Louisell W H (1973) Quantum Statistical Properties of Radiation ( John Wiley and Sons, New York).
- 22. Powel J L and Crasemann B (1961) *Quantum Mechanics* (Addision-Wesley, Massachusetts).
- 23. Bhaumik K and Dutta-Ray B (1975) J. Math. Phys. 16 1131.
- 24. Dutt R, Lakshmanan M, J. Math. Phys. 17 (1976) 482.
- 25. Bradbury T C and Brintzenhoff A (1971) J. Math. Phys. 12 1269.