Quantum Dynamical Entropies for Classical Stochastic Systems

Mark Fannes and Bart Haegeman 1

Instituut voor Theoretische Fysica K.U. Leuven, B-3001 Leuven, Belgium

Abstract

We compare two proposals for the dynamical entropy of quantum deterministic systems (CNT and AFL) by studying their extensions to classical stochastic systems. We show that the natural measurement procedure leads to a simple explicit expression for the stochastic dynamical entropy with a clear informationtheoretical interpretation. Finally, we compare our construction with other recent proposals.

1 Motivation

Dynamical entropy is a standard tool for the study of classical deterministic systems, see, e.g., [Wal]. It measures the marginal amount of uncertainty generated by the dynamics, or, equivalently, the marginal amount of information obtained about the initial condition. Different approaches have been followed to generalize the idea of dynamical entropy to quantum systems, [AliFan1, AliFan2, Acc et al., Con et al., Hud]. Thereby, one encounters, apart from the non-commutativity, still another problem : Not only the dynamics can generate uncertainty, but also the quantum measurements can do so. A good calculation scheme should separate these two contributions.

The latter problem is also present when extending the notion of dynamical entropy to classical stochastic systems. In this case, the different sources of entropy production to separate are the system dynamics and the stochasticity due to the coupling to the unobserved environment. Therefore, from the dynamical entropy point of view, classical stochastic dynamics can be considered as an intermediate case between classical deterministic and quantum dynamics, [Mak].

We will make this link even more explicit by taking two established quantum constructions (CNT [Con et al.] and AFL [AliFan1, AliFan2]) as a starting point and extending them to stochastic systems. As a consequence, next to system and environment, so-called unsharp measurements appear as a third source of dynamical entropy.

¹Aspirant FWO-Vlaanderen

Interestingly, the different quantum constructions lead to clear-cut differences which we can interpret in terms of these three sources. E.g., for the extreme situation of a Bernoulli process, where the stationary completely random state is already reached after a single time step, successive observations of the system do not reveal any information at all on the initial state of our system. The degree of stochasticity of the dynamics can be measured however and, moreover, very unsharp observations of such a process will overestimate this randomness.

Quantum dynamical entropy has recently received new interest in connection with quantum information theory. Both dynamical entropies we will discuss in this paper have been reformulated in this framework, CNT in [Ben] and AFL in [Ali]. The stochasticity we introduce can both model badly isolated information sources or noisy communication channels. Finally, this work can also be considered as a first step in the construction of a dynamical entropy for quantum stochastic systems, as was recently done in [Kos et al.].

2 Preliminaries

Deterministic classic dynamics are given by a transformation T of the phase space X. For stochastic systems one should use stochastic transformations of phase space. It is more convenient to work on the level of the observables, i.e., the functions on the phase space. Such a description allows also to connect with the quantum world by allowing the algebra of observables to become non-commutative. The dynamics is now given by a transformation Θ of a function space on the phase space X. The different concepts needed to introduce dynamical entropy, like partitions, their evolutions and refinements, must be transported from the level of the points of phase space to the level of observables.

2.1 Some notation

Let μ be a probability measure on the set X. Consider a transformation Θ of the algebra of observables, $\Theta: L^{\infty}(X, \mu) \to L^{\infty}(X, \mu)$, which is

- positive, $f \ge 0 \Rightarrow \Theta(f) \ge 0$, for all $f \in L^{\infty}(X, \mu)$,
- unital, $\Theta(\mathbf{1}) = \mathbf{1}$ and
- measure-preserving, $\mu(\Theta(f)) = \mu(f)$, for all $f \in L^{\infty}(X, \mu)$.

The triple (X, μ, Θ) defines a stochastic dynamical system in discrete time.

Example 2.1 (Markov process) Let X be a finite set and $\mu = \{\mu_x | x \in X\}$ a probability measure on X. Let P be a transition matrix satisfying

$$P_{xy} \ge 0, \quad \sum_{y} P_{xy} = 1 \quad \text{and} \quad \sum_{x} \mu_x P_{xy} = \mu_y.$$

The time evolution Θ , given by

$$\Theta(f)(x) := \sum_{y} P_{xy} f(y),$$

defines a stochastic dynamical system. This finite-dimensional example can be generalized considerably. Let (X, S, μ) be a σ -finite probability space. Let P be a measurable function on the product space $X \times X$ satisfying

$$P(x,y) \ge 0, \quad \int d\mu(y)P(x,y) = 1, \ x \in X$$

and

$$\int d\mu(x) \int_{S} d\mu(y) P(x, y) = \mu(S), \ S \in \mathcal{S}.$$

The time evolution Θ is given by

$$\Theta(f)(x) := \int d\mu(y) P(x, y) f(y).$$

Example 2.2 (Deterministic systems) Also deterministic dynamical systems are included in this formalism. They are given by a probability space (X, S, μ) and a transformation $T : X \to X$ which is measure-preserving, $\mu(T^{-1}(S)) = \mu(S), S \in S$. Take then $\Theta(f) := f \circ T$.

The positivity of Θ can be rephrased as $|\Theta(f)| \leq \Theta(|f|)$ (triangle inequality) or as $|\Theta(f)|^2 \leq \Theta(|f|^2)$ (Schwarz inequality). Deterministic systems are then distinguished by the additional property $|\Theta(f)| = \Theta(|f|)$ or by $|\Theta(f)|^2 = \Theta(|f|^2)$. This means exactly that Θ is an endomorphism of $L^{\infty}(X, \mu)$.

We are ready now to construct a partition on the level of the observables. A set of measurable functions $\mathcal{F} = \{f_k \mid k \in K\}$ with K a finite index set, is called a *partition* of unity whenever $f_k \geq 0$ and $\sum_k f_k = \mathbf{1}$. Such a set of functions can be interpreted as a response function for an unsharp measurement. The number $f_k(x)$ equals the probability for the measurement outcome $k \in K$ given the system is located in $x \in X$.

Example 2.3 (Sharp measurements) An important class of partitions of unity are those corresponding to sharp measurements. Let $C = \{C_k \mid k \in K\}$ be a measurable partition of X, i.e., $C_k \subset X$ is measurable, $C_k \cap C_l = C_k \delta_{kl}$ and $\bigcup_k C_k = X$. The set

$$\chi_{\mathcal{C}} := \{ \chi_{C_k} \mid k \in K \},\$$

where χ_C denotes the characteristic function of the set $C \subset X$, is a partition of unity.

Because deterministic dynamics act on the level of points of phase space, sharp measurements suffice in this case. Stochastic dynamics, on the contrary, smooth out sharp measurements. It is then natural to consider unsharp measurements. This implies, however, that a measurement as such, i.e., independent of the dynamics, can contribute to the dynamical entropy. Compared to classical deterministic systems, this is a new phenomenon one should take care off when constructing a dynamical entropy.

2.2 Refined partitions of unity

For two partitions of unity, $\mathcal{F} = \{f_k | k \in K\}$ and $\mathcal{G} = \{g_l | l \in L\}$, let $\mathcal{F} \lor \mathcal{G}$ denote the partition of unity $\{f_k g_l | k \in K, l \in L\}$. Also, define the time evolution of a partition of unity $\mathcal{F}, \Theta(\mathcal{F}) = \{\Theta(f_k) | k \in K\}$, which is again a partition of unity.

The repeated application of the time evolution Θ can be described as a refinement of an initial partition of unity. For deterministic dynamics, for example, the refined partition of unity after n time steps is given by

$$\mathcal{F} \vee \Theta(\mathcal{F}) \vee \ldots \vee \Theta^{n-1}(\mathcal{F}). \tag{1}$$

This can be used as a definition for the evolution of a partition of unity under stochastic dynamics, [Mak]. In this paper, we will use another definition,

$$\mathcal{F}^{(n)}[\Theta] := \mathcal{F} \vee \Theta(\mathcal{F} \vee \ldots \vee \Theta(\mathcal{F})), \tag{2}$$

where the initial partition of unity \mathcal{F} appears *n* times. As Θ is generally only positive and not necessarily an endomorphism, the definitions (1) and (2) coincide for deterministic systems, but differ for stochastic systems. This difference can be illustrated for a Markov process, Example 2.1.

Example 2.4 (Markov process) Let X be the finite state space and $\mathcal{F} = \{f_k\}$ a partition of unity. The element $(k_0, k_1, \ldots, k_{N-1})$ of the refined partition of unity (2) is given by

$$x \mapsto \sum_{(x_0, x_1, \dots, x_{N-1})} f_{k_0}(x) P_{xx_1} f_{k_1}(x_1) P_{x_1 x_2} \dots P_{x_{N-2} x_{N-1}} f_{k_{N-1}}(x_{N-1}),$$

and equals the probability for the measurement outcome $(k_0, k_1, \ldots, k_{N-1})$ given the initial state $x \in X$. A similar interpretation is missing for the refinement (1),

$$x \mapsto \sum_{(x_0, x_1, \dots, x_{N-1})} f_{k_0}(x) P_{xx_1} f_{k_1}(x_1) P_{xx_2}^{(2)} f_{k_2}(x_2) \dots P_{xx_{N-1}}^{(N-1)} f_{k_{N-1}}(x_{N-1}),$$

where $P^{(n)}$ is the n-th matrix power of P.

In quantum systems, partitions are replaced by so-called operational partitions, i.e., sets of observables $\mathcal{X} = \{x_k \mid k \in K\}$ such that $\sum_k x_k^* x_k = \mathbf{1}$. For two such partitions, $\mathcal{X} = \{x_k \mid k \in K\}$ and $\mathcal{Y} = \{y_l \mid l \in L\}$, we can again define $\mathcal{X} \lor \mathcal{Y} = \{x_k y_l \mid k \in K, l \in L\}$ and $\Theta(\mathcal{X}) = \{\Theta(x_k) \mid k \in K\}$. For endomorphisms Θ the refinement of \mathcal{X} can then be defined as in Eq. (1) or, equivalently, Eq. (2). However, for non-endomorphic maps Θ this approach does not work because $\Theta(\mathcal{X})$ will not be an operational partition anymore. Instead, at time *n* the #(K) operators $x_k^* x_k$ should be replaced by

$$x_{k_0}^* \Theta(x_{k_1}^* \dots \Theta(x_{k_{n-1}}^* x_{k_{n-1}}) \dots x_{k_1}) x_{k_0},$$

which are $\#(K)^n$ positive operators summing up to **1**. This is a generalization of (2) rather than of (1).

2.3 Entropy and all that

In this subsection we collect for finite systems the definitions and properties of entropy and of relative entropy which will be needed later [Weh, OhyPet]. These properties naturally extend to infinite systems.

With $\eta : [0,1] \to \mathbb{R}$ the entropy function,

$$\begin{aligned} \eta(x) &:= -x \log x, & x \in (0,1] \\ &:= 0, & x = 0, \end{aligned}$$

we have the following definitions.

Definition 2.5 The entropy (or Shannon entropy) of a probability measure $\mu = \{\mu_i \mid i \in I\}$ with finite index set I, is given by,

$$S(\mu) := \sum_{i \in I} \eta(\mu_i).$$

The quantum entropy (or von Neumann entropy) of a density matrix ρ on a finitedimensional Hilbert space is given by,

$$S_q(\rho) := \operatorname{Tr} \eta(\rho).$$

If we denote by $\operatorname{diag}(\rho)$ the probability measure obtained by restricting the density matrix ρ to its diagonal in a given basis, then $S_q(\rho) \leq S(\operatorname{diag}(\rho))$. Because the entropy function η is concave, the entropies S and S_q are also concave. Moreover, one has,

Proposition 2.6 (Concavity of entropy) Let μ , μ_{α} , $\alpha \in A$ and $\#(A) < \infty$, be probability measures on a finite set I and let $\lambda = \{\lambda_{\alpha} \mid \alpha \in A\}$ be a probability measure such that

$$\mu = \sum_{\alpha \in A} \lambda_{\alpha} \mu_{\alpha},$$

then

$$\sum_{\alpha} \lambda_{\alpha} S(\mu_{\alpha}) \le S(\mu) \le \sum_{\alpha} \lambda_{\alpha} S(\mu_{\alpha}) + S(\lambda).$$

Similarly, for a convex combination of density matrices ρ_{α} on a finite-dimensional Hilbert space,

$$\rho = \sum_{\alpha} \lambda_{\alpha} \rho_{\alpha},$$

it holds that

$$\sum_{\alpha} \lambda_{\alpha} S_q(\rho_{\alpha}) \le S_q(\rho) \le \sum_{\alpha} \lambda_{\alpha} S_q(\rho_{\alpha}) + S(\lambda).$$
(3)

Definition 2.7 Let μ and ν be two probability measures on the set I, $\#(I) < \infty$. The relative entropy of μ with respect to ν is

$$S(\mu \mid \nu) := \sum_{i \in I} \mu_i \log \frac{\mu_i}{\nu_i}.$$

Let ρ_1 and ρ_2 density matrices on a finite-dimensional Hilbert space. The quantum relative entropy of ρ_1 with respect to ρ_2 is

$$S_q(\rho_1 \mid \rho_2) := \operatorname{Tr} \rho_1(\log \rho_1 - \log \rho_2).$$

These relative entropies decrease monotonically under positive transformations.

Proposition 2.8 (Monotonicity of relative entropy) Let $M : \mathcal{C}(J) \to \mathcal{C}(I)$ be a linear positive unital map between the continuous functions on two finite sets I and J. This means, for $f \in \mathcal{C}(J)$, $i \in I$

$$M(f)(i) = \sum_{j \in J} M_{ij} f(j) \quad \text{with} \quad M_{ij} \ge 0, \quad \sum_{j \in J} M_{ij} = 1.$$

The dual of M maps probability measures on I to probability measures on J. For two probability measures μ and ν on I, the relative entropy S satisfies,

$$S(M^*\mu \mid M^*\nu) \le S(\mu \mid \nu).$$

For the quantum case, let $M : \mathcal{B} \to \mathcal{A}$ be a linear completely positive unital map between two finite-dimensional matrix algebras. The dual of M maps states on \mathcal{A} to states on \mathcal{B} . For two density matrices ρ_1 and ρ_2 on \mathcal{A} , the relative entropy S_q satisfies,

$$S_q(M^*\rho_1 \mid M^*\rho_2) \le S_q(\rho_1 \mid \rho_2).$$

As a consequence of this proposition, we have the following lemma.

Lemma 2.9 Let ρ , ρ_{α} , $\alpha \in A$ and $\#(A) < \infty$, be density matrices on a finitedimensional Hilbert space and let $\lambda = \{\lambda_{\alpha} \mid \alpha \in A\}$ be a probability measure such that

$$\rho = \sum_{\alpha \in A} \lambda_{\alpha} \rho_{\alpha},$$

then,

$$S(\operatorname{diag}(\rho)) - \sum_{\alpha} \lambda_{\alpha} S(\operatorname{diag}(\rho_{\alpha})) \le S_q(\rho) - \sum_{\alpha} \lambda_{\alpha} S_q(\rho_{\alpha}).$$

Proof Suppose $\rho, \rho_{\alpha} \in \mathcal{M}_k$ and $\alpha \in A$. Consider $M : \mathcal{M}_k \to \mathcal{M}_k \otimes \mathcal{C}(A)$ defined by $M(A) = A \otimes \mathbf{1}$. This is a completely positive unital map. Then M^* is the partial trace of $\mathcal{M}_k \otimes \mathcal{C}(A)$ to \mathcal{M}_k . Define,

$$\rho_1 = \sum_{\alpha} \lambda_{\alpha} \rho_{\alpha} \otimes \delta_{\alpha} \quad \text{and} \quad \rho_2 = \sum_{\alpha} \lambda_{\alpha} \operatorname{diag}(\rho_{\alpha}) \otimes \delta_{\alpha}.$$

Applying the monotonicity of the quantum relative entropy, Prop. 2.8, we obtain the inequality we are looking for. qed

2.4 Deterministic dynamical systems

The Kolmogorov-Sinai construction for the dynamical entropy of deterministic systems, see, e.g., [Wal], can be cast into this algebraic framework. With the notation of Example 2.2, let ν be a probability measure on X and $\mathcal{F} = \{f_k \mid k \in K\}$ a partition of unity in $L^{\infty}(X, \mu)$. Define a probability measure on K by $\nu \circ \mathcal{F} = \{\nu(f_k) \mid k \in K\}$. The metric or Kolmogorov-Sinai entropy $h^{\text{KS}}[\mu, \Theta]$ for the deterministic dynamics Θ with respect to the invariant measure μ can be written as,

$$h^{\rm KS}[\mu,\Theta] := \sup_{\mathcal{C}} \lim_{N \to \infty} \frac{1}{N} S(\mu \circ \chi_{\mathcal{C}}^{(N)}[\Theta]), \tag{4}$$

where the supremum is over all partitions C of the phase space X, Example 2.3.

3 CNT dynamical entropy

In the following two sections, we will generalize the dynamical entropy (4) for deterministic systems to stochastic systems. Our approach will be as follows. Different quantum dynamical entropies have been proposed in the literature and they all have to handle, albeit implicitly, the uncertainty generated by a quantum measurement. As explained before, a similar problem arises in the construction of a dynamical entropy for classical stochastic systems. We want now to reuse the different quantum solutions to treat the measurement uncertainty in this classical stochastic context.

We start by examining the quantum dynamical entropy proposed by Connes, Narnhofer and Thirring in [Con et al.]. In our language the basic notion is the entropy of a partition of unity with respect to a decomposition of the invariant measure. In particular, let $\mu = \sum_{\alpha} \lambda_{\alpha} \mu_{\alpha}$ be a decomposition of the invariant measure μ as a convex combination of probability measures μ_{α} , $\alpha \in A$ and $\#(A) < \infty$, with coefficients λ_{α} , $\lambda_{\alpha} \geq 0$ and $\sum_{\alpha} \lambda_{\alpha} = 1$. Let $\mathcal{F} = \{f_k\}$ be a partition of unity. We define

$$I[\mu, \{\lambda_{\alpha}\mu_{\alpha}\}, \mathcal{F}] := \sum_{\alpha} \lambda_{\alpha} S(\mu_{\alpha} \circ \mathcal{F} \mid \mu \circ \mathcal{F}) = S(\mu \circ \mathcal{F}) - \sum_{\alpha} \lambda_{\alpha} S(\mu_{\alpha} \circ \mathcal{F}).$$
(5)

To interpret this quantity, note first that a decomposition of μ corresponds to a partition of unity $\mathcal{G} = \{g_{\alpha}\}$ where $\lambda_{\alpha} = \mu(g_{\alpha})$ and $\mu_{\alpha}(f) = \mu(g_{\alpha}f)/\mu(g_{\alpha})$ for $f \in L^{\infty}(X,\mu)$. In other words, the function $g_{\alpha} \in L^{1}(X,\mu)$ is the Radon-Nikodym derivative of μ_{α} with respect to μ . Now \mathcal{G} can be seen as a finite model of X, whereas \mathcal{F} corresponds as usual to a measurement. Define the joint probability distribution $\mu_{\alpha k}^{12} := \mu(g_{\alpha}f_{k})$ with marginals $\mu_{\alpha}^{1} = \mu(g_{\alpha}), \ \mu_{k}^{2} = \mu(f_{k})$. With these definitions,

$$I[\mu, \{\lambda_{\alpha}\mu_{\alpha}\}, \mathcal{F}] = S(\mu^{1}) + S(\mu^{2}) - S(\mu^{12})$$
(6)

is the mutual information of the two marginals, or the average amount of information obtained about an initial condition $g_{\alpha} \in \mathcal{G}$ by performing a measurement $f_k \in \mathcal{F}$.

3.1 CNT construction

The entropic quantity (5) is now used in the definition of the CNT dynamical entropy of a stochastic dynamical system. Multi-index decompositions of the measure μ will be needed, which we write as

$$\mu = \sum_{\overline{\alpha}} \lambda_{\overline{\alpha}} \mu_{\overline{\alpha}},$$

where $\overline{\alpha} = (\alpha_1, \alpha_2, \ldots, \alpha_N)$, $\alpha_n \in A_n$ and $\#(A_n) < \infty$, $\mu_{\overline{\alpha}}$ are probability measures on X and $\lambda_{\overline{\alpha}}$ are the weights. For every $n = 1, 2, \ldots, N$, the marginal of this decomposition over all but the *n*-th index will be written as $\mu = \sum_{\alpha_n} \lambda_{\alpha_n}^{(n)} \mu_{\alpha_n}^{(n)}$. More explicitly,

$$\lambda_{\beta}^{(n)} = \sum_{\overline{\alpha} : \alpha_n = \beta} \lambda_{\overline{\alpha}} \quad \text{and} \quad \mu_{\beta}^{(n)} = \frac{1}{\lambda_{\beta}^{(n)}} \sum_{\overline{\alpha} : \alpha_n = \beta} \lambda_{\overline{\alpha}} \mu_{\overline{\alpha}}.$$

The probability measures $\{\lambda_{\overline{\alpha}} \mid \alpha_n \in A_n, \forall n\}$ and $\{\lambda_{\alpha_n}^{(n)} \mid \alpha_n \in A_n\}$ will be denoted by λ and $\lambda^{(n)}$ respectively.

Definition 3.1 Let (X, μ, Θ) be a stochastic dynamical system. Define,

$$H^{\text{CNT}}[\mu, \{\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_N\}] = \sup_{\mu = \sum_{\overline{\alpha}} \lambda_{\overline{\alpha}} \mu_{\overline{\alpha}}} \left(\sum_{n=1}^N I[\mu, \{\lambda_{\alpha_n}^{(n)} \mu_{\alpha_n}^{(n)}\}, \mathcal{F}_n] - \left(\sum_{n=1}^N S(\lambda^{(n)}) - S(\lambda) \right) \right),$$
(7)

where the supremum is over all finite N-index decompositions. The CNT dynamical entropy of (X, μ, Θ) is given by

$$h^{\text{CNT}}[\mu,\Theta] = \sup_{\mathcal{F}} \lim_{N \to \infty} \frac{1}{N} H^{\text{CNT}}[\mu, \{\mathcal{F},\Theta(\mathcal{F}), \dots, \Theta^{N-1}(\mathcal{F})\}].$$
 (8)

Especially in the quantum case, the optimization problem in (7) is the basic obstacle for calculating this dynamical entropy, see [Ben, Ben et al., Uhl]. We will analyze the supremum for N = 1 and N = 2.

One-time decompositions For the case N = 1, Eq. (7) becomes,

$$H^{\text{CNT}}[\mu, \mathcal{F}] = \sup_{\mu = \sum_{\alpha} \lambda_{\alpha} \mu_{\alpha}} I[\mu, \{\lambda_{\alpha} \mu_{\alpha}\}, \mathcal{F}]$$

= $S(\mu \circ \mathcal{F}) - \inf_{\mu = \sum_{\alpha} \lambda_{\alpha} \mu_{\alpha}} \sum_{\alpha} \lambda_{\alpha} S(\mu_{\alpha} \circ \mathcal{F}).$

The infimum in the second term of the right hand side is a convex optimization problem : find the infimum of the concave entropy functional over the convex domain of finite decompositions of the measure μ . This infimum will then be reached on the set of extremal points of this domain.

Assume that the partition of unity $\mathcal{F} = \{f_k \mid k \in K\}$ consists of simple functions. The result for general f_k will follow by continuity. For such simple functions their exists a partition $\mathcal{C} = \{C_i \mid i \in I\}$ of X, $\#(I) \leq \infty$, such that $f_k = \sum_i f_{ik} \chi_{C_i}$. The measures

 μ , μ_{α} can then be considered as measures on the finite set I, $\mu = \{\mu(C_i) \mid i \in I\}$ and $\mu_{\alpha} = \{\mu_{\alpha}(C_i) \mid i \in I\}.$

Let us now determine the extremal finite decompositions of the measure μ , or, equivalently, the extremal probability measures on $I \times A$ with $\#(A) < \infty$ such that the marginal over the α -index equals μ . We claim that these extremal measures are μ_f , characterized by a map $f: I \to A$ such that $\mu_f(i, \alpha) = \delta_{\alpha, f(i)} \mu(C_i)$. Indeed, μ_f is a probability measure on $I \times A$,

$$\sum_{i,\alpha} \mu_f(i,\alpha) = \sum_{i,\alpha} \delta_{\alpha,f(i)} \mu(C_i) = \sum_i \mu(C_i) = 1.$$

The measure μ_f has μ as marginal over the α -index,

$$\sum_{\alpha \in A} \mu_f(i, \alpha) = \sum_{\alpha \in A} \delta_{\alpha, f(i)} \mu(C_i) = \mu(C_i).$$

Moreover, μ_f is extremal. Suppose we can write μ_f as a convex combination of ν_1 and ν_2 , two probability measures on $I \times A$ with marginal μ ,

$$\mu_f = \frac{1}{2}\nu_1 + \frac{1}{2}\nu_2.$$

Substituting the explicit form for μ_f , one immediately gets $\mu_f = \nu_1 = \nu_2$. Finally, all the extremal points are of this form because every probability measure on $I \times A$ with marginal μ , can be written as a convex combination,

$$\mu_{\alpha}(C_i) = \sum_f c_f \mu_f(i, \alpha),$$

with $c_f \ge 0$ and $\sum_f c_f = 1$.

The infimum will thus be reached on this set of measures μ_f . Moreover, we can restrict our attention to injective maps f. This follows again by the concavity of the entropy functional. The order of the indices α is of no importance, so we can take as optimal decomposition $\mu_{\alpha} = \delta_i$ and $\lambda_{\alpha} = \mu(C_i)$. As a result, for simple functions f_k ,

$$H^{\text{CNT}}[\mu, \mathcal{F}] = S(\mu \circ \mathcal{F}) - \sum_{i} \mu(C_i) S(\{f_{ik} \mid k \in K\}),$$

or, for general f_k ,

$$H^{\text{CNT}}[\mu, \mathcal{F}] = S(\mu \circ \mathcal{F}) - \int d\mu(x) S(\delta_x \circ \mathcal{F}) = \int d\mu(x) S(\delta_x \circ \mathcal{F} \mid \mu \circ \mathcal{F}).$$
(9)

Two-times decompositions In contrast with the case N = 1, the optimization problem (7) is *not* convex for the case N = 2. To see this, suppose the partitions of unity $\mathcal{F} = \{f_k \mid k \in K\}$ and $\mathcal{G} = \{g_l \mid l \in L\}$ consist of simple functions, $f_k = \sum_i f_{ik} \chi_{C_i}$ and $g_l = \sum_j g_{jl} \chi_{D_j}$. Here, $\mathcal{C} = \{C_i\}$ and $\mathcal{D} = \{D_j\}$ are finite partitions of X and so is $\mathcal{C} \vee \mathcal{D} = \{C_i \cap D_j\}$.

Define now a probability measure on the composed system $A \times B \times I \times J \times K \times L$,

$$\mu_{\alpha\beta i j k l}^{1\,2\,3\,4\,5\,6} = \lambda_{(\alpha,\beta)} \mu_{\alpha\beta} (C_i \cap D_j) f_{ik} f_{jl}.$$

This probability measure has marginals

$$\mu_{k}^{5} = \mu(f_{k}), \quad \mu_{l}^{6} = \mu(g_{l}), \quad \mu_{\alpha\beta}^{12} = \lambda_{(\alpha,\beta)},$$
$$\mu_{\alpha k}^{15} = \lambda_{\alpha}^{(1)} \mu_{\alpha}^{(1)}(f_{k}), \quad \mu_{\beta l}^{26} = \lambda_{\beta}^{(2)} \mu_{\beta}^{(2)}(g_{l}).$$

The functional to optimize can then be written as

$$S(\mu^5) + S(\mu^6) + S(\mu^{12}) - S(\mu^{15}) - S(\mu^{26}).$$

The task is now to optimally couple subsystems A and B with I and J. If this optimization problem were convex, the optimal coupling would identify A with I and B with J, as in the case N = 1. This decomposition can yield a negative value for the supremum. However, the supremum has to be positive because the functional is zero for the trivial decomposition, i.e., #(A) = 1 and #(B) = 1. We conclude that this optimization problem is not convex.

3.2 Hudetz construction

The CNT construction seems to be intractable because of the supremum over all multi-index decompositions of the invariant measure, Eq. (7). These multi-index decompositions were introduced to obtain finite-dimensional algebras for the one-time restrictions. For stochastic systems this algebraic structure is absent anyway. The following construction appears to be more natural.

Definition 3.2 Let (X, μ, Θ) be a stochastic dynamical system. Define,

$$H^{\mathrm{Hud}}[\mu, \mathcal{F}] := \sup_{\mu = \sum_{\alpha} \lambda_{\alpha} \mu_{\alpha}} I[\mu, \{\lambda_{\alpha} \mu_{\alpha}\}, \mathcal{F}].$$
(10)

The Hudetz (Hud) dynamical entropy of (X, μ, Θ) is given by

$$h^{\mathrm{Hud}}[\mu,\Theta] := \sup_{\mathcal{F}} \limsup_{N \to \infty} \frac{1}{N} H^{\mathrm{Hud}}[\mu, \mathcal{F}^{(N)}[\Theta]].$$
(11)

The optimization problem at any time N is now the same as the one encountered in the CNT construction for one time decompositon. This supremum was worked out explicitly, Eq. 9,

$$H^{\mathrm{Hud}}[\mu, \mathcal{F}] = S(\mu \circ \mathcal{F}) - \int d\mu(x) \, S(\delta_x \circ \mathcal{F}) = \int d\mu(x) S(\delta_x \circ \mathcal{F} \,|\, \mu \circ \mathcal{F}) \tag{12}$$

Moreover, restricting the supremum to partitions of unity \mathcal{F} which correspond to sharp measurements, leads to the same result.

Proposition 3.3 For a stochastic dynamical system (X, μ, Θ) holds

$$h^{\mathrm{Hud}}[\mu,\Theta] = \sup_{\mathcal{C}} \limsup_{N \to \infty} \frac{1}{N} H^{\mathrm{Hud}}[\mu, \chi_{\mathcal{C}}^{(N)}[\Theta]].$$

Proof We have to show that for every partition of unity $\mathcal{F} = \{f_k\}$ there exist a partition $\mathcal{C} = \{C_i\}$ of X such that,

$$H^{\mathrm{Hud}}[\mu, \mathcal{F}^{(N)}[\Theta]] \le H^{\mathrm{Hud}}[\mu, \chi_{\mathcal{C}}^{(N)}[\Theta]].$$

We consider the case N = 2. The proof for other N is analogous.

Assume that f_k are simple functions. The result for general f_k will follow by continuity. For simple functions f_k there exists a partition $\mathcal{C} = \{C_i\}_i$ of X such that $f_k = \sum_i f_{ik}\chi_{C_i}$ with $M = [f_{ik}]$ a stochastic matrix. Element (k_0, k_1) of the refined partition $\mathcal{F}^{(2)}[\Theta] = \mathcal{F} \vee \Theta(\mathcal{F})$ is then

$$f_{k_0}\Theta(f_{k_1}) = \sum_{i_0i_1} f_{i_0k_0} f_{i_1k_1} \chi_{C_{i_0}}\Theta(\chi_{C_{i_1}})$$

Thus,

$$\mu \circ \mathcal{F}^{(2)}[\Theta] = (M \otimes M)^* (\mu \circ \chi^{(2)}_{\mathcal{C}}[\Theta])$$

with $M \otimes M$ a positive unital transformation. By Prop. 2.8 one obtains,

$$H^{\mathrm{Hud}}[\mu, \mathcal{F}^{(2)}[\Theta]] \le H^{\mathrm{Hud}}[\mu, \chi_{\mathcal{C}}^{(2)}[\Theta]].$$

qed

Referring to the information-theoretic interpretation of $I[\mu, \{\lambda_{\alpha}\mu_{\alpha}\}, \mathcal{F}]$, the quantity $H^{\text{Hud}}[\mu, \mathcal{F}]$ can be seen as the mutual information between the initial state and the measurement outcomes for the best model of the state space X, namely X itself. Therefore, $H^{\text{Hud}}[\mu, \mathcal{F}]$ equals the information obtained about an initial state $x \in X$ by performing a measurement $f_k \in \mathcal{F}$. The Hudetz dynamical entropy $h^{\text{Hud}}[\mu, \Theta]$ equals the average information gain. In this way, out of the three sources of dynamical entropy, only the system dynamics contributes.

Example 3.4 (Deterministic systems) By Prop. 3.3, we can restrict our attention to sharp measurements. Recall that deterministic dynamics transform sharp measurements into sharp ones. Therefore, all the probability measures $\delta_x \circ \mathcal{F}$ appearing in (12) are pure and $H^{\text{Hud}}[\mu, \mathcal{F}] = S(\mu \circ \mathcal{F})$. We conclude that $h^{\text{Hud}} = h^{\text{KS}}$ for deterministic systems.

Example 3.5 (Finite systems) Consider a dynamical system with a finite state space X. Denote the invariant measure by $\mu = \{\mu_x | x \in X\}$. By Eq. 12,

$$H^{\text{Hud}}[\mu, \mathcal{F}] = \sum_{kx} \mu_x f_k(x) \log \frac{f_k(x)}{\mu(f_k)} \le \sum_{kx} \mu_x f_k(x) \log \frac{1}{\mu_x} = S(\mu).$$

This quantity is finite and does not depend on N when \mathcal{F} is replaced by $\mathcal{F}^{(N)}[\Theta]$ in (11). We conclude that $h^{\text{Hud}} = 0$ for finite systems.

Note that the CNT dynamical entropy gives the same results for these two examples. Finally, let us compare the Hudetz dynamical entropy with another definition for the dynamical entropy of stochastic systems. It is closely related to [Mak], but as explained before, we use another refinement of partitions, (2) instead of (1). First, a density matrix is constructed,

$$\left(\rho^{\mathrm{Mak}}[\mu,\mathcal{F}]\right)_{kl} := \mu(\sqrt{f_k f_l}).$$
(13)

The Makarov dynamical entropy is given by

$$h^{\text{Mak}}[\mu,\Theta] := \sup_{\mathcal{F}} \limsup_{N \to \infty} \frac{1}{N} S_q(\rho^{\text{Mak}}[\mu, \mathcal{F}^{(N)}[\Theta]]).$$
(14)

This dynamical entropy leads to the same result as the Hudetz dynamical entropy for the two examples discussed. Moreover, it holds that $h^{\text{Hud}} \leq h^{\text{Mak}}$. Indeed, from Lemma 2.9,

$$S(\operatorname{diag}(\rho)) - \sum_{i} \lambda_i S(\operatorname{diag}(\rho_i)) \le S_q(\rho).$$

This is equivalent with $H^{\text{Hud}}[\mu, \mathcal{F}] \leq S_q(\rho^{\text{Mak}}[\mu, \mathcal{F}])$ for all partitions of unity \mathcal{F} .

4 AFL dynamical entropy

In [AliFan1] another quantum dynamical entropy was proposed, based on an idea of Lindblad, by mapping the evolution of a dynamical system onto a quantum spin chain. For the classical stochastical systems we are interested in, the definition is as follows.

Definition 4.1 Let (X, μ, Θ) be a stochastic dynamical system. Define the density matrix $\rho^{\text{AFL}(N)}$ by

$$\left(\rho^{\text{AFL}(N)}[\mu,\Theta,\mathcal{F}]\right)_{\overline{k},\overline{l}} := \mu\left(\sqrt{f_{k_0}f_{l_0}}\Theta\left(\sqrt{f_{k_1}f_{l_1}}\dots\Theta\left(\sqrt{f_{k_{N-1}}f_{l_{N-1}}}\right)\right)\right).$$
(15)

The AFL dynamical entropy of (X, μ, Θ) is given by

$$h^{\text{AFL}}[\mu,\Theta] := \sup_{\mathcal{F}} \limsup_{N \to \infty} \frac{1}{N} S_q(\rho^{\text{AFL}(N)}[\mu,\Theta,\mathcal{F}]).$$
(16)

Note that the density matrix $\rho^{\text{AFL}(N)}[\mu, \Theta, \mathcal{F}]$ is different from $\rho^{\text{Mak}}[\mu, \mathcal{F}^{(N)}[\Theta]]$ for N > 2. For sharp measurements $\chi_{\mathcal{C}}$ with \mathcal{C} a partition of X, the density matrix $\rho^{\text{AFL}(N)}[\mu, \Theta, \chi_{\mathcal{C}}]$ is diagonal. In that case,

$$S_q(\rho^{\text{AFL}(N)}[\mu,\Theta,\mathcal{F}]) = S(\mu \circ \mathcal{F}^{(N)}[\Theta]).$$

As a consequence,

$$h^{\mathrm{AFL}}[\mu,\Theta] \ge \sup_{\mathcal{F} \text{ sharp}} \limsup_{N \to \infty} \frac{1}{N} S(\mu \circ \mathcal{F}^{(N)}[\Theta])$$

In fact, equality holds.

Proposition 4.2 For a stochastic dynamical system (X, μ, Θ) holds

$$h^{\text{AFL}}[\mu, \Theta] = \sup_{\mathcal{C}} \limsup_{N \to \infty} \frac{1}{N} S(\mu \circ \chi_{\mathcal{C}}^{(N)}[\Theta]).$$

Proof We have to show that for every partition of unity $\mathcal{F} = \{f_k\}$ there exist a partition $\mathcal{C} = \{C_i\}$ of X such that,

$$S(\rho^{\text{AFL}(N)}[\mu,\Theta,\mathcal{F}]) \leq S(\mu \circ \chi_{\mathcal{C}}^{(N)}[\Theta]).$$

We consider the case N = 2. The proof for other N is analogous.

Assume that f_k are simple functions. The result for general f_k will follow by continuity. For such simple functions, there exists a partition of X, $C = \{C_i | i \in I\}$ and $\#(I) < \infty$, such that $f_k = \sum_i f_{ik} \chi_{C_i}$ with $[f_{ik}]$ a stochastic matrix. Component $(k_0, k_1), (l_0, l_1)$ of the density matrix $\rho^{\text{AFL }(2)}[\mu, \Theta, \mathcal{F}]$ is

$$\mu \left(\sqrt{f_{k_0} f_{l_0}} \Theta \left(\sqrt{f_{k_1} f_{l_1}} \right) \right)$$

$$= \mu \left(\sqrt{\sum_{i_0 j_0} f_{i_0 k_0} f_{j_0 l_0} \chi_{C_{i_0}} \chi_{C_{j_0}}} \Theta \left(\sqrt{\sum_{i_1 j_1} f_{i_1 k_1} f_{j_1 l_1} \chi_{C_{i_1}} \chi_{C_{j_1}}} \right) \right)$$

$$= \mu \left(\sqrt{\sum_{i_0} f_{i_0 k_0} f_{i_0 l_0} \chi_{C_{i_0}}} \Theta \left(\sqrt{\sum_{i_1} f_{i_1 k_1} f_{i_1 l_1} \chi_{C_{i_1}}} \right) \right)$$

$$= \sum_{i_0 i_1} \mu \left(\chi_{C_{i_0}} \Theta \left(\chi_{C_{i_1}} \right) \right) \sqrt{f_{i_0 k_0} f_{i_0 l_0}} \sqrt{f_{i_1 k_1} f_{i_1 l_1}}$$

This a convex combination of $\#(I)^2$ vector states. The coefficient of term (i_0, i_1) is $\mu(\chi_{C_{i_0}}\Theta(\chi_{C_{i_1}}))$ and component (k_0, k_1) of the corresponding vector is $\sqrt{f_{i_0k_0}f_{i_1k_1}}$. These vectors are normalized because $[f_{ik}]$ is stochastic. Applying the second inequality in (3) finishes the proof. qed

Prop. 4.2 leads to the following interpretation of $h^{\text{AFL}}[\mu, \Theta]$. It is the average uncertainty on the outcome of sharp measurements. Out of the three sources of dynamical entropy, both the system dynamics and the stochasticity contribute.

Example 4.3 (Deterministic systems) By comparing Eq. 4 and Prop. 4.2, $h^{AFL} = h^{KS}$ for deterministic systems.

Example 4.4 (Finite systems) The dynamical entropy h^{AFL} can be strictly positive for finite systems. In this case, the supremum over all sharp partitions is reached for the extremal partition, i.e., $C = \{\{x\} | x \in X\}$. For a Bernoulli process with probabilities $\{p_x\}$ and $\sum_x p_x = 1$, one obtains $h^{\text{AFL}} = \sum_x \eta(p_x)$, whereas $h^{\text{Hud}} = 0$.

Finally, we compare the AFL dynamical entropy with the other definitions. The given interpretation and the finite case example suggest the inequality $h^{\text{Hud}} \leq h^{\text{AFL}}$. This

can be easily proven by Lemma 2.9. Another definition for the dynamical entropy of stochastic systems was given in [Kos et al.], based on [Acc et al.] for deterministic systems. In our notation it can be written as,

$$h^{\text{KOW}}[\mu, \Theta] = \sup_{\mathcal{F}} \limsup_{N \to \infty} \frac{1}{N} S(\mu \circ \mathcal{F}^{(N)}[\Theta]).$$

By comparing this with Prop. 4.2, $h^{\text{AFL}} \leq h^{\text{KOW}}$. Because unsharp measurements are allowed, they can contribute to the dynamical entropy. The three sources of dynamical entropy are thus taken into account. As a consequence, $h^{\text{KS}} \leq h^{\text{KOW}}$ for deterministic systems where strict inequality can hold. Even stronger, without restricting the set of allowed measurements, this dynamical entropy will always be infinite. Indeed, the partition of unity consisting of k elements $\frac{1}{k}\mathbf{1}$ leads to a dynamical entropy log k. This can grow without bounds.

Acknowledgements It is a pleasure to acknowledge constructive discussions with R. Alicki who pointed out reference [Mak] to us.

References

- [Ali] R. Alicki, Information-theoretical meaning of quantum dynamical entropy, e-print quant-ph/0201012.
- [Ali et al.] R. Alicki, J. Andries, M. Fannes, P. Tuyls, An algebraic approach to the Kolmogorov-Sinai entropy, *Rev. Math. Phys.* 8, 167–184 (1996).
- [AliFan1] R. Alicki, M. Fannes, Defining quantum dynamical entropy, *Lett. Math. Phys.* **32**, 75–82 (1994).
- [AliFan2] R. Alicki, M. Fannes, *Quantum Dynamical Systems*, Oxford University Press, Oxford, 2001.
- [Acc et al.] L. Accardi, M. Ohya, N. Watanabe, Dynamical entropy through quantum Markov chains, Open Sys. & Information Dyn. 4, 71–87 (1997).
- [Ben] F. Benatti, Entropy of a subalgebra and quantum estimation, J. Math. Phys. **37**, 5244–5258 (1996).
- [Ben et al.] F. Benatti, H. Narnhofer, A. Uhlmann, Optimal decomposition of quantum states with respect to entropy, *Rep. Math. Phys.* **38**, 123–141 (1996).
- [Con et al.] A. Connes, H. Narnhofer, W. Thirring, Dynamical entropy of C*-algebra and von Neumann algebras, *Commun. Math. Phys.* **112**, 691–719 (1987).
- [Hud] T. Hudetz, Quantum dynamical entropy revisited, *Banach Center Publ.* **43**, 241–251 (1997).
- [Kos et al.] A. Kossakowski, M. Ohya, N. Watanabe, Quantum dynamical entropy for completely positive map, *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 2, 267–282 (1999).

[Mak]	I. I. Makarov, Dynamical entropy for Markov operators, <i>Journal of Dynamical and Control Systems</i> , 6 , 1–11 (2000).
[OhyPet]	M. Ohya, D. Petz, <i>Quantum entropy and its use</i> , Springer-Verlag, Berlin, 1993.
[Uhl]	A. Uhlmann, Entropy and optimal decomposition of states relative to a maximal commutative subalgebra, <i>Open Sys. & Information Dyn.</i> 5, 209–227 (1998).
[Wal]	P. Walters, An Introduction to Ergodic Theory, Springer-Verlag, Berlin, 1982.
[Weh]	A. Wehrl, General properties of entropy, <i>Rev. Mod. Phys.</i> 50 , 221–260 (1978).